



**HAL**  
open science

# Importance of Choosing Relevant Biological End Points To Predict Nanoparticle Toxicity with Computational Approaches for Human Health Risk Assessment

Valérie Forest, Jean-François Hochepped, Jérémie Pourchez

► **To cite this version:**

Valérie Forest, Jean-François Hochepped, Jérémie Pourchez. Importance of Choosing Relevant Biological End Points To Predict Nanoparticle Toxicity with Computational Approaches for Human Health Risk Assessment. *Chemical Research in Toxicology*, 2019, 32 (7), pp.1320-1326. 10.1021/acs.chemrestox.9b00022 . hal-02189302

**HAL Id: hal-02189302**

**<https://hal.science/hal-02189302v1>**

Submitted on 19 Jul 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **Importance of choosing relevant biological endpoints to predict nanoparticle toxicity with computational approaches for human health risk assessment**

Valérie Forest<sup>†,\*</sup>, Jean-François Hochepped<sup>‡,§</sup>, Jérémie Pourchez<sup>†</sup>.

<sup>†</sup> Mines Saint-Etienne, Univ Lyon, Univ Jean Monnet, INSERM, U 1059 Sainbiose, Centre CIS, F-42023 Saint-Etienne, France.

<sup>‡</sup> MINES ParisTech, PSL Research University, MAT - Centre des matériaux, CNRS UMR 7633, BP 87 91003 Evry, France.

<sup>§</sup> UCP, ENSTA ParisTech, Université Paris-Saclay, 828 bd des Maréchaux, 91762 Palaiseau cedex France.

\* Corresponding author: Valérie Forest:

École Nationale Supérieure des Mines de Saint-Etienne

158 cours Fauriel, CS 62362


42023 Saint-Etienne Cedex 2. FRANCE.

Email address: [vforest@emse.fr](mailto:vforest@emse.fr) - Telephone number: +33477499776

## **Key-words**

Nanoparticles; Cytotoxicity; *In vitro* nanotoxicology; *In silico* nanotoxicology; Predictive models; QSAR.

## Table of Contents Graphic



## **Abstract**

Because it is impossible to assess *in vitro* or *in vivo* the toxicity of all nanoparticles available on the market on a case-by-case basis, computational approaches have been proposed as useful alternatives to predict *in silico* the hazard potential of engineered nanoparticles. Despite promising results, a major issue associated with these mathematical models lies in the *a priori* choice of the physico-chemical descriptors and the biological endpoints. We performed a thorough bibliographic survey on the biological endpoints used for nanotoxicology purposes and compared them between experimental and computational approaches. They were found to be disparate: while conventional *in vitro* nanotoxicology assays usually investigate a large array of biological effects using eukaryotic cells (cytotoxicity, pro-inflammatory response, oxidative stress, genotoxicity), computational studies mostly focus on cell viability and also includes studies on prokaryotic cells. We may thus wonder the relevance of building complex mathematical models able to predict accurately a biological endpoint if this latter is not the most relevant to support human health risk assessment. The choice of biological endpoints clearly deserves to be more carefully discussed. This could bridge the gap between experimental and computational nanotoxicology studies and allow *in silico* predictive models to reach their full potential.

In the field of nanotoxicology a major aim is the investigation of the biological effects induced by engineered nanoparticles to determine their potential impact on the environment and human health. In this context, *in vivo* studies are potentially the most informative as animal models reproduce a physiological integrated response. But because of ethical issues, animal experiments are limited to follow the 3R's rule defined by Russell and Burch <sup>1</sup> aiming to replace, reduce and refine the use of animals for scientific purposes. This trend was followed by regulation <sup>2</sup> which advocates for the reduction of animal models for evaluating nanoparticles toxicity. *In vitro* models have then been developed for human risk assessment. They possess several advantages such as being inexpensive, easy and rapid to perform, but most of all they allow the high throughput screening of biological effects triggered by nanoparticles, also enabling mechanistic studies. In this context, a large panel of assays is usually carried out to determine the cytotoxicity, the pro-inflammatory response, the oxidative stress or the genotoxicity triggered by the contact of nanoparticles with eukaryotic cells <sup>3-6</sup> (some examples are reported in Table 1, even though there are no standardized methods and no international guidelines).

**Table 1** – Biological endpoints commonly evaluated in *in vitro* nanotoxicology studies (not exhaustive).

<b>Toxicity endpoint</b>	<b>Evaluated parameter</b>	<b>Examples of assays</b>
Cell viability	Apoptosis induction	Caspases, TUNEL, AnnexinV
	Cell proliferation	BrdU
	Membrane damages	LDH
	Mitochondrial activity	MTT, WST-1, ATP content
Pro-inflammatory response	Cytokine production	TNF- $\alpha$ , interleukins (IL8, IL6, IL1 $\alpha$ , IL1 $\beta$ , GM-CSF, etc.)



approaches have been applied to nanoparticles such as (Q)SAR ((Quantitative) Structure Activity Relationship)<sup>9-13</sup>, read-across<sup>14-17</sup>, neural network<sup>18</sup> or decision tree<sup>19</sup> classifications. These models can theoretically be built using data obtained through either *in vitro* or *in vivo* assays. But in practice, to be reliable, these models should involve a very large amount of data. And it is thus difficult to get such an amount of data with *in vivo* experiments. Indeed, while *in vivo* experiments are supposed to be closer to “real life” condition because they take into account physiological parameters, they are expensive, time-consuming and limited because of ethical issues. On the contrary, *in vitro* studies seem more suitable as they are cheaper, more rapid and easier to perform. They are thus rather useful for mechanistic studies and as screening tools as they allow to collect more information. This doesn't mean that *in vivo* studies are less qualified to help assess risks, but it argues for considering a multi-step process research. First, *in vitro* assays could be used for the screening of the nanomaterials that are worth investigating further. Based on these data predictive models could be built for risk assessment. And finally particles of interest could be further investigated using *in vivo* assays that can bring complementary information.


*In silico* models are the subject of intensive research and are highly topical issues. An overview of the landscape of the available computational models for nanomaterials has been very recently reported<sup>20</sup>. Although some of these mathematical models have produced promising results, their construction is associated with some challenges and questions. A major issue lies in the *a priori* choice of the physico-chemical descriptors and biological endpoints, which are at the core of the computational modeling. As the QSAR approach remains the most abundantly documented in the literature we will focus our discussion on it thereafter. It was originally developed for risk assessment of chemicals for which many molecular descriptors are available in chemical database or easily calculated with usual softwares. Nanoparticles, because of their specific properties, need to be described by additional parameters accurately experimentally

measured such as size, shape, surface area, surface reactivity, crystalline structure, composition of core and coating, etc.<sup>21</sup>. Therefore, one of the specific features of QSAR modeling for nanotoxicology purposes involves a labor-intensive, time-consuming and costly effort to obtain these “nanodescriptors”. And because descriptors are mathematical representations of relevant properties of nanoparticles, nanoparticle physico-chemical features should be carefully considered, especially the parameters recommended by the ISO/TR13014:2012 standard<sup>22</sup>. Unfortunately, a systematic and comprehensive nanoparticle physico-chemical characterization is not always available in the proposed nanoQSAR models. Furthermore, the reality is much more complex and to take into account the “biological identity” of the nanoparticles (reflecting their complex interactions with biological environments) the presence and nature of the corona formed around the nanoparticles should be introduced in the equation<sup>23–25</sup>.

Regarding the choice of the biological endpoints to be predicted thanks to the model, the question is even more debatable. Indeed, we observed that often, the main biological endpoints that are used in empirical nanotoxicology studies (studies without any modelling purpose) and in studies dedicated to the *in silico* prediction of the nanoparticle toxicity are different. We thus wondered why there is such a discrepancy between the two types of studies. The assays used for empirical nanotoxicology reported in Table 1 can all be a priori used for modelling. But because many other parameters are involved such as the particle type, the cell type, etc... they will not exhibit the same efficiency in their predictive potential for risk assessment. No one could predict which assays (and why) will be more predictive than others. Only the construction of the models will tell if they work or not. To exemplify this point, we performed a literature survey focused on nanoparticles (including nanomaterials and nanostructure) and QSAR model and its variants (because as mentioned before, QSAR approach is among the most abundantly documented and used computational approaches). As illustrated in Figure 1, querying the PubMed database with the search terms “(nanoparticle OR nanomaterials OR nanostructure)



AND (QSAR OR QSPR OR QNAR OR QNTR) revealed that 232 reports have been published on this topic by March 2019 (please refer to Supporting Information for further details). We thoroughly reviewed these articles focusing our attention on studies in which nanoQSAR models were built using *in vitro* toxicological data (because as mentioned before unlike *in vivo* studies, they allow to get enough data to construct reliable models). Based on these criteria, we excluded 167 papers (see Supporting Information), either because they did not include experimental data (52 papers: reviews, commentaries, book chapters, etc.) or because their topic was not directly related to our issue (115 papers with no cytotoxicity data or not related to predictive nanotoxicology for human risk assessment, or based on *in vivo* data, etc.). 65 papers were therefore included in this analysis. Among them, nanoparticle toxicity assessment was performed using prokaryotic cells in 19 papers<sup>10,16–18,26–40</sup>, eukaryotic cells in 40 papers<sup>9,12,41–78</sup> or both in 6 papers<sup>79–84</sup>. Please note that the exclusion of some studies from this bibliographic survey doesn't question their merit or usefulness, it just means they don't match the specific criteria we had previously established.



**Figure 1** – Schematic summary of the literature survey performed querying the PubMed database with the search terms (nanoparticle OR nanomaterials OR nanostructure) AND (QSAR OR QSPR OR QNAR OR QNTR). Of the 232 papers found we excluded 52 papers because they didn't present experimental data and other 115 papers were excluded because they didn't match our criteria (papers with no cytotoxicity data or not related to predictive nanotoxicology for human risk assessment, or based on *in vivo* data, etc.). Of the 65 papers remaining 19 assessed toxicity on prokaryotic cells, 40 on eukaryotic and 6 on both.

When nanoparticle toxicity is investigated using prokaryotic cells, the major biological endpoint evaluated is bacteria viability (often using *Escherichia coli*) by the means of the EC<sub>50</sub> measure<sup>10,26–29,79</sup> (half maximal effective concentration, corresponding to the nanoparticle concentration that reduces bacteria viability by 50%<sup>17</sup>). This cellular model and this parameter are rather useful for ecotoxicology purposes but seem less sensible to infer nanoparticle hazard to human health.

We then analyzed more closely the studies using eukaryotic cells. We especially paid attention to the quantity and quality of the data needed to build the mathematical models, 2 criteria that are commonly acknowledged to be of high importance for the construction of reliable QSAR models. In addition, we examined the number and nature of the selected biological endpoints. Regarding the first criteria about data quantity (*i.e.* the size of the dataset), as mentioned before QSAR models were originally developed for chemicals for which data were easily available and QSAR could be built based on considerably large databases (compounds in the order of hundreds or even thousands). On the contrary, in the case of nanoparticles, due to the difficulty to gather physico-chemical and biological data in standardized conditions, datasets are often very small (units or tens, in the best case)<sup>34</sup>. In our bibliographic survey, we found that most of the studies used a dataset higher than 20 but at the expense of the quality of the data.

Indeed, for standard *in vitro* nanotoxicology, the ISO/TR13014:2012 standard recommends to characterize at least the 8 following nanoparticle physico-chemical features: agglomeration/aggregation state, composition, size, shape, solubility/dispersibility, specific surface area, density of surface groups and surface chemistry<sup>22</sup>. But in computational studies this comprehensive nanoparticle physico-chemical characterization is far from being systematic. Actually, none of the 40 papers experimentally measured these 8 crucial parameters. The nanodescriptors greatly varied depending on the studies, some considered only nanoparticle size and zeta potential while others included much more parameters calculated or experimentally assessed but none characterized the 8 parameters recommended by the ISO/TR13014:2012 standard. The lack of complete characterization may be explained by some metrological issues. First, sample preparation is not trivial for techniques requiring a good dispersion in suspensions. In addition, some questions are still open. For instance, the agglomeration/aggregation state is quite unclear. Agglomeration and aggregation are very different concepts: in agglomerates particles are just gathered by weak interactions and agglomerates size is expected to change a lot during a journey in living organisms, whereas in aggregates particles are cemented by solid bridges and will not budge. So there is a high risk that agglomerates size measurement does not make much sense, unless the medium and sample preparation is representative of biological conditions. As far as size is concerned, even in the case of well dispersed spherical particles - defined by one parameter- the correct measurement of a size distribution is all but trivial, and direct methods as TEM or indirect methods as DLS give different results. Things get even more complicated when shape is not spherical and at least a second dimensional parameter is needed to describe particles. Density of surface groups and surface chemistry may be determined in average, but locally they depend on crystal exposed faces or even edges, which supposes expensive HRTEM studies for a perfect description.

Finally, on the third criteria regarding the biological endpoints, we observed that in the vast majority of the cases (29/40 papers) only one biological endpoint was considered, mainly cell viability. While as mentioned before, in standard *in vitro* nanotoxicology before concluding on the toxicity of a nanoparticle several parameters should be investigated (*i.e.* cytotoxicity, pro-inflammatory response, oxidative stress, genotoxicity, etc.). In this respect, the study from Maher *et al.*<sup>55</sup> was the most complete as in addition to cytotoxicity, reactive oxygen species and pro-inflammatory cytokine production were included as biological endpoints. Similarly, Le *et al.*<sup>47</sup> considered the oxidative stress in their computational study.

In the end, none of the 40 papers met the 3 criteria (dataset>20; physico-chemical characterization as recommended by ISO/TR13014:2012 standard; and more than one biological endpoint studied), 10 met 2, 20 only one and 10 not even one.

Therefore, there is a gap between the nanodescriptors and biological endpoints used in computational approaches and *in vitro* empirical testing for human health risk assessment. Taken together these observations argue for the need to discuss the relevance of the selected biological endpoints. Such discussion could bridge the gap between experimental and computational nanotoxicology studies to get predictive models more useful for human risk assessment. It should be also interesting to consider and include in nanoQSAR alternative biological parameters which could be more relevant to take into account nanoparticle dose-effects such as NOAEL (No Observable Adverse Effect Level, *i.e.* the highest tested nanoparticle dose at which no adverse effect is found where higher doses result in an adverse effect).

It is unanimously acknowledged that many challenges in predictive nanotoxicology are associated with the quantity and quality of the data needed to build the mathematical model. But the nature of the biological endpoints deserves better attention and should be carefully discussed to get more useful and applicable models to support human risk assessment. This

way, mathematical models could reach their full potential, able to predict nanoparticle toxicity avoiding empirical assays, saving time, money and animal experiments.

### **Funding**

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### **Declarations of interest**

The authors declare no competing financial interest.

### **Supporting Information**

Details on the 232 publications found when querying the PubMed database with the search terms (nanoparticle OR nanomaterials OR nanostructure) AND (QSAR OR QSPR OR QNAR OR QNTR).

### **References**

- (1) Russell, W. M. .; Burch, R. . The Principles of Humane Experimental Technique; Methuen, 1959.
- (2) European Union. *Directive 2010/63/EU of the European Parliament and of the Council of 22 September 2010 on the Protection of Animals Used for Scientific Purposes*; 2010.
- (3) Hillegass, J. M.; Shukla, A.; Lathrop, S. A.; MacPherson, M. B.; Fukagawa, N. K.; Mossman, B. T. Assessing Nanotoxicity in Cells in Vitro. *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.* **2010**, 2 (3), 219–231. <https://doi.org/10.1002/wnan.54>.

- (4) Stone, V.; Johnston, H.; Schins, R. P. F. Development of in Vitro Systems for Nanotoxicology: Methodological Considerations. *Crit. Rev. Toxicol.* **2009**, *39* (7), 613–626. <https://doi.org/10.1080/10408440903120975>.
- (5) Khalili Fard, J.; Jafari, S.; Eghbal, M. A. A Review of Molecular Mechanisms Involved in Toxicity of Nanoparticles. *Adv. Pharm. Bull.* **2015**, *5* (4), 447–454. <https://doi.org/10.15171/apb.2015.061>.
- (6) Bahadar, H.; Maqbool, F.; Niaz, K.; Abdollahi, M. Toxicity of Nanoparticles and an Overview of Current Experimental Models. *Iran. Biomed. J.* **2016**, *20* (1), 1–11. <https://doi.org/10.7508/ibj.2016.01.001>.
- (7) Chatterjee, R. Calculating the Costs of Nanohazard Testing. *Environ. Sci. Technol.* **2009**, *43* (10), 3405–3405. <https://doi.org/10.1021/es900758w>.
- (8) Hansch, C.; Maloney, P. P.; Fujita, T.; Muir, R. M. Correlation of Biological Activity of Phenoxyacetic Acids with Hammett Substituent Constants and Partition Coefficients. *Nature* **1962**, *194* (4824), 178–180. <https://doi.org/10.1038/194178b0>.
- (9) Fourches, D.; Pu, D.; Tassa, C.; Weissleder, R.; Shaw, S. Y.; Mumper, R. J.; Tropsha, A. Quantitative Nanostructure-Activity Relationship Modeling. *ACS Nano* **2010**, *4* (10), 5703–5712. <https://doi.org/10.1021/nn1013484>.
- (10) Puzyn, T.; Rasulev, B.; Gajewicz, A.; Hu, X.; Dasari, T. P.; Michalkova, A.; Hwang, H.-M.; Toropov, A.; Leszczynska, D.; Leszczynski, J. Using Nano-QSAR to Predict the Cytotoxicity of Metal Oxide Nanoparticles. *Nat. Nanotechnol.* **2011**, *6* (3), 175–178. <https://doi.org/10.1038/nnano.2011.10>.
- (11) Winkler, D. A.; Mombelli, E.; Pietroiusti, A.; Tran, L.; Worth, A.; Fadeel, B.; McCall, M. J. Applying Quantitative Structure-Activity Relationship Approaches to Nanotoxicology: Current Status and Future Potential. *Toxicology* **2013**, *313* (1), 15–23. <https://doi.org/10.1016/j.tox.2012.11.005>.

- (12) Gajewicz, A.; Schaeublin, N.; Rasulev, B.; Hussain, S.; Leszczynska, D.; Puzyn, T.; Leszczynski, J. Towards Understanding Mechanisms Governing Cytotoxicity of Metal Oxides Nanoparticles: Hints from Nano-QSAR Studies. *Nanotoxicology* **2015**, *9* (3), 313–325. <https://doi.org/10.3109/17435390.2014.930195>.
- (13) Pan, Y.; Li, T.; Cheng, J.; Telesca, D.; Zink, J. I.; Jiang, J. Nano-QSAR Modeling for Predicting the Cytotoxicity of Metal Oxide Nanoparticles Using Novel Descriptors. *RSC Adv.* **2016**, *6* (31), 25766–25775. <https://doi.org/10.1039/C6RA01298A>.
- (14) Gajewicz, A.; Jagiello, K.; Cronin, M. T. D.; Leszczynski, J.; Puzyn, T. Addressing a Bottle Neck for Regulation of Nanomaterials: Quantitative Read-across (Nano-QRA) Algorithm for Cases When Only Limited Data Is Available. *Environ. Sci. Nano* **2017**, *4* (2), 346–358. <https://doi.org/10.1039/C6EN00399K>.
- (15) Gajewicz, A. Development of Valuable Predictive Read-across Models Based on “Real-Life” (Sparse) Nanotoxicity Data. *Environ. Sci. Nano* **2017**, *4* (6), 1389–1403. <https://doi.org/10.1039/C7EN00102A>.
- (16) Gajewicz, A. What If the Number of Nanotoxicity Data Is Too Small for Developing Predictive Nano-QSAR Models? An Alternative Read-across Based Approach for Filling Data Gaps. *Nanoscale* **2017**, *9* (24), 8435–8448. <https://doi.org/10.1039/c7nr02211e>.
- (17) Gajewicz, A.; Cronin, M. T. D.; Rasulev, B.; Leszczynski, J.; Puzyn, T. Novel Approach for Efficient Predictions Properties of Large Pool of Nanomaterials Based on Limited Set of Species: Nano-Read-Across. *Nanotechnology* **2015**, *26* (1), 015701. <https://doi.org/10.1088/0957-4484/26/1/015701>.
- (18) Fjodorova, N.; Novic, M.; Gajewicz, A.; Rasulev, B. The Way to Cover Prediction for Cytotoxicity for All Existing Nano-Sized Metal Oxides by Using Neural Network Method. *Nanotoxicology* **2017**, *11* (4), 475–483. <https://doi.org/10.1080/17435390.2017.1310949>.

- (19) Gajewicz, A.; Puzyn, T.; Odziomek, K.; Urbaszek, P.; Haase, A.; Riebeling, C.; Luch, A.; Irfan, M. A.; Landsiedel, R.; van der Zande, M.; et al. Decision Tree Models to Classify Nanomaterials According to the DF4nanoGrouping Scheme. *Nanotoxicology* **2018**, *12* (1), 1–17. <https://doi.org/10.1080/17435390.2017.1415388>.
- (20) Lamon, L.; Asturiol, D.; Vilchez, A.; Ruperez-Illescas, R.; Cabellos, J.; Richarz, A.; Worth, A. Computational Models for the Assessment of Manufactured Nanomaterials: Development of Model Reporting Standards and Mapping of the Model Landscape. *Comput. Toxicol.* **2018**. <https://doi.org/10.1016/j.comtox.2018.12.002>.
- (21) Bañares, M. A.; Haase, A.; Tran, L.; Lobaskin, V.; Oberdörster, G.; Rallo, R.; Leszczynski, J.; Hoet, P.; Korenstein, R.; Hardy, B.; et al. CompNanoTox2015: Novel Perspectives from a European Conference on Computational Nanotoxicology on Predictive Nanotoxicology. *Nanotoxicology* **2017**, *11* (7), 839–845. <https://doi.org/10.1080/17435390.2017.1371351>.
- (22) ISO/TR 13014:2012. Nanotechnologies - Guidance on physico-chemical characterization of engineered nanoscale materials for toxicologic assessment [http://www.iso.org/iso/catalogue\\_detail?csnumber=52334](http://www.iso.org/iso/catalogue_detail?csnumber=52334) (accessed Nov 28, 2016).
- (23) Walkey, C. D.; Chan, W. C. W. Understanding and Controlling the Interaction of Nanomaterials with Proteins in a Physiological Environment. *Chem. Soc. Rev.* **2012**, *41* (7), 2780–2799. <https://doi.org/10.1039/c1cs15233e>.
- (24) Monopoli, M. P.; Aberg, C.; Salvati, A.; Dawson, K. A. Biomolecular Coronas Provide the Biological Identity of Nanosized Materials. *Nat. Nanotechnol.* **2012**, *7* (12), 779–786. <https://doi.org/10.1038/nnano.2012.207>.
- (25) Forest, V.; Cottier, M.; Pourchez, J. Electrostatic Interactions Favor the Binding of Positive Nanoparticles on Cells: A Reductive Theory. *Nano Today* **2015**, *10* (6), 677–680. <https://doi.org/10.1016/j.nantod.2015.07.002>.



- (26) Kar, S.; Gajewicz, A.; Puzyn, T.; Roy, K.; Leszczynski, J. Periodic Table-Based Descriptors to Encode Cytotoxicity Profile of Metal Oxide Nanoparticles: A Mechanistic QSTR Approach. *Ecotoxicol. Environ. Saf.* **2014**, *107*, 162–169. <https://doi.org/10.1016/j.ecoenv.2014.05.026>.
- (27) Toropova, A. P.; Toropov, A. A.; Rallo, R.; Leszczynska, D.; Leszczynski, J. Optimal Descriptor as a Translator of Eclectic Data into Prediction of Cytotoxicity for Metal Oxide Nanoparticles under Different Conditions. *Ecotoxicol. Environ. Saf.* **2015**, *112*, 39–45. <https://doi.org/10.1016/j.ecoenv.2014.10.003>.
- (28) Mu, Y.; Wu, F.; Zhao, Q.; Ji, R.; Qie, Y.; Zhou, Y.; Hu, Y.; Pang, C.; Hristozov, D.; Giesy, J. P.; et al. Predicting Toxic Potencies of Metal Oxide Nanoparticles by Means of Nano-QSARs. *Nanotoxicology* **2016**, *10* (9), 1207–1214. <https://doi.org/10.1080/17435390.2016.1202352>.
- (29) Pathakoti, K.; Huang, M.-J.; Watts, J. D.; He, X.; Hwang, H.-M. Using Experimental Data of Escherichia Coli to Develop a QSAR Model for Predicting the Photo-Induced Cytotoxicity of Metal Oxide Nanoparticles. *J. Photochem. Photobiol. B* **2014**, *130*, 234–240. <https://doi.org/10.1016/j.jphotobiol.2013.11.023>.
- (30) Zhai, X.; Chen, M.; Lu, W. Predicting the Toxicities of Metal Oxide Nanoparticles Based on Support Vector Regression with a Residual Bootstrapping Method. *Toxicol. Mech. Methods* **2018**, *28* (6), 440–449. <https://doi.org/10.1080/15376516.2018.1449278>.
- (31) Mikolajczyk, A.; Sizochenko, N.; Mulkiewicz, E.; Malankowska, A.; Nischk, M.; Jurczak, P.; Hirano, S.; Nowaczyk, G.; Zaleska-Medynska, A.; Leszczynski, J.; et al. Evaluating the Toxicity of TiO<sub>2</sub>-Based Nanoparticles to Chinese Hamster Ovary Cells and Escherichia Coli: A Complementary Experimental and Computational Approach. *Beilstein J. Nanotechnol.* **2017**, *8*, 2171–2180. <https://doi.org/10.3762/bjnano.8.216>.

- (32) Toropova, A. P.; Toropov, A. A.; Leszczynska, D.; Leszczynski, J. CORAL and Nano-QFAR: Quantitative Feature - Activity Relationships (QFAR) for Bioavailability of Nanoparticles (ZnO, CuO, Co<sub>3</sub>O<sub>4</sub>, and TiO<sub>2</sub>). *Ecotoxicol. Environ. Saf.* **2017**, *139*, 404–407. <https://doi.org/10.1016/j.ecoenv.2017.01.054>.
- (33) Sizochenko, N.; Gajewicz, A.; Leszczynski, J.; Puzyn, T. Causation or Only Correlation? Application of Causal Inference Graphs for Evaluating Causality in Nano-QSAR Models. *Nanoscale* **2016**, *8* (13), 7203–7208. <https://doi.org/10.1039/C5NR08279J>.
- (34) Toropova, A. P.; Toropov, A. A.; Veselinović, A. M.; Veselinović, J. B.; Benfenati, E.; Leszczynska, D.; Leszczynski, J. Nano-QSAR: Model of Mutagenicity of Fullerene as a Mathematical Function of Different Conditions. *Ecotoxicol. Environ. Saf.* **2016**, *124*, 32–36. <https://doi.org/10.1016/j.ecoenv.2015.09.038>.
- (35) Toropov, A. A.; Toropova, A. P. Quasi-SMILES and Nano-QFAR: United Model for Mutagenicity of Fullerene and MWCNT under Different Conditions. *Chemosphere* **2015**, *139*, 18–22. <https://doi.org/10.1016/j.chemosphere.2015.05.042>.
- (36) Speck-Planche, A.; Kleandrova, V. V.; Luan, F.; Cordeiro, M. N. D. S. Computational Modeling in Nanomedicine: Prediction of Multiple Antibacterial Profiles of Nanoparticles Using a Quantitative Structure-Activity Relationship Perturbation Model. *Nanomed.* **2015**, *10* (2), 193–204. <https://doi.org/10.2217/nnm.14.96>.
- (37) Toropov, A. A.; Toropova, A. P. Optimal Descriptor as a Translator of Eclectic Data into Endpoint Prediction: Mutagenicity of Fullerene as a Mathematical Function of Conditions. *Chemosphere* **2014**, *104*, 262–264. <https://doi.org/10.1016/j.chemosphere.2013.10.079>.
- (38) Toropov, A. A.; Toropova, A. P.; Benfenati, E.; Gini, G.; Puzyn, T.; Leszczynska, D.; Leszczynski, J. Novel Application of the CORAL Software to Model Cytotoxicity of

- Metal Oxide Nanoparticles to Bacteria Escherichia Coli. *Chemosphere* **2012**, 89 (9), 1098–1102. <https://doi.org/10.1016/j.chemosphere.2012.05.077>.
- (39) Toropov, A. A.; Toropova, A. P. Quasi-QSAR for Mutagenic Potential of Multi-Walled Carbon-Nanotubes. *Chemosphere* **2015**, 124, 40–46. <https://doi.org/10.1016/j.chemosphere.2014.10.067>.
- (40) Toropova, A. P.; Toropov, A. A. Mutagenicity: QSAR - Quasi-QSAR - Nano-QSAR. *Mini Rev. Med. Chem.* **2015**, 15 (8), 608–621.
- (41) Golbamaki, A.; Golbamaki, N.; Sizochenko, N.; Rasulev, B.; Leszczynski, J.; Benfenati, E. Genotoxicity Induced by Metal Oxide Nanoparticles: A Weight of Evidence Study and Effect of Particle Surface and Electronic Properties. *Nanotoxicology* **2018**, 1–17. <https://doi.org/10.1080/17435390.2018.1478999>.
- (42) Shin, H. K.; Kim, K. Y.; Park, J. W.; No, K. T. Use of Metal/Metal Oxide Spherical Cluster and Hydroxyl Metal Coordination Complex for Descriptor Calculation in Development of Nanoparticle Cytotoxicity Classification Model. *SAR QSAR Environ. Res.* **2017**, 28 (11), 875–888. <https://doi.org/10.1080/1062936X.2017.1400998>.
- (43) Helma, C.; Rautenberg, M.; Gebele, D. Nano-Lazar: Read across Predictions for Nanoparticle Toxicities with Calculated and Measured Properties. *Front. Pharmacol.* **2017**, 8, 377. <https://doi.org/10.3389/fphar.2017.00377>.
- (44) Toropova, A. P.; Toropov, A. A. Nano-QSAR in Cell Biology: Model of Cell Viability as a Mathematical Function of Available Eclectic Data. *J. Theor. Biol.* **2017**, 416, 113–118. <https://doi.org/10.1016/j.jtbi.2017.01.012>.
- (45) Basant, N.; Gupta, S. Modeling Uptake of Nanoparticles in Multiple Human Cells Using Structure-Activity Relationships and Intercellular Uptake Correlations. *Nanotoxicology* **2017**, 11 (1), 20–30. <https://doi.org/10.1080/17435390.2016.1257075>.

- (46) Papa, E.; Doucet, J. P.; Sangion, A.; Doucet-Panaye, A. Investigation of the Influence of Protein Corona Composition on Gold Nanoparticle Bioactivity Using Machine Learning Approaches. *SAR QSAR Environ. Res.* **2016**, *27* (7), 521–538. <https://doi.org/10.1080/1062936X.2016.1197310>.
- (47) Le, T. C.; Yin, H.; Chen, R.; Chen, Y.; Zhao, L.; Casey, P. S.; Chen, C.; Winkler, D. A. An Experimental and Computational Approach to the Development of ZnO Nanoparticles That Are Safe by Design. *Small.* **2016**, *12* (26), 3568–3577. <https://doi.org/10.1002/sml.201600597>.
- (48) Tsiliki, G.; Munteanu, C. R.; Seoane, J. A.; Fernandez-Lozano, C.; Sarimveis, H.; Willighagen, E. L. RRegrs: An R Package for Computer-Aided Model Selection with Multiple Regression Models. *J. Cheminformatics* **2015**, *7*, 46. <https://doi.org/10.1186/s13321-015-0094-2>.
- (49) Manganelli, S.; Leone, C.; Toropov, A. A.; Toropova, A. P.; Benfenati, E. QSAR Model for Predicting Cell Viability of Human Embryonic Kidney Cells Exposed to SiO<sub>2</sub> Nanoparticles. *Chemosphere* **2016**, *144*, 995–1001. <https://doi.org/10.1016/j.chemosphere.2015.09.086>.
- (50) Papa, E.; Doucet, J. P.; Doucet-Panaye, A. Linear and Non-Linear Modelling of the Cytotoxicity of TiO<sub>2</sub> and ZnO Nanoparticles by Empirical Descriptors. *SAR QSAR Environ. Res.* **2015**, *26* (7–9), 647–665. <https://doi.org/10.1080/1062936X.2015.1080186>.
- (51) Liu, R.; Jiang, W.; Walkey, C. D.; Chan, W. C. W.; Cohen, Y. Prediction of Nanoparticles-Cell Association Based on Corona Proteins and Physicochemical Properties. *Nanoscale* **2015**, *7* (21), 9664–9675. <https://doi.org/10.1039/c5nr01537e>.

- (52) Liu, R.; Rallo, R.; Bilal, M.; Cohen, Y. Quantitative Structure-Activity Relationships for Cellular Uptake of Surface-Modified Nanoparticles. *Comb. Chem. High Throughput Screen.* **2015**, *18* (4), 365–375.
- (53) Toropova, A. P.; Toropov, A. A.; Benfenati, E. A Quasi-QSPR Modelling for the Photocatalytic Decolourization Rate Constants and Cellular Viability (CV%) of Nanoparticles by CORAL. *SAR QSAR Environ. Res.* **2015**, *26* (1), 29–40. <https://doi.org/10.1080/1062936X.2014.984327>.
- (54) Toropova, A. P.; Toropov, A. A.; Benfenati, E.; Korenstein, R.; Leszczynska, D.; Leszczynski, J. Optimal Nano-Descriptors as Translators of Eclectic Data into Prediction of the Cell Membrane Damage by Means of Nano Metal-Oxides. *Environ. Sci. Pollut. Res. Int.* **2015**, *22* (1), 745–757. <https://doi.org/10.1007/s11356-014-3566-4>.
- (55) Maher, M. A.; Naha, P. C.; Mukherjee, S. P.; Byrne, H. J. Numerical Simulations of in Vitro Nanoparticle Toxicity - the Case of Poly(Amido Amine) Dendrimers. *Toxicol. Vitro Int. J. Publ. Assoc. BIBRA* **2014**, *28* (8), 1449–1460. <https://doi.org/10.1016/j.tiv.2014.07.014>.
- (56) Toropova, A. P.; Toropov, A. A.; Benfenati, E.; Puzyn, T.; Leszczynska, D.; Leszczynski, J. Optimal Descriptor as a Translator of Eclectic Information into the Prediction of Membrane Damage: The Case of a Group of ZnO and TiO<sub>2</sub> Nanoparticles. *Ecotoxicol. Environ. Saf.* **2014**, *108*, 203–209. <https://doi.org/10.1016/j.ecoenv.2014.07.005>.
- (57) Luan, F.; Kleandrova, V. V.; González-Díaz, H.; Ruso, J. M.; Melo, A.; Speck-Planche, A.; Cordeiro, M. N. D. S. Computer-Aided Nanotoxicology: Assessing Cytotoxicity of Nanoparticles under Diverse Experimental Conditions by Using a Novel QSTR-Perturbation Approach. *Nanoscale* **2014**, *6* (18), 10623–10630. <https://doi.org/10.1039/c4nr01285b>.

- (58) Ehret, J.; Vijver, M.; Peijnenburg, W. The Application of QSAR Approaches to Nanoparticles. *Altern. Lab. Anim. ATLA* **2014**, *42* (1), 43–50.
- (59) Winkler, D. A.; Burden, F. R.; Yan, B.; Weissleder, R.; Tassa, C.; Shaw, S.; Epa, V. C. Modelling and Predicting the Biological Effects of Nanomaterials. *SAR QSAR Environ. Res.* **2014**, *25* (2), 161–172. <https://doi.org/10.1080/1062936X.2013.874367>.
- (60) Kar, S.; Gajewicz, A.; Puzyn, T.; Roy, K. Nano-Quantitative Structure-Activity Relationship Modeling Using Easily Computable and Interpretable Descriptors for Uptake of Magnetofluorescent Engineered Nanoparticles in Pancreatic Cancer Cells. *Toxicol. Vitro Int. J. Publ. Assoc. BIBRA* **2014**, *28* (4), 600–606. <https://doi.org/10.1016/j.tiv.2013.12.018>.
- (61) Toropova, A. P.; Toropov, A. A. Optimal Descriptor as a Translator of Eclectic Information into the Prediction of Membrane Damage by Means of Various TiO<sub>2</sub> Nanoparticles. *Chemosphere* **2013**, *93* (10), 2650–2655. <https://doi.org/10.1016/j.chemosphere.2013.09.089>.
- (62) Toropov, A. A.; Toropova, A. P.; Puzyn, T.; Benfenati, E.; Gini, G.; Leszczynska, D.; Leszczynski, J. QSAR as a Random Event: Modeling of Nanoparticles Uptake in PaCa2 Cancer Cells. *Chemosphere* **2013**, *92* (1), 31–37. <https://doi.org/10.1016/j.chemosphere.2013.03.012>.
- (63) Liu, R.; Rallo, R.; Weissleder, R.; Tassa, C.; Shaw, S.; Cohen, Y. Nano-SAR Development for Bioactivity of Nanoparticles with Considerations of Decision Boundaries. *Small Wein. Bergstr. Ger.* **2013**, *9* (9–10), 1842–1852. <https://doi.org/10.1002/sml.201201903>.
- (64) Epa, V. C.; Burden, F. R.; Tassa, C.; Weissleder, R.; Shaw, S.; Winkler, D. A. Modeling Biological Activities of Nanoparticles. *Nano Lett.* **2012**, *12* (11), 5808–5812. <https://doi.org/10.1021/nl303144k>.

- (65) Sayes, C.; Ivanov, I. Comparative Study of Predictive Computational Models for Nanoparticle-Induced Cytotoxicity. *Risk Anal. Off. Publ. Soc. Risk Anal.* **2010**, *30* (11), 1723–1734. <https://doi.org/10.1111/j.1539-6924.2010.01438.x>.
- (66) Mukherjee, S. P.; Davoren, M.; Byrne, H. J. In Vitro Mammalian Cytotoxicological Study of PAMAM Dendrimers - towards Quantitative Structure Activity Relationships. *Toxicol. Vitro Int. J. Publ. Assoc. BIBRA* **2010**, *24* (1), 169–177. <https://doi.org/10.1016/j.tiv.2009.09.014>.
- (67) Choi, J.-S.; Trinh, T. X.; Yoon, T.-H.; Kim, J.; Byun, H.-G. Quasi-QSAR for Predicting the Cell Viability of Human Lung and Skin Cells Exposed to Different Metal Oxide Nanomaterials. *Chemosphere* **2019**, *217*, 243–249. <https://doi.org/10.1016/j.chemosphere.2018.11.014>.
- (68) Ojha, P. K.; Kar, S.; Roy, K.; Leszczynski, J. Toward Comprehension of Multiple Human Cells Uptake of Engineered Nano Metal Oxides: Quantitative Inter Cell Line Uptake Specificity (QICLUS) Modeling. *Nanotoxicology* **2018**, 1–21. <https://doi.org/10.1080/17435390.2018.1529836>.
- (69) Leone, C.; Bertuzzi, E. E.; Toropova, A. P.; Toropov, A. A.; Benfenati, E. CORAL: Predictive Models for Cytotoxicity of Functionalized Nanozeolites Based on Quasi-SMILES. *Chemosphere* **2018**, *210*, 52–56. <https://doi.org/10.1016/j.chemosphere.2018.06.161>.
- (70) Choi, J.-S.; Ha, M. K.; Trinh, T. X.; Yoon, T. H.; Byun, H.-G. Towards a Generalized Toxicity Prediction Model for Oxide Nanomaterials Using Integrated Data from Different Sources. *Sci. Rep.* **2018**, *8* (1), 6110. <https://doi.org/10.1038/s41598-018-24483-z>.
- (71) Trinh, T. X.; Choi, J.-S.; Jeon, H.; Byun, H.-G.; Yoon, T.-H.; Kim, J. Quasi-SMILES-Based Nano-Quantitative Structure-Activity Relationship Model to Predict the

- Cytotoxicity of Multiwalled Carbon Nanotubes to Human Lung Cells. *Chem. Res. Toxicol.* **2018**, *31* (3), 183–190. <https://doi.org/10.1021/acs.chemrestox.7b00303>.
- (72) Wang, W.; Sedykh, A.; Sun, H.; Zhao, L.; Russo, D. P.; Zhou, H.; Yan, B.; Zhu, H. Predicting Nano-Bio Interactions by Integrating Nanoparticle Libraries and Quantitative Nanostructure Activity Relationship Modeling. *ACS Nano* **2017**, *11* (12), 12641–12649. <https://doi.org/10.1021/acsnano.7b07093>.
- (73) Luan, F.; Tang, L.; Zhang, L.; Zhang, S.; Monteagudo, M. C.; Cordeiro, M. N. D. S. A Further Development of the QNAR Model to Predict the Cellular Uptake of Nanoparticles by Pancreatic Cancer Cells. *Food Chem. Toxicol. Int. J. Publ. Br. Ind. Biol. Res. Assoc.* **2018**, *112*, 571–580. <https://doi.org/10.1016/j.fct.2017.04.010>.
- (74) Cassano, A.; Marchese Robinson, R. L.; Palczewska, A.; Puzyn, T.; Gajewicz, A.; Tran, L.; Manganelli, S.; Cronin, M. T. D. Comparing the CORAL and Random Forest Approaches for Modelling the in Vitro Cytotoxicity of Silica Nanomaterials. *Altern. Lab. Anim. ATLA* **2016**, *44* (6), 533–556.
- (75) Oksel, C.; Winkler, D. A.; Ma, C. Y.; Wilkins, T.; Wang, X. Z. Accurate and Interpretable NanoSAR Models from Genetic Programming-Based Decision Tree Construction Approaches. *Nanotoxicology* **2016**, *10* (7), 1001–1012. <https://doi.org/10.3109/17435390.2016.1161857>.
- (76) Fourches, D.; Pu, D.; Li, L.; Zhou, H.; Mu, Q.; Su, G.; Yan, B.; Tropsha, A. Computer-Aided Design of Carbon Nanotubes with the Desired Bioactivity and Safety Profiles. *Nanotoxicology* **2016**, *10* (3), 374–383. <https://doi.org/10.3109/17435390.2015.1073397>.
- (77) Esposito, E. X.; Hopfinger, A. J.; Shao, C.-Y.; Su, B.-H.; Chen, S.-Z.; Tseng, Y. J. Exploring Possible Mechanisms of Action for the Nanotoxicity and Protein Binding of Decorated Nanotubes: Interpretation of Physicochemical Properties from Optimal QSAR



- Models. *Toxicol. Appl. Pharmacol.* **2015**, 288 (1), 52–62. <https://doi.org/10.1016/j.taap.2015.07.008>.
- (78) Shao, C.-Y.; Chen, S.-Z.; Su, B.-H.; Tseng, Y. J.; Esposito, E. X.; Hopfinger, A. J. Dependence of QSAR Models on the Selection of Trial Descriptor Sets: A Demonstration Using Nanotoxicity Endpoints of Decorated Nanotubes. *J. Chem. Inf. Model.* **2013**, 53 (1), 142–158. <https://doi.org/10.1021/ci3005308>.
- (79) Kleandrova, V. V.; Luan, F.; González-Díaz, H.; Ruso, J. M.; Speck-Planche, A.; Cordeiro, M. N. D. S. Computational Tool for Risk Assessment of Nanomaterials: Novel QSTR-Perturbation Model for Simultaneous Prediction of Ecotoxicity and Cytotoxicity of Uncoated and Coated Nanoparticles under Multiple Experimental Conditions. *Environ. Sci. Technol.* **2014**, 48 (24), 14686–14694. <https://doi.org/10.1021/es503861x>.
- (80) Concu, R.; Kleandrova, V. V.; Speck-Planche, A.; Cordeiro, M. N. D. S. Probing the Toxicity of Nanoparticles: A Unified in Silico Machine Learning Model Based on Perturbation Theory. *Nanotoxicology* **2017**, 11 (7), 891–906. <https://doi.org/10.1080/17435390.2017.1379567>.
- (81) Basant, N.; Gupta, S. Multi-Target QSTR Modeling for Simultaneous Prediction of Multiple Toxicity Endpoints of Nano-Metal Oxides. *Nanotoxicology* **2017**, 11 (3), 339–350. <https://doi.org/10.1080/17435390.2017.1302612>.
- (82) Kar, S.; Gajewicz, A.; Roy, K.; Leszczynski, J.; Puzyn, T. Extrapolating between Toxicity Endpoints of Metal Oxide Nanoparticles: Predicting Toxicity to *Escherichia Coli* and Human Keratinocyte Cell Line (HaCaT) with Nano-QTTR. *Ecotoxicol. Environ. Saf.* **2016**, 126, 238–244. <https://doi.org/10.1016/j.ecoenv.2015.12.033>.
- (83) Sizochenko, N.; Rasulev, B.; Gajewicz, A.; Kuz'min, V.; Puzyn, T.; Leszczynski, J. From Basic Physics to Mechanisms of Toxicity: The “Liquid Drop” Approach Applied

to Develop Predictive Classification Models for Toxicity of Metal Oxide Nanoparticles. *Nanoscale* **2014**, *6* (22), 13986–13993. <https://doi.org/10.1039/c4nr03487b>.

- (84) Toropov, A. A.; Rallo, R.; Toropova, A. P. Use of Quasi-SMILES and Monte Carlo Optimization to Develop Quantitative Feature Property/Activity Relationships (QFPR/QFAR) for Nanomaterials. *Curr. Top. Med. Chem.* **2015**, *15* (18), 1837–1844.

## Supporting Information

### Table of content:

Details on the 232 publications found when querying the PubMed database with the search terms (nanoparticle OR nanomaterials OR nanostructure) AND (QSAR OR QSPR OR QNAR OR QNTR).....S2-S11

	Authors	Journal	Ref in the text	No experimental data (reviews, commentaries)	Papers included in our analysis and performed on:			Papers not included in our analysis for the following reasons:			
					Prokaryotic cells	Eukaryotic cells	Both	No toxicity data	No modeling	<i>In vivo</i> data	Topic not related to human risk assessment
1	Lata and Vikas.	SAR QSAR Environ Res. 2019, 30(2):109-130.	/								x
2	Shchelokov <i>et al.</i>	Langmuir. 2018	/								x
3	Choi <i>et al.</i>	Chemosphere. 2019, 217:243-249.	67			x					
4	Salahinejad and Zolfonoun.	SAR QSAR Environ Res. 2018, 29(12):997-1009.	/								x
5	Möser <i>et al.</i>	Int J Mol Sci. 2018, 19(11), pii: E3482.	/								x
6	Ojha <i>et al.</i>	Nanotoxicology. 2018 :1-21.	68			x					
7	Dabrowska <i>et al.</i>	Int J Mol Sci. 2018, 19(10).	/								x
8	Ahmadi and Akbari.	SAR QSAR Environ Res. 2018, 29(11):895-909.	/								x
9	Tasi <i>et al.</i>	Nanoscale. 2018, 10(44):20863-20866.	/	x							
10	Wang <i>et al.</i>	Chemosphere. 2019, 214:79-84.	/								x
11	Utembe <i>et al.</i>	Environ Toxicol Chem. 2018, 37(12):2972-2988.	/	x							
12	Bhutto <i>et al.</i>	Talanta. 2018, 189:174-181.	/								x
13	Vracko <i>et al.</i>	SAR QSAR Environ Res. 2018, 29(8):567-577.	/				x				
14	Leone <i>et al.</i>	Chemosphere. 2018, 210:52-56.	69			x					
15	Lopalco and Denora.	Methods Mol Biol. 2018, 1800:347-365.	/	x							
16	Golbamaki <i>et al.</i>	Nanotoxicology. 2018, 9:1-17.	41			x					
17	Villaverde <i>et al.</i>	Sci Total Environ. 2018, 634:1530-1539.	/	x							
18	Wang <i>et al.</i>	Anal Bioanal Chem. 2018, 410(18):4379-4386.	/								x
19	Choi <i>et al.</i>	Sci Rep. 2018, 8(1):6110.	70			x					
20	Toropov <i>et al.</i>	Nanomaterials. 2018, 8(4). pii: E243.	/				x				
21	Zhai <i>et al.</i>	Toxicol Mech Methods. 2018, 28(6):440-449.	30		x						

22	Vracko <i>et al.</i>	Curr Comput Aided Drug Des. 2018, 14(1):2-4.	/	x							
23	Gupta and Bassant.	Chemosphere. 2018, 201:361-369.	/								x
24	Roh <i>et al.</i>	Mol Cell. 2018, 69(6):993-1004.e3.	/								x
25	Titma T.	Toxicol In Vitro. 2018, 50:11-21.	/						x		
26	Trinh <i>et al.</i>	Chem Res Toxicol. 2018, 31(3):183-190.	71			x					
27	Gajewicz <i>et al.</i>	Nanotoxicology. 2018,12(1):1-17.	/							x	
28	Shin <i>et al.</i>	SAR QSAR Environ Res. 2017, 28(11):875-888.	42			x					
29	Rajput <i>et al.</i>	Nucleic Acids Res. 2018, 46(D1):D894-D900.	/								x
30	Wang <i>et al.</i>	ACS Nano. 2017, 11(12):12641-12649.	72			x					
31	Mikolajczyk <i>et al.</i>	Beilstein J Nanotechnol. 2017, 8:2171-2180.	31		x						
32	Sizochenko <i>et al.</i>	Nanomaterials (Basel). 2017, 7(10). pii: E330.	/							x	
33	Zhao <i>et al.</i>	ACS Appl Mater Interfaces. 2017, 9(41):35740-35748.	/								x
34	Puzyn <i>et al.</i>	Food Chem Toxicol. 2018, 112:478-494.	/	x							
35	Concu <i>et al.</i>	Nanotoxicology. 2017, 11(7):891-906.	80				x				
36	Johnson <i>et al.</i>	Small. 2017, 13(42).	/								x
37	Bañares <i>et al.</i>	Nanotoxicology. 2017, 11(7):839-845.	21	x							
38	Kovalishyn <i>et al.</i>	Food Chem Toxicol. 2018, 112:507-517.	/	x							
39	Chen <i>et al.</i>	Int J Mol Sci. 2017, 18(7), pii: E1504.	/	x							
40	Helma <i>et al.</i>	Front Pharmacol. 2017, 8:377.	43			x					
41	Burello E.	Toxicol Sci. 2017, 159(2):339-353.	/							x	
42	Boukhvalov and Yoon.	Chem Res Toxicol. 2017, 30(8):1549-1555.	/								x
43	Lai <i>et al.</i>	Environ Sci Pollut Res Int. 2018, 25(4):3060-3077.	/	x							
44	Gajewicz A.	Nanoscale. 2017, 9(24):8435-8448.	16		x						
45	Wang <i>et al.</i>	Expert Opin Drug Discov. 2017, 12(8):769-784.	/	x							





98	Manganelli <i>et al.</i>	Chemosphere. 2016, 144:995-1001.	49			x					
99	Tsiliki <i>et al.</i>	J Cheminform. 2015, 7:46.	48			x					
100	Chen <i>et al.</i>	Altern Lab Anim. 2015, 43(4):221-40.	/	x							
101	Papa <i>et al.</i>	SAR QSAR Environ Res. 2015, 26(7-9):647-65.	50			x					
102	Esposito <i>et al.</i>	Toxicol Appl Pharmacol. 2015, 288(1):52-62.	77			x					
103	Zhang <i>et al.</i>	Anal Chim Acta. 2015, 880:130-5.	/								x
104	Rust <i>et al.</i>	Anal Chem. 2015, 87(14):7250-7.	/								x
105	Huang <i>et al.</i>	Chemosphere. 2015, 138:183-9.	/								x
106	Toropov and Toropova.	Chemosphere. 2015, 139:18-22.	35		x						
107	Melagraki and Afantitis.	Curr Top Med Chem. 2015, 15(18):1827-36.	/								x
108	Toropov <i>et al.</i>	Curr Top Med Chem. 2015, 15(18):1837-44.	84				x				
109	He <i>et al.</i>	Curr Top Med Chem. 2015, 15(18):1887-900.	/	x							
110	Salahinejad M.	Curr Top Med Chem. 2015, 15(18):1868-86.	/	x							
111	Liu <i>et al.</i>	Nanoscale. 2015, 7(21):9664-75.	51			x					
112	Yilmaz <i>et al.</i>	Nanomaterials. 2015, 5(2):778-791.	/					x			
113	Bygd <i>et al.</i>	Biomaterials. 2015, 56:187-97.	/							x	
114	Rabanel <i>et al.</i>	ACS Appl Mater Interfaces. 2015, 7(19):10374-85.	/								x
115	Apul <i>et al.</i>	J Hazard Mater. 2015, 295:138-44.	/								x
116	Toropov <i>et al.</i>	Comb Chem High Throughput Screen. 2015, 18(4):376-86.	/	x							
117	Liu <i>et al.</i>	Comb Chem High Throughput Screen. 2015, 18(4):365-75.	52			x					
118	Castellanos <i>et al.</i>	Nanoscale. 2015, 7(13):5654-64.	/								x
119	Galbiati E <i>et al.</i>	Bioconjug Chem. 2015, 26(4):680-9.	/								x
120	Toropova and Toropov.	Mini Rev Med Chem. 2015, 15(8):608-21.	40		x						





144	Nicolotti <i>et al.</i>	Drug Discov Today. 2014, 19(11):1757-1768.	/	x							
145	Gajewicz <i>et al.</i>	Nanotoxicology. 2015, 9(3):313-25.	12			x					
146	Kar <i>et al.</i>	Ecotoxicol Environ Saf. 2014, 107:162-9.	26		x						
147	Panneerselvam and Choi.	Int J Mol Sci. 2014, 15(5):7158-82.	/	x							
148	Ehret <i>et al.</i>	Altern Lab Anim. 2014, 42(1):43-50.	58			x					
149	Putz and Ori.	Molecules. 2014, 19(4):4157-88.	/								x
150	Delbecq F.	Adv Colloid Interface Sci. 2014, 209:98-108.	/	x							
151	Ma <i>et al.</i>	ACS Comb Sci. 2014, 16(4):155-9.	/								x
152	Winkler <i>et al.</i>	SAR QSAR Environ Res. 2014, 25(2):161-72.	59			x					
153	Andriotis <i>et al.</i>	J Chem Phys. 2014, 140(9):094705.	/								x
154	Zhao <i>et al.</i>	Analyst. 2014, 139(6):1498-503.	/								x
155	Kar <i>et al.</i>	Toxicol In Vitro. 2014, 28(4):600-6.	60			x					
156	Pathakoti <i>et al.</i>	J Photochem Photobiol B. 2014, 130:234-40.	29		x						
157	Cherkasov <i>et al.</i>	J Med Chem. 2014, 57(12):4977-5010.	/	x							
158	Lubinski <i>et al.</i>	SAR QSAR Environ Res. 2013, 24(12):995-1008.	/	x							
159	Ivask <i>et al.</i>	Nanotoxicology. 2014, 8 Suppl 1:57-71.	/	x							
160	Toropov and Toropova.	Chemosphere. 2014, 104:262-4.	37		x						
161	Toropova and Toropov.	Chemosphere. 2013, 93(10):2650-5.	61			x					
162	Sayes <i>et al.</i>	Int J Nanomedicine. 2013, 8 Suppl 1:45-56.	/					x			
163	Wang <i>et al.</i>	Acc Chem Res. 2013, 46(3):761-9.	/						x		
164	Kim <i>et al.</i>	Sci Total Environ. 2014, 470-471:1553-7.	/								x
165	Ahmed <i>et al.</i>	Org Biomol Chem. 2013, 11(35):5798-808.	/								x
166	Athauda <i>et al.</i>	ACS Appl Mater Interfaces. 2013, 5(13):6237-46.	/								x

167	Leifert <i>et al.</i>	Nanoscale. 2013, 5(14):6224-42.	/	x							
168	Kwok <i>et al.</i>	ACS Nano. 2013, 7(5):4668-82.	/								x
169	Li <i>et al.</i>	J Nanosci Nanotechnol. 2013, 13(2):1399-402.	/								x
170	Toropov <i>et al.</i>	Chemosphere. 2013, 92(1):31-7.	62			x					
171	Liu <i>et al.</i>	Small. 2013, 9(9-10):1842-52.	63			x					
172	Shao <i>et al.</i>	J Chem Inf Model. 2013, 53(1):142-58.	78			x					
173	Das <i>et al.</i>	Chem Pharm Bull (Tokyo). 2013, 61(2):125-33.	/								x
174	Winkler <i>et al.</i>	Toxicology. 2013, 313(1):15-23.	/	x							
175	Muthu MS.	Nanomedicine (Lond). 2012, 7(10):1471-3.	/								x
176	Kahru and Ivask.	Acc Chem Res. 2013, 46(3):823-33.	/								x
177	Cohen <i>et al.</i>	Acc Chem Res. 2013, 46(3):802-12.	/	x							
178	Epa <i>et al.</i>	Nano Lett. 2012, 12(11):5808-12.	64			x					
179	Westerhoff and Nowak	Acc Chem Res. 2013, 46(3):844-53.	/					x			
180	Apul <i>et al.</i>	Environ Sci Technol. 2013, 47(5):2295-303.	/					x			
181	Toropov <i>et al.</i>	Chemosphere. 2012, 89(9):1098-102.	38		x						
182	Gajewicz <i>et al.</i>	Adv Drug Deliv Rev. 2012, 64(15):1663-93.	/	x							
183	Ibrahim <i>et al.</i>	Mini Rev Med Chem. 2012, 12(6):447-51.	/								x
184	Saranathan <i>et al.</i>	J R Soc Interface. 2012, 9(75):2563-80.	/								x
185	Combes <i>et al.</i>	Adv Exp Med Biol. 2012, 745:v-xiii, xv, xvii passim.	/	x							
186	Misra <i>et al.</i>	Acta Biomater. 2012, 8(5):1908-17.	/						x		
187	Avila-Salas <i>et al.</i>	J Phys Chem B. 2012, 116(7):2031-9.	/								x
188	Szymański <i>et al.</i>	Int J Mol Sci. 2012, 13(1):427-52.	/	x							
189	Le <i>et al.</i>	Chem Rev. 2012, 112(5):2889-919.	/	x							
190	Ostrowski <i>et al.</i>	J Phys Chem A. 2012, 116(1):631-43.	/								x
191	Carbó-Dorca and Besalú.	SAR QSAR Environ Res. 2011, 22(7-8):661-5.	/	x							
192	Xia <i>et al.</i>	ACS Nano. 2011, 5(11):9074-81.	/								x



