



HAL
open science

p-value presheaves

Erwan Beurier, Dominique Pastor

► **To cite this version:**

Erwan Beurier, Dominique Pastor. p-value presheaves. [Research Report] IMTA-RR-2019-02-SC, IMT Atlantique. 2019. hal-02190029v1

HAL Id: hal-02190029

<https://hal.science/hal-02190029v1>

Submitted on 22 Jul 2019 (v1), last revised 28 Aug 2019 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

IMT Atlantique

Dépt. Signal & Communications
Technopôle de Brest-Iroise - CS 83818
29238 Brest Cedex 3
Téléphone: +33 (0)2 29 00 13 04
Télécopie: +33 (0)2 29 00 10 12
URL: www.imt-atlantique.fr



Collection des rapports de recherche d'IMT Atlantique
IMTA-RR-2019-02-SC

p-value presheaves

Erwan Beurier
IMT Atlantique
Dominique Pastor
IMT Atlantique

Date d'édition : July 21, 2019

Version : 1.0



IMT Atlantique

Bretagne-Pays de la Loire
École Mines-Télécom

Contents

1. Introduction	2
2. Notation	2
3. P-values	2
4. RDT and sheaves	3
5. First results	4
5.1. About p-value sheaves	4
5.2. About sheaf morphisms	6
5.3. About predicates	9
Index	10
References	10

1. Introduction

This report presents a preliminary study of the sheaf-ness properties of the p-value.

2. Notation

$\mathcal{B}(\mathbb{R}^n)$ is the Borel set (the smallest σ -algebra containing all open sets of \mathbb{R}^n).

3. P-values

Definition 3.1 (Size). A *size* is a function $\alpha : \mathcal{B}(\mathbb{R}^n) \rightarrow [0, 1]$.

Definition 3.2 (Retract). Let $\alpha : \mathcal{B}(\mathbb{R}^n) \rightarrow [0, 1]$ be a size.

A *retract* is a function $R : [0, 1] \rightarrow \mathcal{B}(\mathbb{R}^n)$ such that $\alpha \circ R = \text{id}_{[0,1]}$.

In other words, a retract is a right inverse for a size.

Definition 3.3 (Test). A *family of tests*, or simply *test*, is a retract $T : [0, 1] \rightarrow \mathcal{B}(\mathbb{R}^n)$ such that $T(0) = \emptyset$, $T(1) = \mathbb{R}^n$ and $a \leq a' \Rightarrow T(a) \leq T(a')$.

Remark 3.4. If $T : [0, 1] \rightarrow \mathcal{B}(\mathbb{R}^n)$ is a test, then for $a \in [0, 1]$, $T(a)$ is the rejection region of the null hypothesis with size a .

Definition 3.5 (p-value). Let T be a test.

For $x \in \mathbb{R}^n$, the p-value of x is defined as:

$$\text{pval}_T(x) = \inf(\{a \in [0, 1] \mid x \in T(a)\})$$

The p-value of x is the minimum size of the test that puts x into the rejection region.

Definition 3.6 (Topological space \mathcal{T}). For all $a \in [0, 1]$, we define $U_a = [0, a[$ and $\mathcal{T} = \{U_a \mid a \in [0, 1]\}$.

It is easy to see that:

Proposition 3.7. $([0, 1[, \mathcal{T})$ is a topological space.

Remark 3.8. Note that $U_a \cap U_b = U_{\min(a,b)}$ and $\bigcup_{a \in A} U_a = U_{\sup(A)}$.

Definition 3.9 (P-value sheaf). Let $T : [0, 1] \rightarrow \mathcal{B}(\mathbb{R}^n)$ be a test.

The p-value sheaf for T is the following functor:

$$S_T : \begin{cases} \mathcal{T}^{\text{op}} & \longrightarrow & \mathbf{Sets} \\ U_a & \longmapsto & \{x \in \mathbb{R}^n \mid \text{pval}_T(x) \geq a\} \\ U_a \subseteq U_b & \longmapsto & \begin{cases} S_T(U_b) & \longrightarrow & S_T(U_a) \\ x & \longmapsto & x \end{cases} \end{cases}$$

Proposition 3.10. Let $T : [0, 1] \rightarrow \mathcal{B}(\mathbb{R}^n)$ be a test and let S_T be its p-value sheaf.

Then, S_T is an actual sheaf.

Proof. It's easy to see that S_T is a presheaf: it's a functor $\mathcal{T}^{\text{op}} \rightarrow \mathbf{Sets}$. We now have to check the sheaf condition.

Let $A \subset [0, 1[$ and $(U_a)_{a \in A} \in \mathcal{T}^A$ be a cover of $U \in \mathcal{T}$. We have $U = \bigcup_{a \in A} U_a$. Let $(s_a)_{a \in A}$ be a matching family of S_T -sections over $(U_a)_{a \in A}$: $s_a \in S_T(U_a)$ and $s_a|_{U_a \cap U_b} = s_b|_{U_a \cap U_b}$.

By definition of S_T , we have:

$$s_a|_{U_a \cap U_b} = S_T(U_a \cap U_b \subset U_a)(s_a) = s_a$$

which yields, for all $a, b \in A$, $s_a = s_b$. Let $s = s_a$; then s satisfies $\forall a \in A$, $\text{pval}_T(s) \geq a$, so $\text{pval}_T(s) \geq \sup(A)$ and $s \in S_T(U_{\sup(A)}) = S_T\left(\bigcup_{a \in A} U_a\right)$. Consequently, s is the unique gluing of $(s_a)_{a \in A}$, and S_T satisfies the sheaf condition. \square

Consider the sheaf topos based on \mathcal{T} : $\mathbf{Shv}([0, 1[, \mathcal{T})$. Its subobject classifier is:

$$\Omega : \begin{cases} \mathcal{T}^{\text{op}} & \longrightarrow & \mathbf{Sets} \\ U_a & \longmapsto & \{U_{a'} \mid a' \leq a\} \\ U_a \subset U_b & \longmapsto & \begin{cases} \Omega(U_b) & \longrightarrow & \Omega(U_a) \\ U_c & \longmapsto & U_a \cap U_c = U_{\min(a,c)} \end{cases} \end{cases}$$

Let T be a test and S_T its associated p-value sheaf. A sheaf morphism $p : S_T \rightarrow \Omega$ makes the following diagram commute for all $a \leq b$:

$$\begin{array}{ccc} U_a & & S_T(U_b) \xrightarrow{p_{U_b}} \Omega(U_b) \\ \downarrow u & \rightsquigarrow & \downarrow S_T(u) \quad \checkmark \quad \downarrow \Omega(u) \\ U_b & & S_T(U_a) \xrightarrow{p_{U_a}} \Omega(U_a) \end{array}$$

For $x \in S_T(U_b)$, we have:

$$\begin{aligned} p_{U_a} \circ S_T(u)(x) &= \Omega(u) \circ p_{U_b}(x) \\ p_{U_a}(x) &= p_{U_b}(x) \cap U_a \end{aligned}$$

The canonical example is the following natural transformation:

$$p_{U_a} : \begin{cases} S_T(U_a) & \longrightarrow & \Omega(U_a) \\ x & \longmapsto & U_a \end{cases}$$

4. RDT and sheaves

Consider the set-theoretic integer $n \in \mathbb{N}$, $n = \{0, \dots, n-1\}$. Endow n with the discrete topology $(n, \mathcal{P}(n))$. For $I \subset n$, \mathbb{R}^I is the set of functions $I \rightarrow \mathbb{R}$; equivalently, it is the set of I -indexed sets of real numbers.

Let Π_n be the following functor:

$$\Pi_n : \begin{cases} \mathcal{P}(n) & \longrightarrow & \mathbf{Sets} \\ I & \longmapsto & \mathbb{R}^I \\ I \subset J & \longmapsto & \begin{cases} \Pi_n(J) & \longrightarrow & \Pi_n(I) \\ f & \longmapsto & f|_I \end{cases} \end{cases}$$

Proposition 4.1. Π_n is a sheaf.

For $I \subset n$, $\|\cdot\|_{\mathbb{R}^I}$ is the Euclidean norm on \mathbb{R}^I .

Definition 4.2 (τ functor). We define the functor τ by setting:

$$\tau : \begin{cases} \mathcal{P}(n) & \longrightarrow & \mathcal{T} \\ I & \longmapsto & \tau(I) \\ I \subset J & \longmapsto & \tau(I) \subset \tau(J) \end{cases}$$

where \mathcal{T} is the topology defined in Definition 3.6. For $I \in \mathcal{P}(n)$, the unique a such that $\tau(I) = [0, a[$ is denoted τ_I , so that: $\tau(I) = [0, \tau_I[$.

The interest of this definition will tentatively be given later.

Let $f, \theta \in \mathbb{R}^n$. For all $I \in \mathcal{P}(n)$, the proposition: $\|\theta|_I - f|_I\|_{\mathbb{R}^I} \leq \tau_I$ is a proposition the truth of which can be known by simple inspection, provided that θ is known. In practice, we don't have access to θ but to a modified, noisy version $Y = Y(\theta)$ of it, and the question is then, for any given $I \subset n$, whether the proposition $\|\theta|_I - f|_I\|_{\mathbb{R}^I} \leq \tau_I$ is plausible in a certain sense or not, when we observe $Y|_I$.

Definition 4.3 (Optimal pair). Let $f \in \mathbb{R}^n$, we call it model. Let Θ be a random vector whose distribution we don't know, and $X \sim \mathcal{N}(0, C^2)$ is a n -dimensional Gaussian vector. We observe $Y = \Theta + X$.

For all $I \subset n$, there exists a pair $(\alpha_I^{f,\tau}, T_I^{f,\tau})$ [1], called *optimal pair restricted to I* , where:

1. $\alpha_I^{f,\tau} : \mathcal{B}(\mathbb{R}^I) \rightarrow [0, 1]$ is a size
2. $T_I^{f,\tau} : [0, 1] \rightarrow \mathcal{B}(\mathbb{R}^I)$ is a monotone retract of $\alpha_I^{f,\tau}$
3. $T_I^{f,\tau}$ is optimal for testing $\|\Theta|_I - f|_I\|_{\mathbb{R}^I} \leq \tau_I$ against its alternative $\|\Theta|_I - f|_I\|_{\mathbb{R}^I} \geq \tau_I$ when we observe $Y(\Theta)$.

Definition 4.4 (Restricted p-value). We consider the τ functor described above. Let $I \subset n$ and let $(\alpha_I^{f,\tau}, T_I^{f,\tau})$ be an optimal pair restricted to I .

The *I -restricted p-value of x* is the following value:

$$\text{pval}_I^{f,\tau}(y) = \inf \left(\left\{ a \in [0, 1] \mid y \in T_I^{f,\tau}(a) \right\} \right)$$

The *I -restricted p-value sheaf* is the following sheaf:

$$S_I^{f,\tau} : \begin{cases} [0, 1] & \longrightarrow & \mathbb{R}^I \\ U_a & \longmapsto & \left\{ y \in \mathbb{R}^n \mid \text{pval}_I^{f,\tau}(y) \geq a \right\} \\ U_a \subseteq U_b & \longmapsto & \begin{cases} S_T(U_b) & \longrightarrow & S_T(U_a) \\ x & \longmapsto & x \end{cases} \end{cases}$$

The p-value $\text{pval}_I^{f,\tau}(y)$ measures the plausability of proposition $\|\Theta|_I - f|_I\|_{\mathbb{R}^I} \leq \tau_I$ when we observe $y \sim Y = \Theta + X$. The higher $\text{pval}_I^{f,\tau}(y)$, the more plausible the proposition $\|\Theta|_I - f|_I\|_{\mathbb{R}^I} \leq \tau_I$.

5. First results

5.1. About p-value sheaves

It is easy to see that:

Proposition 5.1. (\mathcal{T}, \subset) is a total order with a minimal and maximal element (resp. \emptyset and \mathbb{R}^d).

It is also easy, but less easy, to see that:

Proposition 5.2. \mathcal{T}^{op} has all small limits and colimits.

Proof. Let $D : \mathcal{I} \rightarrow \mathcal{T}^{\text{op}}$ be a diagram in \mathcal{T}^{op} . For all $i \in \mathcal{I}$, define $U_{d_i} = D(i)$. Then:

$$\begin{aligned} \text{Colim}(D) &= \sup_{i \in \text{Ob}_{\mathcal{I}}} D_i = \bigcap_{i \in \text{Ob}_{\mathcal{I}}} U_{d_i} = U_{\inf(\mathcal{I})} \\ \text{Lim}(D) &= \inf_{i \in \text{Ob}_{\mathcal{I}}} D_i = \bigcup_{i \in \text{Ob}_{\mathcal{I}}} U_{d_i} = U_{\sup(\mathcal{I})} \end{aligned}$$

Note that the sup becomes an intersection, and the inf becomes a union, because we are considering \mathcal{T}^{op} and not \mathcal{T} . Also note that the infima and suprema always exist because these are infima and suprema of $\{d_i \mid i \in \text{Ob}_{\mathcal{I}}\}$, which is a subset of \mathbb{R} . \square

Proposition 5.3. For any test T , its p-value sheaf S_T is continuous and cocontinuous.

Proof. Let $D : \mathcal{I} \rightarrow \mathcal{T}^{\text{op}}$ be any (small) diagram in \mathcal{T}^{op} . For all $i \in \mathcal{I}$, define $U_{d_i} = D(i)$.

We only consider the case of a limit; the proof is very similar for colimits.

$$\begin{aligned}
 S_T(\text{Lim}(D)) &= S_T\left(\bigcup_{i \in \mathcal{I}} U_{d_i}\right) \\
 &= S_T(U_{\sup(d_i)}) \\
 &= \left\{x \in \mathbb{R}^d \mid \text{pval}_T(x) \geq \sup_{i \in \mathcal{I}}(d_i)\right\} \\
 &= \left\{x \in \mathbb{R}^d \mid \forall i \in \mathcal{I}, \text{pval}_T(x) \geq d_i\right\} \\
 &= \bigcap_{i \in \mathcal{I}} \left\{x \in \mathbb{R}^d \mid \text{pval}_T(x) \geq d_i\right\} \\
 &= \bigcap_{i \in \mathcal{I}} S_T(U_{d_i})
 \end{aligned}$$

We now have to check that $\bigcap_{i \in \mathcal{I}} S_T(U_{d_i}) \cong \text{Lim}(S_T \circ D)$.

For all $i \rightarrow j \in \mathcal{I}$, we have $S_T(U_{d_i}) \subset S_T(U_{d_j})$. We denote by $\iota_{i,j} = S_T(U_{d_i} \subset U_{d_j}) : S_T(U_{d_i}) \rightarrow S_T(U_{d_j})$ the inclusion mapping between $S_T(U_{d_i})$'s. We also denote by $\iota_i : \bigcap_{i \in \mathcal{I}} S_T(U_{d_i}) \rightarrow S_T(U_{d_i})$ the inclusion mapping of the intersection. For all $i \rightarrow j \in \mathcal{I}$, we have $\iota_{i,j} \circ \iota_i = \iota_j$, so that $\iota = \left(\iota_i : \bigcap_{i \in \mathcal{I}} S_T(U_{d_i}) \rightarrow S_T(U_{d_i})\right)_{i \in \mathcal{I}}$ is a cone to $S_T \circ D$.

Let (A, α) be any cone to $S_T \circ D$. We denote the $S_T(U_{d_i})$ -components of α by $\alpha_i : A \rightarrow S_T(U_{d_i})$. For all $i \rightarrow j \in \mathcal{I}$, and for all $x \in X$, we have $\iota_{i,j} \circ \alpha_i(x) = \alpha_j(x) = \alpha_i(x)$.

$$\begin{array}{ccc}
 S_T(U_{d_i}) & \xleftarrow{\alpha_i} & A \\
 \downarrow \iota_{i,j} & \swarrow \alpha_j & \\
 S_T(U_{d_j}) & &
 \end{array}$$

So in fact, $\alpha_i(X) = \alpha_j(X) \subset S_T(U_{d_i})$, which yields that, for all $i \in \mathcal{I}$, $\alpha_i(X) \subset \bigcap_{i \in \mathcal{I}} S_T(U_{d_i})$.

Let u be such that:

$$u : \begin{cases} X & \longrightarrow \bigcap_{i \in \mathcal{I}} S_T(U_{d_i}) \\ x & \longmapsto \alpha_i(x) \end{cases}$$

for any of the $i \in \mathcal{I}$, because $\alpha_j(x) = \alpha_i(x)$. Then for all $i \in \mathcal{I}$, we have $\iota_i \circ u = \alpha_i$.

It is also easy to check the unicity of that u : suppose $u, u' : X \rightarrow \bigcap_{i \in \mathcal{I}} S_T(U_{d_i})$, then for all $x \in X$, we have:

$$\begin{aligned}
 \iota_i \circ u(x) &= \alpha_i(x) = \iota_i \circ u'(x) \\
 u(x) &= u'(x)
 \end{aligned}$$

which leads to $u = u'$.

Consequently, $\bigcap_{i \in \mathcal{I}} S_T(U_{d_i}) \cong \text{Lim}(S_T \circ D)$. □

5.2. About sheaf morphisms

Let us first study the sheaf morphisms between p-value sheaves $p : S_R \rightarrow S_T$.

Proposition 5.4 (Nesting property). *Let $R, T : [0, 1] \rightarrow \mathcal{B}(\mathbb{R}^n)$ be two tests (of the same size or not). Let $p : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a function.*

The function p defines a natural transformation $\bar{p} : S_R \rightarrow S_T = (p|_{S_R(U_a)})_{a \in [0,1]} \Leftrightarrow$ for all $a \in [0, 1]$, $p(S_R(U_a)) \subset S_T(U_a)$.

In other words, p defines a function between nested open sets, as in Figure 1.




Figure 1: Illustration of the nesting property of p . If $p : \mathbb{R}^d \rightarrow \mathbb{R}^d$ verifies this nesting property, then it defines a natural transformation between two p-value sheaves.

Proof. Easy deduction from the following natural transformation diagram:

$$\begin{array}{ccc}
 U_a & & S_R(U_b) \xrightarrow{p|_{S_R(U_b)}} S_T(U_b) \\
 \downarrow \subset & \rightsquigarrow & \downarrow \subset \quad \checkmark \quad \downarrow \subset \\
 U_b & & S_R(U_a) \xrightarrow{p|_{S_R(U_a)}} S_T(U_a)
 \end{array}$$

□

Corollary 5.5. *If $\bar{p} = (p_{U_a})_{a \in [0,1]}$ is a natural transformation $S_R \rightarrow S_T$, then $p = \bigcup_{a \in A} p_{U_a}$ is a function that verifies the nesting property.*

Proposition 5.6. *For all $a \in [0, 1]$, $S_T(U_a) = \mathbb{R}^d \setminus \left(\bigcup_{b < a} T(b) \right)$.*

Proof. By computation:

$$\begin{aligned}
 S_T(U_a) &= \{x \in \mathbb{R}^d \mid \text{pval}_T(x) \geq a\} \\
 &= \left\{x \in \mathbb{R}^d \mid \inf_{x \in T(b)} (b \in [0, 1]) \geq a\right\} \\
 &= \{x \in \mathbb{R}^d \mid \forall b < a, x \notin T(b)\} \\
 &= \bigcap_{b < a} \mathbb{R}^d \setminus T(b) \\
 &= \mathbb{R}^d \setminus \left(\bigcup_{b < a} T(b)\right)
 \end{aligned}$$

□

Corollary 5.7. For all $a \in [0, 1]$, if $\bigcup_{b < a} T(b) = T(a)$ then $S_T(U_a) = \mathbb{R}^d \setminus T(a)$.

Example 5.8. Note that we are considering very general tests $T : [0, 1] \rightarrow \mathcal{B}(\mathbb{R}^n)$, so $\left(\bigcup_{b < a} T(b)\right)$ has no reason to be equal to $T(a)$. Let us give a counterexample.

Consider the following functions:

$$\begin{aligned}
 f : \begin{cases} [0, 1] & \longrightarrow & [0, 1] \\ x & \longmapsto & \begin{cases} x & \text{if } x < \frac{1}{2} \\ \frac{1}{2}x + \frac{1}{2} & \text{if } x \geq \frac{1}{2} \end{cases} \end{cases} \\
 g : \begin{cases}]0, 1[& \longrightarrow & \mathbb{R} \\ x & \longmapsto & \tan\left(\pi x - \frac{\pi}{2}\right) \end{cases} \\
 T : \begin{cases} [0, 1] & \longrightarrow & \mathcal{B}(\mathbb{R}^n) \\ x & \longmapsto & \begin{cases}]-g \circ f(x), g \circ f(x)[& \text{if } a < 1 \\ \mathbb{R}^d & \text{if } a = 1 \end{cases} \end{cases}
 \end{aligned}$$

The function f is strictly increasing and establishes a bijection $[0, 1] \rightarrow [0, \frac{1}{2}[\cup [\frac{3}{4}, 1]$. Then g is bijective and strictly increasing, so $g \circ f$ is strictly increasing and injective, and finally T is strictly increasing in the Borel set, so it is injective, and has a left inverse $\mathcal{B}(\mathbb{R}^n) \rightarrow [0, 1]$ (which is a size). In other words, T is a test. However:

$$\begin{aligned}
 \bigcup_{a < \frac{1}{2}} T(a) &= \bigcup_{a < \frac{1}{2}}]-g \circ f(a), g \circ f(a)[\\
 &= \left] -\lim_{a < \frac{1}{2}} g \circ f(a), \lim_{a < \frac{1}{2}} g \circ f(a) \right[\\
 &= \left] -g\left(\lim_{a < \frac{1}{2}} f(a)\right), g\left(\lim_{a < \frac{1}{2}} f(a)\right) \right[\\
 &= \left] -g\left(\frac{1}{2}\right), g\left(\frac{1}{2}\right) \right[\\
 &\subsetneq T\left(\frac{1}{2}\right) = \left] -g\left(\frac{3}{4}\right), g\left(\frac{3}{4}\right) \right[
 \end{aligned}$$

Definition 5.9 (Continuous test). Let T be a test. We call T a *continuous test* when, for all $a \in [0, 1]$, we have $\bigcup_{b < a} T(b) = T(a)$.

In the following, we give an example of such a continuous test.

Definition 5.10 (Likelihood ratio). Let $f_0, f_1 : \mathbb{R}^d \rightarrow \mathbb{R}^+$ be two probability density functions, and let $x \in \mathbb{R}^d$.

The *likelihood ratio of x being from $X \sim f_0$ instead of $X \sim f_1$* , written $\Lambda^{f_0, f_1}(x)$ or simply $\Lambda(x)$ when there is no ambiguity, is the following function:

$$\Lambda : \begin{cases} \mathbb{R}^d & \longrightarrow & \mathbb{R}^+ \cup \{\infty\} \\ x & \longmapsto & \begin{cases} \frac{f_0(x)}{f_1(x)} & \text{if } f_1(x) \neq 0 \\ \infty & \text{if } f_1(x) = 0 \text{ and } f_0(x) > 0 \\ 0 & \text{if } f_1(x) = 0 \text{ and } f_0(x) = 0 \end{cases} \end{cases} \quad (1)$$

In the practical case where f_1 is always strictly positive on its domain, Λ has values only in \mathbb{R}^+ , and Λ becomes a measurable function.

Consider a random variable $X \sim f_0$. One can compute $\mathbb{P}[\Lambda(X) \leq k | X \sim f_0] = a$. When $k = 0$, we have $a = 1$. When $k \rightarrow \infty$, we have $a \rightarrow 0$.

Definition 5.11 (NP-threshold function). Let $f_0, f_1 : \mathbb{R}^d \rightarrow \mathbb{R}^+$ be two probability density functions so that $\Lambda^{f_0, f_1} = \Lambda$ is continuous.

The *NP-threshold function*, written k_{NP} or simply k , is the function that assigns to each $a \in [0, 1]$, threshold $k(a)$ such that:

$$a = \int_{\Lambda(x) \leq k(a)} f_0(x) dx = \mathbb{P}[\Lambda(X) \leq k | X \sim f_0]$$

Proposition 5.12. *The threshold function is continuous and decreasing.*

Proof. To be completed. □

Definition 5.13 (Neyman-Pearson test). Let $f_0, f_1 : \mathbb{R}^d \rightarrow \mathbb{R}^+$ be two probability density functions so that $\Lambda^{f_0, f_1} = \Lambda$ is continuous.

The *Neyman-Pearson test* is the following test:

$$\text{NP} : \begin{cases} [0, 1] & \longrightarrow & \mathcal{B}(\mathbb{R}^n) \\ a & \longmapsto & \{x \in \mathbb{R}^d \mid \Lambda(x) \leq k(a)\} \end{cases}$$

Proposition 5.14. *For all $a \in]0, 1[$, we have $\bigcup_{b < a} \text{NP}(b) = \text{NP}(a)$. In other words, NP is a continuous test.*

Proof. This is due to the continuity of the threshold function k , which remains to be proved:

$$\begin{aligned} \bigcup_{b < a} \text{NP}(b) &= \bigcup_{b < a} \{x \in \mathbb{R}^d \mid \Lambda(x) \leq k(b)\} \\ &= \{x \in \mathbb{R}^d \mid \exists b < a, \Lambda(x) \leq k(b)\} \\ &= \left\{x \in \mathbb{R}^d \mid \Lambda(x) \leq \sup_{b < a} (k(b))\right\} \\ &= \{x \in \mathbb{R}^d \mid \Lambda(x) \leq k(a)\} \\ &= \text{NP}(a) \end{aligned}$$

□

Corollary 5.15. $S_{\text{NP}}(U_a) = \mathbb{R} \setminus \text{NP}(a)$.

5.3. About predicates

Definition 5.16 (Predicate). Let S be a sheaf in $\mathbf{Shv}([0, 1[, \mathcal{T})$.

A *predicate* is a sheaf morphism $p : S \rightarrow \Omega$.

Computing predicates

Let $S : \mathcal{T}^{\text{op}} \rightarrow \mathbf{Sets}$ be a sheaf, and let $p_1, p_2 : S \rightarrow \Omega$ be predicates. Let $a \in [0, 1]$. For $x \in S(U_a)$, define $U_{a_1} = p_{1,U_a}(x)$ and $U_{a_2} = p_{2,U_a}(x)$. We have the trivial results:

$$\begin{aligned}(p_1 \wedge p_2)_{U_a}(x) &= U_{\min(a_1, a_2)} \\ (p_1 \vee p_2)_{U_a}(x) &= U_{\max(a_1, a_2)}\end{aligned}$$

Also:

$$\begin{aligned}(p_1 \rightarrow p_2)_{U_a}(x) &= p_{1,U_a}(x) \rightarrow p_{2,U_a}(x) \\ &= \bigcup_{R \cap U_{a_1} \subset U_{a_2}} R \\ &= \bigcup_{\substack{b \in [0, a] \\ U_b \cap U_{a_1} \subset U_{a_2}}} U_b \\ &= \bigcup_{\substack{b \in [0, a] \\ U_{\min(a_1, b)} \subset U_{a_2}}} U_b \\ &= \bigcup_{\substack{b \in [0, a] \\ \min(a_1, b) \leq a_2}} U_b \\ &= U_c\end{aligned}$$

for some $c \in [0, a]$. Let us take a look at that c . We have:

$$c = \sup_{\substack{b \in [0, a] \\ \min(a_1, b) \leq a_2}} (b) = \begin{cases} a & \text{if } a_1 \leq a_2 \\ a_2 & \text{otherwise} \end{cases}$$

The negation $\neg p_1$ corresponds to the special case where $U_{a_2} = U_0 = \emptyset$:

$$\begin{aligned}(\neg p_1)_{U_a}(x) &= p_{1,U_a}(x) \rightarrow \emptyset \\ &= \bigcup_{\substack{b \in [0, a] \\ \min(a_1, b) \leq 0}} U_b \\ &= U_c\end{aligned}$$

With c being:

$$c = \sup_{\substack{b \in [0, a] \\ \min(a_1, b) \leq 0}} (b) = \begin{cases} a & \text{if } a_1 = 0 \\ 0 & \text{if } a_1 > 0 \end{cases}$$

In summary:

Formula	Condition	Result
$p_{1,U_a}(x)$		U_{a_1}
$p_{2,U_a}(x)$		U_{a_2}
$(p_1 \wedge p_2)_{U_a}(x)$		$U_{\min(a_1, a_2)}$
$(p_1 \vee p_2)_{U_a}(x)$		$U_{\max(a_1, a_2)}$
$(p_1 \rightarrow p_2)_{U_a}(x)$	if $a_1 \leq a_2$ if $a_1 > a_2$	U_a U_{a_2}
$(\neg p_1)_{U_a}(x)$	if $a_1 = 0$ if $a_1 > 0$	U_a $U_0 = \emptyset$


References

- [1] D. Pastor and Q.-T. Nguyen, "Random distortion testing and optimality of thresholding tests," *IEEE Transactions on Signal processing*, vol. 61, no. 16, pp. 4161–4171, 2013.

OUR WORLDWIDE PARTNERS UNIVERSITIES - DOUBLE DEGREE AGREEMENTS



3 CAMPUS, 1 SITE



IMT Atlantique Bretagne-Pays de la Loire – <http://www.imt-atlantique.fr/>

Campus de Brest

Technopôle Brest-Iroise
CS 83818
29238 Brest Cedex 3
France
T +33 (0)2 29 00 11 11
F +33 (0)2 29 00 10 00

Campus de Nantes

4, rue Alfred Kastler
CS 20722
44307 Nantes Cedex 3
France
T +33 (0)2 51 85 81 00
F +33 (0)2 99 12 70 08

Campus de Rennes

2, rue de la Châtaigneraie
CS 17607
35576 Cesson Sévigné Cedex
France
T +33 (0)2 99 12 70 00
F +33 (0)2 51 85 81 99

Site de Toulouse

10, avenue Édouard Belin
BP 44004
31028 Toulouse Cedex 04
France
T +33 (0)5 61 33 83 65



IMT Atlantique
Bretagne-Pays de la Loire
École Mines-Télécom

© IMT Atlantique, 2019
Imprimé à IMT Atlantique
Dépôt légal : Juillet 2019
ISSN : 2556-5060