Preface to the Special Section: Dense Water Formations in the Northwestern Mediterranean: From the Physical Forcings to the Biogeochemical Consequences

Pascal Conan, Pierre Testor, Claude Estournel, Fabrizio d'Ortenzio, Mireille Pujo-Pay, Xavier Durrieu de Madron

▶ **To cite this version:**

Pascal Conan, Pierre Testor, Claude Estournel, Fabrizio d'Ortenzio, Mireille Pujo-Pay, et al.. Preface to the Special Section: Dense Water Formations in the Northwestern Mediterranean: From the Physical Forcings to the Biogeochemical Consequences. Journal of Geophysical Research. Oceans, 2018, 123 (10), pp.6983-6995. 10.1029/2018JC014301 . hal-02190724

HAL Id: hal-02190724
https://hal.science/hal-02190724v1

Submitted on 15 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Preface to the Special Section: Dense water formations in the North Western Mediterranean: from the physical forcings to the biogeochemical consequences

Pascal Conan[1], Pierre Testor[2], Claude Estournel[3], Fabrizio D'Ortenzio[4], Mireille Pujo-Pay[1], Xavier Durrieu de Madron[5]

[1] Sorbonne Université, UPMC, Univ. Paris 06, INSU-CNRS, UMR 7621, Laboratoire d'Océanographie Microbienne (LOMIC), Observatoire Océanologique de Banyuls, 66650 Banyuls-sur-Mer, France

[2] Sorbonne Université, UPMC, Univ. Paris 06, CNRS-IRD-MNHN, Laboratoire d'Océanographie et du Climat : Expérimentation et Approches Numériques (LOCEAN), IPSL, 75252 Paris, France

[3] Laboratoire d'Aérologie, Université de Toulouse, CNRS, UPS, 14 avenue Edouard Belin, 31400 Toulouse, France

[4] Sorbonne Université, UPMC Univ. Paris 06, INSU-CNRS, UMR 7093, Laboratoire d'Océanographie de Villefranche, 181 Chemin du Lazaret, 06230 Villefranche-sur-Mer, France

[5] Université de Perpignan Via Domitia, CNRS UMR 5110, Centre d'Etude et de Formation sur les Environnements Méditerranéens, 52 avenue Paul Alduy, 66860 Perpignan, France

Correspondence to: P. Conan; pascal.conan@obs-banyuls.fr

Short Abstract 2

Key Points 3

Program acronyms 3

1. Rationales 4

2. Strategic plan 6

3. Major outcomes 8

3.1. Dynamics of the atmosphere and air-sea fluxes 8

3.2. Interannual variability of dense water formation 9

3.3. Deep water formation during winter 2013 10

3.4. Effects of small scales and intrinsic ocean variability on dense water formation 11

3.5. Effect of dense water formation on biogeochemical budgets and primary production 12

3.6. Zooplankton community response to dense water formation 14

Short Abstract

The northwestern Mediterranean Sea is one of the few sites of open-sea deep convection and dense water formation. The area is characterized by intense air-sea exchanges favored by the succession of strong northerly and northwesterly winds during autumn and winter that eventually induce deep convection episodes and the formation of the 'Western Mediterranean Deep Water'. This region exhibits a significant spring phytoplankton bloom, which appears to be largely influenced in intensity and diversity by the winter mixing.

To understand and resolve the interplay between the atmosphere and ocean, and the impact of ocean circulation and mixing on biogeochemistry, we carried out an unprecedented observational effort during a large experiment between July 2012 and July 2013. A multiplatform approach - combining aircraft, balloons, ships, moorings, floats, and gliders - aimed both at characterizing the dynamics of the atmosphere and at quantifying the physical, biogeochemical and biological properties of the water masses. Beyond a better understanding of the wind dynamics, the interannual variability of the deep convection, and the seasonal variability of the nutrient distribution and plankton structure in the pelagic ecosystem, the experiment provided a reference data set that was used as a benchmark for advancing the modeling of the surface fluxes, convective processes, dense water formation rates, and physical-biogeochemical coupling processes. It also represented an opportunity for complementary investigations such as evaluating model parameterizations or studying the role of sub-mesoscale eddies in dense water spreading and biogeochemistry.

Detailed Abstract

Open sea deep convection and dense water formation constitute an important driving mechanism for the ocean circulation and ventilation. This mechanism also strongly impacts the nutrients distribution and controls the triggering of the blooms of the first producer levels and then, the structure and organization of the whole trophic network. It is particularly sensitive to changes in environmental conditions. It is expected to increase/decrease because of weaker/stronger stratification due to ocean warming and evaporation.

The northwestern Mediterranean Sea is one of the few sites of open-sea deep convection and dense water formation. The area is characterized by intense air-sea exchanges favored by the succession of strong northerly and northwesterly winds during autumn and winter that eventually induce deep convection episodes and the formation of the 'Western Mediterranean Deep Water'. This region exhibits a significant spring phytoplankton bloom, which appears to be largely influenced in intensity and diversity by the winter mixing.

High-resolution, realistic, three-dimensional atmospheric and oceanic models are essential for assessing the intricacy of buoyancy fluxes, horizontal advection, and convective processes, but they need to be validated by data that describe the variability of convection at small spatial scales (a few kilometers) and at high frequency (typically a day), and by accurate concomitant observations throughout the water column and atmosphere, especially during strong wind events. Likewise, the lack of *in situ* observations simultaneously inferring the main physical and biogeochemical variables was a major obstacle to unify a consensual description of the variability of the phytoplankton spring bloom and finally to equilibrate matter fluxes and budgets over an annual cycle.

To understand and resolve the interplay between the atmosphere and ocean, and the impact of ocean circulation and mixing on biogeochemistry, we carried out an unprecedented observational effort during a large experiment between July 2012 and July 2013. A multiplatform approach - combining aircraft, balloons, ships, moorings, floats, and gliders - aimed both at characterizing the

dynamics of the atmosphere and at quantifying the physical, biogeochemical and biological properties of the water masses. Beyond a better understanding of the wind dynamics, the interannual variability of the deep convection, and the seasonal variability of the nutrient distribution and plankton structure in the pelagic ecosystem, the experiment provided a reference data set that was used as a benchmark for advancing the modeling of the surface fluxes, convective processes, dense water formation rates, and physical-biogeochemical coupling processes. It also represented an opportunity for complementary investigations such as evaluating model parameterizations or studying the role of sub-mesoscale eddies in dense water spreading and biogeochemistry.

Key Points

- Dynamics of dense water formation and effects of (sub)mesoscale dynamics and intrinsic ocean variability
- Estimates of dense water formation rate, characteristics, and fate
- Effect of deep water formation on biogeochemical budgets, planktonic biodiversity and trophic levels functioning
- Potential evolution of mechanisms and their impacts on biogeochemistry in the context of global change

Program acronyms

DeWEX:	Deep Water formation Experiment
HyMEX:	Hydrological cycle in the Mediterranean EXperiment
MERMEX:	Marine Ecosystems Response in the Mediterranean Experiment
MISTRALS:	Mediterranean Integrated Studies at Regional And Local Scales
MOOSE:	Mediterranean Ocean Observing System on Environment
NAOS:	Novel Argo Ocean observing System
PERSEUS:	Protection EuRopean Seas and borders through the intelligent Use of surveillance

1. Rationales

The Mediterranean Sea is recognized to be particularly sensitive to climate change [*Adloff et al.*, 2015; *Cacho et al.*, 2002; *Giorgi*, 2006; *Somot et al.*, 2006]. The basin is affected by an increasing anthropogenic pressure, linked to strong economical and touristic activities [*i.e. Attané and Courbage*, 2001]. Its semi-enclosed nature, together with its smaller inertia [*Paluszkiewicz*, 1994] compared to large oceans [*Lacombe and Richez*, 1982], makes it more sensitive to natural variations in the atmosphere-ocean and land-ocean interactions and exchanges, which can, in turn, enhance the effects of climatic perturbations. The Mediterranean Sea is one of the most intricate marine environments on Earth as most of the critical physical processes characterizing the global circulation (dense water formation, thermohaline circulation, sub-basins gyres…) can be observed in a narrow latitudinal belt at mid-latitudes. They act on a large spectrum of spatial and temporal scales [*MERMEX group*, 2011], and induce a pronounced biological heterogeneity [*Conan et al.*, 1999; *Crise et al.*, 1999; *Robinson et al.*, 2001]. It combines a clear West - East gradient in nutrients distribution and oligotrophy [*D'Ortenzio and Ribera d'Alcalà*, 2009; *Pujo-Pay et al.*, 2011; *Ribera d'Alcalà et al.*, 2003] with a high spatio-temporal variability in the dynamics of nutrients and organic matter [*Pujo-Pay and Conan*, 2003; *Pujo-Pay et al.*, 2011; *Santinelli et al.*, 2010] which is propagated through diverse bottom-up and top-down controls within primary, secondary and upper trophic levels.

The Mediterranean Sea can thus be regarded as a reduced model of the global ocean and, in this context, the North Western Mediterranean (NWM hereafter) is consensually assumed as a critical region for the functioning of the whole basin. In this relatively small portion of the Mediterranean Sea, deep convection is regularly observed in winter. If the three phases of convection (preconditioning, vertical mixing and spreading/ restratification) have been defined for several decades [*MEDOC group*, 1970], it appears that these phases overlap and that the physical processes that govern them interact in a wide range of spatial and temporal scales. In particular, small-scale and sub-mesoscale dynamics play a particularly important role, for example convective plumes whose size is less than one kilometer and which are important mixing agents, or sub-mesoscale coherent vortices of radius (~5km) which are involved in the large scale circulation of the newly formed deep water [*Testor and Gascard*, 2003].

Convection induces a redistribution of organic and inorganic matters all over the water column. Moreover, in the same area (although with a larger extension and with a greater interannual variability than the region of deep convection) a large and intense spring bloom is observed [*D'Ortenzio and Ribera d'Alcalà*, 2009; *Mayot et al.*, 2016], presenting a high variability from mesoscale to small scale, often clearly coupled to the physical one [*Diaz et al.*, 2000]. The bloom represents the most important process of the basin in terms of primary and secondary productions

at the origin of large carbon exports to the deep layers [*MERMEX group*, 2011]. In such areas, the intermittent nutrient enrichment promotes a switching between a small-sized microbial community and diatom-dominated populations [*Conan et al.*, 1999; *Siokou-Frangou et al.*, 2010]. In the Ligurian Sea, for example, *Marty and Chiavérini [2010]* found a strong relationship between the depth of the wintertime convection and nutrient enrichment of the surface layer, triggering short and intense diatom production and deep vertical flux before the onset of the stratification and the development of the regular spring bloom [*Stemmann et al.*, 2002]. Higher abundance of zooplankton in the '80s and late '90s was also correlated with dry and cold winters [*Vandromme et al.*, 2011]. Besides, the modeling results of *Auger et al.* [2014] indicated that the total annual phytoplankton biomass is favored by convection because of the reduced zooplankton grazing pressure in winter and early spring while the annual primary production is not affected due to a compensation of its reduction in winter by its increase in spring.

All the above demonstrates that the NWM area concentrates several physical biological and chemical processes that are absolutely critical for the Mediterranean functioning, but that are also dramatically relevant to understand the complex interplay between physical forcing and biogeochemical responses at a synoptic view. However, and despite of several national and international initiatives, the strong imbrications of spatial (from meso- to large-) and temporal (from episodic- to interannual-) scales of the main processes involved limited our comprehension of the functioning of the area and of its role in a Pan-Mediterranean context.

In a review paper devoted to the Mediterranean Sea [*MERMEX group*, 2011], the inadequacy of the observational approaches, which were in the past for the most based on punctual cruises or a small number of fixed observing systems (and as such cannot be extrapolated to a wider spatial context or only under very specific hypotheses) was identified as the critical limitation for our understanding of the region. The Deep Water Formation (DWF hereafter) process generally occurs during a period of less than one month, with a low interannual variability in the timing of the deep convection phase [*Houpert et al.*, 2016]. On the other hand, the DWF is pre-conditioned during at least the whole previous fall [*Madec et al.*, 1996] and possibly previous winter periods. Moreover, the impact of the Levantine Intermediate Water (LIW) on the DWF was acknowledged [*Grignon et al.*, 2010]. The previous observational approaches were inadequate to completely describe the succession of the DWF phases. For the biological and chemical observations, this inadequacy was even more flagrant, with a consequence that, despite of some fixed stations or satellite monitoring, the seasonality of the biogeochemical dynamic of the area was still missing.

A large experiment coupling the exploration of the ocean from the physical and biological point of views to observations of the air-sea interface and atmospheric boundary layer was then

designed to couple the recent technological breakthroughs for small and intelligent autonomous platforms with satellite remote sensing (altimetry, SST, and color) information, with more 'classical' data acquisition by ship and aircraft sampling, and with existing fixed point moorings. The number of oceanic observations was dramatically high compared to previous experiments, and, more importantly, observations were obtained all along a seasonal cycle. These data were directly used in combination with models, before, during and after the observational phase. The strategy was focused on:

- the understanding, in a wide range of spatial and temporal scales of the physical mechanisms and forcing that precondition the convection zone, form dense water and disperse it
- the role of deep winter convection on the biogeochemical properties of the water masses, on the resulting organization of the pelagic ecosystems in spring and on the organic matter export

The modeling activity was central for driving the sampling strategy before the observation phase, for integrating the observations in a complete 4D frame during the observation phase, while favoring the understanding of the integrated system and improving our long-term simulations and scenarii. All these efforts allowed us to obtain a unique and comprehensive data set to validate and improve the physical/ biogeochemical coupled models.

The objective of this special issue is to synthesize the major results of these experiments and related operations. These results bring significant advances in our understanding of the dynamics of DWF and the related functioning of marine ecosystems that have been organized according to six major axes:

- Dynamics of the atmosphere and air-sea fluxes
- Interannual variability of dense water formation
- Deep water formation during winter 2013
- Effects of small scales and intrinsic ocean variability on dense water formation
- Effect of deep water formation on biogeochemical budgets and primary production
- Response of zooplankton community to dense water formation

2. Strategic plan

The experimental strategy was based on pooling the DeWEX [*Testor et al.*, 2018] and HyMeX SOP2 [*Estournel et al.*, 2016b] projects respectively supported by the MerMEX and HyMEX programs (both components of MISTRALS). Several National and European projects (i.e. RemOCEAN ERC, NAOS EquipEx, FP7-GROOM, FP7-JERICO, FP7-PERSEUS), combined with the long lasting monitoring activity

of MOOSE and with some additional surveys (*i.e.* MerMEX WP2-SPECIMED), provided a solid and favorable context contributing to the success of this experiment.

The observational strategy was mainly driven by the need to obtain pertinent observations all along a seasonal cycle. Six cruises were then programmed at key moments of the year 2012-2013 (see figure 1). This intensive ship-based observation effort, carried out from July 2012 to September 2013, was completed by a large variety of tools, for the most robotic or semi-robotic, to obtain a quasi-synoptic view of the area, to interpolate between the cruises and to provide enough observations to initialize models (Fig. 1).

The observation strategies involved then several airborne (aircraft and atmospheric boundary layer balloons), surface (boat, fixed and drifting buoys), and oceanic (physical and bio-optical Argo floats and gliders, mooring lines equipped with CTD sensors, current-meters and sediment traps) platforms (Fig. 2 and 3). Space observations of the sea surface altimetry and ocean color were used to monitor the interannual variability and spatial extent of the convection. The ATR-42 aircraft operated by the "Service des Avions Français Instrumentés pour la Recherche en Environnement (SAFIRE)", investigated the mean and turbulent characteristics of the marine atmospheric boundary layer above the oceanic convection region. Boundary layer pressurized balloons, developed by the "Centre National d'Etudes Spatiales", were used to document the evolution of the thermo-dynamical characteristics along the trajectory of air parcels during northern wind (Mistral) events.

Measurements from research vessels (R/V L'Atalante, R/V Le Suroît, and R/V Tethys II) were intended to be operated over periods of 3 weeks in order to cover most of the NWM during different key phases of the seasonal cycle of the region. The six cruises were carried out in July-August 2012 (summer stratified period), September 2012 (summer-to-fall transition stratified period), February 2013 (pre-bloom, winter mixing period), April 2013 (spring bloom, restratification and spreading period), July 2013 (post bloom and restratified period) and September 2013. The February DeWEX 2013 cruise, which took place during the period of intense mixing, overlapped with the airborne observations. Only during naval operations, a very large set of chemical and biological data were collected, in particular to provide stocks, fluxes, diversities and biological activities. During all the cruises, a CTD carousel composed of 12 Niskin bottles was deployed and discrete samples were collected. 440 surface-to-bottom profiles of CTD and of biogeochemical parameters (chlorophyll-a, dissolved oxygen, turbidity, currents, aggregates and zooplankton images), 9240 water samples for nutrients, dissolved and particulate organic matter, pigments, microbial diversity were collected. During February and April 2013 cruises, primary and bacterial productions were measured, and additional vertical net hauls were performed to collect zooplankton.

Surface and ship-based observations were also collected over the whole NWM. Numerous profiles in the upper layer between the surface and 1000 or 2000 m deep were collected with Argo and BGC-Argo profiling floats and gliders. All these platforms were equipped with CTD sensors, and some platforms also carried biogeochemical sensors (i.e. chlorophyll-a, dissolved oxygen, nitrate, optical backscattering). After interoperability processes, the moorings, profiling floats, and gliders provided an opportunity to characterize the temporal variability of the oceanic conditions throughout the year, and small-scale features, such as sub-mesoscale eddies and convective plumes. Surface drifters (SVP and Marisonde buoys) and fixed meteorological buoys ("Lion" and "Azur Météo-France" buoys) were used to collect air-sea interface parameters. Long mooring lines in the convection region ("Lion") and in the Ligurian Sea ("Dyfamed") provided additional observations of the evolution of the mixed layer depth, water masses, currents, and particle fluxes.

3. Major outcomes

3.1. Dynamics of the atmosphere and air-sea fluxes

The winter-integrated buoyancy loss over the Gulf of Lion was identified as the primary driving factor of the DWF interannual variability [*Somot et al.*, 2016]. At a daily scale, the Atlantic Ridge weather regime was identified as favorable to the Mistral wind and associated strong buoyancy losses and therefore to DWF, whereas the positive phase of the North Atlantic Oscillation was unfavorable. The mesoscale dynamics of the Mistral wind over the Gulf of Lion was detailed in *Drobinski et al. [2017]*. They related boundary layer balloons observations in the marine atmospheric boundary layer with the AROME-WMED weather forecast model and analyzed all the terms of the Lagrangian formulation of the momentum conservation equation to identify forces acting in the injection, ejection and deceleration regions of the Mistral flow.

Several articles have addressed the determination of air/sea fluxes at different spatial and temporal scales. Airborne measurements enabled to characterize the mean and turbulent structure of the marine atmospheric boundary layer [*Brilouet et al.*, 2017]. They showed that a one-dimensional description of the vertical exchanges remains problematic because of the presence in the marine atmospheric boundary layer of organized structures such as two-dimensional rolls favored by strong winds and surface heat fluxes. *Caniaux et al. [2017]* using measurements at the air sea interface and in the water column implemented an inverse method to estimate heat and water fluxes during one year for the NWM basin at a fine scale resolution. They compared air-sea fluxes adjusted using observations, with some operational numerical weather prediction models (such as ARPEGE, NCEP, ERA-INTERIM, ECMWF, and AROME) and concluded that these models were unable to retrieve the mean annual patterns and values of fluxes. *Song and Yu [2017]* performed an analysis

of the whole Mediterranean Sea surface energy budget using nine surface heat flux climatologies obtained from atmospheric re-analyses, satellite or ship observations. They compared them with the heat flux associated to the net transport through the Strait of Gibraltar and to data-assimilated global ocean state estimation.

Uncertainties on heat and water fluxes, in particular related to uncertainties in the parameterization of turbulent fluxes under strong wind conditions, remain an important problem for ocean modeling in the Mediterranean, as shown by the sensitivity study by *Seyfried et al.* [2017] on the modeling of the convection during winter 2012-2013. Unfortunately, no direct observation of the turbulent fluxes was available during the strong wind events of DeWEX/ HyMeX SOP2. However, it can be noted that the uncertainties of turbulent fluxes parameterization are not specific to the Mediterranean.

3.2. Interannual variability of dense water formation

The currently well-accepted hypothesis based on the climate projections, predicts a strong decrease of DWF rate, as surface layers will become too warm to produce dense waters. On the other hand, increasing evaporation should increase the surface water salinity, which could, in turn, raise density and result in convection. So, at least cyclically the Mediterranean should continue to produce dense water through different mechanisms; however, under this scenario, deep water characteristics should be deeply modified, affecting nutrient budgets and the whole ecosystem functioning.

The interannual variability of convection in the Gulf of Lion over the last few decades has not been monitored consistently and uniformly; CTD profiles and short-term mooring lines have been used to document the period from 1970 to 2000 [see *Béthoux et al.*, 2002; *Mertens and Schott*, 1998]. It is only since 2007, with the implementation and simultaneous use, in the framework of MOOSE in particular, of fully instrumented mooring lines and buoy at the Lion site, satellite data, gliders, profiling floats and annual cruises throughout the NWM that a reference dataset has been established to characterize the DWF interannual variability in terms of mixed layer depth, convective surface, DWF rate or water mass characteristics, which allowed to evaluate accurately long-term model simulations.

High frequency temperature, salinity and current measurements, collected at the Lion site [*Durrieu de Madron et al.*, 2017; *Houpert et al.*, 2016], revealed that from 2007 to 2015 bottom reaching convection and production of new WMDW, denser than 29.11 $kg.m^{-3}$, occurred every winter between 2009 and 2013. The overlapping of the mixing and restratification phases was

regularly observed, with a secondary vertical mixing event of 2-4 days long occurring after the beginning of the restratification phase (generally in April). This pattern is retrieved in the simulations produced by [*Léger et al.*, 2016] or *in situ data* detailed by [*Severin et al.*, 2014; *Severin et al.*, 2016].

As expected, the succession of bottom-reaching convection episodes between 2009 and 2013 altered the deep-water masses thermohaline characteristics, which already underwent a significant change after exceptional winter 2005 convection event *[see Herrmann et al., 2010 and references therein]*. The stepwise rise in temperature, salinity and density, observed consecutively to each bottom-reaching convection event, led between 2009 to 2013 to an increasing trend in salinity (+3.3 ± 0.2 10^{-3} a^{-1}) and potential temperature (+3.2 ± 0.5 10^{-3} °C.a^{-1}) for the deep layer [*Houpert et al.*, 2016]. This confirmed the increase in rate observed from literature (+9.2 10^{-4} a^{-1} in salinity and +2 10^{-3} °C a^{-1} in temperature for 1945-2000 period [*Vargas-Yáñez et al.*, 2010]. These increasing trends in the deep layers would result from a heat and salt accumulation during the 1990s in the surface and intermediate layers of the Gulf of Lion, transferred stepwise towards the deep layers when intense convective events occur [like in 1999, 2005 and later; *Somot et al.*, 2016].

The evolution of the deep water thermohaline characteristics due to winter convection depends directly on the volume of newly formed dense water. However, to get such estimates remains a complex issue since there is currently no objective criterion for assessing them. Two different methods were primarily used to estimate the volume of new WMDW, the advantages and disadvantages of each are explained by *Testor et al. [2018]*. First, as the vertical dilution effect of convection reduces the chlorophyll concentration [*Severin et al.*, 2014], the volume of new WMDW can be approximated as the extent of the chlorophyll depleted mixing zone seen by ocean color satellite times the mixed layer depth. Second, the volume of new WMDW can be estimated by quantifying the amount of new water exceeding a given density threshold (e.g. > 29.11 $kg.m^{-3}$, the maximum density of the "old" deep water) over the NWM basin. *Houpert et al. [2016]* estimated that between 2009 and 2012, the new WMDW formation rate varied between 0.91 and 1.25 Sv (when averaged over one year). As for them, *Herrmann et al.* [2017] used altimetry and ocean color satellites data to obtain a wider annual volume range of 0.0 to 2.67 Sv for the 1998-2016 period.

3.3. Deep water formation during winter 2013

The winter buoyancy fluxes in 2012-2013 were sufficient to trigger deep convection and significant production of new WMDW during DeWEX operations. With an observing system simulation experiment approach, *Waldman et al. [2016]* confirmed the ability of the observation network (ship track in Fig.2) to correctly quantify the spatial extent of convection and the seasonal

evolution of dense water volume. Several estimations of DWF rate deduced from observations of the mixed patch volume [*Herrmann et al.*, 2017], dissolved oxygen ventilation of the deep waters [*Coppola et al.*, 2017], and the newly formed dense water volume [*Herrmann et al.*, 2017; *Testor et al.*, 2018] were coherent and ranged between 1.27 and 2.0 Sv. This last method, which appears to be the most robust, was used to quantify the rate of dense water formation for the different models used.

All the observations provided important results for model evaluation and sensitivity studies. Numerical models were generally able to simulate realistic DWF (location, triggering and chronology, characteristics) during winter 2013 [*Estournel et al.*, 2016a; *Lebeaupin Brossier et al.*, 2017; *Léger et al.*, 2016; *Waldman et al.*, 2017a; *Waldman et al.*, 2017b], and calculated DWF rates between 1.33 and 2.59 Sv.

Sensitivity analysis revealed that models are mostly affected by fine-scale ocean structures, such as shelf DWF and export, eddies and fronts at the rim of the convective patch well evidenced by glider observations [*Testor et al.*, 2018]. The resolution of the mesoscale definitively improved the realism of the simulations and their agreement with observations, by increasing the eddy kinetic energy and enhancing the stratifying effect of advection through the mixed patch [*Estournel et al.*, 2016a; *Lebeaupin Brossier et al.*, 2017; *Waldman et al.*, 2017b]. Models also showed a large sensitivity to the initial and boundary conditions (introducing bias on the stratification), the wind intensity, and air-sea fluxes [*Estournel et al.*, 2016a; *Léger et al.*, 2016], which affect the water masses characteristics and the volume of dense water formed during convection.

3.4. Effects of small scales and intrinsic ocean variability on dense water formation

The distinction between the internal (intrinsic) and external (atmospherically-forced) components of the ocean variability is central to understand the forcings modulating DWF intensity and water mass transformations. The large presence of meso- and sub-meso scale eddies, and current instabilities in the NWM basin are sources of ocean intrinsic variability that can retroact on the basin-scale circulation through an inverse energy cascade. The multiple platforms used during the experience allowed identifying new dynamical processes at small scales or better characterizing already known processes (frontal instabilities, convective plumes, vortices). Indeed, *Giordani et al. [2017]* described a symmetric instability at the western edge of the deep convection area when current and dominant northerly wind are in the same direction. This instability results from a cross-front ageostrophic circulation, which subducts surface low-potential vorticity waters on the dense

side of the front and obducts high-potential vorticity waters from the pycnocline on the light side of the front.

In the convection area, *Margirier et al. [2017]* were able from glider navigation data to describe the statistical physical and biogeochemical characteristics of the convective plumes. It appears that these plumes, which had radius of about 350 m and significant vertical velocities (up to 18 $cm.s^{-1}$), covered about one-third of the deep convection area. Vertical velocities are scaled by atmospheric fluxes inducing downward buoyancy fluxes with a vertical diffusion coefficient of 7 $m^2.s^{-1}$. *Bosse et al. [2016]* provided an inventory of the different types of cyclonic and anticyclonic Submesoscale Coherent Vortices (SCVs) observed in the NWM by gliders, cruises, and mooring data during a four-year period (2009-2013). They showed that these energetic features prevailed over the large-scale geostrophic circulation and remained coherent for long periods of time (typically one year). Bottom-reaching convection might favor the formation of cyclonic vortices. Using a realistic high-resolution (1 km) numerical model, *Damien et al. [2017]* simulated the formation and spreading of SCVs during intermediate and deep convection events. They reproduced similar long-lived cyclonic and anticyclonic coherent structures. Both studies showed the prominent role of SCVs in the spreading of the convected waters, their contribution to the ventilation of the deep basin, and their influence on the convection onset the following winter.

Estournel et al. [2016a] discussed the interaction in autumn of the surface and Ekman buoyancy fluxes associated with displacements of the front bounding the convection zone to the south. They show that these processes are important for the convection preconditioning processes. *Waldman et al. [2017a]* showed that intrinsic variability has a strong impact on the timing and geographical extent of a convective event, but has little impact on the rate of convection in the constrained 2012-2013 case. *Waldman et al. [2018]* further showed that intrinsic variability also explains a significant fraction of the deep convection interannual variability, although it has only modest impacts on the long-term mean state.

3.5. Effect of dense water formation on biogeochemical budgets and primary production

The DWF in the NWM impacts the nutrients distribution, matter export and development of the trophic network and it strongly contributes to the modification of the deep stocks of these components at the basin level. Three different areas of the western basin were identified by contrasting vertical mixing regimes: deep convection, shallow convection and stratified area [*Kessouri et al.*, 2018]. Using a 3-D high-resolution coupled hydrodynamic-biogeochemical model these authors showed in all three regions, that the mixed layer deepening during the destratification

process induced an upward nutrients flux, triggering in turn, an autumnal phytoplankton bloom. In contrast at the end of winter, the end of turbulent mixing favored the onset of an intense spring bloom only in the deep convection region. The authors concluded that despite these seasonal variations, annual primary production in all three regions is quite similar, but total organic carbon exported to deep waters was 3 and 8 times higher for moderate and deep convective areas respectively, compared to stratified ones.

The depth reached by the convection directly drives the nutrient stoichiometry of water masses in winter [*Severin et al.*, 2014; *Severin et al.*, 2017], which influences the phytoplankton community structure in spring [*Mayot et al.*, 2017b; *Severin et al.*, 2016], which in return controls the nutrient availability for the rest of the year by direct uptake [*Kessouri et al.*, 2017]. The general emerging pattern is that the spatial (horizontal) extension of the convective area in winter influences the intensity of the resulting bloom in spring (because of variable amount of nutrients injected into the surface layer), but the depth (vertical) reached by the convection (and in particular when deep convection reaches the nepheloïd boundary layer) plays a key role in the bloom diversity (notably due to different nutritional stoichiometry and higher $Si:NO_3$ ratio when convection is deeper) [*Leblanc et al.*, 2018; *Mayot et al.*, 2017b]. These variations could be partly the result of pore water release loaded with nutrients because of sediment resuspension enhanced by the bottom-reaching mixing *[Durrieu de Madron et al., 2017]*.

Moreover, the deep convection events result in a homogenization of hydrological characteristics, but also of the prokaryotic communities over the entire convective cell, resulting in a surprising predominance of typical surface bacteria (such as *Oceanospirillale* and *Flavobacteriale*) [*Severin et al.*, 2016]. However, physical turbulences only were not sufficient to explain this new distribution, but act in synergy with quantity and quality of exported organic matter. Indeed, the authors explained this dominance by the rapid export of fresh and labile organic matter to the deep layers. Finally, the rapid return of specific prokaryotic communities and classic activities in the intermediate layer less than 5 days after the intense mixing (and before hydrological restratification) indicates a marked resilience of the communities, apart from the residual deep mixed water patch [*Severin et al.*, 2017; *Severin et al.*, 2016]. Finally, the heterogeneous surface distribution of nutrients during the winter deep convection event in the NWM was shown to impact the phytoplankton distribution and community structure several weeks after the end of convective events [*Mayot et al.*, 2017b; *Severin et al.*, 2017; *Severin et al.*, 2016].

The multiple platforms of physical, bio-optical and biochemical measurements provided a detailed characterization of the spatial and temporal structuring of the main biogeochemical variables at different scales, from small-scale vortex structures to the basin as a whole. The role of

447 the post convective SCVs on nutrients distribution and phytoplankton communities, as well as on the
448 subsequent enhanced primary production and higher carbon sequestration was revealed by *Bosse et*
449 *al. [2017]*. They showed that SCVs present important dynamical barriers that drastically reduce
450 lateral exchanges between their cores and the surroundings thus enabling them to keep their core
451 characteristics the same for a long time. This suggests that they locally have a great imprint on both
452 physical and biogeochemical cycles. At larger scale, two distinct trophic regimes as “High Bloom”
453 centered on the convection area, and a ‘‘Bloom’’ bioregion located at its periphery were identified
454 [*Mayot et al.*, 2017a; *Mayot et al.*, 2017b], in agreement with regions mentioned above and defined
455 by *Kessouri et al. [2018]*.

456 The interannual variability of the physical and biogeochemical coupling of the NWM was also
457 assessed using satellite ocean color observations [*Mayot et al.*, 2017b] and a 3-D hydrodynamic-
458 biogeochemical coupled model [*Ulses et al.*, 2016]. Both studies showed that phytoplankton biomass
459 during the spring bloom is larger in years associated with intense deep convection events. Modeling
460 gave further evidence that amounts of nutrients annually injected into the surface layer is clearly
461 linked to the intensity of the convection but does not significantly modify the overall primary
462 production budget. Indeed, the primary production is inhibited during severe winters and transiently
463 explodes during the restratification phase, while it increases continuously during both phases when
464 winters are milder. As seen above, the occurrence of a highly productive bloom is also related to an
465 increase in the phytoplankton bloom size area. For the period including winter 2013, the processes
466 responsible for the stoichiometry variability at the seasonal scale were disentangled by using a
467 coupled physical-biogeochemical model able to correctly reproduce the general observed seasonal
468 dynamics of the physical and biogeochemical events [*Kessouri et al.*, 2017]. The dynamics of the
469 nutrients, phyto- and zoo-plankton biomass as well as their interactions during the convection period
470 and the spring bloom were well reproduced. The authors estimated nutrient budgets for the entire
471 year and showed that the convection region represented a sink of inorganic and a source of organic
472 nitrogen and phosphorus. This process can efficiently export organic carbon below the photic layer
473 [*Kessouri et al.*, 2018; *Kessouri et al.*, 2017; *Severin et al.*, 2017].

474

475 **3.6. Zooplankton community response to dense water formation**

476 The deep convection zone is likely an area of both enhanced energy transfer to higher trophic
477 levels and organic matter export [*Kessouri et al.*, 2018; *Severin et al.*, 2017]. The two trophic regimes
478 observed on the primary producer level are also reflected on higher trophic levels. Indeed in winter,
479 low zooplankton abundance and biomass were observed in the deep convection zone, but higher
480 values on its periphery. In spring, this pattern reversed with high biomass dominated by herbivorous

481 species in the deep convection zone, and lower values on the periphery [*Donoso et al.*, 2017]. The
482 potential grazing impact of phytoplankton by zooplankton was estimated to increase by one order of
483 magnitude from winter to spring. In April, all areas except the deep convection zone incurred top-
484 down control by zooplankton [*Siokou-Frangou et al.*, 2010].

485 Despite this significant seasonal variability of zooplankton biomass, the community composition
486 was comparable for both winter mixing and spring bloom periods, typified by high copepod
487 dominance [*Donoso et al.*, 2017]. Using stable isotope mixing models, *Hunt et al.* [2017] estimated
488 that micro particulate organic matter never contributed more than 20% to zooplankton biomass,
489 even in regions where microphytoplankton was plentiful, indicating that a large part of its biomass
490 may have remained ungrazed.

491

492

493 **Acknowledgements**

494 Many people participated in this observational effort: scientists, technicians, crews.... All deserve our
495 grateful thanks. The main efforts were supported by CNRS, French universities (Sorbonne, Toulouse,
496 Marseille, Perpignan…), Météo-France, IFREMER, and CNES through the international
497 metaprogramme MISTRALS dedicated to the understanding of the Mediterranean basin
498 environmental processes (http://www.mistrals-home.org). More specifically by MERMEX-DEWEX,
499 MOOSE and HYMEX French programs.

500

501

502 **References**

503 Adloff, F., S. Somot, F. Sevault, G. Jordà, R. Aznar, M. Déqué, M. Herrmann, M. Marcos, C. Dubois, E.
504 Padorno, E. Alvarez-Fanjul, and D. Gomis, Mediterranean Sea response to climate change in
505 an ensemble of twenty first century scenarios, *Climate Dynamics*, *45* (9), 2775-2802, 2015.
506 Attané, I., and Y. Courbage, La démographie en Méditerranée. Situation et projections, in *Les*
507 *fascicules du Plan Bleu*, pp. 249, 2001.
508 Auger, P.A., C. Ulses, C. Estournel, L. Stemmann, S. Somot, and F. Diaz, Interannual control of
509 plankton communities by deep winter mixing and prey/predator interactions in the NW
510 Mediterranean: Results from a 30-year 3D modeling study, *Progress in Oceanography*, *124*,
511 12-27, 2014.
512 Béthoux, J.-P., X. Durrieu de Madron, F. Nyffeler, and D. Tailliez, Deep water in the western
513 Mediterranean: peculiar 1999 and 2000 characteristics, shelf formation hypothesis,
514 variability since 1970 and geochemical inferences, *Journal of Marine Systems*, *33-34* (C), 117-
515 131, 2002.
516 Bosse, A., P. Testor, L. Houpert, P. Damien, L. Prieur, D. Hayes, V. Taillandier, X. Durrieu de Madron, F.
517 d'Ortenzio, L. Coppola, J. Karstensen, and L. Mortier, Scales and dynamics of Submesoscale

Coherent Vortices formed by deep convection in the northwestern Mediterranean Sea, *Journal of Geophysical Research: Oceans*, *121* (10), 7716-7742, 2016.

Bosse, A., P. Testor, N. Mayot, L. Prieur, F. D'Ortenzio, L. Mortier, H.L. Goff, C. Gourcuff, L. Coppola, H. Lavigne, and P. Raimbault, A submesoscale coherent vortex in the Ligurian Sea: From dynamical barriers to biological implications, *Journal of Geophysical Research: Oceans*, *122* (8), 6196-6217, 2017.

Brilouet, P.-E., P. Durand, and G. Canut, The marine atmospheric boundary layer under strong wind conditions: Organized turbulence structure and flux estimates by airborne measurements, *Journal of Geophysical Research: Atmospheres*, *122* (4), 2016JD025960, 2017.

Cacho, I., J. Grimalt, and M. Canals, *Response of the Western Mediterranean Sea to rapid climatic variability during the last 50,000 years: A molecular biomarker approach*, 253-272 pp., 2002.

Caniaux, G., L. Prieur, H. Giordani, and J.L. Redelsperger, An inverse method to derive surface fluxes from the closure of oceanic heat and water budgets: Application to the north-western Mediterranean Sea, *Journal of Geophysical Research: Oceans*, *122* (4), 2884-2908, 2017.

Conan, P., C.M. Turley, E. Stutt, M. Pujo-Pay, and F. Van Wambeke, Relationship between Phytoplankton Efficiency and the Proportion of Bacterial Production to Primary Production in the Mediterranean Sea, *Aquatic Microbial Ecology*, *17* (2), 131-144, 1999.

Coppola, L., L. Prieur, I. Taupier-Letage, C. Estournel, P. Testor, D. Lefevre, S. Belamari, S. LeReste, and V. Taillandier, Observation of oxygen ventilation into deep waters through targeted deployment of multiple Argo-O2 floats in the north-western Mediterranean Sea in 2013, *Journal of Geophysical Research: Oceans*, *122* (8), 6325-6341, 2017.

Crise, A., J.I. Allen, J. Baretta, G. Crispi, R. Mosetti, and C. Solidoro, The Mediterranean pelagic ecosystem response to physical forcing, *Progress In Oceanography*, *44* (1-3), 219-243, 1999.

D'Ortenzio, F., and M. Ribera d'Alcalà, On the trophic regimes of the Mediterranean Sea: a satellite analysis, *Biogeosciences*, *6* (2), 139-148, 2009.

Damien, P., A. Bosse, P. Testor, P. Marsaleix, and C. Estournel, Modeling postconvective submesoscale coherent vortices in the northwestern Mediterranean Sea, *Journal of Geophysical Research: Oceans*, n/a-n/a, 2017.

Diaz, F., P. Raimbault, and P. Conan, Small-scale study of primary productivity during spring in a Mediterranean coastal area (Gulf of Lions), *Continental Shelf Research*, *20* (9), 975-996, 2000.

Donoso, K., F. Carlotti, M. Pagano, B.P.V. Hunt, R. Escribano, and L. Berline, Zooplankton community response to the winter 2013 deep convection process in the NWMediterranean Sea, *Journal of Geophysical Research: Oceans*, *122* (3), 2319-2338, 2017.

Drobinski, P., B. Alonzo, C. Basdevant, P. Cocquerez, A. Doerenbecher, N. Fourrié, and M. Nuret, Lagrangian dynamics of the mistral during the HyMeX SOP2, *Journal of Geophysical Research: Atmospheres*, *122* (3), 2016JD025530, 2017.

Durrieu de Madron, X., S. Ramondenc, L. Berline, L. Houpert, A. Bosse, S. Martini, L. Guidi, P. Conan, C. Curtil, N. Delsaut, S. Kunesch, J.F. Ghiglione, P. Marsaleix, M. Pujo-Pay, T. Séverin, P. Testor, C. Tamburini, and Antares collaboration, Deep sediment resuspension and thick nepheloid layer generation by open-ocean convection, *Journal of Geophysical Research: Oceans*, *122* (3), 2291-2318, 2017.

Estournel, C., P. Testor, P. Damien, F. D'Ortenzio, P. Marsaleix, P. Conan, F. Kessouri, X. Durrieu de Madron, L. Coppola, J.-M. Lellouche, S. Belamari, L. Mortier, C. Ulses, M.-N. Bouin, and L. Prieur, High resolution modeling of dense water formation in the north-western Mediterranean during winter 2012–2013: Processes and budget, *Journal of Geophysical Research: Oceans*, *121* (7), 5367-5392, 2016a.

Estournel, C., P. Testor, I. Taupier-Letage, M.-N. Bouin, L. Coppola, P. Durand, P. Conan, A. Bosse, P.-E. Brilouet, L. Beguery, S. Belamari, K. Béranger, J. Beuvier, D. Bourras, G. Canut, A. Doerenbecher, X. Durrieu de Madron, F. D'Ortenzio, P. Drobinski, V. Ducrocq, N. Fourrié, H. Giordani, L. Houpert, L. Labatut, C.L. Brossier, M. Nuret, L. Prieur, O. Roussot, L. Seyfried, and S. Somot, HyMeX-SOP2: The Field Campaign Dedicated to Dense Water Formation in the Northwestern Mediterranean, *Oceanography*, *29* (4), 196-206, 2016b.

570 Giordani, H., C. Lebeaupin-Brossier, F. Léger, and G. Caniaux, A PV-approach for dense water
571 formation along fronts: Application to the Northwestern Mediterranean, *Journal of*
572 *Geophysical Research: Oceans*, *122* (2), 995-1015, 2017.
573 Giorgi, F., Climate change hot-spots, *Geophysical Research Letters*, *33* (8), 2006.
574 Grignon, L., D.A. Smeed, H.L. Bryden, and K. Schroeder, Importance of the Variability of Hydrographic
575 Preconditioning for Deep Convection in the Gulf of Lion, NW Mediterranean, *Ocean Science*,
576 *6* (2), 573-586, 2010.
577 Herrmann, M., P.-A. Auger, C. Ulses, and C. Estournel, Long-term monitoring of ocean deep
578 convection using multisensors altimetry and ocean color satellite data, *Journal of Geophysical*
579 *Research: Oceans*, *122* (2), 1457-1475, 2017.
580 Herrmann, M.J., F. Sevault, J. Beuvier, and S. Somot, What induced the exceptional 2005 convection
581 event in the northwestern Mediterranean basin? Answers from a modeling study, *Journal of*
582 *Geophysical Research: Oceans*, *115* (C12), 2010.
583 Houpert, L., X. Durrieu de Madron, P. Testor, A. Bosse, F. D'Ortenzio, M.N. Bouin, D. Dausse, H. Le
584 Goff, S. Kunesch, M. Labaste, L. Coppola, L. Mortier, and P. Raimbault, Observations of open-
585 ocean deep convection in the northwestern Mediterranean Sea: Seasonal and interannual
586 variability of mixing and deep water masses for the 2007-2013 Period, *Journal of Geophysical*
587 *Research: Oceans*, *121* (11), 8139-8171, 2016.
588 Hunt, B.P.V., F. Carlotti, K. Donoso, M. Pagano, F. D'Ortenzio, V. Taillandier, and P. Conan, Trophic
589 pathways of phytoplankton size classes through the zooplankton food web over the spring
590 transition period in the north-west Mediterranean Sea, *Journal of Geophysical Research:*
591 *Oceans*, *122* (8), 6309-6324, 2017.
592 Kessouri, F., C. Ulses, C. Estournel, P. Marsaleix, F. D'Ortenzio, T. Severin, V. Taillandier, and P. Conan,
593 Vertical Mixing Effects on Phytoplankton Dynamics and Organic Carbon Export in the
594 Western Mediterranean Sea, *Journal of Geophysical Research: Oceans*, *0* (0), 2018.
595 Kessouri, F., C. Ulses, C. Estournel, P. Marsaleix, T. Severin, M. Pujo-Pay, J. Caparros, P. Raimbault,
596 O.P.d. Fommervault, F. D'Ortenzio, V. Taillandier, P. Testor, and P. Conan, Nitrogen and
597 phosphorus budgets in the Northwestern Mediterranean deep convection region, *Journal of*
598 *Geophysical Research: Oceans*, *122* (12), 9429-9454, 2017.
599 Lacombe, H., and C. Richez, The regime of the Strait of Gibraltar, in *Hydrodynamics of semi-enclosed*
600 *seas*, edited by J.C.J. Nihoul, pp. 13-73, Elsevier, New York, 1982.
601 Lebeaupin Brossier, C., F. Léger, H. Giordani, J. Beuvier, M.-N. Bouin, V. Ducrocq, and N. Fourrié,
602 Dense water formation in the north-western Mediterranean area during HyMeX-SOP2 in
603 1/36° ocean simulations: Ocean-atmosphere coupling impact, *Journal of Geophysical*
604 *Research: Oceans*, *122* (7), 5749-5773, 2017.
605 Leblanc, K., B. Quéguiner, F. Diaz, V. Cornet, M. Michel-Rodriguez, X. Durrieu de Madron, C. Bowler,
606 S. Malviya, M. Thyssen, G. Grégori, M. Rembauville, O. Grosso, J. Poulain, C. de Vargas, M.
607 Pujo-Pay, and P. Conan, Nanoplanktonic diatoms are globally overlooked but play a role in
608 spring blooms and carbon export, *Nature Communications*, *9* (1), 953, 2018.
609 Léger, F., C. Lebeaupin Brossier, H. Giordani, T. Arsouze, J. Beuvier, M.-N. Bouin, É. Bresson, V.
610 Ducrocq, N. Fourrié, and M. Nuret, Dense water formation in the north-western
611 Mediterranean area during HyMeX-SOP2 in 1/36° ocean simulations: Sensitivity to initial
612 conditions, *Journal of Geophysical Research: Oceans*, *121* (8), 5549-5569, 2016.
613 Madec, G., P. Delecluse, M. Crépon, and F. Lott, Large-Scale Preconditioning of Deep-Water
614 Formation in the Northwestern Mediterranean Sea, *Journal of Physical Oceanography*, *26* (8),
615 1393-1408, 1996.
616 Margirier, F., A. Bosse, P. Testor, B. L'Hévéder, L. Mortier, and D. Smeed, Characterization of
617 Convective Plumes Associated With Oceanic Deep Convection in the Northwestern
618 Mediterranean From High-Resolution In Situ Data Collected by Gliders, *Journal of*
619 *Geophysical Research: Oceans*, *122* (12), 9814-9826, 2017.

Marty, J.C., and J. Chiavérini, Hydrological changes in the Ligurian Sea (NW Mediterranean, DYFAMED site) during 1995–2007 and biogeochemical consequences, *Biogeosciences*, *7* (7), 2117-2128, 2010.

Mayot, N., F. D'Ortenzio, M. Ribera d'Alcalà, H. Lavigne, and H. Claustre, Interannual variability of the Mediterranean trophic regimes from ocean color satellites, *Biogeosciences*, *13* (6), 1901-1917, 2016.

Mayot, N., F. D'Ortenzio, V. Taillandier, L. Prieur, O.P.d. Fommervault, H. Claustre, A. Bosse, P. Testor, and P. Conan, Physical and Biogeochemical Controls of the Phytoplankton Blooms in North Western Mediterranean Sea: A Multiplatform Approach Over a Complete Annual Cycle (2012–2013 DEWEX Experiment), *Journal of Geophysical Research: Oceans*, *122* (12), 9999-10019, 2017a.

Mayot, N., F. D'Ortenzio, J. Uitz, B. Gentili, J. Ras, V. Vellucci, M. Golbol, D. Antoine, and H. Claustre, Influence of the Phytoplankton Community Structure on the Spring and Annual Primary Production in the Northwestern Mediterranean Sea, *Journal of Geophysical Research: Oceans*, *122* (12), 9918-9936, 2017b.

MEDOC group, Observation of formation of deep water in the Mediterranean Sea, 1969, *Nature*, *227*, 1037-1040, 1970.

MERMEX group, Marine ecosystems' responses to climatic and anthropogenic forcings in the Mediterranean, *Progress In Oceanography*, *91* (2), 97-166, 2011.

Mertens, C., and F. Schott, Interannual Variability of Deep-Water Formation in the Northwestern Mediterranean, *Journal of Physical Oceanography*, *28* (7), 1410-1424, 1998.

Paluszkiewicz, T., Deep Convective Plumes in the Ocean, *Oceanography*, *7*, 1994.

Pujo-Pay, M., and P. Conan, Seasonal variability and export of Dissolved Organic Nitrogen in the North Western Mediterranean Sea, *Journal of Geophysical Research*, *108* (C6), 1901-1911, 2003.

Pujo-Pay, M., P. Conan, L. Oriol, V. Cornet-Barthaux, C. Falco, J.F. Ghiglione, C. Goyet, T. Moutin, and L. Prieur, Integrated survey of elemental stoichiometry (C, N, P) from the Western to Eastern Mediterranean Sea, *Biogeosciences*, *8* (4), 883-899, 2011.

Ribera d'Alcalà, M., G. Civitarese, F. Conversano, and R. Lavezza, Nutrient ratios and fluxes hint at overlooked processes in the Mediterranean Sea, *Journal of Geophysical Research*, *108* (C9), 8106, 2003.

Robinson, A.R., G. Wayne, A. Theocharis, and A. Lascaratos, Mediterranean Sea circulation, in *Encyclopedia of Ocean Sciences*, edited by J. Steele, K. Turekian, and S. Thorpe, Academic Press, 2001.

Santinelli, C., L. Nannicini, and A. Seritti, DOC dynamics in the meso and bathypelagic layers of the Mediterranean Sea, *Deep Sea Research Part II: Topical Studies in Oceanography*, *57* (16), 1446-1459, 2010.

Severin, T., P. Conan, X. Durrieu de Madron, L. Houpert, M.J. Oliver, L. Oriol, J. Caparros, J.F. Ghiglione, and M. Pujo-Pay, Impact of open-ocean convection on nutrients, phytoplankton biomass and activity, *Deep Sea Research Part I: Oceanographic Research Papers*, *94* (0), 62-71, 2014.

Severin, T., F. Kessouri, M. Rembauville, E.D. Sánchez-Pérez, L. Oriol, J. Caparros, M. Pujo-Pay, J.-F. Ghiglione, F. D'Ortenzio, V. Taillandier, N. Mayot, X. Durrieu De Madron, C. Ulses, C. Estournel, and P. Conan, Open-ocean convection process: A driver of the winter nutrient supply and the spring phytoplankton distribution in the Northwestern Mediterranean Sea, *Journal of Geophysical Research: Oceans*, *122* (6), 4587-4601, 2017.

Severin, T., C. Sauret, M. Boutrif, T. Duhaut, F. Kessouri, L. Oriol, J. Caparros, M. Pujo-Pay, X.D. de Madron, M. Garel, C. Tamburini, P. Conan, and J.F. Ghiglione, Impact of an intense water column mixing (0-1500m) on prokaryotic diversity and activities during an open-ocean convection event in the NW Mediterranean Sea, *Environ Microbiol*, 2016.

Seyfried, L., P. Marsaleix, E. Richard, and C. Estournel, Modelling deep-water formation in the north-west Mediterranean Sea with a new air–sea coupled model: sensitivity to turbulent flux parameterizations, *Ocean Sci.*, *13* (6), 1093-1112, 2017.

Siokou-Frangou, I., U. Christaki, M.G. Mazzocchi, M. Montresor, M. Ribera d'Alcalá, D. Vaqué, and A. Zingone, Plankton in the open Mediterranean Sea: a review, *Biogeosciences*, *7* (5), 1543-1586, 2010.

Somot, S., L. Houpert, F. Sevault, P. Testor, A. Bosse, I. Taupier-Letage, M.-N. Bouin, R. Waldman, C. Cassou, E. Sanchez-Gomez, X. Durrieu de Madron, F. Adloff, P. Nabat, and M. Herrmann, Characterizing, modelling and understanding the climate variability of the deep water formation in the North-Western Mediterranean Sea, *Climate Dynamics*, 2016.

Somot, S., F. Sevault, and M. Déqué, Transient climate change scenario simulation of the Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation model, *Climate Dynamics*, *27* (7), 851-879, 2006.

Song, X., and L. Yu, Air-sea heat flux climatologies in the Mediterranean Sea: Surface energy balance and its consistency with ocean heat storage, *Journal of Geophysical Research: Oceans*, *122* (5), 4068-4087, 2017.

Stemmann, L., G. Gorsky, J.-C. Marty, M. Picheral, and J.-C. Miquel, Four-year study of large-particle vertical distribution (0–1000m) in the NW Mediterranean in relation to hydrology, phytoplankton, and vertical flux, *Deep Sea Research Part II: Topical Studies in Oceanography*, *49* (11), 2143-2162, 2002.

Testor, P., A. Bosse, L. Houpert, F. Margirier, L. Mortier, H. Legoff, D. Dausse, M. Labaste, J. Karstensen, D. Hayes, A. Olita, A. Ribotti, K. Schroeder, J. Chiggiato, R. Onken, E. Heslop, B. Mourre, F. D'ortenzio, N. Mayot, H. Lavigne, O.d. Fommervault, L. Coppola, L. Prieur, V. Taillandier, X.D.d. Madron, F. Bourrin, G. Many, P. Damien, C. Estournel, P. Marsaleix, I. Taupier-Letage, P. Raimbault, R. Waldman, M.N. Bouin, H. Giordani, G. Caniaux, S. Somot, V. Ducrocq, and P. Conan, Multiscale Observations of Deep Convection in the Northwestern Mediterranean Sea during Winter 2012–2013 Using Multiple Platforms, *Journal of Geophysical Research: Oceans*, *123*, 1745–1776, 2018.

Testor, P., and J.-C. Gascard, Large-Scale Spreading of Deep Waters in the Western Mediterranean Sea by Submesoscale Coherent Eddies, *Journal of Physical Oceanography*, *33* (1), 75-87, 2003.

Ulses, C., P.A. Auger, K. Soetaert, P. Marsaleix, F. Diaz, L. Coppola, M.J. Herrmann, F. Kessouri, and C. Estournel, Budget of organic carbon in the North-Western Mediterranean open sea over the period 2004–2008 using 3-D coupled physical-biogeochemical modeling, *Journal of Geophysical Research: Oceans*, *121* (9), 7026-7055, 2016.

Vandromme, P., L. Stemmann, L. Berline, S. Gasparini, L. Mousseau, F. Prejger, O. Passafiume, J.M. Guarini, and G. Gorsky, Inter-annual fluctuations of zooplankton communities in the Bay of Villefranche-sur-mer from 1995 to 2005 (Northern Ligurian Sea, France), *Biogeosciences*, *8* (11), 3143-3158, 2011.

Vargas-Yáñez, M., P. Zunino, A. Benali, M. Delpy, F. Pastre, F. Moya, M.d.C. García-Martínez, and E. Tel, How much is the western Mediterranean really warming and salting?, *Journal of Geophysical Research: Oceans*, *115* (C4), 2010.

Waldman, R., M. Herrmann, S. Somot, T. Arsouze, R. Benshila, A. Bosse, J. Chanut, H. Giordani, F. Sevault, and P. Testor, Impact of the Mesoscale Dynamics on Ocean Deep Convection: The 2012–2013 Case Study in the Northwestern Mediterranean Sea, *Journal of Geophysical Research: Oceans*, *122* (11), 8813-8840, 2017a.

Waldman, R., S. Somot, M. Herrmann, A. Bosse, G. Caniaux, C. Estournel, L. Houpert, L. Prieur, F. Sevault, and P. Testor, Modeling the intense 2012–2013 dense water formation event in the northwestern Mediterranean Sea: Evaluation with an ensemble simulation approach, *Journal of Geophysical Research: Oceans*, *122* (2), 1297-1324, 2017b.

720 Waldman, R., S. Somot, M. Herrmann, F. Sevault, and P.E. Isachsen, On the Chaotic Variability of
721 Deep Convection in the Mediterranean Sea, *Geophysical Research Letters*, *45* (5), 2433-2443,
722 2018.
723 Waldman, R., S. Somot, M. Herrmann, P. Testor, C. Estournel, F. Sevault, L. Prieur, L. Mortier, L.
724 Coppola, V. Taillandier, P. Conan, and D. Dausse, Estimating dense water volume and its
725 evolution for the year 2012–2013 in the Northwestern Mediterranean Sea: An observing
726 system simulation experiment approach, *Journal of Geophysical Research: Oceans*, *121* (9),
727 6696-6716, 2016.

728

LEGENDS

Figure 1 - Time schedule of the different platforms and means used during the DEWEX experiment between July 2012 and July 2013

Figure 2 - Sketch of sampling strategy and examples of trajectory for airborne and oceanic platforms during the DEWEX experiment in the northwestern Mediterranean

Figure 3 - Side view of the airborne and oceanic platforms used during the DEWEX experiment



729

730 *Figure 1 - Time schedule of the different platforms and means used during the DEWEX experiment*
731 *between July 2012 and July 2013*

732

733



Figure 2 - Sketch of sampling strategy and examples of trajectory for airborne and oceanic platforms during the DEWEX experiment in the northwestern Mediterranean



Figure 3 - Side view of the airborne and oceanic platforms used during the DEWEX experiment