



HAL
open science

Use of organic amendment and endomycorrhizal fungi for steel slag phytostabilization

Mathieu Scattolin, Olivier Faure, Fernando Pereira, Nour-Eddine Menad,
Frédéric Paran, Frédéric Astolfi, Frédéric Gallice

► To cite this version:

Mathieu Scattolin, Olivier Faure, Fernando Pereira, Nour-Eddine Menad, Frédéric Paran, et al.. Use of organic amendment and endomycorrhizal fungi for steel slag phytostabilization. Slag Valorisation Symposium, KU Leuven - Department of Materials Engineering (MTM); EIT RawMaterials – Innovation Hub CLC West, Apr 2019, Mechelen, Belgium. hal-02191545

HAL Id: hal-02191545

<https://hal.science/hal-02191545v1>

Submitted on 23 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

USE OF ORGANIC AMENDMENT AND *ENDOMYCORRHIZAL FUNGI* FOR STEEL SLAG PHYTOSTABILIZATION

Mathieu SCATTOLIN^{1,2}, Olivier FAURE^{1,2}, Fernando PEREIRA^{1,2}, Nour-Eddine MENAD³, Frédéric PARAN^{1,2}, Frédéric ASTOLFI⁴, Frédéric GALLICE^{1,2}

¹ "Association pour la Recherche et le développement des Méthodes et processus INDUSTRIELS" (ARMINES), 75 272 Paris, France

² Mines Saint-Étienne, Centre "Sciences des Processus Industriels et Naturels" (SPIN), Département "Procédés pour l'Environnement et les Géo-ressources" (PEG), 42 023 Saint-Étienne, France

³ "Bureau de Recherches Géologiques et Minières" (BRGM), "Direction Eau, Environnement et Écotechnologies" (D3E), 45 060 Orléans, France

⁴ Industeel France ArcelorMittal, 42 800 Châteauneuf, France

mathieu.scattolin@emse.fr

Abstract

Steel slags are major by-products of the steel and iron industry. Particularly rich in metallic trace elements, these materials are massively stored in slagheaps and constitute as many potential nuisances for the environment and public health. Several studies [(Mendez et Maier, 2008), (Vamerali et al., 2010)] have shown the effectiveness of phytostabilization processes for rehabilitating sites contaminated by high levels of metals, both because of their sustainable and ecological aspects but also due to their low implementation costs. The slagheap studied in this work contains approximately 500 000 tons of slags spread over an area of about 4 ha. A previous study (Bourchardon et al., 2014) carried out on the same site has demonstrated that the use of "Composted Sewage Sludge" (CSS) as an organic amendment favors the installation of metallophytes herbaceous species naturally able to grow on this type of substrate. In this work we evaluated :1) the combined effects of CSS and inoculation of an AMF (Arbuscular Mycorrhizal Fungi) on vegetation in terms of biomass production as well as accumulation of metallic compounds in leaves and 2) the optimal amount of CSS in order to obtain the best rates of mycorrhization.

Keywords: Steel slag, Phytomanagement, Metallic trace elements, AMF.

Introduction

In France, slag deposits resulting from metallurgical activities have been estimated to about 40 millions of tons, spread over several hundred sites. In most cases, these sites are left bare, without any protective measure; they are thus exposed to wind and rainfalls, causing them to erode and leach. In this context, phytostabilization approaches (i.e. implementing a homogenous plant cover without bringing an extra layer of clean soil¹) could be quite efficient for managing sustainably large metallurgical slag dumps².

Recent studies demonstrated that using "Composted Sewage Sludge" (CSS), as a cheap and easily available organic amendment, was an efficient treatment to promote plant development on slags, provided that an assemblage of adapted species (drought tolerant pseudo-metallophyte species) was used³. Nevertheless, decreasing Cr accumulation in plants used for phytostabilizing slag dumpsites remains an important research issue. In this respect, a key-factor to mitigate metal accumulation in plants could rely on stimulating the development of "*Mycorrhizal Fungi*" (AMF) symbioses. In addition to a better tolerance to soil nutrient deficiencies and drought stress, there are a growing number of evidences that AMF could also immobilize some metals in their hyphae (and especially Cr) thereby decreasing the accumulation of toxic metals in plants⁴.

In this work, we hypothesized that using *Rhizophagus irregularis* (Ri), a generalist, easily commercially available AMF which has been used for a long time for agronomical practices, could be as efficient for phytostabilizing metallurgical slags. In order to test this hypothesis, we performed a pot experiment, combining CSS amendment and AMF inoculation, to evaluate: 1) the effects of CSS and Ri inoculation on vegetation in terms of biomass production, Cr accumulation and P concentration in leaves and 2) the optimal amount of CSS in order to obtain the best rates of mycorrhization.

Material and method

Slags used in this study were taken from a metallurgical landfill located at Châteauneuf (Loire, France) of about 4 ha. Pot experiments were performed to assess the combined effects of CSS level and Ri inoculation on plant growth, Cr accumulation and P concentration of two herbaceous species: *Festuca arundinacea* (*Poaceae*) (=F.a.) and *Melilotus officinalis* (*Fabaceae*) (=M.o.). Two CSS levels were compared: 60T·ha⁻¹ (=CSS60) and 120T·ha⁻¹ (=CSS120). AMF inoculation was done for each CSS level at the time of sowing. Thus, the overall experimental design comprised four treatments: CSS60-Ri0; CSS60-Ri; CSS120-Ri0 and CSS120-Ri, completely randomized with ten replicates by treatment. Root colonization was

quantified under optical microscope according to Trouvelot⁵ using: frequency of mycorrhization (F%), percentage of root cortex colonization (M%) and overall abundance of arbuscules (A%). The Kruskal-Wallis tests indicate the significant differences between the treatments (Holm adjustment, $P < 5\%$).

Results and discussion

Shoot dry weights obtained for each species and treatment are shown in Figure 1. Plants grown in CSS60 had very low aerial biomass as compared with those from CSS120. These strong differences could be mostly due to an increasing of water holding capacity, nutrient supply as well as a more pronounced pH decrease in CSS120. Substrate pH around 11 without amendment limit nutrient availability and drive metals speciation and solubility. Plants inoculated with AMF had a slightly higher biomass, but this was not statistically significant.

The most problematic metal on the study site was Cr; its accumulation in leaves is presented in Figure 2a. Results showed that plants grown on CSS60 exhibited higher Cr contents; this could be related to the very small size of these plants. Indeed, for tree species as well as herbaceous species, metal concentrations are generally higher in the early vegetative growth stages. At the opposite, late growth stages present higher biomass production, leading to dilution of accumulated metals.

As one of the main essential plant nutrients, P concentrations in leaves are shown in Figure 2b. P foliar content showed contrasted differences between treatments. The higher P contents were observed on plants growing on CSS120 in the presence of Ri ($P < 5\%$). This is consistent with the general observation that mycorrhizal plants are able to solubilize higher P amounts from soil and improve its concentration in organs.

Indicators of root colonization for both CSS levels are given in Table 1. The frequency of mycorrhization was significantly higher on CSS120 than on CSS60 for both plant species. Likewise, *F. a.* showed the highest frequencies of colonization when grown on CSS120 ($P < 5\%$). Calculation of the M% and A% parameters did not evidence significant differences between species growing on CSS120 and confirmed that the extension of root colonization was globally higher in that condition.




Figure 1: Shoot dry weigh yields of *F. a.* and *M.o.* after 3 months growth. Different letters mean statistical significance (Kruskal-Wallis, $P < 5\%$).




Figure 2: Cr (a) and P (b) concentration in leaves of *F. a.* and *M. o.* after 3 months growth. Different letters mean statistical significance (Kruskal-Wallis, $P < 5\%$).

Table 1: Mycorrhization status of *F. a.* and *M. o.* after 3 months growth.Different letters mean statistical significance (Kruskal-Wallis, $P < 5\%$).

Plant species	Treatment	F%	M%	A%
<i>F. arundinacea</i>	CSS60	47,50 ± 17,85 (bc)	4,90 ± 6,88 (ab)	2,40 ± 3,90 (ab)
	CSS120	94,00 ± 10,20 (a)	14,01 ± 10,29 (a)	3,36 ± 3,87 (a)
<i>M. officinalis</i>	CSS60	27,50 ± 10,90 (c)	0,48 ± 0,43 (b)	0,03 ± 0,04 (b)
	CSS120	58,89 ± 22,83 (b)	17,22 ± 15,85 (a)	5,52 ± 6,38 (a)

Conclusion

In this study, we investigated how CSS amendment and AMF inoculation impact plant development and Cr bioavailability. In addition to increasing water holding capacity and substrate nutrient content, CSS drove the growth of plant species. In addition, CSS amended at $120\text{T}\cdot\text{ha}^{-1}$ led to a lower accumulation of Cr in plants leaves. AMF inoculation slightly increased aerial biomass and improved plant nutrition by stimulating phosphorus uptake.

References

1. P. Kidd, M. Mench, V. Alvarez-Lopez, V. Bert, I. Dimitriou, W. Friesl-Hanl, R. Herzig, J.O. Janssen, A. Kolbas, I. Müller, S. Neu, G. Renella, A. Ruttens, J. Vangronsveld, M. Puschenreiter, "Agronomic practices for improving gentle remediation of trace element-contaminated soils", *International of Phytoremediation*, **17** 1005-1037 (2015).
2. M. Mench, N. Lepp, V. Bert, J.P. Schwitzguébel, S.W. Gawronski, P. Schröder, J. Vangronsveld, "Successes and limitations of phytotechnologies at field scale: outcomes, assessment and outlook from COST action 859", *Journal of Soils and Sediments*, **16** 876-900 (2010).
3. J.L. Bouchardon, O. Faure, G. Lespagnol, J. Moutte, B. Guy, D. Mimoun, D. Graillet, F. Paran, V. Croze, J.D. Athénol, B. Fernando, J. Boisson, S. Perret, A. Hitmi, P. Verney, C. Moussard Gauthier, G. Ledoigt, P. Goupil, C. Sac, M. Mench, "PHYSAFIMM : la phytoStabilisation, méthodologie Applicable aux Friches Industrielles, Métallurgiques et Minières", Rapport final. ADEME, 158 (2014).
4. S. Wu, X. Zhang, Y. Sun, Z. Wu, T. Li, Y. Hu, J. Lv, G. Li, Z. Zhang, J. Zhang, L. Zheng, X. Zhen, B. Chen, "Chromium immobilization by extraradical mycelium of arbuscular mycorrhiza contributes to plant chromium tolerance", *Environmental and Experimental Botany*, **122** 10-18 (2016).
5. A. Trouvelot, J.L. Kough, V. Gianinazzi-Pearson, "Mesure du taux de mycorrhization VA d'un système racinaire. Recherche de méthodes d'estimation ayant une signification fonctionnelle", *Physiological and Genetical Aspects of Mycorrhizae*, 217-221 (1986).