



HAL
open science

Gut microbiome in Chronic Rheumatic and Inflammatory Bowel Diseases: similarities and differences

Fatouma Salem, Nadège Kindt, Julian R. Marchesi, Patrick Netter, Anthony Lopez, Tunay Kokten, Silvio Danese, Jean-Yves Jouzeau, Laurent Peyrin-Biroulet, David Moulin

► To cite this version:

Fatouma Salem, Nadège Kindt, Julian R. Marchesi, Patrick Netter, Anthony Lopez, et al.. Gut microbiome in Chronic Rheumatic and Inflammatory Bowel Diseases: similarities and differences. United European Gastroenterology Journal, 2019, 7 (8), pp.1008-1032. 10.1177/2050640619867555 . hal-02192635

HAL Id: hal-02192635

<https://hal.science/hal-02192635>

Submitted on 24 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Gut microbiome in Chronic Rheumatic and Inflammatory Bowel Diseases: similarities and differences

Fatouma SALEM¹, Nadège KINDT¹, Julian R MARCHESI^{2,3}, Patrick NETTER¹, Anthony LOPEZ^{4,5}, Tunay KOKTEN⁴, Silvio DANESE⁶, Jean-Yves JOUZEAU¹, Laurent PEYRIN-BIROULET^{4,5*} and David MOULIN^{1,7*}

¹ IMoPA, UMR7365 CNRS-Université de Lorraine, Vandœuvre Les Nancy, France

² Division of Integrative Systems Medicine and Digestive Disease, Imperial College London, England

³ School of Biosciences, Museum Avenue, Cardiff University, Cardiff, UK

⁴ NGERE, UMR_ U1256 INSERM-Université de Lorraine, Vandœuvre Les Nancy, France

⁵ Service d'hépatogastroentérologie, CHRU de Nancy, Vandœuvre Les Nancy, France

⁶ Department of Biomedical Sciences, Humanitas University, Rozzano, Milan, Italy.

⁷ CHRU de Nancy, Contrat d'interface, Vandœuvre Les Nancy, France

*Authors contributed equally to this work and are corresponding:

Laurent PEYRIN-BIROULET: peyrinbiroulet@gmail.com, Service d'hépatogastroentérologie, CHRU de Nancy, Vandœuvre Les Nancy, France

David MOULIN: david.moulin@univ-lorraine.fr; IMoPA, UMR7365 CNRS-Université de Lorraine, Vandœuvre Les Nancy, France

Contributors

FS and NK: study concept and design; acquisition, analysis, review and interpretation of data; manuscript preparation and critical revisions. NK: study concept and design; review and interpretation of data; manuscript preparation and critical revision. JRM, PN, AL, TK, SD: review and interpretation of data; manuscript preparation and critical revisions. JYJ, LPB DM: study concept and design; review and interpretation of data; manuscript preparation and critical revisions; study supervision.

Competing interests

The authors declare to have no conflict of interest.

ABSTRACT

Introduction: Inflammatory bowel diseases (IBD), and Chronic Rheumatic Diseases (CRD) are systemic chronic disorders sharing common genetic, immune and environmental factors. About half of patients with IBD develop rheumatism affections and microscopic intestinal inflammation is present in up to half of CRD patients. IBD and CRD patients also share a common therapeutic armamentarium. Disequilibrium in the complex realm of microbes (known as dysbiosis) that closely interact with the gut mucosal immune system, has been associated with both IBD and CRD (Spondyloarthritis and Rheumatoid Arthritis). Whether dysbiosis represents an epiphenomenon or a prodromal feature remains to be determined.

Methods: In an attempt to further interrogate whether specific gut dysbiosis may be the missing link between IBD and CRD in patients developing both diseases, we performed here a systematic literature review focusing on studies looking at bacterial microbiota in CRD and/or IBD patients.

Results: We included 80 studies, with a total of 3799 IBD patients without arthritis, 1084 CRD patients without IBD, 132 IBD patients with arthropathy manifestations and 12 SpA patients with IBD history. Overall, this systematic review indicates that an increase in *Bifidobacterium*, *Staphylococcus*, *Enterococcus*, *Lactobacillus*, *Pseudomonas*, *Klebsiella* and *Proteus* genera, as well as a decrease in *Faecalibacterium*, *Roseburia* genera and species belonging to Verrucomicrobia, Fusobacteria phyla are common features in IBD and CRD patients, whereas dozens of bacterial species are specific features of CRD and IBD.

Conclusion: Further work is needed to understand the functions of bacteria and of their metabolites but also to characterize fungi and viruses that are commonly found in these patients.

Introduction

Inflammatory bowel diseases (IBD), are mainly represented by Crohn's disease (CD) and ulcerative colitis (UC), whereas Chronic Rheumatic Diseases (CRD), encompass Rheumatoid Arthritis and Spondyloarthritis (SpA). These systemic chronic disorders have relapsing and remitting clinical course arising from an interaction between genetic, immune and environmental factors.

CRD and IBD are intercurrent since articular manifestations are observed in up to 40% of IBD patients and intestinal inflammation is often present in CRD subjects ¹. Co-occurring CRD and IBD can be very disabling and are associated with a more severe disease course in IBD patients².

Interestingly, IBD and CRD share common pathophysiology, including common molecular and cellular actors and, consequently, common therapeutic armamentarium. Genetic studies have reinforced the importance of genes and pathways contributing to IBD pathogenesis, such as barrier function, the role of T cell subsets, and cytokine-cytokine receptor signalling ³. In addition, recent studies pointed out new genes and pathways, including autophagy or regulation of interleukin 23 (IL23) signalling, highlighting the importance of host defence pathways, specifically those involved in the management of mycobacteria ⁴. Heredity is also an important feature of CRD and notably in SpA, and several genetic polymorphisms have been shown to influence the disease risk. The most important one is the major histocompatibility complex (MHC) class I allele HLA-B27 ⁵. Remarkably, a large subset of the IBD and CRD susceptibility identified genes are encoding for proteins involved in immune response, and particularly in the IL-23/Th17 pathway of T cell differentiation, which is primarily implicated in response against extracellular pathogens, including bacteria and yeasts, and/or in microbial sensing.

However, the link between pathological gut and joint inflammation in patients with both IBD and CRD is not fully understood. Taken together, these data suggest that the perturbation of the gut microbiome, also called dysbiosis represent an attractive target in this context.

In an attempt to further interrogate whether specific gut dysbiosis may be associated with IBD and CRD and promote pathological inflammation within joint-gut axis, we performed a systematic literature review investigating similarities and differences regarding faecal microbiota in these patients.

Methods

Search strategy and study selection

A systematic literature search was performed according to PRISMA guidelines ⁶. The literature review conducted using PubMed/MEDLINE (from 1950 to December 2018), Web of science (from 1958 to December 2018). Abstracts from annual meetings of national and international gastroenterology and rheumatology conferences (United European Gastroenterology Week [UegW], Digestive Diseases Week [DDW], European Crohn's and Colitis Organization [ECCO], European League Against Rheumatism [EULAR], American College of Rheumatology [ACR]) were searched manually from 2013 to 2018.

The following keywords were searched in various combinations using the boolean terms “AND” and “OR” (“Microbiota”, “Microbiome”, “Gut”, “Gastrointestinal Microbiome”, “Microbiology”, “Colitis”, “Ileitis”, “Intestinal”, “Enteritis”, “Inflammatory Bowel Diseases”, “Crohn Disease”, “Ulcerative Colitis”, “Rheumatoid Arthritis”, “Spondyloarthritis”, “Arthritis”, “Reactive Arthritis”, “Psoriatic Arthritis”, “Rheumatoid Arthritis”, “Infectious Arthritis”, “Ankylosing Spondylitis”, “Mycobiome”, “Fungal Microbiota”, “Intestinal Virome”). This strategy was used both as Medical Subject Headings (MeSH) terms if available and as free text. Searching was limited to publications with human subjects. We only selected English language full text papers and abstracts.

Two authors independently reviewed all articles. Inclusion criteria included the presence of IBD and CRD patient samples and 16S rRNA gene sequencing or metagenomic methods to characterize the gut microbiota. Literature reviews did not include meta-analyses, as well as experimental studies based on *in vitro* findings and animal models.

Study characteristics and outcomes were reported in a Microsoft Excel Office 2016 Professional spread sheets.

Results

6519 were identified (Fig 1) based on defined criteria. After review of the titles and abstracts 5564 papers were excluded. Amongst the remaining studies, another 881 were excluded because they included reviews, data retrieved from studies using animal models and *in vitro* findings. Therefore, 80 studies were included: 56 from IBD patients, with 1 Case-reports³⁹ (Table 1.a, 1.b and 1.c), 21 from CRD patients (RA and SpA) including 5 congress abstracts^{78,80,81,84,85} (Table 2.a and 2.b). Finally, three publications addressed gut microbiota study in IBD patients developing arthropathy^{94,96,97} (Table 3). As microbiota from one individual is different from one sample location to another, table were generated by sample type and are detailed with studied populations characteristics.

1. Literature search results

A. Distinct dysbiosis in IBD and CRD

In order to identify bacterial variations specific of IBD, (i.e. not found in CRD), and vice versa, we adopted two complementary methodologies: we first reviewed bacterial changes reported in studies enrolling IBD patients without information on possible concomitant arthritis, then all studies involving CRD patients without information on possible concomitant IBD. We looked finally at studies comparing gut microbiome in patients with or without IBD-associated CRD.

A.1. Gut bacterial changes reported in IBD patients

Fifty-six studies enrolling 3270 IBD patients from which gut microbiota was mainly analyzed by 16S rRNA gene sequencing or qRNA of DNA extracted from faeces and/or biopsies. A quantitative and qualitative (biodiversity) reduction of the gut microbiome in IBD patients^{7,8} is generally observed.

Firmicutes phyla

A reduction of *Clostridiales* order species from the Firmicutes phylum is observed in the faecal microbiota of IBD and CD patients^{9–11}. Whereas an enrichment of *R.gnavus* is observed in the IBD patients faecal microbiota^{12–14}. This phylogenetic group includes several butyrate-producing bacteria, notably *Faecalibacterium* and *Ruminococcus*, which are among the main members of the *Ruminococcaceae* genera¹⁵. Other bacteria that are considered as ‘beneficial’ for the host have been shown to be quantitatively reduced in the faecal microbiota of these patients. A few studies found a lower number of sequences of the bacterial phylum Firmicutes in the mucosal-associated microbiota (MAM) of CD and UC patients, especially species from the *Lachnospiraceae* genera (*Roseburia* and *coprococcus*)^{12,13,16,17, 18,19,20,21}. Within this phylum, an increase amount of *Streptococcus* genera was observed, in contrast to *Ruminococcaceae* genera (*Faecalibacterium*) that seems to be particularly deficient in CD^{15,20,22–24}. Furthermore, Rehman *et al* demonstrated a population-specific disease-related patterns of Firmicutes phyla, by observing a lower abundance in healthy German samples compared with patients samples, while Lithuanian and Indian patients with CD show the lowest Firmicutes abundances²⁵.

In a recent study using molecular methods of bacterial identification¹⁹, it has been shown that *F. prausnitzii* was one of the most underrepresented species of the *Faecalibacterium* genera in the MAM of patients with IBD (compared with healthy subjects)^{12,13,15,19,22,23,26–29}. Therefore, similar to the results from faecal microbiota studies, a significant decrease of bacteria from the Firmicutes phylum was demonstrated in the MAM of CD patients^{15,22,23}.

A reduction of *Ruminococcaceae*, *Lactobacillaceae*, *Veillonellaceae* and *Erysipelotrichiaceae* genera (*Faecalibacterium*, *Streptococcus*, *Veillonella* and *Catenibacterium* respectively)^{10,13,30–32}, along with *Dialister* genus in CD patients³³, and *Roseburia*, *Clostridium* and

Butyricimonas genus is observed in IBD patients particularly those with UC^{15,21,31,32,34}. A few studies, showed an increased amount of the *Tissierellaceae* family, and a decreased number of *Eubacterium* genera in inflamed colonic mucosa biopsy samples when compared to the non-inflamed sites in UC patients³⁵⁻³⁷. (Fig 2)

Bacteroidetes phyla

Data concerning the Bacteroidetes phylum are more conflicting. Some studies reported a reduction of the *Bacteroides* group in IBD patients especially in CD patients^{9,12,13,20}. In contrast, Andoh and colleagues demonstrated an increase amount of this phyla in the context of IBD³⁸. To note, one study showed an increase of Bacteroidetes phylum in salivary microbiota in UC patients³⁹. Hirano and co-workers showed an enrichment of the *Cloacibacterium* genus, and decreased abundance of *Prevotella* (at both inflamed and non-inflamed mucosal site) and *Butyricimonas* genera at the non-inflamed mucosal site of UC patients compared to the corresponding site in non-IBD controls and in the faecal microbiota of UC patients^{12,34,35,40}. A greater abundance in these two genera was found in the submucosal tissues of patients with CD^{12,34,35,40,41}. As with CD, this strongly suggest a restricted biodiversity in UC and an increased proportion of unusual bacteria^{42,43}. Bacteroidetes show also interesting age-related patterns and population-independent increase in abundance in the standing and active bacteria among healthy subjects and UC patients²⁵. A decrease abundance of *Parabacteroides* genera and *Odoribacteraceae* family in IBD and CD patients respectively is reported^{10,13,15}. Similar to the results from faecal microbiota studies, a significant decrease of bacteria from the phylum Firmicutes was demonstrated in the MAM of patients with CD^{44,45}. A recent study of Walujkar and colleagues revealed significant differences in the MAM of patients manifesting acute exacerbations of UC with increased amount of *Parabacteroides* and *Elizabethkingia* genera in the MAM of UC patients as compared to the same patients during remission stage⁴⁶ (Fig 2).

Actinobacteria phyla

Concerning the Actinobacteria phylum, studies using both culture and recent molecular methods, demonstrated an increase of *Bifidobacterium* genera in the faecal microbiota as well as in the biopsy samples of IBD patients, notably in patients with CD^{12,13,15,19,47}. However, other authors reported an age-related reduction of bacteria of the *Bifidobacterium* genera was

shown in inflamed sites when compared to non-inflamed ones and salivary microbiota of UC patients^{12,13,15,47–49,31,35,39,50}. Walujkar and co-workers showed an increase amount of *Micrococcus* genera in MAM of UC patients when compared to non-IBD subjects⁴⁶ (Fig 2).

Proteobacteria phyla

Published studies display a quantitative alteration of Proteobacteria phylum in IBD especially *Escherichia* and *Shigella* from the *Enterobacteriaceae* family,^{10,12,13,15,24,29,51}. Thus, their increased abundance was reported in the MAM and faecal samples of patients with CD, whether using culture^{24,44} or molecular^{52,17,53} methods. As with CD patients, the MAM of patients with UC contained an abnormally elevated concentration of bacteria, especially anaerobes^{44,45}. A restriction of the MAM biodiversity similar to that observed in patients with CD has been found such as reduction of Firmicutes and an overrepresentation of *Enterobacteriaceae*^{19,25 44 45 54 55 56}. A decreased abundance of the genera *Bilophila* and *Desulfovibrio* was evident at the inflamed site of UC patients compared to the corresponding site of non-IBD controls, whereas a decreased amount of *Bilophila* genera and its species (*B.wadsworthia*) was detected in the faecal microbiota of CD patients^{35,57,58}. Moreover, an age-related reduction of the *Neisseria* genera bacteria was reported in inflamed sites when compared to non-inflamed ones and salivary microbiota of UC patients^{12,13,15,47–49,31,35,39,50}. Walujkar *et al.* suggested an increased abundance of *Stenotrophomonas*, *Ochrobactrum* and *Achromobacter* genera in UC patients as compared to the same patients during remission stage⁴⁶. Finally, Proteobacteria phyla displayed also an age-related patterns²⁵.

(Fig 2)

Other phyla

Finally, a decreased in abundance of Verrucomicrobia (*Akkermansia*) and Fusobacteria (*Leptotrichia*), was reported at the inflamed colonic mucosal sites of CD and UC patients compared to the corresponding site of non-IBD controls. However, further investigation concerning an eventual association between *Leptotrichia* and UC is necessary^{12,31,32,35,59–61}.

In summary, among the 56 available studies on IBD, differential abundance of 40 bacterial species has been reported, 15 were specifically found in CD studies while only 16 species reported in UC studies. These variations mainly concerned Firmicutes, Proteobacteria and Bacteroidetes.

Table 1.a: Bacteria associated with inflammatory bowel disease analysed from biopsy samples.

Author	Methods	Sample origins	Study Cohort	Study Cohort characteristics at the time of sampling			Major findings
				Gender (no. M/no. F)	Mean age (range)	Geo. Origin	
Seksik, P. et al 2005	▪ TTGE of 16S rRNAs	▪ Biopsy samples	▪ 15 CD	(6/9)	▪ 37.6 ▪ (21–63)	France	No bacterial species was found to be specifically associated with CD ulceration, and ulceration did not qualitatively modify the dominant associated microbiota
Ott, S.J. et al., 2004	▪ 16S rDNA based SSCP fingerprint	▪ Biopsy samples	▪ 26 CD ▪ 31 UC ▪ 15 Inflammatory controls ▪ 31 Non-inflammatory controls	▪ (9/17) ▪ (18/13) ▪ (6/9) ▪ (10/21)	▪ 35 (16–56) ▪ 44 (23–74) ▪ 50 (20–82) ▪ 52 (26–74)	N/A	- <i>Bacteroides</i> , <i>Prevotella</i> (↓IBD)
Morgan, X.C. et al., 2012	▪ 16S rRNA-sequencing ▪ WGS	▪ Biopsy samples	▪ 121 CD ▪ 75 UC ▪ 8 Indeterminate ▪ 27 Controls	▪ (49/72) ▪ (38/37) ▪ (3/5) ▪ (12/15)	▪ 38 (35–41) ▪ 42 (38–45) ▪ 27(14–41) ▪ 36 (30–42)	USA	- <i>Prevotella</i> , <i>Streptococcus</i> , <i>Catenibacteria</i> (↓UC) - <i>Roseburia</i> , <i>Ruminococcus</i> (↓CD) - <i>Lactobacillus</i> , <i>Acidaminococcus</i> , <i>Veillonella</i> , <i>Shigella</i> , <i>Aeromonas</i> , <i>Fusobacterium</i> , <i>Shigella</i> (↑CD) - <i>Asteroleplasma</i> , <i>Porphyromonas</i> , <i>Bifidobacterium</i> , <i>Faecalibacterium</i> , <i>Coprococcus</i> (↓IBD)
Ananthakrishnan, AN. et al. 2017	▪ Metagenomic sequencing	▪ Biopsy samples	▪ 42 CD ▪ 43 UC	N/A	N/A	N/A	- <i>Roseburia inulinivorans</i> , <i>Burkholderiales</i> species (↑CD at 14 weeks remission)
Frank, D. N. et al 2007	▪ 16S rRNA sequencing	▪ Biopsy samples	▪ 68 CD ▪ 61 UC ▪ 61 Non-IBD Controls	N/A	▪ 35 (21–49) ▪ 38 (22–54) ▪ 36 (23–49)	N/A	- <i>Bacteroides</i> (<i>B. thetaiotaomicron</i>), <i>Lachnospiraceae</i> (↓IBD) - <i>Actinobacteria</i> , <i>Proteobacteria</i> (↑IBD)
Willing, BP. et al., 2010	▪ T-RFLP ▪ Cloning and 16S rRNA Sequencing	▪ Biopsy from 5 locations between the ileum and rectum	▪ 6 L1-CD ▪ 8 L2-CD ▪ 6 Controls	▪ (3/3) ▪ (6/2) ▪ (3/3)	Born between (1936–1986)	N/A	- <i>F. prausnitzii</i> (↓L1-CD) - <i>E.coli</i> (↑L1-CD)
Png, C. W. et al. 2010	▪ 16S rRNA qPCR ▪ In vitro mucus degradation test	Biopsy samples	▪ 26 CD ▪ 20 UC ▪ 20 Controls	▪ (6 / 20) ▪ (13 / 7) ▪ (9 / 11)	▪ 38 (19 – 74) ▪ 48 (24 – 71) ▪ 53 (22 – 84)	N/A	- <i>R. gnavus</i> <i>R. torques</i> (↑CD/UC) - <i>Akkermansia muciniphila</i> , (↓CD/UC)
Hansen, R. et al 2012	▪ 16S rRNA RT-PCR and pyrosequencing	▪ Colonic mucosa biopsy samples	▪ 13 CD ▪ 12 UC ▪ 12 Controls	▪ (10/3) ▪ (9/3) ▪ (8/4)	▪ 13 (8- 17) ▪ 13 (9 - 16) ▪ 12 (7- 16)	Scotland, UK	- <i>Faecalibacterium</i> (↑CD)
Wang, M. et al 2007	▪ 16S rRNA sequencing	▪ Colonic biopsy samples	▪ 1 UC (colonic microbiota)	▪ (0/1)	▪ 12-year-old	N/A	- <i>Enterobacteriaceae</i> , <i>Bacteroides fragilis</i> , <i>F. prausnitzii</i> -like, <i>Pseudomonas aeruginosa</i> (↑UC)

Rehman, A. et al 2016	16S rRNA pyrosequencing	Mucosal biopsy samples	27 CD (10 Ger.; 8 Lith.; 9 Ind.) 30 UC (10 Ger.; 10 Lith.; 10 Ind.) 30 Controls (10 Ger.; 9 Lith.; 11 Ind.)	Ger. (14/16) Lith. (10/17) Ind. (21/19)	Ger.(16-63) Lith.(19-81) Ind. (17-67)	Germany Lithuania India	- Firmicutes (↓Ger. Controls /CD Lith. Ind.) - Bacteroidetes (↑UC) - Proteobacteria (↑CD Lith./Ind.)
Hirano, A. et al. 2018	16S rRNA sequencing	Mucosal biopsies samples	14 UC 14 Non-IBD (Controls)	(6/8) (8/6)	45 (17-67) 59 (41-73)	N/A	- Cloacibacterium, Neisseria genus, Tissierellaceae family, (↑inflamed site UC compared to non-inflamed site UC) - Prevotella, Eubacterium, Neisseria, Leptotrichia, Bilophila, Desulfovibrio, Butyricimonas (↓UC corresponding site of non-IBD controls). - Prevotella, Butyricimonas (↓UC patients compared with the corresponding site in non-IBD controls)
Chiodini, R. J. et al. 2015	Deep 16S rRNA sequencing	Ilea mucosal and submucosal biopsy samples	20 CD 15 Non-IBD (Controls)	(9/11) (4/11)	41 (24-66) 59 (32-88)	USA	- Desulfovibrionales (↑CD in the subjacent submucosa as compared to the parallel mucosal tissue including) - Ruminococcus spp., Oscillospira spp., Pseudobutyribrio spp., Tumbacillus spp., Propionibacterium spp., Cloacibacterium spp., Proteobacteria (Parasutterella spp., Methylobacterium spp) (↑CD)
Swidsinski, A. et al. 2002	16S rRNA sequences FISH 3 group-specific FISH probes	Colonic biopsy samples	54 CD 119 UC 104 In.C 28 S.I.C 40 Controls	(25/29) (52/67) (46/58) (16/12) (23/17)	35 (17-86) 45 (1786) 46 (19-81) 37(17-70) 50 (26-77)	Berlin , Germany	No principal difference in the composition of the mucosal flora in IBD patients and controls. Species isolated from the washed mucosa were of faecal origin in all groups. Proportion of Enterococci/Streptococci, Clostridia, Peptostreptococci, Eubacteria were lower - Proportion of Collinsella aerofaciens or Propionibacteria higher than usually found in faecal specimens
Swidsinski, A. et al. 2005	FISH 14 group-specific FISH probes	Mucosal Biopsy samples	20 CD 20 UC 20 IBS 10 IBD + antibiotics 20 Controls	(11/9) (9/11) (6/14) (4/6) (7/13)	33 45 48 40 47	N/A	An adherent mucosal biofilm mainly composed of Bacteroides fragilis is a prominent feature in patients with IBD, while biofilm composed of Eubacterium rectale group in IBS.
Walujkar, SA. et al., 2018	16S rRNA gene-based sequencing	Colon biopsy samples	12 UC 7 Non-IBD (Controls)	N/A	(30- 41) (37-54)	Maharashtra, India	Stenotrophomonas, Ochrobactrum, Achromobacter (↑UC)
Kotowski, R. et al. 2007	RISA DNA sequencing	Biopsy samples	13 CD 19 UC 15 Controls	N/A	N/A	Canada	Enterobacteriaceae (↑IBD)
Sokol, H. et al. 2007	TTGE	Biopsies samples	3 Proctitis 7 Left-sided colitis	N/A	N/A	N/A	E.coli subdominant bacteria
Zhang, M. et al. 2007	DGGE analysis	Mucosal biopsy samples	24 UC	(9/15)	40 (16-72)	China	- Lactobacilli, Clostridium leptum subgroup were significantly different

Author	Methods	Sample origins	Study Cohort	Patients characteristics at the time of sampling			Major findings
				Gender (no. M/no. F)	Mean age (range)	Geo. Origin	
Scanlan, PD. et al., 2006	16S rRNA sequencing DGGE	Faecal samples	11 CD (Remission) 5 CD (Relaps) 18 Controls	(7/3) (2/3) (10/8)	40 (25–70) 46 (25–54) 36 (25–51)	N/A	Clostridiales order species
Hourigan, SK. et al., 2015	FISH 5 group-specific FISH sequencing probes	Rectal biopsies Faecal samples	36 CD 46 UC 14 Controls	(1/5) 19/27 (6/8)	51 (16–66) 54 (27–66) 37 (22–69)	N/A N/A	↓ Clostridiales order species -E.coli, Clostridia (↑A-UC) -E. coli (↑CD)
Gevers, D. et al., 2014	16S rRNA-sequencing WGS	Faecal samples	447 CD 221 Controls	N/A	(<17)	North America	- Odoribacter, Roseburia, Faecalibacterium (↓IBD/CD/UC) - Bifidobacterium, (↓IBD, ↑UC) - Coprococcus (↓IBD/CD) - Escherichia, Shigella (↑IBD) - Lactobacillus (↑IBD/CD) - Ruminococcus, Clostridium, Eubacterium

Table 1.b: Bacteria associated with inflammatory bowel disease analysed from faecal samples.

							(↓CD) - <i>Enterococci</i> (↑CD)
<i>Hall, AB. et al., 2017</i>	Metagenomic sequencing	Faecal samples	<ul style="list-style-type: none"> 9 CD 10 UC 1 Indeterminate Colitis 12 Controls (3 with Gastrointestinal symptoms) 	N/A	N/A	N/A	- <i>R.gnavus</i> (↑IBD)
<i>Kaakoush, NO. et al. 2012</i>	High-throughput sequencing of 16S rRNA	Faecal samples	<ul style="list-style-type: none"> 19 L1/L4 CD 21 Controls 	<ul style="list-style-type: none"> (12/7) (13/8) 	<ul style="list-style-type: none"> 12 (11-15) 10 (9-14) 	Sydney, Australia	- <i>Oscillospira</i> (↓CD)
<i>Aomatsu, T. et al 2012</i>	16S rRNA Sequencing T-RFLP analysis	Faecal samples	<ul style="list-style-type: none"> 10 CD 14 UC 27 Controls 	<ul style="list-style-type: none"> (4/6) (5/9) (12/15) 	<ul style="list-style-type: none"> (8-18) (8-15) (1-5) 	N/A	<ul style="list-style-type: none"> - <i>Parabacteroides</i>, <i>Bacteroides</i>, <i>Roseburia</i>, <i>Coprococcus</i>, <i>Blautia</i>, <i>Dorea</i>, <i>Ruminococcus</i>, <i>Oscillospira</i>, <i>Eubacteria</i>, <i>Dialister</i>, <i>Sutterella</i>, <i>Bilophila</i> (↓CD) - <i>Lactobacillus</i>, <i>Streptococcus</i>, <i>Enterococcus</i>, <i>Gemella</i>, <i>Haemophilus</i> (spp.), <i>Eikenella</i> (↑CD) - <i>Bacteroides</i> (↓IBD)
<i>Machiels, K. et al 2014</i>	DGGE of 16S rRNA Metabolites quantification by gas chromatography–mass spectrometry	Faecal samples	<ul style="list-style-type: none"> 127 UC 87 Controls 	<ul style="list-style-type: none"> (74/53) (39/48) 	<ul style="list-style-type: none"> 43 (32-55) 42 (30-53) 	Belgium	<ul style="list-style-type: none"> <i>Roseburia</i> (<i>R. hominis</i>), <i>Clostridium</i>, <i>Butyricimonas</i>, <i>F. prausnitzii</i> (↓IBD : UC)
<i>Duboc, H. et al 2013</i>	16S rRNA qPCR	Faecal samples	<ul style="list-style-type: none"> 7 A-CD 5 R-CD 16 A-UC 14 R-UC 29 Controls 	<ul style="list-style-type: none"> (3/4) (2/3) (7/9) (9/5) (11/18) 	<ul style="list-style-type: none"> 38 (19-57) 42 (23-61) 36 (22-50) 38 (26-50) 35 (21-49) 	N/A	<ul style="list-style-type: none"> - <i>Clostridium</i> (<i>C. leptum</i>), <i>Blautia</i> (<i>B. coccoides</i>) (↓IBD) - <i>F. prausnitzii</i> (↓CD) - <i>Escherichia</i> (<i>E. coli</i>) (↑IBD)
<i>Fujimoto, T. et al 2012</i>	16S rRNA qPCR T-RFLP	Faecal samples	<ul style="list-style-type: none"> 47 CD 20 Controls 	<ul style="list-style-type: none"> (31/16) (14/6) 	<ul style="list-style-type: none"> 36 (26-45) 45 (28/62) 	Japon	<i>F. prausnitzii</i> (↓CD)
<i>Pascal, V et al. 2017</i>	16S rDNA sequencing	Faecal samples	<ul style="list-style-type: none"> Spanish cohort (34 CD, 33 UC, 111 Controls) Belgian cohort (53 CD) 	<ul style="list-style-type: none"> (21/13) (25/28) 	<ul style="list-style-type: none"> 34 (18–58) 41 (27–53) 	<ul style="list-style-type: none"> Spain Belgium 	<ul style="list-style-type: none"> - <i>Faecalibacterium</i>, <i>Peptostreptococcaceae</i>, <i>Anaerostipes</i>, <i>Methanobrevibacter</i>, <i>Christensenellaceae</i>, <i>Collinsella</i> (↓CD) - <i>Fusobacterium</i>, <i>Escherichia</i> (↑CD)
<i>Swidsinski, A. et al 2008</i>	Fluorescence in situ hybridization (FISH)	Faecal samples	<ul style="list-style-type: none"> 82 CD 105 UC 32 Controls 	N/A	<ul style="list-style-type: none"> 34.8 (17-78) 41.2(18-84) 40 (18-60) 	Germany	<ul style="list-style-type: none"> - <i>F.prausnitzii</i>, (↓CD/↑UC) - <i>Enterobacteriaceae</i> (↑CD/UC) - <i>Enterobacteriaceae</i> (↓CD/↑UC) - <i>Eubacterium hallii</i>, <i>E. cylindroides</i> bacteria (↓CD) - <i>Bifidobacteria</i>, <i>Atopobium</i> (↑UC)
<i>Sokol, H. et al. 2009</i>	16S rRNA	Faecal samples	<ul style="list-style-type: none"> 22 A-CD 10 R-CD 13 A-UC 	<ul style="list-style-type: none"> (7/15) (4/6) (8/5) 	<ul style="list-style-type: none"> 37 (34 - 41) 40 (35 - 44) 40 (37 - 44) 	N/A	- <i>F.prausnitzii</i> (↓R-IBD/ IC/ A-CD/A-UC)

			<ul style="list-style-type: none"> ▪ 4 R-UC ▪ 8 IC ▪ 27 Controls 	<ul style="list-style-type: none"> ▪ (1/3) ▪ (5/3) ▪ (11/16) 	<ul style="list-style-type: none"> ▪ 35 (31 – 40) ▪ 34 (29 - 39) ▪ 36 (35- 37) 		- <i>Bifidobacterium</i> (↓IC)
Sabino, J. et al 2016	16S rDNA sequencing	Faecal samples	<ul style="list-style-type: none"> ▪ 18 PSC only ▪ 27 PSC-UC ▪ 21 PSC-CD ▪ 30 CD ▪ 13 UC ▪ 52 Controls 	<ul style="list-style-type: none"> ▪ (10/8) ▪ (20/7) ▪ (18/3) ▪ (15/15) ▪ (4/9) ▪ (49/3) 	<ul style="list-style-type: none"> ▪ Median age 49 (15.25) ▪ Median age 43 (14) ▪ Median age 49 (17) ▪ Median age 52 (14.25) ▪ Median age 50 (28) ▪ Median age 51.5 (17) 	Belgium	<i>Enterococcus</i> , <i>Fusobacterium</i> , <i>Lactobacillus</i> (↑PSC only/ PSC-UC/PSC-CD)
Bajer, L. et al. 2017	16S rRNA Sequencing	Faecal samples	<ul style="list-style-type: none"> ▪ 32 PSC-IBD ▪ 31 Controls 	<ul style="list-style-type: none"> ▪ (17/15) ▪ (13/18) 	<ul style="list-style-type: none"> ▪ 40 (20-71) ▪ 44 (22-72) 	Prague	- <i>Rothia</i> , <i>R. mucilaginosa</i> , <i>Fusobacteriaceae</i> (↑PSC-IBD) - <i>Adlercreutzia</i> , <i>Ruminococcus</i> (↓PSC-IBD) - <i>Butyricoccus pullicaecorum</i> sp (↓UC)
Eeckhaut, V. et al. 2013	16S rRNA sequencing Genus-specific qPCR	Faecal samples	<ul style="list-style-type: none"> ▪ 51 CD ▪ 91 UC ▪ 88 Controls 	<ul style="list-style-type: none"> ▪ (23/21) ▪ (54/37) ▪ (39/49) 	<ul style="list-style-type: none"> ▪ Median age 39 ▪ Median age 44 ▪ Median age 41 	N/A	<i>Butyricoccus</i> (↓CD/UC)
Knoll, R. L. et al. 2016	Metagenomic analysis	Faecal samples	<ul style="list-style-type: none"> ▪ 6 CD ▪ 6 UC ▪ 12 Controls 	<ul style="list-style-type: none"> ▪ (3/3) ▪ (2/4) ▪ (6/6) 	<ul style="list-style-type: none"> ▪ (11-17) ▪ (11-16) ▪ (8-20) 	N/A	- <i>F. prausnitzii</i> , <i>E. rectale</i> (↓CD/UC) - <i>E.coli</i> , <i>F.nucleatum</i> , <i>E. coli</i> , <i>F. nucleatum</i> (↑IBD)
Andoh, A. et al. 2011	16S rRNA sequencing T-RFLP PCR T-RFL	Faecal samples	<ul style="list-style-type: none"> ▪ 31 CD ▪ 31 UC ▪ 30 Controls 	<ul style="list-style-type: none"> ▪ (16/15) ▪ (15/16) ▪ (12/18) 	<ul style="list-style-type: none"> ▪ 30 ▪ 33 ▪ 35 	N/A	- <i>Clostridium</i> (↓IBD)
Sokol, H. et al. 2006	16S rDNA and rRNA PCR TTAG	Faecal samples	<ul style="list-style-type: none"> ▪ 9 UC ▪ 9 Controls 	<ul style="list-style-type: none"> ▪ (5/4) ▪ (6/3) 	<ul style="list-style-type: none"> ▪ 39 (25-69) ▪ 43 (23-69) 	N/A	- <i>Clostridium coccooides</i> (↓UC)
Sokol, H. et al. 2006	FISH 6 group-specific FISH probes Flow cytometry	Faecal samples	<ul style="list-style-type: none"> ▪ 13 CD ▪ 13 UC ▪ 5 IC ▪ 13 Controls 	<ul style="list-style-type: none"> ▪ (2/11) ▪ (7/6) ▪ (2/3) ▪ (7/6) 	<ul style="list-style-type: none"> ▪ 37(24–50) ▪ 41(28–54) ▪ 29(25–33) ▪ 40(25–56) 	N/A	- <i>C. coccooides</i> (↓UC) - <i>C. leptum</i> (↓CD) - <i>Bacteroides</i> (↑IC)
Giaffer, MH. et al. 1991	16S rRNA quantitative and semi-quantitative bacterial culture techniques	Faecal samples	<ul style="list-style-type: none"> ▪ 22 A-CD ▪ 20 Quiescent CD ▪ 18 A-UC ▪ 19 Quiescent UC ▪ 21 Controls 	<ul style="list-style-type: none"> ▪ (6/16) ▪ (5/15) ▪ (8/10) ▪ (7/12) ▪ (11/10) 	<ul style="list-style-type: none"> ▪ 38 ▪ 50 ▪ 37 ▪ 50 ▪ 35 	N/A	<i>Lactobacillus</i> , <i>Bifidobacteria</i> (↓CD)
Seksik, P. et al 2003	16S rDNA quantitative dot blot hybridization TTGE of 16S rDNA	Faecal samples	<ul style="list-style-type: none"> ▪ 8 A-CD ▪ 13 R-CD ▪ 16 Controls 	<ul style="list-style-type: none"> ▪ (1/7) ▪ (3/6) ▪ (7/9) 	<ul style="list-style-type: none"> ▪ 35 (16–68) ▪ 47 (32–62) 	N/A	- <i>Enterobacteria</i> (↑CD)
Schwiertz, A. et al 2010	16S rRNA sequencing	Faecal samples	<ul style="list-style-type: none"> ▪ 21 A-CD ▪ 19 R-CD ▪ 13 A-UC ▪ 16 R-UC ▪ 25 Controls 	N/A	14 (5-19)	N/A	- <i>Bifidobacteria</i> (↓IBD) - <i>Faecalibacterium</i> (↓CD)

Thorildsen, L. T. et al. 2013	<ul style="list-style-type: none"> ▪ 16S rRNA sequencing ▪ MCR 	<ul style="list-style-type: none"> ▪ Faecal samples 	<ul style="list-style-type: none"> ▪ 30 CD ▪ 33 UC ▪ 3 IBDU ▪ 33 Non-IBD 	<ul style="list-style-type: none"> ▪ (10/20) ▪ (17/16) ▪ (1/2) ▪ (14/19) 	<ul style="list-style-type: none"> ▪ 33 (21-53) ▪ 34 (17-62) ▪ 42(35-53) ▪ 33 (20-56) 	Norway	<ul style="list-style-type: none"> - <i>Escherichia</i> (↑CD) - <i>Shigella</i> (↑IBD/CD)
Martinez-Medina, M. et al. 2006	<ul style="list-style-type: none"> ▪ 16S rRNA gene sequencing ▪ PCR-DGGE ▪ BLAST database 	<ul style="list-style-type: none"> ▪ Faecal samples 	<ul style="list-style-type: none"> ▪ 19 CD ▪ 2 UC ▪ 1 Ischemic colitis ▪ 15 Controls 	<ul style="list-style-type: none"> ▪ (9/10) ▪ (1/1) ▪ (0/1) ▪ (5/11) 	<ul style="list-style-type: none"> ▪ (33-41) ▪ (29-34) ▪ 27 ▪ (43-50) 	N/A	<ul style="list-style-type: none"> - <i>Clostridium</i> spp - <i>Ruminococcus</i> - <i>Escherichia coli</i> (↑CD) - <i>γ-proteobacteria</i> occasionally, in CD mucosal microbiota
Jia, W. et al. 2012	<ul style="list-style-type: none"> ▪ DNA 454 sequencing ▪ DGGE ▪ In-depth sequencing ▪ NGS 	<ul style="list-style-type: none"> ▪ Faecal samples 	<ul style="list-style-type: none"> ▪ 20 CD ▪ 14 UC ▪ 21 IBS ▪ 18 Controls 	N/A	N/A	England	<ul style="list-style-type: none"> - <i>B. wadsworthia</i>, - <i>Desulfovibrio piger</i> (↑CD/UC/IBS)
Vigsnaes, L. k. et al. 2012	<ul style="list-style-type: none"> ▪ DGGE 	<ul style="list-style-type: none"> ▪ Faecal samples 	<ul style="list-style-type: none"> ▪ 6 R-UC ▪ 6 UC ▪ 6 Controls 	N/A	N/A	Danemark	<ul style="list-style-type: none"> - <i>Lactobacillus</i> spp. and <i>Akkermansia</i> (<i>A. muciniphila</i>) (↓UC)
Michail S. et al., 2012	<ul style="list-style-type: none"> ▪ PCR of bacterial 16S rRNA ▪ Microarray hybridization 	<ul style="list-style-type: none"> ▪ Faecal samples 	<ul style="list-style-type: none"> ▪ 27 UC ▪ 26 Controls 	<ul style="list-style-type: none"> ▪ (17/10) ▪ (14/12) 	<ul style="list-style-type: none"> ▪ (10-17) ▪ (10-16) 	N/A	<ul style="list-style-type: none"> - <i>Clostridia</i> (↓UC) - <i>γ-proteobacteria</i> (↑UC)
Papa, E. et al., 2012	<ul style="list-style-type: none"> ▪ DNA 454 pyrosequencing ▪ Sanger sequencing 	<ul style="list-style-type: none"> ▪ Faecal samples 	<ul style="list-style-type: none"> ▪ 23 CD ▪ 43 UC ▪ 1 IBDU ▪ 24 Controls 	<ul style="list-style-type: none"> ▪ (13/10) ▪ (21/22) ▪ (1/0) ▪ (10/14) 	<ul style="list-style-type: none"> ▪ 15 (3-20) ▪ 14 (4-24) ▪ 10 (3-17) ▪ 14 	N/A	
Varela, E. et al. 2013	<ul style="list-style-type: none"> ▪ qPCR 	<ul style="list-style-type: none"> ▪ Faecal samples 	<ul style="list-style-type: none"> ▪ 116 R-UC ▪ 29 First degree relatives ▪ 31 Controls 	<ul style="list-style-type: none"> ▪ (55/61) ▪ (13/16) ▪ (17/14) 	<ul style="list-style-type: none"> ▪ 40 (32-46) ▪ 37 (27-54) ▪ 32 (23-41) 	Spain	<ul style="list-style-type: none"> - <i>F.prausnitzii</i> (↓UC/relatives/↑R-UC)

Table 1.c : Bacteria associated with inflammatory bowel disease analysed from both faecal and biopsy samples.

Author	Methods	Sample origins	Study Cohort	Patients characteristics at the time of sampling			Major findings	A/R-CD / UC = Active / Remission Crohn Disease / Ulcerative Colitis ; CD= Crohn disease, DGG = Denaturing gradient gel electrophoresis; FISH = Fluorescence in situ hybridization; IBD= Inflammatory bowel disease
				Gender (no. M/no. F)	Mean age (range)	Geo. Origin		
Willing, BP. et al., 2010	16S rRNA-sequencing	Faecal samples Mucosal samples	15 L1 12 L2 2 L3 15 UC 35 Controls	(7/8) (6/6) (0/2) (7/8) (10/25)	53 (20–70) 47 (20–70) 46 (42–49) 54 (30–69) 52 (30–70)	Swedish	- <i>Bacteroides</i> (↑IBD) - <i>Prevotella</i> (↓UC) - <i>Lactobacillus</i> , <i>R.gnavus</i> , <i>Veillonella</i> (↑CD) - <i>Faecalibacterium</i> (↓CD)	
Sokol, H. et al. 2008	qPCR of <i>F. prau</i>	Mucosal biopsy and Faecal samples	98 CD	N/A	N/A	N/A	- <i>F. prausnitzii</i> , <i>C. leptum</i> group (↓L1-CD)	
Chen, L. et al. 2014	16S rRNA 454-pyrosequencing	Biopsies different locations (Ileum, Cecum and Rectum) Faecal samples	26 CD 46 UC 21 Controls	(17/9) 30/11 (10/11)	30 (18-46) 42 (19-70) 28 (22-40)	China	- <i>Faecalibacterium</i> (↓CD/↑UC) - The abundance of the genus <i>Escherichia-Shigella</i> (↑CD/UC) - <i>Enterococcus</i> , (↑IBD).	
Vermeiren, J. et al. 2012	M-SHIME in vitro dynamic gut model DGGE of 16S rRNA	Luminal and mucosal biopsy samples Faecal samples	6 UC 6 Controls	N/A	41 (33-78) 27 (25-34)	N/A	- <i>Clostridium</i> cluster XIVa, <i>Roseburia</i> spp., members of the <i>C. coccoides/E. rectale</i> group, <i>F. prausnitzii</i> , a species of the <i>C. leptum</i> group, <i>Bacteroides/Prevotella</i> (↓UC)	
Wang, W. et al., 2014	16S rRNA-sequencing	Faecal samples Biopsy samples	25 CD 41 UC 21 Controls	(12/9) 30/11 N/A	30 (17-51) 43 (19-74) N/A	China	- <i>Lactobacillus</i> (↑IBD)	

e; IBDU = Inflammatory bowel disease unclassified; IBS = Irritable bowel syndrome; IC = Infectious Colitis; L1-CD= Ileum localized CD(Montreal classification); L1/L4 CD = Ileum localized CD with upper-gut involvement (Montreal classification); L2-CD= CD with primarily Colonic involvement (Montreal classification); L3-CD= Ileocolonic Crohn's Disease (Montreal classification); MCR= Multivariate curve resolution; N/A = Not available; NGS = Next generation sequencing; PCR= Polymerase Chain Reaction; PSC = Primary sclerosing cholangitis; qPCR = quantitative Polymerase Chain Reaction; RISA = Ribosomal intergenic spacer analysis; RT-PCR = Reverse Transcription - Polymerase Chain Reaction; SSCP = Single Strand Conformation Polymorphism; S.L.c = Self-limiting colitis (s.l.c); T-RFLP = Terminal restriction fragment length polymorphism; TTGE= Temporal temperature gradient gel electrophoresis; UC = Ulcerative Colitis; In.C = Indeterminate colitis; WGS = Whole Genome Shotgun

A.2. Gut bacterial changes reported in chronic rheumatic diseases patients

A total of twenty-one studies, enrolling 993 CRD patients analyzing the gut microbiota by 16S rRNA gene sequencing from faeces. Breban *et al.* have demonstrated that β -diversity analysis, which evaluates the shared diversity between different microbiomes in terms of various ecological distances, showed a microbiota composition significantly different between the RA, SpA and healthy subjects (HS) groups. Both SpA and RA patients differed from HSs as well as SpA from RA patients. This study showed also that α -diversity, which evaluates the species' richness and evenness within the microbiota, assessed by the number of observed species was significantly decreased in both SpA and RA patients, as compared with HSs^{62,63}. In ankylosing spondylitis (AS) patients, the diversity of the gut microbiome was similar to HSs at the genus level but was significantly higher in the controls at the species level⁶⁴.

Firmicutes phyla

Concerning the Firmicutes phylum, several bacteria from the *Lachnospiraceae* family, including *Ruminococcus* (*R. gnavus* sp.), *Dorea*, *Coprococcus* and *Blautia* genera are overabundant in SpA⁶². Increased amount of several *Blautia* and *Ruminococcus* could characterize HLA-B27⁺ siblings⁶². Likewise, inflamed ileal biopsies of SpA patients revealed an increase in the *Dialister* genus which could be a microbial marker of disease activity^{65,66}. In contrast, SpA patients seemed to present a decreased amount of *Roseburia* species⁶². Concerning RA patients, a fewer Firmicutes of the *Ruminococcaceae* family but an increase in *Lactobacillus* species and *Faklamia* have been observed^{62,67}. A study by Picchianti-Diamanti *et al.* characterized the gut microbiota of RA patients on different immunosuppressants treatment strategies (ETN, MTX, or ETN plus MTX) and compared it with that of treatment-naïve patients. The drop in Proteobacteria caused by ETN which in general are abundant in both intestinal and extra-intestinal inflammatory diseases⁶⁸. Moreover, a decrease in *Clostridiaceae* was observed upon ETN treatment which were previously found enriched in patients with RA and IBD-associated arthropathy⁶⁹. In patients treated with MTX, analysis revealed a significant decrease in Enterobacteriales⁶⁷.

Liu *et al.* reported that RA patients, compared to HSs, exhibited an increased bacterial diversity within *Lactobacillus* community with increase in *L.salivarius* and *L.iners*^{62,70,71} for instance. The analysis of faeces from RA patients have demonstrated the presence of a large cluster including Firmicutes bacteria belonging to *Lachnospiraceae* and *Clostridiaceae* (*Clostridium*) family, as well as small clusters containing strains from the *Lactobacillus* and

Ruminococcus genera^{70–73}. In the RA patients' gut, a decrease of bacteria from the *Veillonellaceae* family was observed^{72,74}. In contrast to SpA patients, PsA patients showed depletion in *Coprococcus*, *Ruminococcus*, *Clostridium* and *Pseudobutyrvibrio* compared to HSs^{62,74–76}. Finally, SpA patients exhibited a decreased fecal abundance of *F.prausnitzii* compared to HSs. This bacterium may be, at least in part, responsible for the pathogenesis of SpA^{66,77,78}.

Bacteroidetes phyla

There is a significant enrichment of the *Prevotellaceae* species, and more particularly of *Prevotella copri*, within the Bacteroidetes phylum, in intestinal microbiota of patients with new-onset RA, compared to chronic RA patients and HSs^{79–81}. This bacterium is relatively scarce in the general population. In addition, *Bacteroides* genera counts were lower in the same group, while being higher in SpA patients^{79,66,72}. However, *P. copri* decreased in the gut of RA patients along with disease chronicity⁸⁰. Breban *et al.* also demonstrated that SpA and RA patients have decreased populations of *Prevotellaceae* and *Paraprevotellaceae* genera compared to HSs⁶². However, in AS patients, *Prevotellaceae* are more abundant in terminal ileal biopsy samples⁷⁷. Furthermore, a quantitative metagenomics study has shown that the microbial communities in the AS cases were characterized by a higher abundance of *Prevotellaceae* genera (*Prevotella copri*) compared to HSs⁶⁴. Other bacteria from the Bacteroidetes phylum, such as *Porphyromonas*, were shown to be decreased in RA patients while being increased in terminal biopsies of AS patients^{77,82}.

Actinobacteria phyla

Regarding the Actinobacteria phylum, which is a low-abundant one, patients with RA or SpA had a higher amount of bacteria from the *Coriobacteriaceae* family and especially of the *Bifidobacterium* genus, including *B. bifidum* species than HSs^{62,66}. However, RA patients are also characterized by an increase of *Corynebacterium* species⁶². The metagenomic analysis and 16S sequencing have additionally brought into light the presence of the bacteria *Gordonibacter pamelaee*, *Eggerthella lenta* and *Collinsella* in RA patients^{63,64,72}. The latter could contribute to the increased permeability of the gut and enhanced production of pro-inflammatory cytokines⁶⁶. In SpA patients, an overabundance of *Collinsella*, *Rothia* and *Actinomyces* genera was reported^{63,64,76}.

Proteobacteria phyla

The Proteobacteria phylum is more abundant in RA patients than in HSs, concerning more specifically the *Klebsiella* and *Bilophila* genera from *Enterobacteriaceae*,

Author	Methods	Samples origins	Study Cohort	Patients characteristics at the time of sampling			Major findings
				Gender (no. M/no. F)	Mean age (range)	Geo. Origin	

Desulfovibrionaceae and *Succinivibrionaceae* families⁶². In SpA patients there is a decrease of *Citrobacter*, *Enterobacter* and *Erwinia* genera^{71,73,76}. The latter was particularly reduced in the HLA-A24 positive group of patients. In contrast, an overabundance of *Neisseria* genera was reported SpA patients⁶⁴.

Other phyla

Finally, other phyla as Synergistetes, Tenericutes, Fusobacteria and Verrucomicrobia were also retrieved to be increased or decreased in RA and SpA patients^{12,62,64,75,83}. (Fig 2)

In summary, among the available studies on CRD (n= 21), 33 bacterial species were reported in CRD, among those 17 were specifically reported in SpA studies while only 9 species reported in RA studies. Variations mainly concerned *Firmicutes*, *Bacteroidetes* and *Actinobacteria* phyla.

Table 2.a: Bacteria associated with chronic rheumatoid diseases analysed from faecal samples.

Breban, M. et al. 2017	16S rRNA gene sequencing	Faecal samples	86 SpA patients (74 SpA, 12 SpA+ IBD history) 28 RA 69 Controls	(41/46) (6/22) (26/43)	(35-63) (54-76) (27-63)	France	- <i>Klebsiella</i> , <i>Desulfovibrionaceae</i> (<i>bilophila</i>), <i>Succinivibrionaceae</i> , <i>Synergistetes</i> , <i>Tenericutes</i> (↑RA) - <i>Bifidobacterium</i> (↓RA/↑SpA) - <i>Paraprevotella</i> (↓SpA) - <i>Coriobacteriaceae</i> , <i>Ruminococcus</i> , <i>coprococcus</i> , <i>Dorea</i> , <i>Blautia</i> (↑SpA)
Picchianti - Diamanti, A. et al. 2018	NGS 16S Rrna	Faecal samples	11 RA treatment naïve patients 11 RA received MTX 10 RA received ETN 10 RA received ETN+MTX 10 Controls	(1/10) (2/9) (1/9) (2/8) N/A	56 63 60 65 N/A	Finland	- <i>Lactobacillaceae</i> , <i>Lactobacillus</i> (↑RA) - <i>Faecalibacterium</i> (↓RA) - <i>Cyanobacteria</i> phylum, <i>Nostocophycideae</i> , <i>Nostocales</i> group (↑RA-ETN) - <i>Deltaproteobacteria</i> (↑RA-ETN/UC) - <i>Clostridiaceae</i> upon (↓RA-ETN) - <i>Enterobacteriales</i> (↓RA- MTX) - <i>Eggerthella</i> (↑RA) - <i>Collinsella</i> (↑RA/SpA)
Chen, J. et al. 2016	16S rRNA sequencing	Faecal samples	40 RA patients, 32 Controls	(12/28) (6/26)	56 53	USA	- <i>Collinsella</i> , <i>Prevotella copri</i> (↑RA/SpA) - <i>Actinobacteria</i> , <i>Neisseria</i> , <i>rothia</i> , <i>Actinomyces</i> (↑SpA) - <i>Fusobacteria</i> , <i>Citrobacter</i> , <i>Verrucomicrobia</i> (↓SpA)
Wen, C. et al. 2017	Deep shotgun sequencing	Faecal samples	97 AS 114 Controls	(57/40) (72/42)	(14-71) (23-70)	China	- <i>Collinsella</i> , <i>Prevotella copri</i> (↑RA/SpA) - <i>Actinobacteria</i> , <i>Neisseria</i> , <i>rothia</i> , <i>Actinomyces</i> (↑SpA) - <i>Fusobacteria</i> , <i>Citrobacter</i> , <i>Verrucomicrobia</i> (↓SpA)
Stoll, ML. et al. 2018	16S rRNA sequencing Shotgun sequencing	Faecal samples	30 ERA 19 Controls 11 SpA 10 Controls	(19/11) (13/6) (4/7) (3/7)	14 (11-17) 14 (11-17) 52 (45-60) 47 (39-56)	USA	- <i>Bifidobacterium</i> , <i>Actinobacteria</i> , <i>Lachnospiracea</i> (↑RA/SpA) - <i>F. Prausnitzii</i> (↓RA/SpA)
Liu, X. et al. 2013	16S sequencing	Faecal samples	15 RA 15 Controls	(3/12) (5/10)	48 41	China	<i>Lactobacillus</i> genera (<i>Lactobacillus salivrius</i> , <i>L. Iners</i> , <i>L. ruminis</i>) (↑RA)
Maeda, Y. et al. EULAR 2012	RT-qPCR bacterial rRNA-targeted	Faecal samples	37 RA patients 59 Controls	(12/25) (6/53)	60 (49-71) 35 (25-45)	Japan	<i>L. ruminis</i> , <i>L.fermentum</i> , <i>L. reuteri</i> , <i>Enterococcus</i> (↑RA)
Scher, JU. et al. 2015	16S rRNA sequencing	Faecal samples	16 SpA 17 Controls	(7/9) (7/10)	47 43	USA	<i>Verrucomicrobia</i> , <i>Pseudobutyrvibrio</i> (↓SpA)
Manasson . et al., 2018	16S rRNA sequencing	Faecal samples	32 ReA 32 Controls	N/A	(18-55)	USA	- <i>Rikenellaceae</i> (↑SpA) - <i>Pseudomonas</i> (↑RA/SpA)
Stoll, ML. et al. 2015	16S rRNA sequencing	Faecal samples	12 recent onset ERA 21 Controls	ACR meeting Abstract			<i>F. Prausnitzii</i> (↓RA/SpA)
Scher, JU. et al. 2013	16S rRNA sequencing Shotgun sequencing	Faecal samples	44 NORA 26 CRA 16 PsA 28 Controls	(11/33) (3/23) (7/9) (7/21)	43 50 47 43	USA	<i>Prevotella copri</i> (↑RA/SpA)
Maeda, Y. et al. 2016	16S rRNA sequencing Shotgun sequencing	Faecal samples	17 RA 14 Controls	(3/14) (0/14)	64 (51-69) 53 (44-70)	Japan	<i>Prevotella copri</i> (↑RA/SpA)
Vahtovuori, J. et al. 2008	Flow cytometry 16S rRNA hybridization DNA-staining	Faecal samples	51 RA	(9/42)	57 (44- 70)	Finland	<i>Porphyromonas</i> (↓RA/SpA)
Stoll, M. L. et al. 2014	16S rRNA sequencing	Faecal samples	25 ERA 13 Controls	(14/11) (6/7)	13 (7-19) 13 (6-18)	USA	- <i>F. prausnitzii</i> (↓ERA) - <i>Clostridium leptum</i> group (↓AS)
Stebbing, S. et al.	DGGE	Faecal samples	15 AS 15 Controls	N/A	N/A	N/A	<i>Klebsiella pneumoniae</i> , <i>Bacteroides vulgatus</i>

Table 2.b: Bacteria associated with chronic rheumatoid diseases analysed from biopsy samples.

Table 2.c: Bacteria associated with chronic rheumatoid diseases analysed from faecal and other origin samples.

Author	Methods	Samples origins	Study Cohort	Patients characteristics at the time of sampling			Major findings
				Gender (no. M/no. F)	Mean age (range)	Geo. Origin	
Tito, RY. et al. 2017	16S rRNA sequencing	Biopsy samples ileal and colonic	27 SpA 15 Controls	(13/14) N/A	(10-50) N/A	Belgium	<i>Dialister</i> (↑SpA)
Costello, ME. et al. 2016	16S rRNA sequencing	Intestinal biopsy	10 HLA-B27+ 85 HLA-B27-	ACR meeting Abstract			<i>Veillonellaceae</i> (↓RA/SpA)
Costello, ME. et al. 2013	16S sequencing	Terminal ileal Biopsy	N/A AS N/A CD N/A Controls	ACR meeting Abstract			- <i>Porphyromonas</i> , <i>F. Prausnitzii</i> (↓RA/SpA) - <i>Ruminococc</i> (↑SpA)

ACR = American College of Rheumatology ; AS = Ankylosing spondylitis; CRA = Chronic, treated rheumatoid arthritis; DMARD = Disease-modifying antirheumatic drug; ETN = Etanercept; EULAR = European League Against Rheumatism; ERA = Entesitis-related arthritis; IBD = Inflammatory bowel disease; NORA = new onset untreated rheumatoid arthritis; MTX = Methotrexate; PsA = Psoriasis arthritis; RA = Rheumatoid arthritis; ReA = Reactive arthritis; SpA = Spondylo-arthritis; ; UA = Undifferentiated Arthritis; UC = Ulcerative Colitis.

Author	Methods	Samples origins	Study Cohort	Patients characteristics at the time of sampling			Major findings
				Gender (no. M/no.F)	Mean age (range)	Geo. Origin	
Zhang, X. et al. 2015	Metagenomic sequencing	Faecal samples Dental samples Salivary	115 RA (21 DMARD) 97 Controls	(31/84) (28/69)	50 (27-74) 43 (19-68)	China	- <i>Collinsella</i> , <i>Eggerthella</i> , <i>Gordonibacter pamelaeeae</i> , <i>Clostridium</i> , <i>Lachnospiraceae</i> (↑RA) - <i>Veillonellaceae</i> (↓RA/SpA)
Benham, H. et al. 2016	16S rRNA sequencing Tongue and faecal swabs	Tongue and faecal swabs	116 RA 63 First-Degree Relatives 43 Controls	ACR meeting Abstract			- <i>Enterococcus</i> (↑RA) - <i>Pseudomonas</i> (↑RA/SpA)

A.3.
Diff
eren
ces
bet
wee
n
IBD

and CRD gut microbiota

Three studies enrolling a total of 554 patients, directly compared 356 IBD patients without known arthropathy, and a total of 132 IBD with joint extra-intestinal-manifestation (EIM) patients were analysed (Table 3). One study indirectly compared three cohorts of patients,

SpA patients without IBD history (n=74) as well as SpA patients with an IBD history (n=12), and RA patients (n=28) compared with HCs (n= 69) (Table 3)⁶².

Firmicutes phyla

Amongst included studies, some pointed out important differences, including variable amount of several Firmicutes genera. For instance, the overabundance of *Veillonella* observed in CD patients contrasted with its paucity in CRD (RA, SpA) patients. Conversely, the *Eubacterium*, *Clostridium*, *Ruminococcus* and *Coprococcus* genera, that were increased in CRD (RA, SpA) patients, were decreased in patients with CD^{15,12,13, 19,72,77,79,80,82,83,84}. Variation of the *Ruminococcus* genus is the most surprising since a paradoxical overabundance, especially of *R. gnavus*, has been reported in SpA patients. This increased abundance correlated positively with SpA activity whatever patients IBD history, even though IBD was inactive at the time of sampling in most of them^{12,62}. In IBD, *R. gnavus* was mostly associated with the gut mucosa, which conferred to this mucolytic bacteria a possible role in the triggering or maintenance of inflammation^{12,32}. Whether its lonely increase could be linked to specific genetic predispositions to SpA warrants more investigation. As for the *Dialister* genera, belonging to the same bacterial family, an increased number of sequences was observed in SpA groups whereas a decrease was found in CD patients⁶². In UC patients with a joint EIM, the *Staphylococcus* genus was found more frequently in stool cultures⁸⁵.

Bacteroidetes phyla

Variations in Bacteroidetes phylum concerned mainly two genera: *Bacteroides*, which was in increased amount in SpA patients and in reduced amount in RA and IBD groups and *Prevotella* which showed a high abundance in CRD (RA and SpA) patients and was lowered in UC patients^{15,20, 63,64,66,79,80 84, 86}.

Proteobacteria phyla

In the Proteobacteria phylum, the genus *Bilophila* was overabundant in RA and SpA patients while being found in reduced amounts in CD patients^{12,51,62,87,88}. Dorofeyev *et al.* showed a significant abundance of *Enterobacter*, *Klebsiella* and *Proteus* genera in stools cultures from UC patients with a joint EIM, compared to HSs and UC patients without EIM⁸⁵. In contrast, in UC a decreased amount of *Neisseria* was observed^{12,13,15,47–49,31,35,39,50}. However, metagenomics studies of gut microbiome in patients with enteropathic arthritis are still lacking. Using qPCR, a relative overabundance of the *Enterobacteriaceae* family, concomitant to a reduction of the *Clostridia* group *XIVa* cluster, was reported in the gut microbiota in IBD patients with joint manifestations. As a whole, the *Enterobacteriaceae*

family seemed to be increased in the gut of IBD patients and this tendency is even more pronounced in those with arthropathy⁸⁹.

Actinobacteria phyla

Concerning the Actinobacteria phylum, an overabundance of *Gordonibacter pamelaee*, *Eggerthella lenta* and *Collinsella* was observed in RA patients^{63,64, 72,90}. However, an increase of *Micrococcus* genera was also characterized in MAM UC patients⁴⁶. In SpA patients, an overabundance of *Collinsella*, *Rothia* and *Actinomyces* genera was reported^{63,64,76}.

Other phyla

Finally, the *Fusobacterium* phylum is more abundant in CD patients and less abundant in SpA patients⁶². In contrast, amounts of the *Tenericutes* phylum are increased in SpA patients^{10,62,64}.

Taken together, when considering all available studies (n=80), 40 bacterial species were reported only in IBD patients and 33 bacterial species were reported only in CRD subjects (Fig 2). Main variations were mostly observed in the *Firmicutes* phylum.

B. Similarities regarding bacterial microbiome in IBD and CRD

When comparing studies on IBD patients without known CRD *versus* studies on CRD patients without known IBD, we first observed that some dysbiotic changes share similarities between chronic IBD and chronic joint diseases, among which a lower microbial diversity and a diminished abundance of the Firmicutes phylum.

Firmicutes phyla

Amongst the Firmicutes genera, a common decreased amount was described for *Faecalibacterium* and *Roseburia* species in both IBD subtypes (CD and UC), as well as in SpA and RA patients^{12,13,15,20,50,62,84}. A few studies using bacterial culture, in addition to recent molecular methods, have demonstrated an increase amount of *Lactobacillus* and *Enterococcus* in the faecal microbiota of IBD patients especially those with CD and RA patients, although others demonstrated a reduction of *Lactobacillus* in CD patients^{7,12,15, 13, 47, 30,48,49,52,70,72,84}. An overabundance of *staphylococcus* was observed in UC patients with arthritis when compared with patients without EIM and healthy population.

Proteobacteria phyla

In the Proteobacteria phylum, an overabundance of several genera was observed, such as *Klebsiella* and *Proteus* in all UC patients with arthritis. These facultative microbiota were significantly higher in these patients than in the HSs and UC patients without EIM^{46,62,85,91,92}. An increase of *Pseudomonas* was recently showed by Walujkar *et al* in the MAM of UC patients as compared to the same patients during remission stage⁴⁶, as well as showed by Manasson *et al.* and Benham *et al* in patients with SpA or RA.^{73,76}

Actinobacteria phyla

Concerning the Actinobacteria phylum, an overabundance of *Bifidobacterium* was reported in SpA patients, especially those with enthesitis-related arthritis (ERA), and in IBD patients notably in patients with CD^{12,13,15,19,47 50,62,64,66,72, 82, 83}.

Other phyla

Finally, a common decrease of Verrucomicrobia and Fusobacteria belonging species was reported in both CD and UC patients compared to non-IBD controls^{10,12,31,32,35,59–61}, and in RA and SpA patients^{12,62,64,75,83}.

In summary, variations of species belonging to Firmicutes, Proteobacteria, Actinobacteria, Verrucomicrobia and Fusobacteria phyla represent the main common trait between IBD and CRD gut microbiota. A figure depicting similarities and differences observed in bacterial species amounts in biopsy and faeces from IBD and CRD patients is proposed (Fig 2).

Table 3: Bacteria associated with inflammatory bowel disease and chronic rheumatic diseases.

Author	Methods	Samples origins	Population studied	Patients characteristics at the time of sampling			Major findings
				Gender (no. M/no. F)	Mean age (range)	Geo. Origin	
Muniz-Pedrogo, D.A. et al. 2018	16S rRNA sequencing	Faecal samples	<ul style="list-style-type: none"> ▪ 25 IBD-A ▪ 66 IBD-N, ▪ 25 RA ▪ 64 Controls 	<ul style="list-style-type: none"> ▪ (11/14) ▪ (26/40) ▪ (10/15) ▪ (27/37) 	<ul style="list-style-type: none"> ▪ 49 ▪ 49 ▪ 52 ▪ 50 	N/A	<i>Escherichia</i> (↑IBD)
Dorofeyev, A.E. et al. 2009	Culture dependent techniques	<ul style="list-style-type: none"> ▪ Biopsies samples ▪ Faecal samples 	<ul style="list-style-type: none"> ▪ 131 Distal UC ▪ 102 Left-sided UC ▪ 86 Pancolitis ▪ 95 UC+ joint EIM 	<ul style="list-style-type: none"> ▪ (147/172) ▪ Idem ▪ Idem ▪ N/A 	<ul style="list-style-type: none"> ▪ (40-47) ▪ Idem ▪ Idem ▪ N/A 	N/A	<ul style="list-style-type: none"> - <i>Bifidobacteria</i>, <i>lactobacilli</i> and <i>Escherichia coli</i> (↓ UC) - Facultative flora (↑UC) - <i>Staphylococcus</i>, <i>Klebsiella</i> and <i>Proteus</i> were found more often in stool cultures (↑UC+ joint EIM)
Kabeerdoss, J. et al. 2014	16S rRNA sequencing	Faecal samples	<ul style="list-style-type: none"> ▪ 12 IBD + Arthropathy ▪ 12 IBD 	N/A	N/A	N/A	<ul style="list-style-type: none"> - <i>Enterococcaceae</i>, <i>Enterococcus</i> and <i>Enterococcus faecium</i> (↑IBD+ arthropathy)

EIM: Extra-intestinal manifestation; IBD= Inflammatory bowel disease; IBD-A/N = IBD-associated/ without arthropathy; N/A = Not available; RA = Rheumatoid arthritis.

Conclusion and perspectives

To our knowledge, this is the first systematic review concerning evidence regarding the gut microbiota in IBD and CRD patients. Our analysis highlights the general finding that microbiota favouring proteolytic-fuelled fermentation and lactic acid-producing bacteria, are increased in both CRD and IBD inflammatory conditions while those producing butyrate are generally decrease in both diseases. Secondly, variations of gut microbiota composition in IBD patients mainly concerned *Firmicutes*, *Proteobacteria* and *Bacteroidetes*. Within the Firmicute phylum variations of species as *Roseburia*, *coprococcus*, *F. prausnitzii* and *Streptococcus* genera, was observed either in the mucosal-associated microbiota (MAM) of CD patients or UC patients. In terms of Proteobacteria phylum published data display a quantitative alteration in IBD CD and UC patients compared to control groups especially of *Escherichia*, *Shigella*, *Bilophila*, *Desulfovibrio*, *Neisseria*, *Stenotrophomonas*, *Ochrobactrum* and *Achromobacter* genera. Concerning the Bacteroidetes, variations of *Cloacibacterium*, *Prevotella*, *Butyricimonas*, *Parabacteroides*, *Elizabethkingia* genera and *Odoribacteraceae* family in IBD, CD and UC patients are observed.

However, in CRD patients, variations are mainly observed in Firmicutes, Bacteroidetes and Actinobacteria phyla. Alterations of gut microbiota observed in the Firmicutes phyla included *Ruminococcus* (*R. gnavus* sp.), *Dorea*, *Coprococcus*, *Blautia*, and *Dialister* genus in RA and SpA patients. In addition alterations of *Roseburia*, *Lactobacillus*, *Faklamia*, *Staphylococcus*, *Clostridium*, *Pseudobutyrvibrio*, *F.prausnitzii* species and *Veillonellaceae* family was observed in patients compared to healthy subjects. There is a significant variations of species within the Bacteroidetes phylum, particularly of, *Bacteroides*, *Prevotellaceae* (*P.copri*) *Paraprevotellaceae* and *Porphyromonas* genera in RA and SpA patients compared to HSs. Regarding the Actinobacteria phylum, which is a low-abundant one, in patients with RA or SpA variations of the *Bifidobacterium* genus, including among others *B. bifidum* species, *Gordonibacter pamelaee*, *Eggerthella lenta*, *Collinsella*, *Rothia* and *Actinomyces* genera was reported compared to control groups.

Another major finding of this study, is the reduction of bacterial diversity, observed in both CRD and IBD and the presence of common bacterial phyla changes. We can mention an increased abundance in *Lactobacillus*, *Enterococcus*, *Staphylococcus*, *Bifidobacterium*,

Klebsiella, *Pseudomonas* and *Proteus* genera in both CRD and IBD, whereas *Faecalibacterium*, *Roseburia* genera and Verrucomicrobia, Fusobacteria phyla are decreased in both diseases.

Interestingly, experimental studies have confirmed the role of *Faecalibacterium* in immune controlled in both type of affections. First, Hablot and colleagues suggested that experimental Dextran Sulfate Sodium (DSS) induced colitis could altered the gut microbiota of mice with arthritis compared to mice with colitis alone and thus could delayed the appearance of “pro-arthritis” bacteria⁹³. This delay is associated with a difference of microbiota composition between mice with arthritis and colitis and mice with colitis only. Members of the Firmicutes phylum are mainly affected; *Lactobacillus* genus and Clostridiales order are more present in mice with arthritis and colitis compared to mice with only colitis. Several studies showed that species from *Lactobacillus* are beneficial in DSS-induced colitis^{94,95}. Thereby, *Lactobacillus* *sp* increase in arthritis + colitis group might play a role in the subclinical improvement as observed by the decrease in fecal lipocalin-2 level. A difference of the fecal microbiota composition is also observed between arthritis and arthritis + colitis groups. At arthritis and colitis onset, *Lactobacillaceae*, and notably *Lactobacillus R.gnavus* and S24_7 species belonging to Bacteroidales are more present in mice with arthritis and colitis compared to arthritis group. Interestingly, these groups of bacteria had been shown to be more present in mice with higher susceptibility to arthritis development^{93,96}.

Viladomiu and colleagues recently identified an enrichment of IgA-coated *E. coli* in CD-SpA with an adherent-invasive *E. coli* (AIEC) pathotype. Experimental models highlight two features of the host-pathogen interaction that must be considered to understand the specificity of pathogenetic mechanisms, namely, host susceptibility and strain variability⁹⁷. CD SpA-derived AIEC protect against acute injury and death from DSS induced colitis in WT mice. Resident microbiota, including AIEC, induce colonic ROR γ t/Foxp3⁺ CD4⁺ T cells, which play an important role in restraining inflammatory colitis⁹⁸. Consistent with a higher Enterobacteriaceae in 6-month-old infants correlated with better nutritional status⁹⁹. Thus, in situations of nutritional sufficiency or immunocompetence, the response to Enterobacteriaceae may have coevolved to protect the host; however, persistent nutritional deficiency⁹⁹ or genetic susceptibility (modeled in IL-10-deficient and K/BxN mice) evokes maladaptive responses, which, in turn, promote more severe inflammatory Th17 disease. Likewise, this data link the shared genetic susceptibility in the IL23R locus in both CD and SpA¹⁰⁰ with increased systemic *E. coli* sero-reactivity and Th17 inflammatory cytokines.

These results highlight the functional implication of IgA-coated *E. coli* enriched in CD-SpA and identify a Th17 immunophenotype characteristic of this EIM. This mechanistic link between intestinal microbiota and systemic inflammation may underlie the clinical efficacy of sulfasalazine in peripheral joint symptoms ¹⁰¹. While anti-TNF α therapy improves axial symptoms in patients with active CD ¹⁰², this data also highlights the overactivation of the IL-23/IL-17 pathway in CD patients with peripheral symptoms.

This review displays several methodological and theoretical limitations. First, heterogeneity of studied populations (in terms of age, gender and origins) and microbiota analysing methodology deeply impact gut microbiota picture. The purpose of our study, i.e. to identify similarities and differences between gut microbiome in IBD and in CRD patients, is challenging considering also the relatively small number of studies in CRD compared to IBD. Indeed, first studies analysing gut microbiota in IBD were published in 2005, whereas gut microbiota in CRD has been explored a decade later. Since the first studies, more than 4000 IBD patients have been analysed whereas only 300 for CRD.

Secondly, inconsistencies may exist among the findings from available studies due to the heterogeneity in sample size, biopsy location, local inflammation and types of samples (biopsies vs stool) that may influence the microbiota composition. Furthermore, complexity of the microbiota must be put into perspective along with current technological limitations (analysing DNAs encoding 16S RNA gene still provides only an incomplete picture of bacterial populations and some study presented here used culture dependent determination methodology).

Despite these considerations and in an effort to synthesize already published data we provide detailed tables by clinical condition and sample type as well as a figure providing an overview of the data available (figure 2).

Finally, information on the possible concomitant arthritis and IBD was not provided in some of the 80 included studies involving IBD and CRD patients. It is thus impossible to rule out the presence of subclinical joint-gut inflammation in these patients.

We can mention also the absence of healthy controls groups in certain studies or the incomplete description of clinical situation of patients (for instance patients with IBD history without information on disease activity or medication or faeces consistency score at time of sampling) that could influence gut microbiota ¹⁰³.

Bacteria are not the only component of gut microbiota, fungi and virus may have a role in both diseases' initiation or severity. Bacteria and fungi could compete for the same substrates or produce synergistically metabolites that could affect host immunity and metabolism. Only a few studies on intestinal fungal microbiota and its relationship with IBD have been conducted. Much evidence has shown that fungi and their communities may be involved in the pathogenesis of IBD, especially CD¹⁰⁴. To date fungal microbiota implication in CRD has not been explored.

The enteric virome is known to be altered in patients with IBD, with specific changes assessed between UC and CD. Enormous numbers of candidate viruses have been thought to be triggering factor of arthritis, particularly of RA, but most of the evidence implicating viruses in the pathogenesis of CRD are circumstantial and inconclusive. Tantalizing observations have often been based on *in vitro* or animal studies, case reports, or studies with small sample sizes, cross-sectional designs, or without control groups.

The description of the viral, fungal, bacterial metagenomes in patients suffering from IBD and or CRD shall provide a better understanding of the interactions between the microbiome and host immunity within the joint-gut axis. The identification of specific species in well-defined categories of patients can provide valuable information, which can be translated into prognostic, diagnostic or therapeutic tools that are critically lacking for these diseases. Furthermore, such studies hold great promise for the development of future strategies aiming at early detection of relapse and at controlling/manipulating the microbiome to reduce the burden of these ailments.

In conclusion a total of 80 studies investigated bacterial microbiome in patients with IBD and/or CRD. These studies showed that some bacterial taxons seem specifically imbalanced in IBD (n=40) and CRD (n=33), while increased abundance in Firmicutes genera *Lactobacillus* and *Staphylococcus*, Actinobacteria *Bifidobacterium*, and Proteobacteria genera such as *Pseudomonas*, *Klebsiella* and *Proteus*. Whereas, Firmicutes phyla *Faecalibacterium*, *Roseburia* genera and Verrucomicrobia phylum are decreased in both CRD and IBD. Large and well-designed prospective studies are eagerly awaited to further elucidate the role of gut microbiome in promoting pathological inflammation within joint-gut axis.

Figures



Figure 1: Flow-diagram of identified studies.

Figure 2 Salem et al.




Figure 2 : Similarities and differences regarding gut bacteria between IBD and CRD patients.

Genera colours represents Phylum: **Bacteroidetes**, **Firmicutes**, **Actinobacteria**, **Proteobacteria**, **Fusobacteria**, **Tenericutes**, **Synergistetes**.

↑/↓= increase / decrease in patients with IBD or CRD

References

1. Klingberg E, Strid H, Ståhl A, et al. A longitudinal study of fecal calprotectin and the development of inflammatory bowel disease in ankylosing spondylitis. *Arthritis Res Ther*; 19. Epub ahead of print 2017. DOI: 10.1186/s13075-017-1223-2.
2. Peyrin-Biroulet L, Ferrante M, Magro F, et al. Results from the 2nd Scientific Workshop of the ECCO. I: Impact of mucosal healing on the course of inflammatory bowel disease. *J Crohns Colitis* 2011; 5: 477–483.
3. Lees CW, Barrett JC, Parkes M, et al. New IBD genetics: common pathways with other diseases. *Gut* 2011; 60: 1739–1753.
4. Jostins L, Ripke S, Weersma RK, et al. Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* 2012; 491: 119–124.
5. Atar D, Birkeland KI, Uhlig T. ‘Treat to target’: moving targets from hypertension, hyperlipidaemia and diabetes to rheumatoid arthritis. *Ann Rheum Dis* 2010; 69: 629–630.
6. PRISMA, <http://prisma-statement.org/prismastatement/Checklist.aspx> (accessed 29 June 2018).
7. Seksik P, Lepage P, de la Cochetiere M-F, et al. Search for Localized Dysbiosis in Crohn’s Disease Ulcerations by Temporal Temperature Gradient Gel Electrophoresis of 16S rRNA. *J Clin Microbiol* 2005; 43: 4654–4658.
8. Ott SJ, Musfeldt M, Wenderoth DF, et al. Reduction in diversity of the colonic mucosa associated bacterial microflora in patients with active inflammatory bowel disease. *Gut* 2004; 53: 685–693.
9. Scanlan PD, Shanahan F, O’Mahony C, et al. Culture-Independent Analyses of Temporal Variation of the Dominant Fecal Microbiota and Targeted Bacterial Subgroups in Crohn’s Disease. *J Clin Microbiol* 2006; 44: 3980–3988.
10. Hourigan SK, Chen LA, Grigoryan Z, et al. Microbiome changes associated with sustained eradication of *Clostridium difficile* after single faecal microbiota transplantation in children with and without inflammatory bowel disease. *Aliment Pharmacol Ther* 2015; 42: 741–752.
11. Duvallet C, Gibbons SM, Gurry T, et al. Meta-analysis of gut microbiome studies identifies disease-specific and shared responses. *Nat Commun*; 8. Epub ahead of print December 2017. DOI: 10.1038/s41467-017-01973-8.
12. Willing BP, Dicksved J, Halfvarson J, et al. A Pyrosequencing Study in Twins Shows That Gastrointestinal Microbial Profiles Vary With Inflammatory Bowel Disease Phenotypes. *Gastroenterology* 2010; 139: 1844-1854.e1.
13. Gevers D, Kugathasan S, Denson LA, et al. The Treatment-Naive Microbiome in New-Onset Crohn’s Disease. *Cell Host Microbe* 2014; 15: 382–392.
14. Hall AB, Yassour M, Sauk J, et al. A novel Ruminococcus gnavus clade enriched in inflammatory bowel disease patients. *Genome Med*; 9. Epub ahead of print December 2017. DOI: 10.1186/s13073-017-0490-5.
15. Morgan XC, Tickle TL, Sokol H, et al. Dysfunction of the intestinal microbiome in inflammatory bowel disease and treatment. *Genome Biol* 2012; 13: R79.
16. Ananthakrishnan AN, Luo C, Yajnik V, et al. Gut Microbiome Function Predicts Response to

- Anti-integrin Biologic Therapy in Inflammatory Bowel Diseases. *Cell Host Microbe* 2017; 21: 603-610.e3.
17. Kaakoush NO, Day AS, Huinao KD, et al. Microbial Dysbiosis in Pediatric Patients with Crohn's Disease. *J Clin Microbiol* 2012; 50: 3258–3266.
 18. Gophna U, Sommerfeld K, Gophna S, et al. Differences between Tissue-Associated Intestinal Microfloras of Patients with Crohn's Disease and Ulcerative Colitis. *J Clin Microbiol* 2006; 44: 4136–4141.
 19. Frank DN, Amand ALS, Feldman RA, et al. Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases. *Proc Natl Acad Sci* 2007; 104: 13780–13785.
 20. Aomatsu T, Imaeda H, Fujimoto T, et al. Terminal Restriction Fragment Length Polymorphism Analysis of the Gut Microbiota Profiles of Pediatric Patients with Inflammatory Bowel Disease. *Digestion* 2012; 86: 129–135.
 21. Machiels K, Joossens M, Sabino J, et al. A decrease of the butyrate-producing species *Roseburia hominis* and *Faecalibacterium prausnitzii* defines dysbiosis in patients with ulcerative colitis. *Gut* 2014; 63: 1275–1283.
 22. Willing B, Halfvarson J, Dicksved J, et al. Twin studies reveal specific imbalances in the mucosa-associated microbiota of patients with ileal Crohn's disease. *Inflamm Bowel Dis*; 15: 653–660.
 23. Sokol H, Pigneur B, Watterlot L, et al. *Faecalibacterium prausnitzii* is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients. *Proc Natl Acad Sci* 2008; 105: 16731–16736.
 24. Pascal V, Pozuelo M, Borrueal N, et al. A microbial signature for Crohn's disease. *Gut* 2017; 66: 813–822.
 25. Rehman A, Rausch P, Wang J, et al. Geographical patterns of the standing and active human gut microbiome in health and IBD. *Gut* 2016; 65: 238–248.
 26. Duboc H, Rajca S, Rainteau D, et al. Connecting dysbiosis, bile-acid dysmetabolism and gut inflammation in inflammatory bowel diseases. *Gut* 2013; 62: 531–539.
 27. Fujimoto T, Imaeda H, Takahashi K, et al. Decreased abundance of *Faecalibacterium prausnitzii* in the gut microbiota of Crohn's disease. *J Gastroenterol Hepatol*; 28: 613–619.
 28. Swidsinski A, Loening-Baucke V, Vaneechoutte M, et al. Active Crohn's disease and ulcerative colitis can be specifically diagnosed and monitored based on the biostructure of the fecal flora. *Inflamm Bowel Dis* 2008; 14: 147–161.
 29. Sokol H, Seksik P, Furet JP, et al. Low counts of *Faecalibacterium prausnitzii* in colitis microbiota. *Inflamm Bowel Dis*; 15: 1183–1189.
 30. Sabino J, Vieira-Silva S, Machiels K, et al. Primary sclerosing cholangitis is characterised by intestinal dysbiosis independent from IBD. *Gut* 2016; 65: 1681–1689.
 31. Bajer L, Kverka M, Kostovcik M, et al. Distinct gut microbiota profiles in patients with primary sclerosing cholangitis and ulcerative colitis. *World J Gastroenterol* 2017; 23: 4548–4558.
 32. Png CW, Lindén SK, Gilshenan KS, et al. Mucolytic Bacteria With Increased Prevalence in IBD Mucosa Augment *In Vitro* Utilization of Mucin by Other Bacteria. *Am J Gastroenterol* 2010; 105: 2420–2428.

33. Borody TJ, Paramsothy S, Agrawal G. Fecal Microbiota Transplantation: Indications, Methods, Evidence, and Future Directions. *Curr Gastroenterol Rep*; 15. Epub ahead of print 2013. DOI: 10.1007/s11894-013-0337-1.
34. Eeckhaut V, Machiels K, Perrier C, et al. Butyricicoccus pullicaecorum in inflammatory bowel disease. *Gut* 2012; gutjnl-2012-303611.
35. Hirano A, Umeno J, Okamoto Y, et al. Comparison of the microbial community structure between inflamed and non-inflamed sites in patients with ulcerative colitis. *J Gastroenterol Hepatol*; 0. Epub ahead of print 02 2018. DOI: 10.1111/jgh.14129.
36. Vermeiren J, Van den Abbeele P, Laukens D, et al. Decreased colonization of fecal *Clostridium coccoides*/*Eubacterium rectale* species from ulcerative colitis patients in an in vitro dynamic gut model with mucin environment. *FEMS Microbiol Ecol* 2012; 79: 685–696.
37. Knoll RL, Forslund K, Kultima JR, et al. Gut microbiota differs between children with Inflammatory Bowel Disease and healthy siblings in taxonomic and functional composition: a metagenomic analysis. *Am J Physiol-Gastrointest Liver Physiol* 2016; 312: G327–G339.
38. Andoh A, Imaeda H, Aomatsu T, et al. Comparison of the fecal microbiota profiles between ulcerative colitis and Crohn's disease using terminal restriction fragment length polymorphism analysis. *J Gastroenterol* 2011; 46: 479–486.
39. Said HS, Suda W, Nakagome S, et al. Dysbiosis of Salivary Microbiota in Inflammatory Bowel Disease and Its Association With Oral Immunological Biomarkers. *DNA Res* 2014; 21: 15–25.
40. Allen TD, Lawson PA, Collins MD, et al. *Cloacibacterium normanense* gen. nov., sp. nov., a novel bacterium in the family Flavobacteriaceae isolated from municipal wastewater. *Int J Syst Evol Microbiol* 2006; 56: 1311–1316.
41. Chiodini RJ, Dowd SE, Chamberlin WM, et al. Microbial Population Differentials between Mucosal and Submucosal Intestinal Tissues in Advanced Crohn's Disease of the Ileum. *PLOS ONE* 2015; 10: e0134382.
42. Sokol H, Lepage P, Seksik P, et al. Temperature Gradient Gel Electrophoresis of Fecal 16S rRNA Reveals Active *Escherichia coli* in the Microbiota of Patients with Ulcerative Colitis. *J Clin Microbiol* 2006; 44: 3172–3177.
43. Sokol H, Seksik P, Rigottier-Gois L, et al. Specificities of the Fecal Microbiota in Inflammatory Bowel Disease. *Inflamm Bowel Dis* 2006; 12: 106–111.
44. Swidsinski A, Ladhoff A, Pernthaler A, et al. Mucosal flora in inflammatory bowel disease. *Gastroenterology* 2002; 122: 44–54.
45. Swidsinski A, Weber J, Loening-Baucke V, et al. Spatial Organization and Composition of the Mucosal Flora in Patients with Inflammatory Bowel Disease. *J Clin Microbiol* 2005; 43: 3380–3389.
46. Walujkar SA, Kumbhare SV, Marathe NP, et al. Molecular profiling of mucosal tissue associated microbiota in patients manifesting acute exacerbations and remission stage of ulcerative colitis. *World J Microbiol Biotechnol* 2018; 34: 76.
47. Wang W, Chen L, Zhou R, et al. Increased Proportions of Bifidobacterium and the Lactobacillus Group and Loss of Butyrate-Producing Bacteria in Inflammatory Bowel Disease. *J Clin Microbiol* 2014; 52: 398–406.
48. GIAFFER MH, HOLDSWORTH CD, DUERDEN BI. The assessment of faecal flora in patients with inflammatory bowel disease by a simplified bacteriological technique. *J Med Microbiol*

1991; 35: 238–243.

49. Seksik P, Rigottier-Gois L, Gramet G, et al. Alterations of the dominant faecal bacterial groups in patients with Crohn's disease of the colon. *Gut* 2003; 52: 237–242.
50. Schwartz A, Jacobi M, Frick J-S, et al. Microbiota in Pediatric Inflammatory Bowel Disease. *J Pediatr* 2010; 157: 240-244.e1.
51. Thorkildsen LT, Nwosu FC, Avershina E, et al. Dominant Fecal Microbiota in Newly Diagnosed Untreated Inflammatory Bowel Disease Patients. *Gastroenterology Research and Practice*. Epub ahead of print 2013. DOI: 10.1155/2013/636785.
52. Martinez-Medina M, Aldeguer X, Gonzalez-Huix F, et al. Abnormal microbiota composition in the ileocolonic mucosa of Crohn's disease patients as revealed by polymerase chain reaction-denaturing gradient gel electrophoresis. *Inflamm Bowel Dis* 2006; 12: 1136–1145.
53. Kotlowski R, Bernstein CN, Sepehri S, et al. High prevalence of Escherichia coli belonging to the B2+D phylogenetic group in inflammatory bowel disease. *Gut* 2007; 56: 669–675.
54. Sokol H, Lepage P, Seksik P, et al. Molecular comparison of dominant microbiota associated with injured versus healthy mucosa in ulcerative colitis. *Gut* 2007; 56: 152–154.
55. Zhang M, Liu B, Zhang Y, et al. Structural Shifts of Mucosa-Associated Lactobacilli and Clostridium leptum Subgroup in Patients with Ulcerative Colitis. *J Clin Microbiol* 2007; 45: 496–500.
56. Mylonaki M, Rayment NB, Rampton DS, et al. Molecular Characterization of Rectal Mucosa-Associated Bacterial Flora in Inflammatory Bowel Disease. *Inflamm Bowel Dis* 2005; 11: 481–487.
57. Earley H, Lennon G, Balfe A, et al. A Preliminary Study Examining the Binding Capacity of Akkermansia muciniphila and Desulfovibrio spp., to Colonic Mucin in Health and Ulcerative Colitis. *PLOS ONE* 2015; 10: e0135280.
58. Jia W, Whitehead RN, Griffiths L, et al. Diversity and distribution of sulphate-reducing bacteria in human faeces from healthy subjects and patients with inflammatory bowel disease. *FEMS Immunol Med Microbiol* 2012; 65: 55–68.
59. Vignæs L k., Brynskov J, Steenholdt C, et al. Gram-negative bacteria account for main differences between faecal microbiota from patients with ulcerative colitis and healthy controls. *Benef Microbes* 2012; 3: 287–297.
60. Michail S, Durbin M, Turner D, et al. Alterations in the gut microbiome of children with severe ulcerative colitis. *Inflamm Bowel Dis* 2012; 18: 1799–1808.
61. Papa E, Docktor M, Smillie C, et al. Non-invasive mapping of the gastrointestinal microbiota identifies children with inflammatory bowel disease. *PloS One* 2012; 7: e39242.
62. Breban M, Tap J, Leboime A, et al. Faecal microbiota study reveals specific dysbiosis in spondyloarthritis. *Ann Rheum Dis* 2017; 76: 1614–1622.
63. Chen J, Wright K, Davis JM, et al. An expansion of rare lineage intestinal microbes characterizes rheumatoid arthritis. *Genome Med* 2016; 8: 43.
64. Wen C, Zheng Z, Shao T, et al. Quantitative metagenomics reveals unique gut microbiome biomarkers in ankylosing spondylitis. *Genome Biol* 2017; 18: 142.
65. Tito RY, Cypers H, Joossens M, et al. Brief Report: Dialister as a Microbial Marker of Disease Activity in Spondyloarthritis. *Arthritis Rheumatol*; 69: 114–121.

66. Stoll ML, Weiss PF, Weiss JE, et al. Age and fecal microbial strain-specific differences in patients with spondyloarthritis. *Arthritis Res Ther* 2018; 20: 14.
67. Picchianti-Diamanti A, Panebianco C, Salemi S, et al. Analysis of Gut Microbiota in Rheumatoid Arthritis Patients: Disease-Related Dysbiosis and Modifications Induced by Etanercept. *Int J Mol Sci* 2018; 19: 2938.
68. Rizzatti G, Lopetuso LR, Gibiino G, et al. Proteobacteria: A Common Factor in Human Diseases. *BioMed Res Int*; 2017. Epub ahead of print 2017. DOI: 10.1155/2017/9351507.
69. Muniz-Pedrogo DA, Chen J, Hillmann BM, et al. Mo1934 - Gut Microbial Markers of Arthritis Including Inflammatory Bowel Disease Associated Arthropathy. *Gastroenterology* 2018; 154: S-856.
70. Liu X, Zou Q, Zeng B, et al. Analysis of Fecal Lactobacillus Community Structure in Patients with Early Rheumatoid Arthritis. *Curr Microbiol* 2013; 67: 170–176.
71. Maeda Y, Matsushita M, Katayama M, et al. SAT0079 The analysis of fecal microbiota in rheumatoid arthritis patients compared to healthy volunteers using bacterial RRNA-targeted reverse transcription-quantitative PCR. *Ann Rheum Dis* 2013; 71: 496–496.
72. Zhang X, Zhang D, Jia H, et al. The oral and gut microbiomes are perturbed in rheumatoid arthritis and partly normalized after treatment. *Nat Med* 2015; 21: 895–905.
73. Benham H, MARADANA M, LAKIS V Anne. Distinct Oral and Fecal Community Profiles Enriched in Opportunistic Pathogens in RA Patients and First Degree Relatives Are Influenced By Environmental Risk Factors, Including Smoking, Dental History and Lung Infection. *ACR Meeting Abstracts*, <https://acrabstracts.org/abstract/distinct-oral-and-fecal-community-profiles-enriched-in-opportunistic-pathogens-in-ra-patients-and-first-degree-relatives-are-influenced-by-environmental-risk-factors-including-smoking-dental-history/> (2016).
74. Costello ME, Asquith M, kim-Anh LC. HLA-B27 and Ankylosing Spondylitis Have Shared Effects on the Gut Microbiome. *ACR Meeting Abstracts*, <https://acrabstracts.org/abstract/hla-b27-and-ankylosing-spondylitis-have-shared-effects-on-the-gut-microbiome/> (accessed 13 April 2019).
75. Scher JU, Ubeda C, Artacho A, et al. Decreased bacterial diversity characterizes the altered gut microbiota in patients with psoriatic arthritis, resembling dysbiosis in inflammatory bowel disease. *Arthritis Rheumatol Hoboken NJ* 2015; 67: 128–139.
76. Manasson J, Shen N, Garcia Ferrer HR, et al. Gut Microbiota Perturbations in Reactive Arthritis and Postinfectious Spondyloarthritis. *Arthritis Rheumatol Hoboken NJ* 2018; 70: 242–254.
77. Costello, ME. Evidence Of a Microbial Signature In The Intestinal Microbiome In Ankylosing Spondylitis. - UQ eSpace, <https://espace.library.uq.edu.au/view/UQ:315298> (accessed 22 May 2018).
78. Stoll ML, Weiss P F, Weiss JE. Enteric Flora in Newly Diagnosed Spondyloarthritis: A Collaborative Study. *ACR Meeting Abstracts: abstract Nombre: 2805*.
79. Scher JU, Sczesnak A, Longman RS, et al. Expansion of intestinal Prevotella copri correlates with enhanced susceptibility to arthritis. *eLife*; 2. Epub ahead of print 5 November 2013. DOI: 10.7554/eLife.01202.
80. Maeda Y, Kurakawa T, Umemoto E, et al. Dysbiosis Contributes to Arthritis Development via Activation of Autoreactive T Cells in the Intestine. *Arthritis Rheumatol Hoboken NJ* 2016; 68: 2646–2661.
81. Takahashi N, Ishihara K, Kimizuka R, et al. The effects of tetracycline, minocycline,

- doxycycline and ofloxacin on *Prevotella intermedia* biofilm. *Oral Microbiol Immunol*; 21: 366–371.
82. Vahtovuo J, Munukka E, Korkeamäki M, et al. Fecal Microbiota in Early Rheumatoid Arthritis. *J Rheumatol* 2008; 35: 1500–1505.
83. Stoll ML, Kumar R, Morrow CD, et al. Altered microbiota associated with abnormal humoral immune responses to commensal organisms in enthesitis-related arthritis. *Arthritis Res Ther* 2014; 16: 486.
84. Varela E, Manichanh C, Gallart M, et al. Colonisation by *Faecalibacterium prausnitzii* and maintenance of clinical remission in patients with ulcerative colitis. *Aliment Pharmacol Ther*; 38: 151–161.
85. Dorofeyev AE, Vasilenko IV, Rassokhina OA. Joint extraintestinal manifestations in ulcerative colitis. *Dig Dis Basel Switz* 2009; 27: 502–510.
86. Costello M-E, Ciccia F, Willner D, et al. Brief Report: Intestinal Dysbiosis in Ankylosing Spondylitis. *Arthritis Rheumatol*; 67: 686–691.
87. Siala M, Gdoura R, Fourati H, et al. Broad-range PCR, cloning and sequencing of the full 16S rRNA gene for detection of bacterial DNA in synovial fluid samples of Tunisian patients with reactive and undifferentiated arthritis. *Arthritis Res Ther* 2009; 11: R102.
88. Stebbings S, Munro K, Simon MA, et al. Comparison of the faecal microflora of patients with ankylosing spondylitis and controls using molecular methods of analysis. *Rheumatology* 2002; 41: 1395–1401.
89. J. Kabeerdoss, S. Pugazhendhi, A. Balekuderu, et al. Gut microbiome in inflammatory bowel disease patients with and without arthropathy: Poster Sessions WED-136. *FEBS J* 2014; 281: 65–784.
90. Würdemann D, Tindall BJ, Pukall R, et al. *Gordonibacter pamelaiae* gen. nov., sp. nov., a new member of the Coriobacteriaceae isolated from a patient with Crohn's disease, and reclassification of *Eggerthella hongkongensis* Lau et al. 2006 as *Paraeggerthella hongkongensis* gen. nov., comb. nov. *Int J Syst Evol Microbiol* 2009; 59: 1405–1415.
91. Rothfuss KS, Stange EF, Herrlinger KR. Extraintestinal manifestations and complications in inflammatory bowel diseases. *World J Gastroenterol WJG* 2006; 12: 4819–4831.
92. Danese S, Semeraro S, Papa A, et al. Extraintestinal manifestations in inflammatory bowel disease. *World J Gastroenterol* 2005; 11: 7227–7236.
93. Hablot J, Peyrin-Biroulet L, Kokten T, et al. Experimental colitis delays and reduces the severity of collagen-induced arthritis in mice. *PLoS ONE*; 12. Epub ahead of print 19 September 2017. DOI: 10.1371/journal.pone.0184624.
94. Ahl D, Liu H, Schreiber O, et al. *Lactobacillus reuteri* increases mucus thickness and ameliorates dextran sulphate sodium-induced colitis in mice. *Acta Physiol* 2016; 217: 300–310.
95. Chen L, Zou Y, Peng J, et al. *Lactobacillus acidophilus* suppresses colitis-associated activation of the IL-23/Th17 axis. *J Immunol Res* 2015; 2015: 909514.
96. Liu X, Zeng B, Zhang J, et al. Role of the Gut Microbiome in Modulating Arthritis Progression in Mice. *Sci Rep* 2016; 6: 30594.
97. Viladomiu M, Kivolowitz C, Abdulhamid A, et al. IgA-coated *E. coli* enriched in Crohn's disease spondyloarthritis promote TH17-dependent inflammation. *Sci Transl Med*; 9. Epub ahead of print 8 February 2017. DOI: 10.1126/scitranslmed.aaf9655.

98. Esen Sefik. Individual intestinal symbionts induce a distinct population of ROR γ + regulatory T cells, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4700932/> (accessed 27 September 2018).
99. Andrew L. Kau. Functional characterization of IgA-targeted bacterial taxa from malnourished Malawian children that produce diet-dependent enteropathy, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4423598/> (accessed 27 September 2018).
100. Burton PR, Clayton DG, Cardon LR, et al. Association scan of 14,500 nonsynonymous SNPs in four diseases identifies autoimmunity variants. *Nat Genet* 2007; 39: 1329–1337.
101. DANIEL O. CLEGG. Comparison of sulfasalazine and placebo in the treatment of psoriatic arthritis. A department of veterans affairs cooperative study - Clegg - 1996 - Arthritis & Rheumatism - Wiley Online Library, <https://onlinelibrary.wiley.com/doi/abs/10.1002/art.1780391210>.
102. Generini S. Infliximab in spondyloarthritis associated with Crohn's disease: an open study on the efficacy of inducing and maintaining remission of musculoskeletal and gut manifestations. *Ann Rheum Dis* 2004; 63: 1664–1669.
103. Vandeputte D, Falony G, Vieira-Silva S, et al. Stool consistency is strongly associated with gut microbiota richness and composition, enterotypes and bacterial growth rates. *Gut* 2016; 65: 57–62.
104. Standaert–Vitse A, Jouault T, Vandewalle P, et al. Candida albicans Is an Immunogen for Anti–Saccharomyces cerevisiae Antibody Markers of Crohn's Disease. *Gastroenterology* 2006; 130: 1764–1775.