



HAL
open science

Association Between Microscopic Lesions at Ileal Resection Margin and Recurrence After Surgery in Patients With Crohn's Disease

Nassim Hammoudi, Dominique Cazals-Hatem, Claire Auzolle, Charlotte Gardair, Marjolaine Ngollo, Hugo Bottois, Stéphane Nancey, Benjamin Pariente, Anthony Buisson, Xavier Tréton, et al.

► To cite this version:

Nassim Hammoudi, Dominique Cazals-Hatem, Claire Auzolle, Charlotte Gardair, Marjolaine Ngollo, et al.. Association Between Microscopic Lesions at Ileal Resection Margin and Recurrence After Surgery in Patients With Crohn's Disease. *Clinical Gastroenterology and Hepatology*, 2019, 10.1016/j.cgh.2019.04.045 . hal-02193633

HAL Id: hal-02193633

<https://hal.science/hal-02193633v1>

Submitted on 21 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Association Between Microscopic Lesions at Ileal Resection Margin and**
2 **Recurrence After Surgery in Patients With Crohn's Disease**

3
4 Nassim Hammoudi^{*1,2}, Dominique Cazals-Hatem^{*3}, Claire Auzolle^{1,2}, Charlotte
5 Gardair⁴, Marjolaine Ngollo¹, Hugo Bottois¹, Stéphane Nancey⁵, Benjamin Pariente⁶,
6 Anthony Buisson⁷, Xavier Treton⁸, Mathurin Fumery⁹, Madeleine Bezault¹, Philippe
7 Seksik¹⁰, Lionel Le Bourhis¹, REMIND study group investigators, Jean-François
8 Flejou¹¹, Matthieu Allez^{1,2}

9 * Both authors contributed equally

10
11 Affiliations:

12 1, INSERM UMRS 1160, Université Paris Diderot, Sorbonne Paris-Cité University,
13 Paris, France. 2, Gastroenterology Department, AP-HP, Hôpital Saint-Louis, Paris,
14 France. 3 Pathology Department, AP-HP, Hôpital Beaujon, Clichy, France, 4,
15 Pathology Department, AP-HP, Hôpital Saint-Louis, Paris, France. 5,
16 Gastroenterology Department, Lyon Sud Hospital, Hospices Civils de Lyon, Lyon,
17 France. 6 Gastroenterology Department, Hôpital Claude Huriez, University of Lille 2,
18 Lille, France. 7 Université Clermont Auvergne, Inserm U1071, M2iSH, 3iHP, CHU
19 Clermont-Ferrand, Service de Médecine de l'Appareil Digestif, Clermont-Ferrand,
20 France. 8 Hôpital Beaujon, Service de Gastroentérologie, MICI et Assistance
21 Nutritive, Clichy, France. 9 Hepatogastroenterology Department, Amiens University
22 Hospital, Amiens, France. 10 Sorbonne Universités, Department of
23 Gastroenterology, Saint Antoine Hospital, AP-HP, Ecole normale supérieure, PSL
24 Research University, CNRS, INSERM, Laboratoire des Biomolécules (LBM), Paris,

25 France. 11 Pathology Department, AP-HP, Hôpital Saint-Antoine, Faculté de
26 Médecine Sorbonne Université, Paris, France.

27

28 Word count:

29 Abstract: 274

30 Manuscript (including references and figure legends): 4010

31 Manuscript: 2951 / References: 26, 629 words / Figures and legends: 430

32 Pages: 26 / Tables: 3 / Figures: 3

33

34 Corresponding Author: Prof Matthieu Allez

35 Département de Gastroentérologie

36 Hôpital Saint-Louis

37 1 avenue Claude Vellefaux

38 75010 Paris, France

39 Email : matthieu.allez@aphp.fr

40

41 *Conflict of interest statement:* NH declares no competing interest. DCH declares no
42 competing interest. CA declares no competing interest. CG declares no competing
43 interest. MN declares no competing interest. HB declares no competing interest. SN
44 received honoraria from MSD, Abbvie, Takeda, Janssen, HAC Pharma, Tillots,
45 Ferring, and Novartis. BP received honoraria from AbbVie, MSD, Takeda, Janssen,
46 Bioagaran, and Ferring. AB received honoraria from MSD, Abbvie, Ferring, Takeda,
47 Vifor Pharma, Sanofi-Aventis, Hospira, and Janssen. XT received honoraria from
48 Abbvie, MSD, Takeda, Ferring, Norgine, and Janssen. MF received honoraria from
49 AbbVie, MSD, Takeda, Janssen, Pfizer, Ferring, and Boehringer. MB declares no

50 competing interest. LLB declares no competing interest. PS received honoraria from
51 Takeda, MSD, Biocodex, Ferring and Abbvie and financial support from Takeda. JFF
52 declares no competing interest. MA received honoraria from Abbvie, MSD, Janssen,
53 Takeda, Pfizer, Novartis, Ferring, Tillots, Celgene, and Genentech/Roche.

54 *Guarantor of the article:* Matthieu Allez

55 *Author contributions:* NH: histological analysis, interpretation of data and drafting of
56 the manuscript; DCH: histological analysis, design, supervision of histological
57 analysis and critical review of the manuscript; CA: statistical analysis and critical
58 review of the manuscript; CG: histological analysis and critical review of the
59 manuscript; HB, MN, SN, BP, AB, XT, MF, PS, LLB: data collection and critical
60 review of the manuscript; MB: management of the study; JFF: design of histological
61 analysis, critical review of the manuscript; MA: conception, design and supervision of
62 the study, analysis of data, drafting and critical review of the manuscript. All authors
63 reviewed and approved the final version of the manuscript.

64

65 **ABSTRACT**
66

67 **BACKGROUND AND AIMS:** Different types of histologic lesions at the ileal margin,
68 detected by histology, have been associated with increased rates of recurrence after
69 ileocaecal surgery in patients with Crohn's disease (CD). We aimed to characterize
70 histologic features of the ileal margin and to evaluate their association with disease
71 recurrence. **METHODS:** We collected histologic data from 211 patients with ileal or
72 ileocolonic CD who underwent ileocolonic resections at hospitals in France from
73 September 2010 through December 2016. Ileal margins were analyzed. Early
74 endoscopic recurrence was defined by a Rutgeerts score of ≥ 2 or more, 6 months
75 after surgery. We also collected data from 10 adults with healthy ileum who
76 underwent ileocecal resection for colonic tumors (controls). Clinical relapse was
77 defined by CD-related symptoms confirmed by imaging, endoscopy, therapy
78 intensification, CD-related complication, or subsequent surgery. **RESULTS:** Six
79 months after surgery, 49% of patients had endoscopic recurrence; 5 years after
80 surgery, 57% of patients had clinical relapse. Ileal margins were macroscopically
81 affected in 20.9% of patients. CD transmural lesions at the margin (defined by
82 mucosal ulceration or cryptitis, submucosal fibrosis and lymphoplasmocytic infiltrate
83 of the subserosa) were observed in 13.6% of patients. Endoscopic recurrence was
84 observed in 75% of patients with CD transmural lesions vs 46% of patients without
85 ($P=.005$). In multivariate analysis, CD transmural lesions at the margin were
86 independently associated with early endoscopic recurrence (OR, 3.83; 95% CI, 1.47-
87 11.05; $P=.008$) and clinical recurrence (OR 2.04; 95% CI, 1.09-3.99; $P=.026$).
88 **CONCLUSION:** In patients with CD, transmural lesions at the ileal margin were
89 associated with an increased risk of post-operative recurrence. Histologic features of
90 the ileal margin should be included in making decisions about post-operative therapy.

91

92 **KEYWORDS**

93

94 REMIND trial

95

96 Outcome

97

98 Prognostic Factor

99

100 IBD Surgery

101

102

103 **ABBREVIATIONS**

104

105 CD: Crohn's Disease

106

107 HES: Hematosin Eosin Saffron

108

109 IQR: Inter Quartile Range

110

111 MP: Myenteric Plexitis

112

113 OR: Odds Ratio

114 INTRODUCTION

115

116 Despite recent advances in the medical management of Crohn's Disease
117 (CD), more than two thirds of patients still require intestinal resection during the
118 course of their disease (1-3). Furthermore, surgery is not curative and post-operative
119 recurrence occurs in the majority of patients. Early post-operative endoscopic
120 recurrence, which is mainly observed at the anastomosis or in the neo-terminal
121 ileum, precedes clinical recurrence and its severity predicts the long-term clinical
122 outcome (4). Several risk factors have been reported to increase post-operative
123 recurrence, including active smoking, penetrating disease, history of perianal
124 disease, previous intestinal surgery and male gender (5-7). There is still a need for
125 factors that can accurately risk-stratify patients (8).

126 Macroscopically involved ileal margin could increase post-operative
127 recurrence risk but this assumption was pushed aside in 1996 by Fazio et al with the
128 results of a randomized controlled trial showing no benefit of extended resection
129 (clearance of 12 cm versus 2 cm of normal ileal margin) (9). Retrospective studies
130 showed conflicting results regarding the impact of CD microscopic lesions on the ileal
131 margin (mainly granuloma or active inflammation) on post-operative recurrence (10-
132 13). Several publications have described an association between plexitis and
133 endoscopic, clinical or surgical recurrence. Plexitis is classically defined by the
134 presence of inflammatory cells in the myenteric or submucosal plexi (14). There were
135 diverse definitions of plexitis and various endpoints were used (15-21). More
136 recently, a cohort analysis of consecutive CD patients undergoing a first ileocaecal
137 resection in two specialized centers found that microscopic involvement of the ileal
138 margin, with the presence of granuloma, ulcers or transmural inflammation, was an
139 independent risk factor of clinical and surgical recurrence (22).

140 The impact of CD lesions at the proximal ileal margin on recurrence is, thus,
141 still unclear. Our aims were to characterize the microscopic involvement of the
142 proximal ileal margin and to study its predictive value on rates of endoscopic
143 recurrence 6 months after surgery, clinical relapse and surgery during the long-term
144 follow-up. We performed this study on a well-characterized prospective multicenter
145 cohort of CD patients followed from surgery (7).

146 **MATERIALS AND METHODS**

147

148 ***Post-operative cohort and patient selection***

149 The present work was undertaken on a prospective multicenter study
150 performed by the REMIND group that aimed at identifying predictors of early post-
151 operative endoscopic recurrence (7). Patients were included between September
152 2010 and December 2016. Inclusion criteria were: age >18 years, ileal or ileocolonic
153 CD and indication of CD-related intestinal surgery (ileocolonic resection). A post-
154 operative treatment was recommended according to a pre-established algorithm,
155 based on the following risk factors: current smoking, previous bowel resection,
156 penetrating phenotype and active perianal disease (detailed in 7). All patients had, 6
157 months after surgery, a colonoscopy to assess the endoscopic recurrence according
158 to the Rutgeerts score (23). From this cohort, we selected patients who had a
159 paraffin-embedded ileal margin sample available for histological analysis. Margins
160 with poor quality paraffin embedding were excluded. All patients provided an
161 informed written consent. The study was approved by AFSAPS (IDRCB: 2009-
162 A00205-52), the French Ethic Committee - Hôpital Saint-Louis (CPP 2009/17) and
163 declared to clinicaltrials.gov (NCT03458195).

164 To test CD-related histological lesions' specificity, 10 control adults with
165 healthy ileum who underwent ileocaecal resection for colonic tumor constituted a
166 control group for the histological analysis. All control patients provided an approval
167 consent form.

168

169

170

171 ***Data collection***

172 Clinical parameters were collected prospectively at time of surgery and at
173 post-operative endoscopy. They included demographical data (gender, age at
174 diagnosis, age at surgery, disease duration at surgery, smoking status at surgery),
175 clinical data, complication and surgical indication, anoperineal lesions, previous
176 history of intestinal resection, extra-digestive manifestations, treatment history
177 including previous exposure to immunosuppressants (azathioprine, 6-MP or
178 methotrexate) and anti-TNF agents.

179 An endoscopy was performed 6 months after surgery. Endoscopic data were
180 collected prospectively. All lesions were reported in all segments, including the
181 anastomotic region and the neo-terminal ileum. The physician who performed the
182 colonoscopy evaluated the Rutgeerts score for each patient. Two physicians (MA
183 and CA) checked the colonoscopy's report blinded to histological analysis and post-
184 operative treatment.

185 Long-term data of patients with a follow-up of at least 18 months were
186 collected retrospectively. They included clinical data (smoking behavior, type and
187 date of clinical symptoms, type and date of CD related complications: intra-abdominal
188 abscess, occlusive manifestation, anoperineal lesion and death), treatment data
189 (modification or optimization with initiation and cessation date for each drug),
190 endoscopic data (date and results of colonoscopy performed during the follow-up),
191 imaging data (date and results of magnetic resonance enterography and/or CT scan
192 performed during the follow-up) and surgical data (date and type of subsequent
193 surgery performed).

194

195

196 ***Histopathological analysis***

197 Paraffin blocks of the ileal margin were cut into a full-thickness 3- μ m section
198 and stained with hematein eosin saffron (HES). Each section was jointly analyzed by
199 two expert pathologists (DCH and CG) blinded to the baseline detailed clinical data
200 and to the post-operative outcomes of the patients. An analysis grid was built to
201 assess CD-related lesions affecting each intestinal wall layer (mucosa, submucosa,
202 and subserosa/muscularis).

203 Each of these lesions was considered as binary variable except for plexitis, for
204 which three degrees were defined according to Ferrante et al (15). Ileal margins
205 from the control group (10 patients with healthy ileum without CD) were similarly
206 examined for the same histologic lesions.

207 For macroscopic evaluation, we collected the length of ileal resection for each
208 patient (in centimeters) and the distance from the ileal margin to the first macroscopic
209 CD mucosal ulceration noticed by pathologist on opened specimen (in centimeters).
210 The margin was defined as “macroscopically affected” if CD’s mucosal ulcerations
211 were noticed on the ileal margin.

212

213 ***Evaluated outcomes***

214 Post-operative endoscopic recurrence was evaluated 6 months after surgery.
215 Endoscopic recurrence was defined as a Rutgeerts score \geq i2.

216 Clinical recurrence was defined by: (i) CD-related clinical manifestations
217 confirmed either by endoscopy (Rutgeerts score \geq i2 or severe colonic lesions in one
218 of the follow-up colonoscopies), imaging (active lesions of the small bowel or colon
219 on an examination performed by the center referring radiologist) or therapeutic
220 intensification (treatment optimization or drug switch), (ii) CD-related complications

221 (intra-abdominal abscess or occlusive manifestation) or (iii) CD-related subsequent
222 surgery.

223 Surgical recurrence was defined by a further CD-related surgery at the
224 anastomotic site (subsequent ileocolonic resection or stricturoplasty).

225

226 ***Statistical analyses***

227 Descriptive statistics were calculated for all quantitative variables as medians
228 with their respective interquartile ranges (IQR). Correlation between histology and
229 recurrence, for each item of the analysis grid, was evaluated by both univariate and
230 multivariate analyses. Chi-2 and Logistic regression were respectively used for binary
231 variables. For continuous data (lengths in centimeters), Mann-Whitney test was used.
232 Sensitivity, specificity, positive and negative predictive values were calculated using
233 standard formulas. Spearman's rank correlation test was used when necessary to
234 evaluate the correlation between variables.

235 For both clinical and surgical recurrence, Kaplan-Meier survival curves and
236 crude log rank tests were calculated for all variables (histological items and clinical
237 data at baseline) before survival multivariate analysis. A multivariate Cox model
238 combining demographical data (gender and age) and variables significantly
239 associated with clinical or surgical recurrence in univariate analyses was built.

240 All statistical analyses were two-tailed and a p value of less than 0.05 was
241 considered as statistically significant. To perform these analyses, R software ® (3.2.2
242 version) and PrismGraphPad ® were used.

243 **RESULTS**

244 ***Patients' characteristics at surgery and during follow-up***

245 Two hundred and twenty five patients were included (7). Ileal margins were
246 not technically exploitable for 14. In total, ileal margins were analyzed in 211 patients
247 and 206 ileal margins had a complete transmural histological evaluation
248 **(Supplementary Figure 1)**.

249 Median age at surgery was 36.3 years (IQR: 30.7-47.5) and median disease
250 duration was 6.3 years (IQR: 1.8-11.9). Ninety-two (44%) patients were male, 67
251 (32%) active smokers and 38 (18%) had a previous ileocaecal resection in their past
252 medical history (**Table 1**). Post-operative treatment was given to 102 (48%) patients,
253 including an anti-TNF treatment in 52 (25%) patients.

254 Post-operative endoscopy was performed in all patients with a median time
255 between resection and colonoscopy of 6.8 months. One hundred and four patients
256 (49%) had an endoscopic recurrence (i.e. Rutgeerts score \geq i2) and 43 (20%)
257 patients had severe lesions (i.e. i3 or i4). One hundred and seven (51%) patients had
258 no endoscopic recurrence including 68 (32%) without any lesion (i0).

259 Long-term follow-up (more than 18 months) was available in 168 (75%)
260 patients. Baseline characteristics, post-operative management and endoscopic
261 recurrence rates of these 168 patients did not differ significantly from the whole
262 cohort.

263 Median follow-up was 3.8 years (IQR: 2.55-5.41). Seventy-five (45%) patients
264 experienced clinical relapse during the follow-up. Cumulative clinical recurrence rate
265 was 5%, 37% and 57% at 1, 3 and 5 years respectively. At the end of the follow-up, 4
266 (2%) patients underwent a CD-related subsequent surgery (3 ileocolonic resections,
267 1 stricturoplasty).

268

269 ***Histopathological analysis of ileal margin***

270 Median ileal resection length was 19 cm (IQR: 12-27). The ileal margin was
271 considered as macroscopically affected by the pathologist in 44 (20.9%) patients
272 (**Figure 1**). Among them, 41 (93.2%) presented with at least one mucosal lesion in
273 histology. The median distance between macroscopical ulceration (noticed by the
274 pathologist in 178 patients) and the ileal margin was 4 cm (IQR: 1-8).

275 Microscopically, grade 1 MP was found on the ileal margin of the 10 control
276 patients revealing CD-unrelated lesion (discarded from the final grid). Among the 211
277 patients studied, 65 (30.8%) had a normal ileal margin and 146 (69.2%) had at least
278 one CD lesion (**Table 2**). The most affected layer of the bowel wall was the mucosa
279 (111 patients (52.6%)). Sixty patients (28.4%) had mucosal ulceration or cryptitis
280 indicative of CD activity. The submucosal layer was affected in 104 (49.3%) patients,
281 16 (7.8%) with neutrophils infiltrate (all associated with active mucosal lesion). The
282 most frequent submucosal lesions were lymphoplasmocytic infiltrate (40%) and
283 fibrosis (33.8%). All patients with submucosal plexitis (38.6%) had concomitant
284 lymphoplasmocytic infiltrate. The subserosa/muscularis (plexitis included) were
285 affected in 92 (44.7%) patients. The subserosa was affected in 64 (31.1%) patients
286 (**Supplementary Figure 2**).

287 Patients with at least one lesion of each layer frequently presented grade 2 or
288 3 MP (respectively 21 (55%) patients and 12 (33%)). Interestingly, plexitis (either
289 submucosal or myenteric) was the only lesion found in 6 patients (3%).

290 The presence of microscopic lesions was inversely correlated with the
291 distance between ileal margin and macroscopic ulceration: patients with transmural
292 lesions had a shorter disease-free margin when compared to patients with 2, 1 or

293 none of the three histological features defining transmural impairment (Spearman's
294 rank correlation test: $r=-0.476$; $p<0.001$) (**Supplementary Figure 3**).

295

296 ***Correlation with endoscopic recurrence***

297 The ileal resection length, the macroscopic involvement of the ileal margin and
298 the distance between the ileal margin and the first macroscopic CD ulceration were
299 not correlated with endoscopic recurrence ($p=0.96$, $p=0.31$ and $p=0.43$ respectively).

300 In univariate analysis, two histological lesions were significantly associated
301 with endoscopic recurrence: submucosal fibrosis (recurrence rate: 59.1% vs. 44.6%,
302 OR=1.80, 95%CI [1.01-3.21], $p=0.04$) and lymphoplasmocytic infiltrate of the
303 subserosa (recurrence rate: 60.3% vs. 44,6%, OR=1.89, 95%CI [1.01-3.51], $p=0.04$)
304 (**Table 2**). A trend towards recurrence was also found in patients displaying mucosal
305 ulceration or cryptitis indicative of active CD (recurrence rate: 58.3% vs. 45.7%,
306 OR=1.66, 95%CI [0.91-3.07], $p=0.09$) (**Table 2**). Submucosal or myenteric plexitis
307 were not associated with endoscopic recurrence.

308 Based on this univariate analysis, we defined *a posteriori* the item “CD
309 transmural lesions” by the presence of mucosal ulceration or cryptitis, submucosal
310 fibrosis and lymphoplasmocytic infiltrate of the subserosa. This histological feature
311 was observed in 28 patients (13.6%) out of the 206 samples in which these three
312 lesions could be assessed. These 28 patients had a significant increased risk of
313 endoscopic recurrence when compared to those without this histological feature
314 ($p=0.005$) (**Figure 2a**). Severe recurrence (i.e. score i3 or i4) was also more frequent
315 in patients with CD transmural lesions on margin ($p=0.03$) (**Figure 2a**). The sensitivity
316 and specificity of the presence of CD transmural lesions for the risk of endoscopic

317 recurrence was 20% and 93% respectively with an accuracy of 63%. Consequently,
318 the positive and negative predictive values were of 75% and 53% respectively.

319 Only 8 of the 28 patients who had CD transmural lesions received post-
320 operative anti-TNF. Only 2 were i0 and 6 had a recurrence (\geq i2). In patients with no
321 post-operative biotherapy, CD transmural lesions were also associated with
322 endoscopic recurrence ($p=0.03$, **figure 2b**).

323 We integrated the selected lesions indicative of CD transmural lesions in a
324 multivariate analysis with the clinical parameters associated with an increased risk of
325 endoscopic recurrence in our cohort (i.e. male gender, previous intestinal surgery,
326 active smoker at surgery and post-operative preventive anti-TNF treatment (7)). CD
327 transmural lesions remained independently associated with endoscopic recurrence
328 (OR: 3.83, 95%CI [1.47-11.05], $p=0.008$) (**Table 3**).

329

330 ***Correlation with long-term clinical outcome***

331 In univariate analysis, clinical parameters significantly associated with long-
332 term clinical recurrence were: active smoking at time of surgery ($p=0.007$), previous
333 intestinal surgery ($p=0.012$), and presence of extra-intestinal manifestations at
334 surgery ($p=0.049$). CD transmural lesions at the ileal margin were also associated
335 with a significantly poorer outcome ($p=0.022$) (**Figure 3**).

336 In multivariate analysis, the four factors identified in univariate analysis,
337 including CD transmural lesions at ileal margin, remained independently associated
338 with clinical recurrence (**Table 3**).

339

340 **DISCUSSION**

341 This prospective study is, to our knowledge, the largest analyzing the impact
342 of macroscopic and microscopic CD involvement of the ileal resection margin on
343 endoscopic and clinical outcomes after ileocaecal resection. CD transmural lesions at
344 ileal margin were strongly correlated with both early post-operative endoscopic
345 recurrence and long-term clinical outcome independent of other risk factors.
346 Conversely, we didn't find any correlation between reported macroscopic ulceration
347 noticed on the ileal margin and recurrence.

348 About one third of our patients had a microscopically normal ileal margin while
349 the macroscopic evaluation didn't notice mucosal ulcerations in 80% of patients. This
350 difference highlights the need for a complete microscopic evaluation of the bowel wall
351 to fully assess tissue injury.

352 A recent literature review emphasized the crucial need of new studies
353 assessing the involvement of both macroscopic and microscopic lesions in CD
354 recurrence after surgery (24). Two recent publications found an association between
355 microscopic lesions at ileal margins and recurrence in both adult and pediatric
356 patients (22, 25). However, no detail on the type of lesions and their location in the
357 intestinal wall were provided. Herein, two experts pathologists assessed each CD
358 histological lesions, including plexitis, using a reproducible grid. Thus, in contrast to
359 recent literature on the subject (15-21), we considered all CD-related lesions
360 potentially found in the whole bowel wall through a complete and reproducible
361 analysis rather than focusing on plexitis alone. In our cohort, plexitis, whatever its
362 severity, was not associated with recurrence. Interestingly, most of the patients with
363 grade 3 MP had significant lesions in other layers, while few patients (n=6) had
364 plexitis only.

365 We hypothesized that some histological lesions could be more relevant than
366 others and selected them from our statistical analysis. In univariate analysis, two
367 lesions were significantly associated with endoscopic recurrence - fibrosis of the
368 submucosa and lymphoplasmocytic infiltrate of the subserosa - reflecting sequelae
369 of previous CD flares or persistent chronic inflammation. Moreover, mucosal lesions
370 such as ulceration or cryptitis were also associated with a trend towards recurrence.
371 These 3 lesions had a similar impact on post-operative recurrence (odds ratios for
372 ulceration/cryptitis, fibrosis of the submucosa and lymphoplasmocytic infiltrate of the
373 subserosa were 1.66 [0.91-3.07], 1.80 [1.01-3.21] and 1.89 [1.01-3.51] respectively).
374 Thus, we used the combination of these 3 CD-related lesions, each affecting one
375 layer of the bowel wall, to define intense and active CD transmural involvement of the
376 ileal margin, which was strongly associated with recurrence. This finding highlights
377 that, rather than isolated microscopic lesions, deep CD-related tissue injury of the
378 ileal margin could influence recurrence risk. These patients could benefit from an
379 immediate post-operative treatment.

380 No association was found between macroscopic examination of the opened
381 surgical specimen (length of ileal resection and distance from margin to first
382 macroscopic ulceration) and rates of clinical or endoscopic recurrence. These data
383 are in alignment with a previous study using frozen section examination of resection
384 margins during surgery to achieve microscopical clearance of CD lesions (26). This
385 approach did not improve short and long-term patient's clinical outcomes. Our study
386 also does not support extensive resections. We believe that surgeons should still
387 perform limited resections.

388 CD is characterized by discontinuous patchy or extensive transmural
389 inflammation with skip lesions. Recurrence remains frequent despite normal

390 resection margins. Indeed, 46% of patients with normal ileal margin experienced
391 endoscopic recurrence, which was even severe in 10% (i3 or i4). Histological
392 evaluation of the margin has therefore a low sensitivity for prediction of post-
393 operative recurrence.

394 Our study has several limitations. First, the two pathologists analyzed jointly
395 the histological slides and we did not assess inter-observer variability. Second, there
396 was no endoscopic central reading, which could have led to misclassifications
397 between i1 and i2. However, two physicians checked the colonoscopies report
398 blinded of treatment and patients outcomes, which could have limited this potential
399 bias. Third, long-term clinical outcomes were collected retrospectively. Moreover, in
400 this large prospective cohort, some patients were lost to follow-up. However, no
401 difference was found between the 168 patients with long-term data and the whole
402 cohort. Finally, we did not assess the colonic margin, in which Inflammation could
403 potentially impact post-operative recurrence risk in a similar way.

404

405 In conclusion, we show in this prospective multicenter cohort study that CD
406 transmural histological lesions at the ileal margin of ileocaecal resection specimens
407 in CD were associated with an increased risk of endoscopic and clinical recurrence.
408 Pathologists should systematically detail in their reports the analysis of the ileal
409 margin of resection notifying the presence of CD transmural lesions
410 (ulceration/erosion, fibrosis and lymphoplasmocytic infiltrate). A systematic post-
411 operative treatment could be proposed to patients with transmural histological
412 lesions, even in absence of other risk factors. The better post-operative treatment to
413 initiate to these patients remains to be determined; ClinicalTrials.gov number,
414 NCT03458195.

415 REMIND STUDY GROUP INVESTIGATORS
416

417 Saint-Louis, Paris: Pierre Cattan, Mircea Chirica, Nicolas Munoz-Bongrand, H el ene
418 Corte, Nathan Beaupel, Jonathan Catry, Jean-Marc Gornet, Clotilde Baudry, Nelson
419 Lourenco, Mariane Maillet, My-Linh Tran-Minh, Victor Chardiny, C eline Grand, Brice
420 Gergaud, Joelle Bonnet, Leila Chedouba, Andr ee Nisard; Saint-Antoine, Paris:
421 Laurent Beaugerie, Harry Sokol, Anne Bourrier, Isabelle Nion-Larmurier, Julien
422 Kirchgesner, Elodie Quevrain, Loic Brot, Najim Chafai, Jeremie H. Lefevre,
423 Emmanuel Tiret, Magali Svrcek; Beaujon, Clichy: Nathalie Guedj, Yves Paris, Leon
424 Magiorri, Marianne Ferron, Yoram Bouhnik, Olivier Corcos, Carmen Stefanescu;
425 Lariboisi ere, Paris: Philippe Marteau, Xavier Dray, Ulrika Chaput, Rachid Kaci;
426 Clermont-Ferrand: Anne Dubois, Gilles Bommelaer, Marion Goutte, Nicolas Barnich,
427 Dilek Coban, Catherine Godfraind, Juliette Joubert Zakeyh ; Lille: Pierre
428 Desreumaux, Maria Nachury, Coralie Sommeville, Florence Renaud; Amiens: Jean-
429 Louis Dupas, Julien Loreau, Franck Brazier, Denis Chatelain, Christophe Attencourt,
430 Charles Sabbagh, Martine Leconte; Lyon: Gilles Boschetti, Bernard Flouri e, Yves
431 Fran ois, Eddy Cotte, Anne-Laure Charlois, Peggy Falgon, Helena Hadjisavvas,
432 Driffa Moussata, Marion Chauvenet, Sarah Boyer, Alexandra Traverse-Glehen; Nice:
433 Xavier Hebuterne, J erome Filippi, Paul Hofmann, Amine Rahili, St ephanie
434 Patouraux. Statistiques, PARCC Paris: Xavier Jouven

435

436 ACKNOWLEDGMENTS

437 We thank the Association Francois Aupetit, Helmsley Charitable Trust, MSD France,
438 and La Fondation pour la Recherche M edicale (DEA20170637874) for their funding
439 grants.

440 We also acknowledge Nathalie Colnot for her technical support.

441 **REFERENCES**

- 442 **1**-Binder V, Hendriksen C, Kreiner S. Prognosis in Crohn's disease--based on results
443 from a regional patient group from the county of Copenhagen.*Gut*.1985;26(2):146-
444 150.
- 445 **2**-Cosnes J, Nion-Larmurier I, Beaugerie L et al. Impact of the increasing use of
446 immunosuppressants in Crohn's disease on the need for intestinal
447 surgery.*Gut*.2005;54(2):237-241.
- 448 **3**-Peyrin-Biroulet L, Harmsen WS, Tremaine WJ, et al. Surgery in a Population-
449 Based Cohort of Crohn's Disease From Olmsted County, Minnesota (1970-
450 2004).*Am J Gastroenterol*.2012;107(11):1693-1701.
- 451 **4**-Rutgeerts P, Geboes K, Vantrappen G, et al. Natural history of recurrent Crohn's
452 disease at the ileocolonic anastomosis after curative surgery.*Gut*.1984;25(6):665-
453 672.
- 454 **5**-Gionchetti P, Dignass A, Danese S, et al. 3rd European Evidence-based
455 Consensus on the Diagnosis and Management of Crohn's Disease 2016: Part 2:
456 Surgical Management and Special Situations.*J Crohn's Colitis*.2017;11(2):135-149.
- 457 **6**-Buisson A, Chevaux J-B, Allen PB, et al. Review article: the natural history of
458 postoperative Crohn's disease recurrence.*Aliment Pharmacol Ther*.2012;35(6):625-
459 633.
- 460 **7**-Auzolle C, Nancey S, Tran-Minh ML, et al. Male gender, active smoking and
461 previous intestinal resection are risk factors of postoperative endoscopic recurrence
462 in Crohn's disease: results from a prospective cohort study.*Aliment Pharmacol*
463 *Ther*.2018;48(10):1156-1157
- 464 **8**-Singh S. Editorial: predicting recurrence of Crohn's disease after surgical resection-
465 close to a crystal ball.*Aliment Pharmacol Ther*.2018;48(10):1156-7.

466 **9**-Fazio VW, Marchetti F, Church M, et al. Effect of resection margins on the
467 recurrence of Crohn's disease in the small bowel. A randomized controlled trial.*Ann*
468 *Surg.*1996;224(4):563-71-3.

469 **10**-Pennington L, Hamilton SR, Bayless TM, et al. Surgical management of Crohn's
470 disease. Influence of disease at margin of resection.*Ann Surg.*1980;192(3):311-318.

471 **11**-Cooper JC, Williams NS. The influence of microscopic disease at the margin of
472 resection on recurrence rates in Crohn's disease.*Ann R Coll Surg*
473 *Engl.*1986;68(1):23-26.

474 **12**-Kotanagi H, Kramer K, Fazio VW, et al. Do microscopic abnormalities at resection
475 margins correlate with increased anastomotic recurrence in Crohn's disease?
476 Retrospective analysis of 100 cases.*Dis Colon Rectum.*1991;34(10):909-916.

477 **13**-Heimann TM, Greenstein AJ, Lewis B, et al. Prediction of early symptomatic
478 recurrence after intestinal resection in Crohn's disease.*Ann Surg.*1993;218(3):294-8-
479 9.

480 **14**-Geboes K, Rutgeerts P, Ectors N, et al. Major histocompatibility class II
481 expression on the small intestinal nervous system in Crohn's
482 disease.*Gastroenterology.*1992;103:439-447.

483 **15**-Ferrante M, de Hertogh G, Hlavaty T, et al. The Value of Myenteric Plexitis to
484 Predict Early Postoperative Crohn's Disease
485 Recurrence.*Gastroenterology.*2006;130(6):1595-1606.

486 **16**-Ng SC, Lied GA, Kamm MA, et al. Predictive value and clinical significance of
487 myenteric plexitis in Crohn's disease.*Inflamm Bowel Dis.*2009;15(10):1499-1507.

488 **17**-Sokol H, Polin V, Lavergne-Slove A, et al. Plexitis as a predictive factor of early
489 postoperative clinical recurrence in Crohn's disease.*Gut.*2009;58(9):1218-1225.

490 **18**-Bressenot A, Chevaux J-B, Williet N, et al. Submucosal Plexitis as a Predictor of
491 Postoperative Surgical Recurrence in Crohn's Disease.*Inflamm Bowel*
492 *Dis.*2013;19(8):1654-1661.

493 **19**-Misteli H, Koh CE, Wang LM, et al. Myenteric plexitis at the proximal resection
494 margin is a predictive marker for surgical recurrence of ileocaecal Crohn's
495 disease.*Colorectal Dis.*2015;17(4):304-310.

496 **20**-Decousus S, Boucher A-L, Joubert J, et al. Myenteric plexitis is a risk factor for
497 endoscopic and clinical postoperative recurrence after ileocolonic resection in
498 Crohn's disease.*Dig Liver Dis.*2016;48(7):753-758.

499 **21**-Lemmens B, de Buck van Overstraeten A, Arijis I, et al. Submucosal Plexitis as a
500 Predictive Factor for Postoperative Endoscopic Recurrence in Patients with Crohn's
501 Disease Undergoing a Resection with Ileocolonic Anastomosis: Results from a
502 Prospective Single-centre Study.*J Crohn's Colitis.*2017;11(2):212-220.

503 **22**-de Buck van Overstraeten A, Eshuis EJ, Vermeire S, et al. Short- and medium-
504 term outcomes following primary ileocaecal resection for Crohn's disease in two
505 specialist centres.*Br J Surg.*2017;104(12):1713-1722.

506 **23**-Rutgeerts P, Geboes K, Vantrappen G, et al. Predictability of the postoperative
507 course of Crohn's disease.*Gastroenterology.*1990;99(4):956-963.

508 **24**-Patel K V., Darakhshan AA, Griffin N, et al. Patient optimization for surgery
509 relating to Crohn's disease.*Nat Rev Gastroenterol Hepatol.*2016;13(12):707-719.

510 **25**-Diederer K, de Ridder L, van Rheenen P, et al. Complications and Disease
511 Recurrence After Primary Ileocecal Resection in Pediatric Crohn's Disease.*Inflamm*
512 *Bowel Dis.*2017;23(2):272-282.

513 **26**-Hamilton SR, Reese J, Pennington L, et al. The role of resection margin frozen
514 section in the surgical management of Crohn's disease.*Surg Gynecol*
515 *Obstet.*1985;160(1):57-62.

516

517

518

519

520 **TABLE LEGENDS**

521

522 **Table 1: Patients' characteristics at time of surgery**

523

524 **Table 2: Pathological data after examination of the ileal margin of the 211**
525 **patients and correlation of each histological lesion with endoscopic**
526 **recurrence.**

527 The margin is "affected" if at least one histological lesion is observed in one of the
528 bowel wall layer (as defined below). Margin macroscopically affected is defined as
529 mucosal ulceration observed by the pathologist on the ileal margin when opening the
530 surgical specimen

531 Granuloma=nest of at least 5 epithelioid cells in the wall; Eosinophilic infiltrate=nest
532 of at least 50 eosinophils in the lamina propria at x400 magnification; Villous
533 atrophy=extensive subtotal / total villous atrophy

534 Pyloric Metaplasia=metaplastic glands of the ileal mucosa

535 Ulceration/Cryptitis=acute inflammation (neutrophils) in epithelium or crypt

536 Neutrophilic infiltrate=nest of at least 50 neutrophils in the submucosae at x400

537 magnification; Lymphoplasmocytic infiltrate=nest of at least 50 lymphocytes or

538 plasma cells at x400 magnification; Plexitis=inflammatory cells around or inside

539 submucosal or myenteric plexus as defined previously (14) (Grade 1 Myenteric

540 Plexitis (MP: at least 1 but less than 4 immune cells within or contiguous to the

541 plexus, Grade 2 MP: at least 4 but less than 10 immune cells and Grade 3 MP: at

542 least 10 immune cells);

543 Nervous hyperplasia=enlarged nervous bundles visible at x100 magnification;

544 Fibrosis=significant collagen deposit in intestinal layer visible at x100 magnification.

545

546

547 **Table 3: Correlation between clinical factors, CD transmural lesions of the ileal**
548 **margin and both endoscopic and clinical recurrences.**

549 Multivariate analysis. Significance was reached if $p < 0.05$.

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577 **FIGURES LEGENDS**

578

579

580 **Figure 1: Pathological analysis of the whole ileal margin on surgical specimen**

581 **and on section stained by hematein-eosin-saffron. (A)-Macroscopic view of an**

582 **opened ileocaecal resection specimen with mucosal ulcerations visible on ileal**

583 **proximal margin (arrow) (B)-Microscopic view of the whole ileal margin section**

584 **showing a bowel wall with CD-related lesions affecting the 3 layers (mucosal,**

585 **submucosal and subserosal layers in circles). (C)-Fibrosis observed in the**

586 **submucosa. (D)-Lymphoplasmocytic infiltrate observed in the subserosa.**

587

588 **Figure 2: Endoscopic recurrence rates depending on selected**

589 **histopathological lesions at ileal margin (ulceration, fibrosis in submucosa,**

590 **lymphoplasmocytic infiltrate in subserosa).**

591 **No lesion: patients with normal ileal margin. CD lesions but not transmural: patients**

592 **with ileal margin affected but without CD transmural lesions defined by the presence**

593 **of mucosal ulceration or cryptitis, submucosal fibrosis and lymphoplasmocytic**

594 **infiltrate of the subserosa. CD transmural lesions: patients with these three lesions on**

595 **ileal margin. A: Whole cohort. B: Patients with no post-operative biotherapy.**

596

597 **Figure 3: Clinical recurrence-free survival depending on selected**

598 **histopathological lesions at ileal margins. Survival curves were built using**

599 **Kaplan-Meier method and crude log rank tests were calculated. Significance was**

600 **reached if $p < 0.05$.**

601

602




603

Patients' characteristics at time of surgery	Patients n=211 (%)
Men	92 (44%)
Median age (y, IQR)	36.3 (30.7 – 47.5)
Age at Crohn's Disease diagnosis (Montreal classification)	19 (9%)
- ≤16 years (A1)	170 (81%)
- 17 - 40 years (A2)	22 (10%)
- >40 years (A3)	
Median disease duration (y, IQR)	6.3 (1.8 – 11.9)
Median time between resection and colonoscopy (mo, IQR)	6.8 (6.1 – 8.4)
Smoking	67 (32%)
- Active smoker	13 (6%)
- Smoking cessation at surgery	131 (62%)
- Non smoker	
Previous intestinal resection	38 (18%)
- <10 years	14 (7%)
Number of previous resection(s)	173 (82%)
- 0	27 (13%)
- 1	7 (3%)
- 2	4 (2%)
- 3	
Surgical indication	115 (55%)
- Stricturing complication	83 (39%)
- Penetrating complication	13 (6%)
- Failure of drug therapy	
Disease location (Montreal classification)	132 (63%)
- Ileal (L1)	79 (37%)
- Ileocolonic (L3)	50 (24%)
- Anoperineal lesion	
Extra-digestive symptoms	40 (19%)
- Joint manifestations	12 (6%)
- Skin manifestations	4 (2%)
- Eye manifestations	
Previous exposure to anti-TNF therapy	134 (64%)
Anti-TNF therapy within 3 months before surgery	107 (51%)
Previous exposure to thiopurines	145 (69%)
Thiopurines within 3 months before surgery	60 (28%)

MICROSCOPIC DATA	Histological lesions	Prevalence (affected/analyzed)	Endoscopic recurrence (%)	<i>p</i>
Normal margin		65/211 (30.8%)	46.2	0.54
Margin affected		146/211 (69.2%)	50.7	0.54
<i>Mucosal layer</i>		111/211 (52.6%)	51.4	0.58
	Granuloma	18/211 (8.5%)	55.6	0.58
	Eosinophilic Infiltrate	68/182 (37.4%)	48.5	0.88
	Ulceration/Cryptitis	60/211 (28.4%)	58.3	0.09
	Villous Atrophy	20/211 (9.5%)	55.0	0.59
	Pyloric Metaplasia	43/211 (20.5%)	51.2	0.78
<i>Submucosal layer</i>		104/210 (49.3%)	52.9	0.30
	Neutrophilic Infiltrate	16/210 (7.6%)	62.5	0.27
	Lymphoplasmocytic Infiltrate	84/210 (40.0%)	53.6	0.31
	Plexitis	81/210 (38.6%)	48.1	0.79
	Nervous Hyperplasia	17/210 (8.1%)	52.9	0.75
	Fibrosis	71/210 (33.8%)	59.1	0.04
<i>Subserosal and muscularis layer</i>		92/211 (43.6%)	50.5	0.78
	Lymphoplasmocytic Infiltrate	58/206 (28.2%)	60.3	0.04
	Fibrosis	22/206 (10.7%)	59.1	0.33
	Myenteric Plexitis Grade 2	74/211 (35.1%)	50.0	0.77
	Myenteric Plexitis Grade 3	38/211 (18.0%)	50.0	0.92
MACROSCOPIC DATA				


Margin macroscopically affected	44/211 (20.9%)	56.1	0.31
	Median in non recurrent patients (cm)	Median in recurrent patients (cm)	p
Length of ileal resection	19.0	18.8	0.96
Distance between first macroscopic ulceration and ileal margin	4.0	4.0	0.43

	Correlation with endoscopic recurrence (multivariate analysis)			Correlation with clinical long term recurrence (multivariate analysis)		
	<i>Odds Ratio</i>	<i>95% CI</i>	<i>p</i>	<i>Odds Ratio</i>	<i>95% CI</i>	<i>p</i>
Age	0.98	0.95 - 1.01	0.082	1.01	0.99 - 1.03	0.320
Gender (male)	2.31	1.26 - 4.32	0.008	1.39	0.85 - 2.28	0.189
Previous surgery	5.46	2.22 - 14.57	< 0.001	1.86	1.01 - 3.27	0.046
Active smokers	2.78	1.47 - 5.39	< 0.001	1.69	1.15 - 2.95	0.011
Postoperative anti-TNF	0.42	0.21 - 0.84	0.016	NA	NA	NA
Extradigestive manifestations	NA	NA	NA	2.11	1.19 - 3.63	0.010
CD transmural lesions	3.83	1.47 - 11.05	0.008	2.04	1.09 - 3.99	0.026





A

Endoscopic Recurrence Rate

**B**

Endoscopic Recurrence Rate





n: Without CD transmural lesions	147	143	110	72	39	22	10	5	2
n: With CD transmural lesions	21	21	13	6	3				

Need to Know

Background: We aimed to characterize histologic features of the ileal margin in patients undergoing surgery for Crohn's disease (CD) and to evaluate their association with recurrence.

Findings: In patients with CD, transmural lesions at the ileal margin were associated with increased risk of post-operative disease recurrence.

Implications for Patient Care: Histologic features of the ileal margin should be included when physicians make decisions about post-operative therapy.

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 1: Flow chart

Supplementary Figure 2: Ven diagram: Distribution of the whole cohort of patients (n = 211) according to the histological data: 28.9% of patients presented an ileal margin affected with CD-related lesions in the 3 layers of the bowel wall. 30.8% had a normal ileal margin.

Supplementary Figure 3: Distance between first macroscopical ulceration and ileal margin correlated to histological data. No lesion: patients with normal ileal margin. One layer: lesions affecting one layer of the intestinal wall. Two layers: lesions affecting at least two layers of the intestinal wall. CD transmural lesions: presence of mucosal ulceration or cryptitis, submucosal fibrosis and lymphoplasmocytic infiltrate in the subserosa. Red points stand for patients who experienced endoscopic recurrence and blue points for patients who did not. Correlation test performed using Spearman's rank correlation test.