



HAL
open science

Volatile composition of six horsetails: prospects and perspectives

Françoise Fons, Didier Froissard, Jean-Marie Bessière, Alain Fruchier, Bruno Buatois, Sylvie Rapior

► **To cite this version:**

Françoise Fons, Didier Froissard, Jean-Marie Bessière, Alain Fruchier, Bruno Buatois, et al.. Volatile composition of six horsetails: prospects and perspectives. *Natural Product Communications*, 2013, 8 (4), pp.509-512. 10.1177/1934578X1300800426 . hal-02194690

HAL Id: hal-02194690

<https://hal.science/hal-02194690>


Submitted on 25 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



Volume 8. Issue 4. Pages 427-552. 2013
ISSN 1934-578X (printed); ISSN 1555-9475 (online)
www.naturalproduct.us

EDITOR-IN-CHIEF**DR. PAWAN K AGRAWAL**

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us

EDITORS**PROFESSOR ALEJANDRO F. BARRERO**

Department of Organic Chemistry,
University of Granada,
Campus de Fuente Nueva, s/n, 18071, Granada, Spain
afbarre@ugr.es

PROFESSOR ALESSANDRA BRACA

Dipartimento di Chimica Bioorganica e Biofarmacia,
Università di Pisa,
via Bonanno 33, 56126 Pisa, Italy
braca@farm.unipi.it

PROFESSOR DEAN GUO

State Key Laboratory of Natural and Biomimetic Drugs,
School of Pharmaceutical Sciences,
Peking University,
Beijing 100083, China
gda5958@163.com

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimaki@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry
University of Wollongong
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry,
Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry
The University of Alabama in Huntsville
Huntsville, AL 35809, USA
wsetzer@chemistry.uah.edu

PROFESSOR YASUHIRO TEZUKA

Institute of Natural Medicine
Institute of Natural Medicine, University of Toyama,
2630-Sugitani, Toyama 930-0194, Japan
tezuka@inm.u-toyama.ac.jp

PROFESSOR DAVID E. THURSTON

Department of Pharmaceutical and Biological Chemistry,
The School of Pharmacy,
University of London, 29-39 Brunswick Square,
London WC1N 1AX, UK
david.thurston@pharmacy.ac.uk

HONORARY EDITOR**PROFESSOR GERALD BLUNDEN**

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Berhanu M. Abegaz
Gaborone, Botswana

Prof. Viqar Uddin Ahmad
Karachi, Pakistan

Prof. Øyvind M. Andersen
Bergen, Norway

Prof. Giovanni Appendino
Novara, Italy

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Lee Banting
Portsmouth, U.K.

Prof. Julie Banerji
Kolkata, India

Prof. Anna R. Bilia
Florence, Italy

Prof. Maurizio Bruno
Palermo, Italy

Prof. César A. N. Catalán
Tucumán, Argentina

Prof. Josep Coll
Barcelona, Spain

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Ana Cristina Figueiredo
Lisbon, Portugal

Prof. Cristina Gracia-Viguera
Murcia, Spain

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. Kurt Hostettmann
Lausanne, Switzerland

Prof. Martin A. Iglesias Arteaga
Mexico, D. F., Mexico

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Vladimir I Kalinin
Vladivostok, Russia

Prof. Niel A. Koorbanally
Durban, South Africa

Prof. Karsten Krohn
Paderborn, Germany

Prof. Chiaki Kuroda
Tokyo, Japan

Prof. Hartmut Laatsch
Gottingen, Germany

Prof. Marie Lacaille-Dubois
Dijon, France

Prof. Shoei-Sheng Lee
Taipei, Taiwan

Prof. Francisco Macias
Cadiz, Spain

Prof. Imre Mathe
Szeged, Hungary

Prof. Ermino Murano
Trieste, Italy

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Luca Rastrelli
Fisciano, Italy

Prof. Monique Simmonds
Richmond, UK

Dr. Bikram Singh
Palampur, India

Prof. John L. Sorensen
Manitoba, Canada

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Sylvia Urban
Melbourne, Australia

Prof. Karen Valant-Vetschera
Vienna, Austria

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2013 subscription price: US\$2,395 (Print, ISSN# 1934-578X); US\$2,395 (Web edition, ISSN# 1555-9475); US\$2,795 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Volatile Composition of Six Horsetails: Prospects and Perspectives

Françoise Fons^{a*}, Didier Froissard^b, Jean-Marie Bessière^c, Alain Fruchier^d, Bruno Buatois^c and Sylvie Rapior^a^aLaboratoire de Botanique, Phytochimie et Mycologie, Faculté de Pharmacie (Université Montpellier 1), UMR 5175 CEFE, B.P. 14 491, 15 avenue Charles Flahault, F-34093 Montpellier Cedex 5, France^bLaboratoire de Botanique, Faculté de Pharmacie de Limoges, 2 rue du Docteur Raymond Marcland, F-87025 Limoges Cedex, France^cCentre d'Ecologie Fonctionnelle et Evolutive – Plate-forme d'analyses chimiques en écologie, UMR 5175 CEFE, 1919 Route de Mende, F-34293 Montpellier Cedex 5, France^dENSCM, UMR 5253, 8 Rue de l'Ecole Normale, F-34296 Montpellier Cedex 5, France

francoise.fons@univ-montp1.fr

Received: November 20th, 2012; Accepted: February 18th, 2013

Six horsetails were investigated for volatile organic compounds (VOC) by GC-MS using organic solvent extraction. Seventy-five VOC biosynthesized from the shikimic, lipidic and terpenic pathways including isoprenoid derivatives were detected from these putative natural resources. *E. palustre* var. *americana* contained mainly lipidic derivatives, i.e., 1-octen-3-ol (mushroom-like odor), (*E*)-2-hexenoic acid (fruity odor) and (*E*)-2-hexenal (green odor). Many isoprenoid flavour precursors, i.e., 3-oxo- α -ionol (spicy odor) and (*E,E*)-pseudoionone (balsamic odor), as well as odorous benzenic derivatives, i.e., phenylethanal (hyacinth, lilac note) and 2-phenylethanol (rose odor) contributed to the odor of *E. arvense*. The volatile pattern of *E. telmateia* is dominated by high amounts of isoprenoids and benzenic derivatives. The complex volatile profiles of *E. hyemale* and *E. ramosissimum* are based on ferulic acid isomers, along with either (*E*)-2-heptenal (green vegetable-like odor) or 4-vinylguaiaicol (spicy clove smoky odor) for *E. hyemale* and *E. ramosissimum*, respectively. The broad spectrum of *E. scirpioides* shows the lowest VOC content with high amount of isoprenoids (46.9%), mainly ionone derivatives. *Equisetum* resources are of great interest as bioactive litter and new potential functional feed ingredients.

Keywords: Horsetails, *Equisetum*, Volatile Organic Compounds, (*E*)-2-Heptenal, 4-Vinylguaiaicol, Ferulic acid isomers, (*E,E*)-Pseudoionone, 3-Oxo- α -ionol.

Horsetails have a subcosmopolitan distribution. The fifteen well-known *Equisetum* species always grow in wet sites and form large long-lived populations via highly developed rhizomes [1a,1b]. In the new phylogenetic classification, they belong to Monilophytes, where they are closer to ferns than in the previous classifications [1c,1d]. *Equisetum* species biosynthesize a great number of various phytochemicals, which may possess health-promoting properties. Horsetails have been frequently used in medicine since Greek antiquity to treat wounds and inflammations of the skin and to cure fractures, thanks to the action of silicic acid and various identified minerals [2a]. Recently *E. arvense*, commonly used as a traditional Polish medicine, was investigated for silicon from whole plant and tea bags [2b]. Nowadays several species of horsetails are still used in traditional medicine around the world, i.e., *E. debile* against jaundice and hepatitis, as well as for treatment of urinary tract problems, including kidney stones [2c,2d], *E. giganteum* as a diuretic [2e], and *E. telmateia* as an antiseptic [2f]. *E. ramosissimum*, *E. arvense* and *E. telmateia* were investigated for antioxidant activity, mainly due to polyphenols [2g-2k], and *E. hyemale* for both antioxidant and anti-inflammatory properties [2l]. Recently, anti-cancer activity of *E. hyemale* extract was demonstrated on murine leukemia L1210 cells [2m]. *E. arvense* and *E. palustre* were recently utilized in cosmetics to regenerate the skin and stabilize the connective tissue due to flavonoid and saponin derivatives [3a,3b]. Nowadays *Equisetaceae* extracts are patented for uses in skin treatments and capillary preparations for hair problems [3c,3d].

However, very few data have been reported on volatile organic compounds (VOC) of horsetails and their biological activities. Two

research teams published on wild *Equisetum* essential oils from China and Serbia, respectively [4a-4d]. Within the framework of our investigations for natural volatiles in mushrooms and plants [4e-4j], especially in ferns in the Monilophytes group, six wild horsetails were investigated for VOC.

Fresh aerial parts of six *Equisetum* species were investigated for their volatile profiles using GC-MS: *E. arvense* L., *E. palustre* var. *americana* Vict. and *E. telmateia* Ehrh. from the *Equisetum* subgenus; *E. hyemale* L., *E. ramosissimum* Desf. and *E. scirpioides* Michx. from the *Hippochaete* subgenus. The main volatile components of *Equisetum* sp. were identified by comparison with the National Institute of Standards and Technology Mass Spectral Library [5a], retention indices [5b] as reported in the literature for the eleven French ferns previously studied [4i,4j].

Seventy-five VOC biosynthesized from the shikimic, lipidic and terpenic pathways were detected and sixty-five identified from the concentrated diethyl ether extracts of the six horsetails (Table 1).

Twenty-one VOC participated in the general odor of *E. arvense*. This plant contained the highest biodiversity of isoprenoid flavor precursors (75.8%), i.e., 3-hydroxy-7,8-epoxy- β -ionol (17.5%), (*E,E*)-pseudoionone (13.1%) and 3-oxo- α -ionol (9.8%; Table 1) [6a,6b], as well as odorous benzenic derivatives, i.e., phenylethanal (3.8%; hyacinth, lilac note), 2-phenylethanol (3.8%; rose odor), benzaldehyde (2.2%; bitter almond odor) and homovanillic acid (3.8%). The last compound is well-known as one of the quercetin metabolites from *E. arvense* extract when administered to humans [6c].

Table 1: Percentage of volatile organic compounds^a in fresh aerial part of horsetails.

Compounds	RI ^b	<i>Equisetum arvense</i>	<i>Equisetum palustre</i> var. <i>americana</i>	<i>Equisetum telmateia</i>	<i>Equisetum hyemale</i>	<i>Equisetum ramosissimum</i>	<i>Equisetum scirpioides</i>
Aromatic compounds		18.0	15.3	26.3	25.5	76.6	34.8
Benzaldehyde	962	2.2	1.2				0.5
Benzylalcohol	1044			4.4			
Phenylethanal	1046	3.8	1.0	1.4		1.9	6.5
Acetophenone	1068		3.5				
2-Phenylethanol	1117	3.8	2.0			6.5	1.8
Benzoic acid	1187		1.2				2.9
Benzothiazole	1228		0.5				0.8
Phenylacetic acid	1266		1.5			1.9	1.2
4-Vinylguaiaicol	1316	2.2		18.1	5.9	19.6	2.3
Isovanillin	1404	2.2	1.0	2.4	0.7		
(Z)-3-Hexenyl benzoate	1552		0.3				3.5
Homovanillic acid	1561	3.8					15.2
(Z)-Ferulic acid	1777				8.6	18.7	
(E)-Ferulic acid	1889				10.3	28.0	
Esculetine	1890		3.3				
Lipid derivatives		6.2	58.8	3.7	69.7	20.2	18.3
(E)-2-Hexenal	851		6.6				
(Z)-3-Hexenol	855		2.5				
(E)-2-Hexenol	866		2.0				
Hexanol	870		1.0				
(E)-2-Heptenal	956		2.1		22.4	2.3	4.1
1-Octen-3-one	977		0.8				
1-Octen-3-ol	984		12.4				
2,3-Octanedione	985	1.6	0.8	1.4			0.3
3-Octanol	1000		1.3				
(E,Z)-2,4-Heptadienal	1001				1.7		3.8
Hexanoic acid	1009	2.2			0.7	0.9	1.2
(E,E)-2,4-Heptadienal	1013				1.0		0.9
(Z)-3-Hexenoic acid	1017	1.1	9.9	2.3			
(E)-2-Hexenoic acid	1037	1.3	10.7				
NI ^c	1041				4.5		
NI ^c	1064				3.8		
(E)-2-Octenol	1072		1.3				
Allyl 3-oxo-butanoate	1075				2.1		
Heptanoic acid	1096		0.3		3.1	1.9	
6-Hydroxy-6-methylheptan-2-one	1109				6.2		
Allylheptanoate	1123				2.8		
(Z)-3-Nonenol	1158		0.7				
(E)-2-Nonenal	1162		0.8				
Ethylmethylmaleimide	1242		0.7		1.0		1.2
NI ^c	1256				3.1		
(E)-2-Decenal	1263					7.1	1.1
Methylvinylmaleimide	1273		0.3		0.3		0.6
Nonanoic acid	1283		1.3			1.9	
(E,Z)-2,4-Decadienal	1296					0.9	1.1
2,4,4-Trimethyl-2-butenolide	1298				2.4		
NI ^c	1301				3.8		
NI ^c	1316				6.2		
(E,E)-2,4-Decadienal	1321					2.3	4.1
NI ^c	1348				2.2		
2,4-Decanedione	1373				2.4		
Lauric acid	1570		1.3			2.8	
Myristic acid	1773		2.0				
Terpenes		0.0	0.7	2.9	0.0	0.0	0.0
Linalool	1102		0.7				
β -Caryophyllene	1422						
Isoprenoid derivatives		75.8	25.2	70.6	2.4	3.3	46.9
1,2-Dimethylcyclohexanol	1113		0.7				
1-Acetyl-2,2-dimethylcyclohexane	1257				2.4		
α -Ionone	1433		0.2	1.4			
Dihydroactinidiolide	1539	4.4					
(E,Z)-Pseudoionone	1596	4.9	2.1				
NI ^c	1603		1.2				
(E,E)-Pseudoionone	1611	13.1	1.8	2.3			
NI ^c	1617	0.6	1.3				
3-Oxo-7,8-dihydro- α -ionone	1628		1.2	5.6			
3-Hydroxy- β -ionol	1640	8.2	1.5				
3-Hydroxy- α -ionone	1645		1.8	10.3			5.0
NI ^c	1648		3.3	15.0		2.3	7.0
4-Hydroxy- β -ionol ^d	1653	2.2	1.3				
3-Oxo- α -ionol	1658	9.8		6.8			
3-Hydroxy-7,8-dihydro- β -ionol	1659	4.4					
3-Hydroxy- β -ionone	1668						13.5
3-Hydroxy-7,8-epoxy- β -ionol	1678	17.5	4.3	25.4		0.9	15.5
4-Oxo- β -ionone	1680	4.4					
4-Hydroxy-7,8-dihydro- β -ionol	1694	6.5	4.6	3.8			
3-Hydroxy-7,8-epoxy- α -ionone	1755						2.9
NI ^c	1790						2.9

^a Relative percentage of the VOC based on the GC-MS chromatographic area; ^b RI = Retention Indices on SLBTM-5MS column (Supelco); ^c NI = not identified; ^d = tentatively identified.

The volatile profile of *E. palustre* var. *americana* showed forty-four VOC with mostly lipidic derivatives (58.8%), i.e., 1-octen-3-ol (12.4%; mushroom-like odor), (*E*)-2-hexenoic acid (10.7%; fruity odor), as well as odorous compounds such as (*E*)-2-hexenal (6.6%; green odor), (*Z*)-3-hexenol (2.5%; grass, fresh odor) [6d], hexanol (1%; balsamic odor), and (*E*)-2-nonenal (0.8%; fatty, green odor) [6e]. Only one terpenic derivative was identified, i.e., linalool (0.7%; floral, orange odor). The 1-octen-3-ol responsible for the mushroom-like odor has also been reported from many mushrooms [6f,6g], ferns [4i,4j] and Angiosperms [6h]. (*E*)-2-Hexenoic acid (10.7%), with a fruity, herbal odor, has also been found in both ferns [4i,4j] and mushrooms [4h].

The volatile pattern of *E. telmateia* based on fifteen VOC was dominated by a large number of isoprenoid derivatives (70.6%), i.e., 3-hydroxy-7,8-epoxy- β -ionol (25.4%) and 3-hydroxy- α -ionone (10.3%), as well as one benzenic derivative, i.e., 3-methoxy-4-hydroxystyrene also called 4-vinylguaiaicol (18.1%; spicy clove smoky odor).

Ionol and ionone derivatives were identified from quince leaves [7a] and ferns [4i,4j]; α -ionone shows a more floral *smell* than β -ionone. Isoprenoids, widely produced by animals, microbes and plants, are of great interest in the industrial production of pharmaceuticals, colorants and flavors. Research has focused on their transformation into functional isoprenoids [7b].

The major VOC from *E. hyemale* derived from the lipidic pathway (69.7%), such as (*E*)-2-heptenal (22.4%; green vegetable-like odor with fatty undertone), and the shikimic pathway (25.5%), i.e., (*E*)- and (*Z*)-ferulic acid isomers (10.3% and 8.6%, respectively), 4-vinylguaiaicol (5.9%) and isovanillin (0.7%, vanilla odor). Ferulic acid derivatives are converted into 4-vinylguaiaicol by decarboxylation, then oxidized to vanillin [7c].

Cinnamic acid derivatives have received much attention in medicinal research, more recently as synthetic antitumor agents [7d,7e].

E. ramosissimum showed the highest amount of shikimic derivatives (76.6%), i.e., (*E*)-ferulic acid (28.0%), 4-vinylguaiaicol (19.6%), (*Z*)-ferulic acid (18.7%), and 2-phenylethanol (6.5%; rose odor). Lipid derivatives, such as (*E*)-2-decenal (7.1%), also contributed to the complex odor of *E. ramosissimum*.

The volatile profiles of *E. ramosissimum* and *E. hyemale*, both belonging to the *Hippochaete* subgenus, showed similarities, with both producing the same main shikimic compounds (ferulic acid isomers and 4-vinylguaiaicol). 2-Phenylethanol and (*E*)-2-decenal were previously detected in several ferns (*Adiantum capillus-veneris*, *Blechnum spicant* and *Asplenium trichomanes*) [4i,4j]. In addition, (*E*)-2-decenal, a natural plant and mushroom VOC with a plastic oxidized mutton fatty odor was an abundant key component of the strong "stink bug" odor with (*E,Z*)-2,4-decadienal and (*E,E*)-2,4-decadienal [7f]; these three VOC were also found in *A. capillus-veneris*, *E. ramosissimum* and *E. scirpioides*.

E. scirpioides showed the lowest VOC content with a broad spectrum based on twenty-five VOC. High amounts of isoprenoids (46.9%), mainly ionone derivatives, were identified as 3-hydroxy- β -ionone (13.5%) or 3-hydroxy-7,8-epoxy- β -ionol (15.5%). Lower amounts of aromatics (34.8%), i.e., homovanillic acid (15.2%) and phenylethanol (6.5%; hyacinth, lilac note) [4f,4h], already found in *Gymnocarpium dryopteris* and *Phegopteris connectilis* [4j], as well

as lipid derivatives, i.e., (*E*)-2-heptenal (4.1%), (*E,E*)-2,4-decadienal (4.1%; orange-like odor) and (*E,Z*)-2,4-heptadienal (3.8%) complete the volatile pattern of *E. scirpioides*.

Among the six *Equisetum* species investigated for volatiles belonging either to *Hippochaete* or *Equisetum* subgenera, the quantitatively major volatiles were aromatic compounds, i.e., 4-vinylguaiaicol, homovanillic acid, and (*Z*)- and (*E*)-ferulic acid isomers, which were particularly present in the *Hippochaete* subgenus. Ferulic acid, a precursor of vanillin with antioxidant activity, may be effective in preventing cancer [7g]. 3-Hydroxy-7,8-epoxy- β -ionol and 3-hydroxy- β -ionone were the isoprenoid derivatives detected in the highest amount. Another megastigmane glucoside, i.e., (6*R*,9*S*)-3-oxo- α -ionol 9-*O*- β -D-glucopyranoside, was previously reported in *E. debile* [7h]. On the other hand, lipids (C₆-C₁₀) constituted the most diversified VOC content in horsetails with more than 20% of (*E*)-2-heptenal in *E. hyemale*.

Conclusion

Equisetum resources have great potential as new bioactive litter and potential functional feed ingredients due to the importance of their biologically active constituents, for example silicon, phenolic derivatives and volatile organic compounds [2a-2b,4b,8a,8b].

In addition, each investigated *Equisetum* species synthesizes numerous volatiles known to exhibit insecticidal, pesticidal and/or arthropod repellent properties. These plants may be useful sources of chemicals for the control of micro-organisms of medical, veterinary and agricultural importance. Except *E. palustre*, which is a well-known livestock poison, containing the alkaloid palustrine, *Equisetum* natural resources are of major importance as new potential functional feed ingredients and bioactive litter.

Experimental

Plant material: Fresh aerial parts of *Equisetum arvense* L., *E. palustre* var. *americana* Vict., *E. ramosissimum* Desf., and *E. scirpioides* Michx. were collected on 25th November, 2009 from the Botanical Garden of Limoges. *E. telmateia* Ehrh and *E. hyemale* L. were obtained on 31st August, 2010, from the Botanical Garden of Strasbourg. Voucher specimens are deposited at the Laboratory of Botany (Faculty of Pharmacy, Limoges, France).

Plant part and GC-MS analyses: Fresh aerial parts of *Equisetum* were cubed and extracted with diethyl ether (Carlo Erba, 6 ppm BHT). After one-week of maceration at room temperature, the concentrated organic extracts were used for Gas Chromatography-Mass Spectrometry (GC-MS) analyses, as reported in the literature [4i,4j]. The analyses were performed at the Platform for Chemical Analyses in Ecology of the "SFR 119 Montpellier Environnement Biodiversité".

VOC were identified by comparison with National Institute of Standards and Technology Mass Spectral Library [5a], retention indices listed in the literature [5b], and our own data bank. Internal standards (*n*-alkanes) were used as reference points in the calculation of relative retention indices.

Acknowledgments - The authors greatly thank the Botanical Garden of Limoges for providing *Equisetum arvense*, *E. palustre* var. *americana*, *E. ramosissimum* and *E. scirpioides*; and the Botanical Garden of Strasbourg for providing *E. hyemale* and *E. telmateia*.

References

- [1] (a) Prelli R, Boudry M. (2001) *Les fougères et plantes alliées de France et d'Europe occidentale*. Belin Publisher, Paris; (b) Kenrick P, Crane PR. (1997) The origin and early evolution of plants on land. *Nature*, **389**, 33-39; (c) Pryer KM, Schneider H, Smith AR, Cranfill R, Wolf PG, Hunt JS, Sipes SD. (2001) Horsetails and ferns are a monophyletic group and the closest living relatives to seed plants. *Letters to Nature*, **409**, 618-622; (d) Prelli R. (2010) La classification phylogénétique des ptéridophytes actuelles. *Journal de Botanique de la Société Botanique de France*, **49**, 37-47.
- [2] (a) Sandhu NS, Kaur S, Chopra D. (2010) *Equisetum arvense*: pharmacology and phytochemistry - a review. *Asian Journal of Pharmaceutical and Clinical Research*, **3**, 146-150; (b) Pajchel L, Nykiel P, Kolodziejski W. (2011) Elemental and structural analysis of silicon forms in herbal drugs using silicon-29 MAS NMR and WD-XRF spectroscopic methods. *Journal of Pharmaceutical and Biomedical Analysis*, **56**, 846-850; (c) Abbasi AM, Khan MA, Ahmad M, Zafar M, Khan H, Muhammad N, Sultana S. (2009) Medicinal plants used for the treatment of jaundice and hepatitis based on socio-economic documentation. *African Journal of Biotechnology*, **8**, 1643-1650; (d) Chauhan P, Kumar D, Kasana MS. (2009) Medicinal plants of Muzaffargarh district used in treatment of urinary tract and kidney stones. *Indian Journal of Traditional Knowledge*, **8**, 191-195; (e) Hernández MP, Colares MN, Civitella SM. (2009) Plantas utilizadas en medicina popular en un sector del Partido de Berisso, Buenos Aires, Argentina. *Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas*, **8**, 435-444; (f) Uzun E, Sariyar G, Adersen A, Karakoc B, Otük G, Oktayoglu E, Pirildar S. (2004) Traditional medicine in Sakarya province (Turkey) and antimicrobial activities of selected species. *Journal of Ethnopharmacology*, **95**, 287-296; (g) Čanadanović-Brunet JM, Četković GS, Djilas SM, Tumbas VT, Savatović SS, Mandić AI, Markov SL, Cvetković DD. (2009) Radical scavenging and antimicrobial activity of horsetail (*Equisetum arvense* L.) extracts. *International Journal of Food Science and Technology*, **44**, 269-278; (h) Cetojević-Simin DD, Čanadanović-Brunet JM, Bogdanović GM, Djilas SM, Četković GS, Tumbas VT, Stojiljković BT. (2010) Antioxidative and antiproliferative activities of different horsetail (*Equisetum arvense* L.) extracts. *Journal of Medicinal Food*, **13**, 452-459; (i) Štajner D, Popović BM, Čanadanović-Brunet J, Boža P. (2006) Free radical scavenging activity of three *Equisetum* species from Fruškagora mountain. *Fitoterapia*, **77**, 601-604; (j) Štajner D, Popović BM, Čanadanović-Brunet, Anackov G. (2009) Exploring *Equisetum arvense* L., *Equisetum ramosissimum* L. and *Equisetum telmateia* L. as sources of natural antioxidants. *Phytotherapy Research*, **23**, 546-550; (k) Correia H, González-Paramás A, Amaral MT, Santos-Buelga C, Batista MT. (2005) Characterization of polyphenols by HPLC-PAD-ESI/MS and antioxidant activity in *Equisetum telmateia*. *Phytochemical Analysis*, **16**, 380-387; (l) Park EY, Jeon H. (2008) Antioxidant and anti-inflammatory activities of *Equisetum hyemale*. *Natural Product Sciences*, **14**, 239-243; (m) Li H, Wang P, Liu Q, Cheng X, Zhou Y, Xiao Y. (2012) Cell cycle arrest and cell apoptosis induced by *Equisetum hyemale* extract in murine leukemia L1210 cells. *Journal of Ethnopharmacology*, **144**, 322-327.
- [3] (a) Yesilada E, Gurbuz I. (2010) Evaluation of the antiulcerogenic activity profile of a flavonol diglucoside from *Equisetum palustre* L., *Journal of Ethnopharmacology*, **131**, 17-21; (b) Ozay Y, Ozyurt S, Guzel S, Cimbiz A, Olgun EG, Cayci MK. (2010) Effects of *Equisetum arvense* ointment on dermal wound healing in rats. *Wounds: A Compendium of Clinical Research and Practice*, **22**, 261-267; (c) Fukuda R, Kiden A. (2001) Hair preparations containing *Equisetum arvense* extracts for prevention of gray hair. *Japanese Kokai Tokkyo Koho*, N° JP2001288044 (10 pp); (d) Cockbill S. (2010) Commentary: Effects of *Equisetum arvense* ointment on dermal wound healing in rats. *Wounds: A Compendium of Clinical Research and Practice*, **22**, 268-269.
- [4] (a) Zhao C, Zeng Y, Wan M, Li R, Liang Y, Li C, Zeng Z, Chau FT. (2009) Comparative analysis of essential oils from eight herbal medicines with pungent flavor and cool nature by GC-MS and chemometric resolution methods. *Journal of Separation Science*, **32**, 660-670; (b) Radulović N, Stojanović G, Palić R. (2006) Composition and antimicrobial activity of *Equisetum arvense* L. essential oil. *Phytotherapy Research*, **20**, 85-88; (c) Radulović N, Stojanović G, Milovanović V, Doković D, Radelović V. (2008) Volatile constituents of *Equisetum fluviatile* L. *Journal of Essential Oil Research*, **20**, 437-441; (d) Milovanović V, Radulović N, Mitić V, Palić R, Stojanović G. (2008) Chemical composition of the essential oils of *Equisetum palustre* L. and *Equisetum telmateia* Ehrh. *Journal of Essential Oil Research*, **20**, 310-314; (e) Breheret S, Talou T, Rapior S, Bessièrre JM. (1999) Geosmin, a sesquiterpenoid compound responsible for the musty-earthy odor of *Cortinarius herculeus*, *Cystoderma amianthinum*, and *Cy. carcharias*. *Mycologia*, **91**, 117-120; (f) Rapior S, Kanska G, Guillot J, Andary C, Bessièrre JM. (2000) Volatile composition of *Laetiporus sulphureus*. *Cryptogamie, Mycologie*, **21**, 67-72; (g) Rapior S, Fons F, Bessièrre JM. (2000) The fenugreek odor of *Lactarius helvus*. *Mycologia*, **92**, 305-308; (h) Fons F, Rapior S, Eyssartier G, Bessièrre JM. (2003) Les substances volatiles dans les genres *Cantharellus*, *Craterellus* et *Hydnum*. *Cryptogamie, Mycologie*, **24**, 367-376; (i) Fons F, Froissard D, Bessièrre JM, Buatois B, Rapior S. (2010) Biodiversity of volatile organic compounds from five French ferns. *Natural Product Communications*, **5**, 1655-1658; (j) Froissard D, Fons F, Bessièrre JM, Buatois B, Rapior S. (2011) Volatiles of French ferns and "fougère" scent in perfumery. *Natural Product Communications*, **6**, 1723-1726.
- [5] (a) National Institute of Standard and Technology. (2005) *PC version of the NIST / EPA / NIH Mass Spectra Database*, Gaithersburg, Maryland, USA; (b) Adams RP. (2007) *Identification of essential oil components by gas chromatography / mass spectroscopy*. 4th Ed., Allured, Carol Stream, IL, USA.
- [6] (a) Humpf HU, Schreier P. (1992) 3-Hydroxy-5,6-epoxy- β -ionol β -D-glucopyranoside and 3-hydroxy-7,8-dihydro- β -ionol β -D-glucopyranoside: New C₁₃ norisoprenoid glucoconjugates from sloe tree (*Prunus spinosa* L.) leaves. *Journal of Agricultural and Food Chemistry*, **40**, 1898-1901; (b) Herderich M, Neubert C, Winterhalter P, Schreier P, Skouroumounis GK. (1992) Identification of C₁₃-norisoprenoid flavour precursors in starfruit (*Averrhoa carambola* L.). *Flavour and Fragrance Journal*, **7**, 179-185; (c) Graefe E, Veit M. (1999) Urinary metabolites of flavonoids and hydroxycinnamic acids in humans after application of a crude extract from *Equisetum arvense*. *Phytomedicine*, **6**, 239-246; (d) Mannschreck A, von Angerer E. (2011) The scent of roses and beyond: Molecular structures, analysis, and practical applications of odorants. *Journal of Chemical Education*, **88**, 1501-1506; (e) Kula J, Sadoswka H. (1993) Unsaturated aliphatic C₉-aldehydes as natural flavorants. *Perfumer and Flavorist*, **18**, 23-25; (f) Rapior S, Breheret S, Talou T, Pélissier Y, Bessièrre JM. (2002) The anise-like odor of *Clitocybe odora*, *Lentinellus cochleatus* and *Agaricus essettei*. *Mycologia*, **94**, 373-376; (g) Rapior S, Breheret S, Talou T, Pélissier Y, Milhau M, Bessièrre JM. (1998) Volatile components of fresh *Agrocybe aegerita* and *Tricholoma sulfureum*. *Cryptogamie, Mycologie*, **19**, 15-23; (h) Fons F, Rapior S, Gargadennec A, Andary C, Bessièrre JM. (1998) Volatile components of *Plantago lanceolata* (Plantaginaceae). *Acta botanica Gallica*, **145**, 265-269.
- [7] (a) Lutz-Röder A, Schneider M, Winterhalter P. (2002) Isolation of two new ionone glucosides from quince (*Cydonia oblonga* Mill.) leaves. *Natural Product Letters*, **16**, 119-122; (b) Misawa N. (2011) Pathway engineering for functional isoprenoids. *Current Opinion in Biotechnology*, **22**, 627-633; (c) Tsujiyama SI, Ueno M. (2008) Formation of 4-vinylguaiaicol as an intermediate in bioconversion of ferulic acid by *Schizophyllum commune*. *Biosciences, Biotechnology and Biochemistry*, **72**, 212-215; (d) De P, Baltas M, Bedos-Belval F. (2011) Cinnamic acid derivatives as anticancer agents - A review. *Current Medicinal Chemistry*, **18**, 1672-1703; (e) Ou S, Kwok K-C. (2004) Ferulic acid: pharmaceutical functions, preparation and applications in foods. *Journal of the Science of Food and Agriculture*, **84**, 1261-1269; (f) Ho HY, Kou R, Tseng HK. (2003) Semicemicals from the predatory stink bug *Eocanthe conofurcellata* (Wolff): components of metathoracic gland, dorsal abdominal gland, and sternal gland secretions. *Journal of Chemical Ecology*, **29**, 2101-2114; (g) Srinivasan M, Sudheer AR, Menon VP. (2007) Ferulic acid: therapeutic potential through its antioxidant property. *Journal of Clinical Biochemistry and Nutrition*, **40**, 92-100; (h) Kanchanapoom T, Otsuka H, Ruchirawat S. (2007) Megastigmane glucosides from *Equisetum debile* and *E. diffusum*. *Chemical Pharmaceutical Bulletin*, **55**, 1277-1280.
- [8] (a) Chigarina KM, Alaverdiev IM, Alaverdieva SI, Abdrashitov MA, Ivanyuk TB, Arakcheeva GA, Troshina LN. (2001) Antibacterial toilet soaps containing triclosan and extracts of burdock, calendula and horsetail. Patent No. RU2163255; (b) Zhou Q, Xu T, Zhang GR, Gu DX, Zhang WQ. (2003) Repellent effects of herbivore-induced rice volatiles on the brown planthopper *Nilaparvata lugens*. *Acta Entomologica Sinica*, **46**, 739-744.

Volatile Composition of Six Horsetails: Prospects and Perspectives Françoise Fons, Didier Froissard, Jean-Marie Bessière, Alain Fruchier, Bruno Buatois and Sylvie Rapior	509
Chemical Compositions of the Rhizome, Leaf and Stem Oils from Malaysian <i>Hornstedtia leonurus</i> Nor Akmalazura Jani, Hasnah Mohd. Sirat, NorAzah Mohamad Ali and Azrina Aziz	513
Effect on Emotional Behavior and Stress by Inhalation of the Essential oil from <i>Chamaecyparis obtusa</i> Hikaru Kasuya, Erika Hata, Tadaaki Satou, Masaki Yoshikawa, Shinichiro Hayashi, Yoshinori Masuo and Kazuo Koike	515
Chemical Composition and Antibacterial Activity of Rhizome Oils from Five <i>Hedychium</i> Species Ratchuporn Suksathan, Siriwoot Sookkhee, Somboon Anuntalabhochai and Sunee Chansakaow	519
Chemical Composition and Antimicrobial Activity of Three Essential Oils from <i>Curcuma wenyujin</i> Jingjing Zhu, Agnieszka D. Lower-Nedza, Meng Hong, Song Jiec, Zhimin Wang, Dong Yingmao, Christine Tschiggerl, Franz Bucar and Adelheid H. Brantner	523
Essential Oil Composition and Antimicrobial Activity of Aerial Parts and Ripe Fruits of <i>Echinophora spinosa</i> (Apiaceae) from Italy Daniele Fraternali, Salvatore Genovese and Donata Ricci	527
Composition and <i>in vitro</i> Anticancer Activities of the Leaf Essential Oil of <i>Neolitsea variabilissima</i> from Taiwan Yu-Chang Su, Kuan-Ping Hsu, Eugene I-Chen Wang and Chen-Lung Ho	531
<u>Review/Account</u>	
Natural Products from Marine Algae of the Genus <i>Osmundaria</i> (Rhodophyceae, Ceramiales) Kelvin Osako and Valéria Laneuville Teixeira	533
Phenols, Alkaloids and Terpenes from Medicinal Plants with Antihypertensive and Vasorelaxant Activities. A Review of Natural Products as Leads to Potential Therapeutic Agents Francesco Maione, Carla Cicala, Giulia Musciacco, Vincenzo De Feo, Anibal G. Amat, Armando Ialenti and Nicola Mascolo	539
Diosmin – Isolation Techniques, Determination in Plant Material and Pharmaceutical Formulations, and Clinical Use Anna Bogucka – Kocka, Michał Woźniak, Marcin Feldo, Janusz Kocki and Katarzyna Szewczyk	545

Natural Product Communications

2013

Volume 8, Number 4

Contents

<u>Original Paper</u>	<u>Page</u>
Anti-melanogenesis Constituents from the Seaweed <i>Dictyota coriacea</i> Ryeo Kyeong Ko, Min-Chul Kang, Sang Suk Kim, Tae Heon Oh, Gi-Ok Kim, Chang-Gu Hyun, Jin Won Hyun and Nam Ho Lee	427
Methyl Carnosate, an Antibacterial Diterpene Isolated from <i>Salvia officinalis</i> Leaves Elisa Klimati, Fabio Mastrogiovanni, Maria Valeri, Laura Salvini, Claudia Bonechi, Nilufar Zokirzhonovna Mamadalieva, Dilfuza Egamberdieva, Anna Rita Taddei and Antonio Tiezzi	429
Cytotoxicity of Meroterpenoids from <i>Sargassum siliquastrum</i> against Human Cancer Cells Jung Im Lee, Myoung K. Kwak, Hee Y. Park and Youngwan Seo	431
Isolation of Methyl 27-caffeoyloxyoleanolate – A New Oleanane Triterpenoid from the Roots of <i>Hibiscus vitifolius</i> Duraismy Ramasamy and Ariamuthu Saraswathy	433
Synthesis and Cytotoxic Activity of New Betulin and Betulinic Acid Esters with Conjugated Linoleic Acid (CLA) Barbara Tubek, Pawel Mitula, Natalia Niezgoda, Katarzyna Kempinska, Joanna Wietrzyk and Czeslaw Wawrzenczyk	435
Analysis of Pyrrolizidine Alkaloids and Evaluation of Some Biological Activities of Algerian <i>Senecio delphinifolius</i> (Asteraceae) Soukaina Tidjani, Philippe N. Okusa, Amar Zellagui, Laetitia Moreno Y Banuls, Caroline Stévigny, Pierre Duez and Salah Rhouati	439
Berbanine: a New Isoquinoline-isoquinolone Alkaloid from <i>Berberis vulgaris</i> (Berberidaceae) Anna Hošťálková, Zdeněk Novák, Milan Pour, Anna Jirošová, Lubomír Opletal, Jiří Kuneš and Lucie Cahliková	441
Dicentrine Production in Callus and Cell Suspension Cultures of <i>Stephania venosa</i> Tharita Kitisripanya, Jukrapun Komaikul, Nirachara Tawinkan, Chuennapha Atsawinkowit and Waraporn Putalun	443
New Flavan and Alkyl α,β-Lactones from the Stem Bark of <i>Horsfieldia superba</i> Nabil Ali Al-Mekhlafi, Khozirah Shaari, Faridah Abas, Ethyl Jeyaseela Jeyaraj, Johnson Stanslas, Shaik Ibrahim Khalivulla and Nordin H. Lajis	447
New Flavonol Triglycosides from the Leaves of Soybean Cultivars Yoshinori Murai, Ryoji Takahashi, Felipe Rojas Rodas, Junichi Kitajima and Tsukasa Iwashina	453
Melitidin: A Flavanone Glycoside from <i>Citrus grandis</i> ‘Tomentosa’ Wei Zou, Yonggang Wang, Haibin Liu, Yulong Luo, Si Chen and Weiwei Su	457
Two New Chalcones from the Flowers of <i>Clerodendrum inerme</i> Shaik Khadar Shahabuddin, Rachakunta Munikishore, Golakoti Trimurtulu, Duvvuru Gunasekar, Alexandre Deville and Bernard Bodo	459
A Novel Phenolic Compound from <i>Phyllanthus emblica</i> Gaimei She, Ruiyang Cheng, Lei Sha, Yixia Xu, Renbin Shi, Lanzhen Zhang and Yajian Guo	461
Anti-austeric Activity of Phenolic Constituents of Seeds of <i>Arctium lappa</i> Yasuhiro Tezuka, Keiichi Yamamoto, Suresh Awale, Feng Li, Satoshi Yomoda and Shigetoshi Kadota	463
Bioactive Lignans from the Leaves and Stems of <i>Schisandra wilsoniana</i> Guang-Yu Yang, Rui-Rui Wang, Zhong-Hua Gao, Yin-Ke Li, Liu-Meng Yang, Xiao-Nian Li, Shan-Zhai Shang, Yong-Tang Zheng, Wei-Lie Xiao and Han-Dong Sun	467
Antioxidative / Acetylcholinesterase Inhibitory Activity of Some Asteraceae Plants Ivana Generalić Mekinić, Franko Burčul, Ivica Blažević, Danijela Skroza, Daniela Kerum and Višnja Katalinić	471
Antioxidant and Antimicrobial Activities, and Phenolic Compounds of Selected <i>Inula</i> species from Turkey Alper Gökbulut, Onural Özhan, Basri Satılmış, Kadir Batçoğlu, Selami Günel and Engin Şarer	475
Two New Dihydrostilbenoid Glycosides Isolated from the Leaves of <i>Litsea coreana</i> and their Anti-inflammatory Activity Wenjian Tang, Weili Lu, Xiaoqing Cao, Yilong Zhang, Hong Zhang, Xiongwen Lv and Jun Li	479
Inhibitory Activity of Benzophenones from <i>Anemarrhena asphodeloides</i> on Pancreatic Lipase Yang Hee Jo, Seon Beom Kim, Jong Hoon Ahn, Qing Liu, Bang Yeon Hwang and Mi Kyeong Lee	481
Identification and Quantification of Furanocoumarins in Stem Bark and Wood of Eight Algerian Varieties of <i>Ficus carica</i> by RP-HPLC-DAD and RP-HPLC-DAD-MS Samia Rouaiguia-Bouakkaz, Habiba Amira-Guebailia, Céline Rivière, Jean-Claude Delaunay, Pierre Waffo-Téguo and Jean-Michel Mérillon	485
UPLC-Q-TOF/MS Coupled with Multivariate Statistical Analysis as a Powerful Technique for Rapidly Exploring Potential Chemical Markers to Differentiate Between <i>Radix Paeoniae Alba</i> and <i>Radix Paeoniae Rubra</i> Nian-cui Luo, Wen Ding, Jing Wu, Da-wei Qian, Zhen-hao Li, Ye-fei Qian, Jian-ming Guo and Jin-ao Duan	487
Antimicrobial Activity of Crude Methanolic Extract from <i>Phyllanthus niruri</i> Darah Ibrahim, Lim Sheh Hong and Ninthianantham Kuppan	493
Cellulose Contents of Some Abundant Indian Seaweed Species Arup K. Siddhanta, Sanjay Kumar, Gaurav K. Mehta, Mahesh U. Chhatbar, Mihir D. Oza, Naresh D. Sanandiya, Dharmesh R. Chejara, Chirag B. Godiya and Stalin Kondaveeti	497
Anti-inflammatory Potential of Silk Sericin Pornanong Aramwit, Pasarapa Towiwat and Teerapol Srichana	501
Composition of Essential Oil from Aerial and Underground Parts of <i>Geum rivale</i> and <i>G. urbanum</i> Growing in Poland Aleksandra Owczarek, Jan Gudej and Agnieszka Kice	505

Continued Inside backcover