



HAL
open science

DNA-Templated [2+2] Photocycloaddition: A Straightforward Entry into the Aplysinopsin Family of Natural Products

Nicolas Duchemin, Adam Skiredj, Justine Mansot, Karine Leblanc,
Jean-Jacques Vasseur, Mehdi Beniddir, Laurent Evanno, Erwan Poupon,
Michaël Smietana, Stelios Arseniyadis

► To cite this version:

Nicolas Duchemin, Adam Skiredj, Justine Mansot, Karine Leblanc, Jean-Jacques Vasseur, et al.. DNA-Templated [2+2] Photocycloaddition: A Straightforward Entry into the Aplysinopsin Family of Natural Products. *Angewandte Chemie International Edition*, 2018, 57 (36), pp.11786-11791. 10.1002/anie.201806357 . hal-02197152

HAL Id: hal-02197152

<https://hal.science/hal-02197152>

Submitted on 31 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DNA-Templated [2+2] Photocycloaddition: A Straightforward Entry into the Aplysinopsin Family of Natural Products**

Nicolas Duchemin,[†] Adam Skiredj,[†] Justine Mansot, Karine Leblanc, Jean-Jacques Vasseur, Mehdi A. Beniddir, Laurent Evanno,^{*} Erwan Poupon,^{*} Michael Smietana,^{*} and Stellios Arseniyadis^{*}

Abstract: Biosynthetic considerations inspired us to harness the template properties offered by DNA to promote a [2+2] photo-induced cycloaddition. The method was developed based on the dimerization of (*E*)-aplysinopsin, which was previously shown to be unproductive in solution. In sharp contrast, exposure of this tryptophan-derived olefin to light in the presence of salmon testes DNA (st-DNA) reproducibly afforded the corresponding homo-dimerized spiro-fused cyclobutane in excellent yields. DNA provides unique templating interactions enabling a singular mimic of the solid-state aggregation necessary for the [2+2] photo-cycloaddition to occur. This method was ultimately used to promote the prerequisite dimerizations leading to both dictazole B and tubastrindole B, thus constituting the first example of a DNA-mediated transformation to be applied to the total synthesis of a natural product.

The importance of light-induced reactions in biosynthetic transformations is undisputable as showcased by the thousands of examples found in nature.^[1] Interestingly however, despite the significant efforts made in the field,^[2] photochemical reactions still remain underused in natural product synthesis mainly due to the difficulty in taming the high-energy intermediates involved during the process. Exploiting supramolecular interactions and confinement to modify the excited-state behaviour of molecules has emerged as a particularly appealing strategy.^[3] The use of cucurbituril, fullerenes and, more recently, cyclodextrins, just to name a few, has allowed a preorganization of the reactants and ultimately a control of the photochemical process.^[4]

Since Roelfes' and Feringa's seminal paper published a bit over a decade ago,^[5] the field of DNA-based asymmetric catalysis has been thriving resulting in the development of a number of catalytic transformations by several groups^[6] including ours.^[7] The field

recently culminated with the development of a [2+2] photo-catalysed cycloaddition^[8] involving a modified three-way junction DNA bearing a covalently-attached photo-sensitizer. This elegant approach prompted us to explore yet another interesting feature offered by DNA, namely its template properties, to promote a photo-dimerization process perhaps otherwise impossible. As a model reaction, we decided to study the photo-induced cycloaddition of (*E*)-aplysinopsin monomers in solution. Indeed, in our quest to validate the biosynthetic hypotheses of various structurally intriguing cyclobutane-containing natural products, we recently showed that the (*E*)-aplysinopsin monomers could undergo a [2+2] photo-cycloaddition to produce the dictazole-type spiro-fused cyclobutanes, however this reaction was only possible under solvent-free conditions (Figure 1A).^[9] In comparison, the reactions performed in solution failed to provide any of the desired product. To circumvent this lack of reactivity in solution, we envisioned a DNA-templated strategy^[10] where the compartmentalization properties offered by DNA could allow to bypass the inability of the monomers to self-organize in solution and ultimately promote the title cycloaddition. Most importantly, if successful, this strategy could be extended to the synthesis of a number of cyclobutane-containing natural products and analogues thereof (Figure 1B).

Our first set of experiments rapidly revealed that DMF (3.75 v/v%) was necessary to solubilize (*E*)-aplysinopsin **1a** (1.5 mM) in the aqueous buffer solution (MOPS, 20 mM, pH 6.5) containing st-DNA (3 mM bp). In a typical reaction, the mixture was stirred by inversion in quartz tubes under a UV-B enriched lamp (ReptiSun 10.0 ZOOMED[®] UV-B) for 4 days and the product distribution was determined by HPLC-UV-MS analysis (Table 1). Interestingly, under these initial conditions, we were able to observe the formation of the desired dimerized product **2a** (Table 1, entry 2), while a control experiment run in the absence of DNA (Table 1, entry 3) only produced the formylindole derivative thus demonstrating the crucial role played by DNA in this reaction. It is also worth pointing out that increasing the percentage of DMF or the concentration of the substrate had a detrimental effect on the conversion of **1a** to **2a** (Table 1, entries 4-5). With these preliminary results in hand, we next evaluated the photophysical features offered by the complexing properties of a series of water-compatible Lewis acids (not shown in Table 1, see ESI for details). Interestingly, the use of La(OTf)₃ appeared to improve the yield to up to 56%. However, similar yields could also be obtained by simply improving the solubility of the starting material by adding a DNA-compatible co-solvent such as DMSO^[11] (25 v/v%) instead of the Lewis acid (Table 1, entries 6-8).

The head-to-tail dimerization resulting from the topochemical control of the DNA-templated photodimerization was also confirmed by ¹H and ¹³C NMR analysis after running the reaction on a 0.15 mmol preparative scale.^[12] Interestingly, the homodimerized product **2a** was obtained in up to 65% yield (40% isolated yield after preparative HPLC) after running the reaction for 48 h (Scheme 1A).^[13] The latter was accompanied with 11% of **2a'** (8% isolated yield) bearing the dictazole A-type *syn* stereochemistry

[*] N. Duchemin, Dr. S. Arseniyadis
Queen Mary University of London
School of Biological and Chemical Sciences
Mile End Road, London, E1 4NS (UK)
E-mail: s.arseniyadis@qmul.ac.uk


A. Skiredj, K. Leblanc, Dr. M. A. Beniddir, Dr. L. Evanno,
Prof. Dr. E. Poupon
Laboratoire de Pharmacognosie associé au CNRS
UMR 8076 BioCIS, Université Paris-Sud
5 rue Jean-Baptiste Clément
92296 Châtenay-Malabry (France)
E-mail: erwan.poupon@u-psud.fr
E-mail: laurent.evanno@u-psud.fr

J. Mansot, Dr. J.-J. Vasseur, Prof. Dr. Michael Smietana
Institut des Biomolécules Max Mousseron, UMR 5247
Université de Montpellier, ENSCM
Place Eugène Bataillon, 34095 Montpellier (France)
E-mail: michael.smietana@umontpellier.fr

[†] These authors contributed equally

[**] Supporting information for this article is available on the WWW

A. Aplysinopsins family: dictazoles as synthetic gateways



B. This work: DNA-templated total synthesis of dictazoles, biomimetic entry into the "aplysinopsins' cascade"




Figure 1. DNA-templated photoinduced [2+2] cycloaddition of aplysinopsins

resulting from the dimerization of (*E*)-aplysinopsin with its (*Z*) isomer formed *in situ*. Hence, the optimized DNA-based photodimerization conditions led to an unprecedented 76% combined yield, affording 20 mg of the *spiro*-fused cyclobutane in a single batch. In contrast, the solid-state protocol required combining 14 batches, each on a 0.025 mmol scale, to isolate roughly the same amount of product (40% conversion, 14% isolated yield).

In order to rationalize the intricate interaction between the aplysinopsin analogues and DNA, we performed circular dichroism (CD) experiments. Indeed, non-covalent ligands that bind to DNA *via* intercalation, groove binding or sometimes external binding, display characteristic CD signals.^[14] In addition, while single binding modes are rarely observed with neutral ligands, the presence of cationic charges further increases the number of possible modes of interaction. As shown in Figure 2A, st-DNA exhibits a typical B-type CD spectrum characterized by a negative and a positive Cotton effect at $\lambda = 245$ and 278 nm respectively. Upon addition of compound **1a** into a solution containing st-DNA (30 μ M), no CD activity could be observed when $R < 1$ ($R = [\mathbf{1a}]/[\text{DNA}]$). Interestingly, when the concentration of **1a** was increased from $R = 1$ to $R = 2$ while keeping the concentration of st-DNA constant, a new biphasic CD signal with a strong positive (340 nm) and a broad negative (390 nm) maximum and a zero cross-over at 360 nm was observed (Figure 2A). The appearance of this bisignate shape was concomitant with a small decrease of intensity of st-DNA bands attributed to a small change in the winding of the nucleobase pairs around the DNA helix. The splitting of the ICD signal into a positive and a negative band is characteristic of an exciton coupling effect which arises when two or more dissymmetrically arranged chromophores interact with each other through space. It is generally taken as evidence for the formation of dimers or higher order complexes and characteristic of groove binding or external stacking.^[15-17] Obviously, as compound **1a** is an achiral molecule, the presence of a CD signal in solution is characteristic of a chiral organisation. More specifically, the presence of exciton-coupled bisignate bands strongly supports groove binding, ligand-ligand binding or electrostatic interactions.^[18] Interestingly, upon increasing the ionic strength of the buffer from 0 to 1 M NaCl, new CD spectral features were observed (Figure 2B). While the st-DNA bands in the 200-300 nm region were not affected, the original exciton-coupled CD signals observed at lower ionic strength entirely disappeared and were replaced by a broad negative band at

$\lambda = 400$ nm. This suggests that at low ionic strength, electrostatic and groove binding modes majorly contribute to the CD spectra, while at higher ionic strength, phosphodiester are totally paired with sodium cation and therefore groove binding predominates.^[19]

The DNA-templated formation of helical stacked aggregates of cationic organic molecules is a well-studied phenomenon.^[20,21] It has notably been shown that electronic couplings occurs in H-aggregates (parallel stacking) and J-aggregates (head-to-tail stacking). By analogy with well-studied cationic groove binders such as distamycin, we thus hypothesized that two aplysinopsin molecules could stack head-to-tail with charged groups positioned at opposite ends.^[22,23] This spatial arrangement would allow the photoinduced [2+2] cycloaddition to occur.

To corroborate the importance of these multiple interaction modes, two control experiments were performed. First, to probe the importance of the groove binding interactions, we attempted to form a 2:1 complex between **1a** and the biologically relevant pyrophosphate (PPI) to mimic DNA's phosphate backbone.^[24] In that case, irradiation of the mixture failed to provide the desired cycloaddition product thus confirming not only that an electrostatically-induced sandwich-like dimer cannot homodimerize but also that the groove binding aggregation is necessary. Second, to evaluate the influence of the electrostatic interactions, we set up the

Table 1. Systematic study

Reaction scheme showing the synthesis of dictazole B (**2a**) from aplysinopsin (**1a**) using st-DNA, MOPS buffer (pH 6.5), and light ($\lambda = 280-315$ nm).

Entry	st-DNA (mM)	1a (mM)	Co-solvent/Additive	Yield ^a (%)
1	3	1.5	-	0
2	3	1.5	DMF (3.75 v/v%)	23
3	0	1.5	DMF (3.75 v/v%)	0
4	3	3.0	DMF (3.75 v/v%)	17
5	3	4.5	DMF (3.75 v/v%)	3
6	3	1.5	DMSO (3.75 v/v%)	28
7	3	1.5	DMSO (10 v/v%)	39
8	3	1.5	DMSO (25 v/v%)	55

Conditions: All reactions were ran in 2 mL quartz tubes: a 6 mM base pair solution of st-DNA in 20 mM MOPS (500 μ L, pH 6.5) was slowly added to a 3 mM solution of (*E*)-aplysinopsin **1a** in a 1:1 MOPS (20mM, pH 6.5)/DMSO solution (500 μ L) at rt under artificial light during 96 h. ^a All yields were determined by HPLC analysis (reproducibility $\pm 3\%$; average over three independent runs).




Figure 2. (A) CD spectra changes of st-DNA (30 μM) in MOPS buffer (20 mM, pH 6.5, 0 M NaCl) at 5 °C upon addition of compound **1a** (0-70 μM), (B) CD spectra changes of st-DNA (30 μM) in MOPS buffer (20 mM, pH 6.5, 1 M NaCl) at 5 °C upon addition of compound **1a** (0-60 μM). (C) Rationalization of the DNA-templated [2+2] cycloaddition and representative HPLC chromatogram.

homophotodimerization of the 3'-deimino-3'-oxoaplysinopsin **1b** bearing a neutral urea moiety instead of the guanidium present in **1a** (Scheme 1, B). Here again, no pseudodictazole resulting from the [2+2] cycloaddition was observed thus supporting the importance of the electrostatic binding between the aplysinopsin monomers and the polyanionic backbone of DNA.

Based on these results, we propose a plausible cascade of events (Figure 2C) featuring: **1** the electrostatic and groove binding of a first leading monomer, **2** the aggregation and groove binding of a second monomer in a head-to-tail conformation to avoid charge repulsion and **3** the key [2+2] cycloaddition. This pathway should therefore not only allow the homodimerization of a positively charged monomer, but also a potential heterodimerization with a second monomer which would aggregate in the DNA groove. If confirmed, this would allow a straightforward entry into the natural hetero-dimer dictazole B (**2d**).

To support this scenario, we logically started by evaluating the dimerization of the brominated analogue **1c** on a preparative scale. Interestingly, the use of a 3 mM base pair concentration of st-DNA in 25 v/v% DMSO at room temperature afforded the expected pseudo-dictazole **2c** albeit in only 14% isolated yield (Scheme 1, C). In this case, the reaction mainly led to the recovery of the starting monomer and only traces of the brominated formyl indole. This lower yield compared to the one obtained for the homodimerization of **1a** can be explained by the fast disassembly of the resulting cyclobutane product **2c** within a few hours in solution, as experimentally observed on a pure sample.


We then naturally sought to apply our optimized conditions to a 1:1 mixture of **1c** and **1d** hoping to favour the formation of the natural heterodimer, dictazole B (Scheme 1, D). Indeed, this structurally fascinating framework featuring a four-membered ring spirocyclic scaffold and isolated from the sponge *Smenospongia cerebriformis*,^[25] belongs to the aplysinopsin family of indole marine natural products which also includes aplysinopsin (**1a**) and its deimino and brominated analogues **1b**, **1c** and **1d**, (Figure 1B),^[26] and the tetrahydrocarbazole cycloaplysinopsins; tubastrindole B (**3a**)

and dictazoline C (**3b**).^[25,27-30] We recently suggested a potential biosynthetic scenario for these natural products featuring a unified "aplysinopsin cascade", where the dictazoles are the central dimers *en route* to the cycloaplysinopsins.^[31] While this hypothesis still remains questionable as far as biosynthetic considerations are concerned, this DNA-templated approach allows a straightforward and particularly efficient entry into this family of natural products.

To our delight, the condensation of the two monomers **1c** and **1d** led to the formation of the corresponding heterodimer, which spectral data matched those reported in the literature for natural dictazole B (**2d**).^[9] Even more surprising was the efficacy of the reaction, as the natural product was obtained in an improved 16% isolated yield in a single run (0.15 mmol scale),^[32] which compared favourably with the 3.4% previously obtained after combining 28 batches, each on a 0.025 mmol scale. This could potentially be explained by the postulated competitive disassembly of **2c** in solution during the course of the reaction, which would favour the predominant formation of the more stable heterodimer **2d**. Finally, the enantioselectivity obtained for dictazole B, although modest (7% ee), further confirmed the implication of DNA during the photocycloaddition process.

Although J-aggregates are usually assumed to form in the minor groove of DNA,^[33] it is not possible at this time to unambiguously assert the exact nature of the groove binding mode of the aplysinopsin derivatives. Nevertheless, these results highlight the different roles played by both the leading (positively charged) and following (neutral) monomers during the aggregation process, which results in the [2+2] cycloaddition.

Hence, starting from three different natural aplysinopsin-derived monomers, we were able to access four cyclobutane-centered skeletons, including natural dictazole B. Interestingly, the homodimer **2a** could also be further converted to yet another natural product, tubastrindole B, through a ring-expansion cascade involving the formation of Mancini's intermediate (Scheme 1), thus showcasing the possible generalization of this approach to the entire aplysinopsin family of natural products.^[34]



Scheme 1. Expanding the method to the aplysinopsin series and cycloaplysinopsin counterparts after ring expansion

In summary, we've developed an unprecedented DNA-templated [2+2] photocycloaddition process and demonstrated its synthetic utility by applying it to the synthesis of spiro-fused cyclobutane-containing compounds including the natural heterodimer dictazole B. The latter was obtained in 16% isolated yield, which significantly improved the results obtained under the standard solvent-free conditions. Moreover, by applying solid-state supramolecular principles ascertained by CD spectroscopy, we have shown that DNA could be a useful template to build aggregates with photoreactive units. Most importantly, we believe that this method is not limited to the aplysinopsins. Indeed, by fine-tuning the specific interaction between DNA and any given monomer, we should be able to extend the scope of the method and thus pave the way to new target-oriented syntheses.

Acknowledgements

The authors gratefully thank the Agence Nationale de la Recherche for financial support (D-CYSIV project; ANR-2015-CE29-0021-01).

Received: ((will be filled in by the editorial staff))

Published online on ((will be filled in by the editorial staff))

Keywords: DNA · [2+2] photocycloaddition · templated synthesis · synthesis · Aplysinopsin · Dictazole B

[1] a) N. Hoffmann, *Chem. Rev.* **2008**, *108*, 1052; b) S. Thompson; A. G. Coyne, P. C. Knipe, M. D. Smith, *Chem.*

Soc. Rev. **2011**, *40*, 4217; c) K. Glusac, *Nat. Chem.* **2016**, *8*, 734; d) T. Bach, J. P. Hehn, *Angew. Chem. Int. Ed.* **2011**, *50*, 1000.

[2] a) T. P. Nicholls, D. Leonori, A. C. Bissember, *Nat. Prod. Rep.* **2016**, *33*, 1248; b) B. S. Matsuura, P. Kölle, D. Trauner, R. Vivie-Riedle, R. Meier, *ACS Central Sci.* **2017**, *3*, 39; c) D. Stichnoth, P. Kölle, T. J. Kimbrough, E. Riedle, R. Vivie-Riedle, D. Trauner, *Nat. Commun.* **2014**, *6*, 5597; d) T. Bach, J. P. Hehn, *Angew. Chem. Int. Ed.* **2011**, *50*, 1000; e) F. Mayr, C. Wiegand, T. Bach, *Chem. Commun.* **2014**, *50*, 3355; f) S. Poplata, A. Tröster, Y. Q. Zhou, T. Bach, *Chem. Rev.* **2016**, *116*, 9748.

[3] a) B. Bibal, C. Mongin, D. M. Bassani, *Chem. Soc. Rev.* **2014**, *43*, 4179; b) J. Svoboda, B. König, *Chem. Rev.* **2006**, *106*, 5413.

[4] a) V. Ramamurthy, J. Sivaguru, *Chem. Rev.* **2016**, *116*, 9914; b) S. J. Barrow, S. Kasera, M. J. Rowland, J. del Barrio, O. A. Scherman, *Chem. Rev.* **2015**, *115*, 12320; c) K. I. Assaf, W. M. Nau, *Chem. Soc. Rev.* **2015**, *44*, 394; d) B. C. Pemberton, N. Barooah, D. K. Srivatsava, J. Sivaguru, *Chem. Commun.* **2010**, *46*, 225; e) A. T. Mansour, J. Buendia, J. Xie, F. Brisset, S. Robin, D. Naoufal, O. Yazbeck, D. J. Aitken, *J. Org. Chem.* **2017**, *82*, 9832.

[5] G. Roelfes, B. L. Feringa, *Angew. Chem. Int. Ed.* **2005**, *44*, 3230.

[6] a) N. Duchemin, I. Heath-Apostolopoulos, M. Smetana, S. Arseniyadis, *Org. Biomol. Chem.* **2017**, *15*, 7072; b) A. Rioz-Martinez, J. Oelerich, N. Ségaud, G. Roelfes, *Angew. Chem. Int. Ed.* **2016**, *55*, 14136; c) A. Rioz-Martinez, G. Roelfes, *Curr. Opin. Chem. Biol.* **2015**, *25*, 80; d) S. Park, H. Sugiyama, *Molecules* **2012**, *17*, 12792; e) A. J. Boersma, R. P. Megens, B. L. Feringa, G. Roelfes, *Chem. Soc. Rev.*

- 2010, 39, 2083; f) S. Park, H. Sugiyama, *Angew. Chem. Int. Ed.* **2010**, 49, 3870; g) S. K. Silverman, *Angew. Chem. Int. Ed.* **2010**, 49, 7180.
- [7] a) J. Wang, E. Benedetti, L. Bethge, S. Vonhoff, S. Klussmann, J.-J. Vasseur, J. Cossy, M. Smietana, S. Arseniyadis, *Angew. Chem. Int. Ed.* **2013**, 52, 11546; b) E. Benedetti, N. Duchemin, L. Bethge, S. Vonhoff, S. Klussmann, J.-J. Vasseur, J. Cossy, M. Smietana, S. Arseniyadis, *Chem. Commun.* **2015**, 51, 6076; c) K. Amirbekyan, N. Duchemin, E. Benedetti, R. Joseph, A. Colon, S. A. Markarian, L. Bethge, S. Vonhoff, S. Klussmann, J. Cossy, J.-J. Vasseur, S. Arseniyadis, M. Smietana, *ACS Catal.* **2016**, 6, 3096; d) N. Duchemin, E. Benedetti, L. Bethge, S. Vonhoff, S. Klussmann, J.-J. Vasseur, J. Cossy, M. Smietana, S. Arseniyadis, *Chem. Commun.* **2016**, 52, 8604.
- [8] a) N. Gaß, J. Gebhard, H.-A. Wagenknecht, *ChemPhotoChem* **2017**, 1, 48; b) U. Reisacher, L. Antusch, R. Hofsäb, C. Schwechheimer, B. Lehmann, H.-A. Wagenknecht, *Curr. Opin. Chem. Biol.* **2017**, 40, 119.
- [9] A. Skiredj, M. A. Beniddir, D. Joseph, K. Leblanc, G. Bernadat, L. Evanno, E. Poupon, *Angew. Chem. Int. Ed.* **2014**, 53, 6419.
- [10] A. Garcia-Fernández, R. P. Megens, L. Villarino, G. Roelfes, *J. Am. Chem. Soc.* **2016**, 138, 16308.
- [11] The use of DMSO was shown not to perturb the double-helix structure of DNA, see: a) J. F. Escara, J. R. Hutton, *Biopolymers*, **1980**, 19, 1315-1327; b) S. A. Markarian, A. M. Asatryan, K. R. Grigoryan, H. R. Sargsyan *Biopolymers*, **2006**, 82, 1-5 and references therein.
- [12] Reaction run on a 0.15 mmol scale in a crystallizer dish with slow magnetic stirring and direct irradiation.
- [13] A similar yield could be obtained using a 1.3 mM base pair concentration of st-DNA after 24 h.
- [14] M. Misiak, F. Mantegazza, G. L. Beretta, *Curr. Pharm. Des.* **2016**, 22, 6596.
- [15] a) N. C. Garbett, P. A. Ragazzon, J. B. Chaires, *Nature Protocols*, **2007**, 2, 3166; b) B. Nordén, T. Kurucsev, *J. Mol. Recognit.*, **1994**, 7, 141; c) Y. Y. Mao, K. Y. Liu, L. Chen, X. H. Cao, T. Yi, *Chem. Eur. J.* **2015**, 21, 16623.
- [16] W. C. Tse, D. L. Boger, *Acc. Chem. Res.* **2004**, 37, 61.
- [17] a) A. L. Stadler, B. R. Renikuntla, D. Yaron, A. S. Fang, B. A. Armitage, *Langmuir* **2011**, 27, 1472; b) B. A. Armitage, in *DNA Binders, Related Subjects*, Eds. M. J. Waring, J. B. Chaires, 2005, vol. 253, pp. 55; c) K. C. Hannah, B. A. Armitage, *Acc. Chem. Res.* **2004**, 37, 845.
- [18] a) K. A. Schug, W. Lindner, *Chem. Rev.* **2005**, 105, 67; b) K. Ohara, M. Smietana, J.-J. Vasseur, *J. Am. Soc. Mass Spectrom.* **2006**, 17, 283; c) K. Amirbekyan, J. Mansot, K. Ohara, S. A. Markarian, J.-J. Vasseur, M. Smietana, *Tetrahedron Lett.* **2018**, 59, 293.
- [19] M. Wang, G. L. Silva, B. A. Armitage, *J. Am. Chem. Soc.* **2000**, 122, 9977.
- [20] L. Di Bari, G. Pescitelli, P. Salvadori, *J. Am. Chem. Soc.* **1999**, 121, 7998.
- [21] B. L. Cannon, D. L. Kellis, L. K. Patten, P. H. Davis, J. Lee, E. Graugnard, B. Yurke, W. B. Knowlton, *J. Phys. Chem. A* **2017**, 121, 6905.
- [22] T. Šmidlehner, I. Piantanida, G. Pescitelli, *Beilstein J. Org. Chem.* **2018**, 14, 84.
- [23] S. M. Nelson, L. R. Ferguson, W. A. Denny, *Mutat. Res.* **2007**, 623, 24.
- [24] S. Nishizawa, Y. Kato, N. Teramae, *J. Am. Chem. Soc.* **1999**, 121, 9463.
- [25] a) E. M. Boyd, J. J. Sperry, *Chem. N. Z.* **2010**, 74, 109; b) J. Dai, J.; Jimenez, J. I.; Kelly, M.; Williams, P. G. *J. Org. Chem.* **2010**, 75, 2399.
- [26] R. Kazlauskas, P. T. Murphy, R. J. Quinn, R. J. Wells, *Tetrahedron Lett.* **1977**, 18, 61.
- [27] A. Skiredj, M. A. Beniddir, D. Joseph, G. Bernadat, L. Evanno, E. Poupon, *Synthesis* **2015**, 47, 2367.
- [28] a) I. Mancini, G. Guella, H. Zibrowius, F. Pietra, *Tetrahedron* **2003**, 59, 8757; b) M. Meyer, F. Delberghe, F. Liron, M. Guillaume, A. Valentin, M. Guyot, *Nat. Prod. Res.* **2009**, 23, 178.
- [29] a) T. Isagawa, M. Miyazaki, H. Okamura, M. Nakatani, M. Doe, K. Takemura, *Tetrahedron Lett.* **2003**, 44, 2533; b) T. Isagawa, M. Miyazaki, Y. Yokogawa, H. Okamura, M. Nakatani, M. Doe, Y. Morimoto, K. Takemura, *Heterocycles* **2008**, 75, 2023; c) W. Balansa, R. Islam, D. F. Gilbert, F. Fontaine, X. Xiao, H. Zhang, A. M. Piggott, J. W. Lynch, R. J. Capon, *Bioorg. Med. Chem.* **2013**, 21, 4420.
- [30] J. Dai, J. I. Jimenez, M. Kelly, S. Barnes, P. Lorenzo, P. G. Williams, *J. Nat. Prod.* **2008**, 71, 1287.
- [31] A. Skiredj, M. A. Beniddir, D. Joseph, K. Leblanc, G. Bernadat, L. Evanno, E. Poupon, *Org. Lett.* **2014**, 16, 4980.
- [32] Under these conditions, we were also able to isolate 4% of **2c** resulting from the homodimerization of **1c**.
- [33] Losytskyy, M. Y.; Yashchuk, V. M.; *Fluorescent J-Aggregates and their Biological Applications in Advanced Fluorescence Reporters in Chemistry and Biology II*. Springer Series on Fluorescence (Methods and Applications), Ed. A. Demchenko, vol 9. Springer, Berlin, Heidelberg (2010).
- [34] M. A. Beniddir, L. Evanno, D. Joseph, A. Skiredj, E. Poupon, *Nat. Prod. Rep.* **2016**, 33, 820.

