



HAL
open science

Quantum optimal transport is cheaper

Emanuele Caglioti, François Golse, Thierry Paul

► **To cite this version:**

Emanuele Caglioti, François Golse, Thierry Paul. Quantum optimal transport is cheaper. 2019. hal-02214344v2

HAL Id: hal-02214344

<https://hal.science/hal-02214344v2>

Preprint submitted on 8 Aug 2019 (v2), last revised 4 May 2020 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

QUANTUM OPTIMAL TRANSPORT IS CHEAPER

EMANUELE CAGLIOTI, FRANÇOIS GOLSE, AND THIERRY PAUL

ABSTRACT. We compare bipartite (Euclidean) matching problems in classical and quantum mechanics. The quantum case is treated in terms of a quantum version of the Wasserstein distance. We show that the optimal quantum cost can be cheaper than the classical one. We treat in detail the case of two particles: the equal mass case leads to equal quantum and classical costs. Moreover, we show examples with different masses for which the quantum cost is strictly cheaper than the classical cost.

CONTENTS

1. Introduction	1
2. The equal mass case	6
3. The unequal mass case	10
4. Concluding remarks on quantum optimal transport	12
References	14

1. INTRODUCTION

The paradigm of modern optimal transport theory uses extensively the 2-Wasserstein distance between two probability measures μ, ν on \mathbf{R}^n , defined as

$$(1) \quad W_2(\mu, \nu)^2 := \inf_{\Pi \text{ coupling of } \mu \text{ and } \nu} \int |x - y|^2 \Pi(dx, dy).$$

We have called coupling (or transport plans) of the two probabilities μ and ν any probability measure $\Pi(dx, dy)$ on $\mathbf{R}^n \times \mathbf{R}^n$ whose marginals on the first and the second factors are μ and ν resp., i.e.

$$(2) \quad \int_{\mathbf{R}^n \times \mathbf{R}^n} a(x) \Pi(dx, dy) = \int_{\mathbf{R}^n} a(x) \mu(dx), \quad \int_{\mathbf{R}^n \times \mathbf{R}^n} b(y) \Pi(dx, dy) = \int_{\mathbf{R}^n} b(y) \nu(dy)$$

for all test (i.e. continuous and bounded) functions a and b .

Restricting the definition of W_2 to couplings of the form

$$(3) \quad \Pi = \delta(y - T(x)) \mu(dy)$$

where T is a transformation of \mathbf{R}^d such that ν is the image $T_{\#}\mu$ of μ by T , one sees that the minimization problem in the definition of $W_2(\mu, \nu)$ contains the (quadratic) Monge problem:

$$(4) \quad M(\mu, \nu)^2 := \inf_{T_{\#}\mu = \nu} \int_{\mathbf{R}^n} (x - T(x))^2 \mu(dx).$$

There is a converse result due to Knott, Smith and Brenier: under certain restrictions on the regularity of μ , any optimal coupling for the minimization problem defined by (1) is of the form (3) for some transport map T (see e.g. [1] Section 1 for some details and Theorem 2.12 in [2] for an extensive study).

Associated to W_2 is the bipartite matching problem which can be described as follows. Let us consider M material points on the real line $\{x_i\}_{i=1,\dots,M}$ with $x_i < x_{i+1}$, and with masses $\{m_i\}_{i=1,\dots,M}$, and on the other hand N points $\{y_j\}_{j=1,\dots,N}$ with $y_j < y_{j+1}$, and with masses $\{n_j\}_{j=1,\dots,N}$. We normalize the total mass as follows:

$$\sum_{i=1}^M m_i = \sum_{j=1}^N n_j = 1.$$

The bipartite problem consists in finding a coupling matrix $(p_{i,j})_{i=1,\dots,M,j=1,\dots,N}$ satisfying

$$\sum_{j=1}^N p_{i,j} = m_i, \quad \sum_{i=1}^M p_{i,j} = n_j, \quad p_{i,j} \geq 0 \text{ for each } i, j$$

which minimizes the quantity $\sum_{i,j} p_{i,j} |x_i - y_j|^2$.

That is to say, we define the optimal transport cost as

$$C_c := \inf_{\substack{p_{i,j} \geq 0 \\ \sum_{j=1}^N p_{i,j} = m_i, \sum_{i=1}^M p_{i,j} = n_j}} \sum_{i,j} p_{i,j} |x_i - y_j|^2.$$

It is natural to associate to the sets $\{x_i\}_{i=1,\dots,M}$ and $\{m_i\}_{i=1,\dots,M}$, and to the sets $\{y_j\}_{j=1,\dots,N}$ and $\{n_j\}_{j=1,\dots,N}$ the following discrete probability measures

$$\mu := \sum_{i=1}^M m_i \delta_{x_i}, \quad \nu := \sum_{j=1}^N n_j \delta_{y_j}.$$

It is easy to see that any optimal coupling of μ, ν for W_2 takes the form

$$\Pi = \sum_{i,j} p_{i,j} \delta_{x_i} \otimes \delta_{y_j}, \quad i.e. \quad \Pi(x, y) = \sum_{i,j} p_{i,j} \delta(x - x_i) \delta(y - y_j),$$

so that

$$C_c = W_2(\mu, \nu).$$

A general review of the bipartite problem is out of the scope of the present paper, and the reader is referred to the seminal work [3], the thesis [4] which contains an extensive bibliography, and [5] for a lucid presentation of the mathematical theory pertaining to this problem. Let us describe the simplest case $M = N = 2$.

In the case of equal masses, that is $m_1 = m_2 = n_1 = n_2 = \frac{1}{2}$, the optimal coupling is shown to be diagonal, in the sense that the mass $\frac{1}{2}$ is transported from the point x_1 to the point y_1 , and likewise for x_2 and y_2 . Thus

$$\Pi_{op} = \frac{1}{2} \delta_{x_1} \otimes \delta_{y_1} + \frac{1}{2} \delta_{x_2} \otimes \delta_{y_2},$$




FIGURE 1. equal masses




FIGURE 2. different masses

or equivalently

$$\Pi_{op}(x, y) = \frac{1}{2}\delta(x - x_1)\delta(y - y_1) + \frac{1}{2}\delta(x - x_2)\delta(y - y_2),$$

and therefore

$$C_c = \frac{1}{2}(x_1 - y_1)^2 + \frac{1}{2}(x_2 - y_2)^2.$$

In the case of unequal masses, let us consider the example where $m_1 = \frac{1-\eta}{2}$ and $m_2 = \frac{1+\eta}{2}$ for some $0 < \eta < 1$, while $n_1 = n_2 = \frac{1}{2}$. In this case, one shows that the optimal transport moves the mass $\frac{1}{2}$ from x_2 to y_2 , moves the remaining amount of the mass at x_2 , i.e. $\frac{\eta}{2}$, from x_2 to y_1 , and finally moves the mass $\frac{1-\eta}{2}$ from x_1 and y_1 . Therefore, the optimal coupling in this case is

$$\Pi_{op}(x, y) = \frac{1-\eta}{2}\delta(x - x_1)\delta(y - y_1) + \frac{\eta}{2}\delta(x - x_2)\delta(y - y_1) + \frac{1}{2}\delta(x - x_2)\delta(y - y_2),$$

so that

$$C_c = \frac{1-\eta}{2}(x_1 - y_1)^2 + \frac{\eta}{2}(x_2 - y_1)^2 + \frac{1}{2}(x_2 - y_2)^2.$$

A quantum analogue to the Wasserstein distance has been recently introduced in [6] according to the following general fact.

When passing from classical to quantum mechanics,

1. functions on phase-space should be replaced by operators on the Hilbert space of square integrable functions on the underlying configuration space, and
2. integration (over phase space) of classical functions should be replaced by the trace of the corresponding operators. Moreover,
3. coordinates q of the configuration space should be replaced by the multiplication operator \hat{q} by the q variable, while momentum coordinates p should be replaced by the operator $\hat{p} = -i\hbar\nabla$.

These considerations are in full accordance with the definition of quantum density matrices as self-adjoint positive operators of trace 1 on $\mathfrak{H} := L^2(\mathbf{R}^d)$. They are also consistent with the definition of couplings Q of two density matrices R and S as density matrices on $\mathfrak{H} \otimes \mathfrak{H}$ (identified with $L^2(\mathbf{R}^{2d})$) whose marginals (defined consistently again as partial traces on the two factors of $\mathfrak{H} \otimes \mathfrak{H}$) are equal to R and S . In other words

$$\text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((A \otimes I_{\mathfrak{H}})Q) = \text{trace}_{\mathfrak{H}}(AR), \quad \text{trace}_{\mathfrak{H} \otimes \mathfrak{H}}((I_{\mathfrak{H}} \otimes B)Q) = \text{trace}_{\mathfrak{H}}(BS)$$

for all bounded operators A, B on \mathfrak{H} , by analogy with (2).

Moreover they lead naturally to the following definition of the analogue of the Wasserstein distance between two quantum densities R and S . Consistently with (1) expressed on the phase-space \mathbf{R}^{2d} , therefore with $n = 2d$, we define $MK_2 \geq 0$ by

$$MK_2(R, S)^2 := \inf_{Q \text{ coupling of } R \text{ and } S} \text{trace}(CQ),$$

with

$$C := (\hat{p} \otimes I - I \otimes \hat{p})^2 + (\hat{q} \otimes I - I \otimes \hat{q})^2 - 2d\hbar.$$

In other words, expressed as an operator on $L^2(\mathbf{R}^d, dx) \otimes L^2(\mathbf{R}^d, dy)$,

$$C = (x - y)^2 - \hbar^2(\nabla_x - \nabla_y)^2 - 2d\hbar = -4\hbar^2\nabla_{x-y}^2 + (x - y)^2 - 2d\hbar.$$

The operator $\frac{1}{2}(C + 2d\hbar)$ is a quantum harmonic oscillator in the variable $(x - y)/\sqrt{2}$, and in particular $C \geq 0$. In fact, C is the antinormal (anti-Wick) ordering quantization of the classical cost function (on phase space) $(p_x - p_y)^2 + (x - y)^2$. As a consequence of this, $\text{trace}(CQ) = \int ((p_x - p_y)^2 + (x - y)^2) \widetilde{W}[Q](x, y, p_x, p_y) dx dy dp_x dp_y$, where $\widetilde{W}[Q]$ is the Husimi function of Q (whose definition is recalled below). Since $\widetilde{W}[Q]$ is the only *positive* classical function on phase space associated to a positive operator (at variance with the Wigner function or other symbols), this choice of quantization makes the definition (5) the closest to (1).

The quantity MK_2 is not a distance as shown in [6] p. 171. Nevertheless, it was established in [6] the two following links between MK_2 and W_2^1 . First, for any pair of

¹Note the unessential difference with the definition of the cost C in [6, 7, 8] created by the shift $-2d\hbar$ and accounts for a shift by $2d\hbar$ in the two next formulas.

density matrices R and S , the Husimi functions $\widetilde{W}[R]$ and $\widetilde{W}[S]$ of R and S satisfy

$$W_2(\widetilde{W}[R], \widetilde{W}[S])^2 \leq MK_2(R, S)^2 + 4d\hbar.$$

On the other hand, if R and S are Töplitz operators of symbols μ and ν ,

$$(5) \quad MK_2(R, S)^2 \leq W_2(\mu, \nu)^2.$$

Let us recall that a Töplitz operator T (or positive quantization, or anti-Wick ordering quantization) of symbol a probability measure τ is²

$$T := \int_{\mathbf{R}^{2d}} |q, p\rangle \langle q, p| \tau(dq, dp),$$

where $|q, p\rangle$ is a coherent state at point (q, p) i.e.

$$\langle x|q, p\rangle := (\pi\hbar)^{-d/4} e^{-(x-q)^2/2\hbar} e^{ipx/\hbar}.$$

We also recall the definition of the Husimi function of a density matrix R :

$$\widetilde{W}[R](q, p) := (2\pi\hbar)^{-d} \langle q, p|R|q, p\rangle.$$

The functional MK_2^2 (more precisely $MK_2^2 + 2d\hbar$ with the definition chosen in the present paper) has been systematically used and extended in [6, 7, 8] in order to study various problems, such as the validity of the mean-field limit uniformly in \hbar , the semiclassical approximation of quantum dynamics, and the problem of metrizing of the set of quantum densities in the semiclassical regime.

The quantum bipartite problem can be therefore stated as follows, in close analogy with the classical picture introduced earlier.

One considers two density matrices built in terms of the positions and masses already used for the classical bipartite problem, in the following way

$$R = \sum_{i=1}^M m_i |x_i, 0\rangle \langle x_i, 0|, \quad S = \sum_{j=1}^N n_j |y_j, 0\rangle \langle y_j, 0|.$$

Indeed, it is natural to associate coherent states to material points, as they saturate the Heisenberg uncertainty inequalities. Moreover, one sees that R and S are precisely the Töplitz operators of symbols μ and ν respectively.

The quantum bipartite problem consists then in finding an optimal coupling of R and S for $MK_2(R, S)$ and the optimal quantum cost defined as

$$C_q := MK_2(R, S).$$

Since R and S are Töplitz operators, we know from (5) that

$$C_q \leq C_c.$$

The question we address in this paper is whether there exist pairs of density matrices for which

$$C_q < C_c.$$

²Here also, we use a different normalization than the one in [6, 7, 8], since we deal exclusively with density matrices. With the present normalization, one has $\text{trace } T = \int_{\mathbf{R}^{2d}} \tau(dq, dp)$.

In other words, we address the question of whether quantum optimal transportation can be cheaper than its classical analogue.

We shall study the two cases introduced at the beginning of this section and described in Figures 1 and 2. For the sake of simplicity, we shall take $x_1 = -x_2 = -a, y_1 = -y_2 = -b$, with $a < b$ in the equal mass case, and $a = b$ in the unequal mass case.

In the equal mass case, studied in Section 2, both classical and quantum transport are achieved without splitting mass for each particle: the two costs are shown to be equal (see (13)), and an optimal quantum coupling is the Töplitz quantization to the optimal classical coupling.

In Section 3 we study the case of different masses and construct a family of examples for which the optimal quantum cost is strictly cheaper than the classical one (see (21)).

We also show in Section 4 that the optimal quantum coupling cannot be the Töplitz quantization of any classical coupling: in particular the optimal quantum transport is different from the natural quantization of the underlying classical one. In fact the quantum optimal transport in the latter case does not correspond to the classical optimal transport and involves strictly quantum effects.

2. THE EQUAL MASS CASE

For $a, b > 0$ we will transport a superposition of two density matrices which are pure states associated to two coherent states of null momenta localized at $+a$ and $-a$ towards a similar density matrix associated to the points $(\pm b, 0)$ in phase space. In other words, we consider the coherent states denoted $|c\rangle$ for simplicity (instead of $|c, 0\rangle$), i.e. $\langle x|c\rangle := (\pi\hbar)^{-1/4}e^{-(x-c)^2/2\hbar}$ and consider the two density matrices

$$R := \frac{1}{2}(|a\rangle\langle a| + |-a\rangle\langle -a|), \quad S := \frac{1}{2}(|b\rangle\langle b| + |-b\rangle\langle -b|).$$

Define

$$\lambda := \langle a|-a\rangle = e^{-a^2/\hbar}, \quad \mu := \langle b|-b\rangle = e^{-b^2/\hbar},$$

and consider the two pairs of orthogonal vectors

$$(6) \quad \phi_{\pm} := \frac{|a\rangle \pm |-a\rangle}{\sqrt{2(1 \pm \lambda)}}, \quad \psi_{\pm} := \frac{|b\rangle \pm |-b\rangle}{\sqrt{2(1 \pm \mu)}}.$$

Hence

$$R = \alpha_+|\phi_+\rangle\langle\phi_+| + \alpha_-|\phi_-\rangle\langle\phi_-|, \quad S = \beta_+|\psi_+\rangle\langle\psi_+| + \beta_-|\psi_-\rangle\langle\psi_-|,$$

with

$$\alpha_+ := \frac{1}{2}(1 + \lambda), \quad \alpha_- = \frac{1}{2}(1 - \lambda), \quad \beta_+ := \frac{1}{2}(1 + \mu), \quad \beta_- = \frac{1}{2}(1 - \mu).$$

In the whole present paper, we will only use couplings of R and S that act from the four-dimensional linear span of $\phi_{\pm} \otimes \psi_{\pm}$ to itself. Therefore, in order to compute $\text{trace}(CQ)$ for such couplings, we need to project the cost operator C on the basis

$\{\phi_+ \otimes \psi_+, \phi_+ \otimes \psi_-, \phi_- \otimes \psi_+, \phi_- \otimes \psi_-\}$. This is a tedious but straightforward computation which results in the following 4×4 matrix:

$$(7) \quad C = \begin{pmatrix} a^2 \frac{1-\lambda}{1+\lambda} + b^2 \frac{1-\mu}{1+\mu} & 0 & 0 & -2ab \frac{\lambda^2 + \mu^2 - \lambda^2 \mu^2 - \lambda \mu}{\sqrt{(1-\lambda^2)(1-\mu^2)}} \\ 0 & a^2 \frac{1-\lambda}{1+\lambda} + b^2 \frac{1+\mu}{1-\mu} & -2ab \frac{\lambda^2 + \mu^2 - \lambda^2 \mu^2 + \lambda \mu}{\sqrt{(1-\lambda^2)(1-\mu^2)}} & 0 \\ 0 & -2ab \frac{\lambda^2 + \mu^2 - \lambda^2 \mu^2 + \lambda \mu}{\sqrt{(1-\lambda^2)(1-\mu^2)}} & a^2 \frac{1+\lambda}{1-\lambda} + b^2 \frac{1-\mu}{1+\mu} & 0 \\ -2ab \frac{\lambda^2 + \mu^2 - \lambda^2 \mu^2 - \lambda \mu}{\sqrt{(1-\lambda^2)(1-\mu^2)}} & 0 & 0 & a^2 \frac{1+\lambda}{1-\lambda} + b^2 \frac{1+\mu}{1-\mu} \end{pmatrix},$$

abbreviated for simplicity as

$$(8) \quad C = \begin{pmatrix} \mathcal{A} & 0 & 0 & \gamma \\ 0 & \mathcal{B} & \delta & 0 \\ 0 & \delta & \mathcal{C} & 0 \\ \gamma & 0 & 0 & \mathcal{D} \end{pmatrix}.$$

As a warm up in order to find an ansatz for the general case, let us first neglect the contributions of λ, μ , exponentially small in the Planck constant. In this case $\alpha_{\pm} = \beta_{\pm} = \frac{1}{2}$, and the cost is equal to

$$C_0 = \begin{pmatrix} a^2 + b^2 & 0 & 0 & -2ab \\ 0 & a^2 + b^2 & -2ab & 0 \\ 0 & -2ab & a^2 + b^2 & 0 \\ -2ab & 0 & 0 & a^2 + b^2 \end{pmatrix}.$$

On the other hand, one has

$$Q_0 := \begin{pmatrix} \frac{1}{4} & 0 & 0 & \frac{1}{4} \\ 0 & \frac{1}{4} & \frac{1}{4} & 0 \\ 0 & \frac{1}{4} & \frac{1}{4} & 0 \\ \frac{1}{4} & 0 & 0 & \frac{1}{4} \end{pmatrix} \geq 0,$$

since the spectrum of Q_0 is easily shown to be $\{0, \frac{1}{2}\}$ by using the elementary formula

$$(9) \quad \det \begin{pmatrix} \bar{a} & 0 & 0 & \gamma \\ 0 & \bar{b} & \delta & 0 \\ 0 & \delta & \bar{c} & 0 \\ \gamma & 0 & 0 & \bar{d} \end{pmatrix} = (\bar{a}\bar{d} - \gamma^2)(\bar{b}\bar{c} - \delta^2) \text{ for all } \bar{a}, \bar{b}, \bar{c}, \bar{d}, \gamma, \delta.$$

Moreover, one easily checks that $\text{trace}_2 Q_0 = R$ and $\text{trace}_1 Q_0 = S$ so that Q_0 is a coupling of R and S .

Another easy computation shows that

$$\text{trace}(CQ_0) = (a - b)^2.$$

Therefore

$$MK_2(R, S)^2 \leq (a - b)^2 = W_2\left(\frac{1}{2}(\delta_{-a} + \delta_a), \frac{1}{2}(\delta_{-b} + \delta_b)\right)^2.$$

For the “true” case $\lambda, \mu \neq 0$, we make the following ansatz on the coupling Q

$$Q = Q_0 + \frac{1}{4} \begin{pmatrix} p + \lambda + \mu & 0 & 0 & u \\ 0 & -p + \lambda - \mu & v & 0 \\ 0 & v & -p - \lambda + \mu & 0 \\ u & 0 & 0 & p - \lambda - \mu \end{pmatrix}, \quad p, u, v \in \mathbf{R}.$$

Straightforward computations show that

$$\text{trace } Q = \text{trace } Q_0 = 1, \quad \text{trace}_2 Q = \text{trace}_2 Q_0 = R, \quad \text{trace}_1 Q = \text{trace}_1 Q_0 = S.$$

Using again (9) shows that

$$Q \geq 0 \iff -1 + \sqrt{(\lambda + \mu)^2 + (1 + u)^2} \leq p \leq 1 - \sqrt{(\lambda - \mu)^2 + (1 + v)^2}.$$

Therefore, assuming that p, u, v satisfy this constraint, Q is a coupling of R and S .

Denoting $U := 1 + u$ and $V := 1 + v$, we compute $W := \text{trace}(CQ)$ by using (7) and (8):

$$\begin{aligned} 4W &= 2\gamma U + 2\delta V + p(\mathcal{A} + \mathcal{B} - \mathcal{C} - \mathcal{D}) + \mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} \\ &\quad + (\lambda + \mu)(\mathcal{A} - \mathcal{D}) + (\lambda - \mu)(\mathcal{B} - \mathcal{C}) \\ &= 2\gamma U + 2\delta V + p(\mathcal{A} - \mathcal{B} - \mathcal{C} + \mathcal{D}) + W' = 2\gamma U + 2\delta V + W', \end{aligned}$$

with

$$\begin{aligned} W' &:= \mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} + \lambda(\mathcal{A} + \mathcal{B} - \mathcal{C} - \mathcal{D}) + \mu(\mathcal{A} - \mathcal{B} + \mathcal{C} - \mathcal{D}) \\ &= 4\left(a^2 \frac{1 + \lambda^2}{1 - \lambda^2} + b^2 \frac{1 + \mu^2}{1 - \mu^2}\right) - 8a^2 \frac{\lambda^2}{1 - \lambda^2} - 8b^2 \frac{\mu^2}{1 - \mu^2} \\ &= 4(a^2 + b^2). \end{aligned}$$

Since W is linear in U, V , we minimize $\gamma U + \delta V$ by taking

$$U = \sqrt{(p + 1)^2 - (\lambda + \mu)^2} \quad \text{and} \quad V = \sqrt{(p - 1)^2 - (\lambda - \mu)^2},$$

and, since $\delta \leq \gamma$, we conclude that

$$4W = 2T + W',$$

where

$$\begin{aligned} T &= - \max_{-1 + \lambda - \mu \leq p \leq 1 - (\lambda - \mu)} \left(-\gamma \sqrt{(p + 1)^2 - (\lambda + \mu)^2} - \delta \sqrt{(p - 1)^2 - (\lambda - \mu)^2} \right) \\ &= \frac{-2ab}{\sqrt{(1 - \lambda^2)(1 - \mu^2)}} \max_{-1 + \lambda - \mu \leq p \leq 1 - (\lambda - \mu)} \left((1 - \lambda\mu) \sqrt{(p + 1)^2 - (\lambda + \mu)^2} \right. \\ &\quad \left. + (1 + \lambda\mu) \sqrt{(p - 1)^2 - (\lambda - \mu)^2} \right). \end{aligned}$$

One can check that the max is attained for $p = \lambda\mu \rightarrow 0$ as $\hbar \rightarrow 0$, and that

$$T = - \frac{4ab}{\sqrt{(1 - \lambda^2)(1 - \mu^2)}} \sqrt{1 + \lambda^2\mu^2 - \lambda^2 - \mu^2} = -4ab.$$

Eventually, we arrive at the same result as in the semiclassical regime $\lambda = \mu = 0$, viz.

$$(10) \quad MK_2(R, S)^2 \leq (a - b)^2.$$

Since R and S are Töplitz operator, the inequality (10) was already known by using (5). Nevertheless we gave this explicit computation as we believe the result to be valid for more general density matrices.

In order to get a lower bound for $MK_2(R, S)$, we shall use a dual version of the definition of MK_2 , proved in [1], that is a quantum version of the Kantorovitch duality theorem for W_2 (see [2, 9]):

$$MK_2(R, S)^2 = \sup_{\substack{A=A^*, B=B^* \text{ bounded operators on } \mathfrak{H} \\ \text{such that } A \otimes I + I \otimes B \leq C}} \text{trace}(RA + SB).$$

We make the following diagonal ansatz on A and B :

$$A = \begin{pmatrix} \alpha_1 & 0 \\ 0 & \alpha_2 \end{pmatrix} \quad B = \begin{pmatrix} \beta_1 & 0 \\ 0 & \beta_2 \end{pmatrix},$$

so that

$$A \otimes I = \begin{pmatrix} \alpha_1 & 0 & 0 & 0 \\ 0 & \alpha_1 & 0 & 0 \\ 0 & 0 & \alpha_2 & 0 \\ 0 & 0 & 0 & \alpha_2 \end{pmatrix} \quad \text{and} \quad I \otimes B = \begin{pmatrix} \beta_1 & 0 & 0 & 0 \\ 0 & \beta_2 & 0 & 0 \\ 0 & 0 & \beta_1 & 0 \\ 0 & 0 & 0 & \beta_2 \end{pmatrix}.$$

Hence

$$A \otimes I + I \otimes B - C := \begin{pmatrix} a & 0 & 0 & -\gamma \\ 0 & b & -\delta & 0 \\ 0 & -\delta & c & 0 \\ -\gamma & 0 & 0 & d \end{pmatrix},$$

and, according to (8),

$$\bar{a} = \alpha_1 + \beta_1 - \mathcal{A}, \quad \bar{b} = \alpha_1 + \beta_2 - \mathcal{B}, \quad \bar{c} = \alpha_2 + \beta_1 - \mathcal{C}, \quad \bar{d} = \alpha_2 + \beta_2 - \mathcal{D}.$$

Notice that

$$\bar{a} + \bar{d} = \bar{b} + \bar{c}.$$

Using (9) to compute the characteristic polynomial of $A \otimes I + I \otimes B - C$, we find that

$$(11) \quad A \otimes I + I \otimes B \leq C \iff \bar{a} + \bar{d} \leq -\sqrt{(\bar{a} - \bar{d})^2 + 4\gamma^2} \quad \text{and} \quad b + c \leq -\sqrt{(\bar{b} - \bar{c})^2 + 4\delta^2}.$$

Moreover,

$$\begin{aligned} \text{trace}(AR + BS) &= \frac{1}{2}(\alpha_1 + \alpha_2 + \beta_1 + \beta_2) + \frac{\lambda}{2}(\alpha_1 - \alpha_2) + \frac{\mu}{2}(\beta_1 - \beta_2) \\ &= \frac{1}{4}(\bar{a} + \bar{b} + \bar{c} + \bar{d}) + \frac{1}{4}(\bar{a} + \bar{b} - \bar{c} - \bar{d})\lambda + \frac{1}{4}(\bar{a} - \bar{b} + \bar{c} - \bar{d})\mu + a^2 + b^2. \end{aligned}$$

Let us denote

$$x := \bar{a} + \bar{d} = \bar{b} + \bar{c},$$

so that

$$(12) \quad \text{trace}(AR + BS) = \frac{1}{2}x + \frac{1}{4}(\lambda + \mu)(\bar{a} - \bar{d}) + \frac{1}{4}(\lambda - \mu)(\bar{b} - \bar{c}) + a^2 + b^2.$$

The constraints (11) are expressed as

$$\begin{aligned} x = \bar{a} + \bar{d} &\leq -\sqrt{(\bar{a} - \bar{d})^2 + 4\gamma^2}, \\ x = \bar{b} + \bar{c} &\leq -\sqrt{(\bar{b} - \bar{c})^2 + 4\delta^2}. \end{aligned}$$

Without loss of generality we assume that $\lambda \geq \mu$, that is to say $a < b$. Since the right hand side of (12) is linear in x , in $(\bar{a} - \bar{d})$, and in $(\bar{b} - \bar{c})$, one has to saturate the constraints to maximize $\text{trace}(AR + BS)$. In other words, we must take

$$\bar{a} - \bar{d} = \sqrt{x^2 - 4\gamma^2}, \quad \text{and} \quad \bar{b} - \bar{c} = \sqrt{x^2 - 4\delta^2}.$$

Since $\delta \leq \gamma \leq 0$, this amounts to computing

$$\max_{x \leq 2\delta} f(x), \quad \text{with } f(x) := \frac{x}{2} + \frac{1}{4}(\lambda + \mu)\sqrt{x^2 - 4\gamma^2} + \frac{1}{4}(\lambda - \mu)\sqrt{x^2 - 4\delta^2}.$$

We check that $f'(x)$ is an increasing function of x^2 , so that the maximum of $f(x)$ for $x \leq 2\delta$ is attained at

$$f'(x) = 0 \iff x = -\frac{4ab(1 - \lambda^2\mu^2)}{(1 - \lambda^2)(1 - \mu^2)}, \quad \text{which implies } f(x) = -2ab.$$

We conclude from (12) that

$$MK_2(R, S)^2 \geq \text{trace}(AR + BS) \geq a^2 + b^2 - 2ab = (a - b)^2.$$

Together with (10), this implies that

$$MK_2(R, S)^2 = (a - b)^2 = W_2\left(\frac{1}{2}(\delta_{-a} + \delta_a), \frac{1}{2}(\delta_{-b} + \delta_b)\right)^2.$$

Therefore,

$$(13) \quad C_q = C_c,$$

so that the classical and the quantum optimal transport costs are equal in this case.

3. THE UNEQUAL MASS CASE

In this section, we construct a family of density matrices R and S for which the quantum cost of optimal transport is smaller than the classical analogous cost.

With the same notations as in previous section, we set

$$R := \frac{1+\eta}{2}|a\rangle\langle a| + \frac{1-\eta}{2}| -a\rangle\langle -a|, \quad S := \frac{1}{2}|a\rangle\langle a| + \frac{1}{2}| -a\rangle\langle -a|, \quad 0 < \eta < 1.$$

In other words, we consider the same situation as in the previous section with $a = b$, but with different masses for the quantum density matrix R .

In the orthonormal basis $\{\phi_+, \phi_-\}$, the density matrix R takes the form

$$R = \begin{pmatrix} \frac{1+\eta}{2} & \frac{\eta}{2}(1 - \lambda^2) \\ \frac{\eta}{2}(1 - \lambda^2) & \frac{1-\eta}{2} \end{pmatrix},$$

while S is the same as before.

We define the “quantized classical” coupling as

$$(14) \quad Q_c := \frac{1}{2}|a; a\rangle\langle a; a| + \frac{1-\eta}{2}| - a; -a\rangle\langle -a; -a| + \frac{\eta}{2}|a; -a\rangle\langle a; -a|,$$

with the obvious notation

$$\langle a; b| := \langle a| \otimes \langle b|; \quad |a; b\rangle := |a\rangle \otimes |b\rangle.$$

Obviously $Q_c \geq 0$ by construction, and

$$\text{trace}_2(Q_c) = \frac{1}{2}|a\rangle\langle a| + \frac{\eta}{2}|a\rangle\langle a| + \frac{1-\eta}{2}| - a\rangle\langle -a| = R, \quad \text{while } \text{trace}_1(Q_c) = S.$$

Viewed as a matrix in the basis $\{\phi_+ \otimes \psi_+, \phi_+ \otimes \psi_-, \phi_- \otimes \psi_+, \phi_- \otimes \psi_-\}$,

$$(15) \quad Q_c = \begin{pmatrix} \frac{1}{4}(1+\lambda)^2 & 0 & \frac{1}{4}\eta\sqrt{1-\lambda}(1+\lambda)^{\frac{3}{2}} & \frac{1}{4}(-1+\eta)(-1+\lambda^2) \\ 0 & \frac{1}{4}(1-\lambda^2) & \frac{1}{4}(-1+\eta)(-1+\lambda^2) & \frac{1}{4}\eta(1-\lambda)^{\frac{3}{2}}\sqrt{1+\lambda} \\ \frac{1}{4}\eta\sqrt{1-\lambda}(1+\lambda)^{\frac{3}{2}} & \frac{1}{4}(-1+\eta)(-1+\lambda^2) & \frac{1}{4}(1-\lambda^2) & 0 \\ \frac{1}{4}(-1+\eta)(-1+\lambda^2) & \frac{1}{4}\eta(1-\lambda)^{\frac{3}{2}}\sqrt{1+\lambda} & 0 & \frac{1}{4}(-1+\lambda)^2 \end{pmatrix}.$$

With (7), we easily compute

$$(16) \quad \text{trace}(CQ_c) = 2\eta a^2 = W_2\left(\frac{1+\eta}{2}\delta_a + \frac{1-\eta}{2}\delta_{-a}, \frac{1}{2}\delta_a + \frac{1}{2}\delta_{-a}\right)^2.$$

Indeed, let us recall the classical optimal transport from R to S in this case: first, one “moves” the amount of mass $\frac{1}{2}$ from a in R to a in S . The amount of mass $\frac{\eta}{2}$ remaining at a in R is transported to $-a$ in S , and the outstanding amount of mass $\frac{1-\eta}{2}$, located at $-a$ in R , is “transported” to $-a$ in S (see Figure 2).

For each $\epsilon > 0$, set

$$(17) \quad Q_\epsilon := Q_c + \epsilon Q_q,$$

with

$$Q_q := \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & -1 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix}$$

One easily checks that

$$\text{trace}_1(Q_q) = \text{trace}_2(Q_q) = \text{trace}(Q_q) = 0,$$

so that

$$(18) \quad \text{trace}_1(Q_\epsilon) = S, \quad \text{and} \quad \text{trace}_2(Q_\epsilon) = R, \quad \text{so that } \text{trace}(Q_\epsilon) = 1.$$

The characteristic polynomial of Q_c is found to be of the form

$$\det(Q_c - tI) = tP_3(t),$$

where P_3 is a cubic polynomial satisfying

$$P_3(0) = -\frac{\eta}{8}(1-\eta)(1-\eta^2) < 0.$$

Therefore the spectrum of Q_c is $\{0, \lambda_1 > 0, \lambda_2 > 0, \lambda_3 > 0\}$ since $Q_c = Q_c^* \geq 0$. One can also check that

$$(19) \quad \det(Q_\epsilon - tI)|_{t=0} = \det Q_\epsilon = \epsilon\eta\lambda^2(1-\eta)(1-\lambda^4) + O(\epsilon^2) > 0 \quad \text{for } 0 < \epsilon \ll 1,$$

together with

$$\frac{d}{dt} \det(Q_c - tI)|_{t=0} := P_3(0) < 0.$$

Hence there exists D (independent of ϵ) such that

$$(20) \quad \frac{d}{dt} \det(Q_\epsilon - tI)|_{t=0} \leq D < 0 \quad \text{for } 0 < \epsilon \ll 1.$$

Both (19) and (20) clearly imply that $\det(Q_\epsilon - tI)$ has a positive zero that is ϵ -close to 0, and three other roots which are ϵ -close to λ_1 , λ_2 and $\lambda_3 > 0$ respectively. Therefore, $Q_\epsilon = Q_\epsilon^* > 0$ for $0 < \epsilon \ll 1$, and (18) implies that Q_ϵ is a coupling of R and S .

Another elementary computation shows that

$$\text{trace}(CQ_q) = -\frac{8\eta^2\lambda^2}{1-\lambda^2},$$

so that

$$\begin{aligned} MK_2(R, S)^2 &\leq \text{trace}(CQ_\epsilon) = \text{trace}(CQ_c) - \epsilon \frac{8\eta^2\lambda^2}{1-\lambda^2} \\ &< W_2\left(\frac{1+\eta}{2}\delta_a + \frac{1-\eta}{2}\delta_{-a}, \frac{1}{2}\delta_a + \frac{1}{2}\delta_{-a}\right)^2, \end{aligned}$$

for each ϵ satisfying $0 < \epsilon \ll 1$, according to formula (16). In other words,

$$(21) \quad C_q < C_c,$$

the quantum cost is (strictly) below the classical cost.

4. CONCLUDING REMARKS ON QUANTUM OPTIMAL TRANSPORT

The result of Section 2 shows that, in the equal mass case, an optimal coupling is given by the following matrix in the basis $\{\phi_+ \otimes \psi_+, \phi_+ \otimes \psi_-, \phi_- \otimes \psi_+, \phi_- \otimes \psi_-\}$:

$$Q = \frac{1}{4} \begin{pmatrix} 1 + \lambda\mu + \lambda + \mu & 0 & 0 & \sqrt{(1 + \lambda\mu)^2 - (\lambda + \mu)^2} \\ 0 & 1 - \lambda\mu + \lambda - \mu & \sqrt{(1 - \lambda\mu)^2 - (\lambda - \mu)^2} & 0 \\ 0 & \sqrt{(1 - \lambda\mu)^2 - (\lambda - \mu)^2} & 1 - \lambda\mu - \lambda + \mu & 0 \\ \sqrt{(1 + \lambda\mu)^2 - (\lambda + \mu)^2} & 0 & 0 & 1 + \lambda\mu - \lambda - \mu \end{pmatrix}.$$

Using (6) and with the same notation as in (14), the optimal coupling Q can be put in the form

$$(22) \quad Q = \frac{1}{2} (|a; b\rangle\langle a; b| + |-a; -b\rangle\langle -a; -b|).$$

In other words, Q is the Töplitz operator of symbol

$$\Pi(q, p; q', p') = \frac{1}{2} \delta_{(-a,0)}(q, p) \delta_{(-b,0)}(q', p') + \frac{1}{2} \delta_{(a,0)}(q, p) \delta_{(b,0)}(q', p').$$

Likewise, we recall that R is the Töplitz operator of symbol

$$\mu(q, p) = \frac{1}{2} (\delta_{(-a,0)}(q, p) + \delta_{(a,0)}(q, p)),$$

while S is the Töplitz operator of symbol

$$\nu(q, p) = \frac{1}{2} (\delta_{(-b,0)}(q, p) + \delta_{(b,0)}(q, p)).$$

Therefore,

$$\begin{aligned}
 \Pi(q, p; q', p') &= \frac{1}{2}((\delta_{(-a,0)}(q, p) + \delta_{(a,0)}(q, p))\delta((q', p') - \Phi(q, p))) \\
 (23) \qquad \qquad &= \mu(q, p)\delta((q', p') - \Phi(q, p)),
 \end{aligned}$$

where Φ is any map satisfying $\Phi(a, 0) = (b, 0)$ and $\Phi(-a, 0) = (-b, 0)$.

The second equality in (23) says the following: in the equal mass case, in agreement with (3), an optimal quantum coupling Q is the Töplitz operator of symbol the classical optimal coupling associated to the optimal transport map

$$((-a, 0), (a, 0)) \mapsto ((-b, 0), (b, 0)).$$

In the unequal mass case treated in Section 3, the coupling Q_c defined by (14) is also a Töplitz operator, with symbol

$$\begin{aligned}
 \Pi_c(q, p; q', p') &= \frac{1}{2}\delta_{(a,0)}(q, p)\delta_{(a,0)}(q', p') \\
 &+ \frac{1-\eta}{2}\delta_{(-a,0)}(q, p)\delta_{(-a,0)}(q', p') + \frac{\eta}{2}\delta_{(a,0)}(q, p)\delta_{(-a,0)}(q', p').
 \end{aligned}$$

This expression is easily interpreted as the optimal coupling associated to the “transport” introduced in Section 1, Figure 2, exactly as in the equal mass case. But, as explained in the previous section, Q_c cannot be an optimal coupling, since the coupling Q_ϵ defined by (17) leads to a strictly lower quantum cost.

We did not compute any optimal coupling in this situation. Observe however that, thanks to (15) and (6) specialized to $a = b$ (so that $\lambda = \mu$), one can expand Q_q in the form

$$Q_q = \sum_{i,j,k,l=\pm 1} q_{i,j,k,l} |ia; ja\rangle \langle ka; la|.$$

The contribution of the “diagonal” terms $q_{i,j,i,j}$ defines a Töplitz operator, unlike the off-diagonal terms such as $q_{1,1,-1,1} = \frac{-4\lambda}{(1-\lambda^2)^2} \neq 0$ for instance.

In general, when R and S are Töplitz operators of symbols μ and ν satisfying $MK_2(R, S) < W_2(\mu, \nu)$, no optimal coupling Q_{op} of R and S can be a Töplitz operator: if such was the case, the Töplitz symbol of Q_{op} would be a coupling of μ and ν with classical transport cost $MK_2(R, S) < W_2(\mu, \nu)$, which is impossible. The presence of nonclassical off-diagonal terms in Q_{op} , such as $q_{1,1,-1,1} = \frac{-4\lambda}{(1-\lambda^2)^2} \neq 0$ in the example discussed above, are precisely the reason why quantum optimal transport can be cheaper in this case than classical optimal transport.

Finally, observe that both $q_{1,1,-1,1}$ and $W_2(\frac{1+\eta}{2}\delta_a + \frac{1-\eta}{2}\delta_{-a}, \frac{1}{2}\delta_a + \frac{1}{2}\delta_{-a})^2 - \text{trace}(CQ_\epsilon)$ are exponentially small as $\hbar \rightarrow 0$, but of course are not small for $\hbar = 1$.

Acknowledgements. This work has been partially carried out thanks to the support of the LIA AMU-CNRS-ECM- INdAM Laboratoire Ypatie des Sciences Mathématiques (LYSM).

REFERENCES

- [1] E. Caglioti, F. Golse, T. Paul: *Towards optimal transport for quantum densities*, preprint hal-01963667, 2018.
- [2] C. Villani: “Topics in Optimal Transportation”, Amer. Math. Soc., Providence (RI), 2003.
- [3] M. Mézard, G. Parisi: *The Euclidean Matching Problem*, J. Phys. France **49** (1988), 2019–2025.
- [4] G. Sicuro: ”The Euclidean Matching Problem”, Springer-Verlag, Berlin, Heidelberg, 2017.
- [5] H. Brezis: *Remarks on the Monge-Kantorovich problem in the discrete setting*, C. R. Acad. Sci. Paris, Sér. I **356** (2018), 207–213.
- [6] F. Golse, C. Mouhot, T. Paul: *On the Mean-Field and Classical Limits of Quantum Mechanics*, Commun. Math. Phys. **343** (2016), 165–205.
- [7] F. Golse, T. Paul: *The Schrödinger Equation in the Mean-Field and Semiclassical Regime*, Arch. Rational Mech. Anal. **223** (2017), 57–94.
- [8] F. Golse, T. Paul: *Wave Packets and the Quadratic Monge-Kantorovich Distance in Quantum Mechanics*, Comptes Rendus Mathématique **356** (2018), 177–197.
- [9] C. Villani: “Optimal Transport. Old and New”, Springer-Verlag, Berlin, Heidelberg, 2009.

(E.C.) SAPIENZA UNIVERSITÀ DI ROMA, DIPARTIMENTO DI MATEMATICA, PIAZZALE ALDO MORO 5, 00185 ROMA
E-mail address: `caglioti@mat.uniroma1.it`

(F.G.) CENTRE DE MATHÉMATIQUES LAURENT SCHWARTZ, ECOLE POLYTECHNIQUE, 91128 PALAISEAU CEDEX, FRANCE
E-mail address: `francois.golse@polytechnique.edu`

(T.P.) CENTRE DE MATHÉMATIQUES LAURENT SCHWARTZ, CNRS, ECOLE POLYTECHNIQUE, 91128 PALAISEAU CEDEX, FRANCE
E-mail address: `thierry.paul@polytechnique.edu`