



Study of the relationships between homogenization of standardized milk and milk fat and meltability and stretchability of Emmental cheese

Jean-René Kerjean, Romain Richoux, Marie-Hélène Famelart

► To cite this version:

Jean-René Kerjean, Romain Richoux, Marie-Hélène Famelart. Study of the relationships between homogenization of standardized milk and milk fat and meltability and stretchability of Emmental cheese. 25 th Congrès FIL/IDF, Sep 1998, Aarhus, Denmark. 1998. hal-02244968

HAL Id: hal-02244968

<https://hal.science/hal-02244968>

Submitted on 1 Aug 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Study of the Relationships between Homogenization of Standardized Milk and Milk Fat and Meltableity and Stretchability of Emmental Cheese.

Jean-René Kerjean*, Romain Richoux*, Marie-Hélène Famelart*

* ITC Ouest, 73 rue de Saint-Brieuc, BP 6224, F-35062 Rennes, France

° INRA, LRTL, 65 rue de Saint-Brieuc, F-35042 Rennes, France

Melting and stretching abilities of cheese are of great significance since 75 % of French Emmental production are consumed in a cooked form. The fat content and fat globule membrane state are expected to play a key role in cooking properties of cheese (1). The aim of this work was to study the influence of milk fat homogenization on the melting and stretching of Emmental cheese.







Material and methods

Small-scale Emmental cheese (800 g) were made following a standardized technology. Various amounts (0, 25, 50, 75, 100 %) of homogenized (6.0 Mpa, 50 °C), protein/fat standardized milk were added to the cheese milk. Alternatively, the fat of the cheese milk was adjusted with the same amounts (0, 25, 50, 75 or 100 % of fat recovered in cheese) of homogenized dairy cream. Melting, free-oil (Schreiber test) and stretching (original method, see Kerjean *et al.* this symposium) were measured after 7, 21 and 49 days ripening.

Results

Homogenized cheese milk induced poor melting abilities and low free oil formation in cheese. Swiss-type cheese made with 100 % homogenized milk did not melt at all.

The homogenization of the dairy cream increased dramatically the melting abilities of cheese. A strong correlation ($r=0.958$) was found between the amount of homogenized cream and the average diameter of melted cheese.



Conclusion

This work confirms the importance of the size and the state of milk fat globule membrane on the melting abilities of cheese.

References

M.H. Tunick, E. Malin, P.W. Smith, J.J. Shieh, B.C. Sullivan, K.L. Mackey, V.H. Holstinger (1993). Proteolysis of rheology of low fat and full-fat mozzarella cheese prepared from homogenized milk. *J. Dairy Sci.*, 76, 3621-3628.

Despite strong technological corrections MNFS (moisture in non-fat substance) increased from 52.3 % in control cheeses to 55.2 % with 100 %-homogenized cream. However, MNFS did not differ statistically (54.8 %) in the range [50 %-100 %] homogenized cream but meltableity increased from 2.93 cm to 3.53 cm. Introducing MNFS data did not improve the correlation with meltableity. Also the sole humidity difference do not explain the increase of meltableity.

Free-oil was significantly reduced by increasing homogenized cream addition. Stretching was improved for the early stage of ripening (day 7). The cream fat content (30 or 42 %) did not have any effect on the melting of cheese.