

Chiral Magnetic Josephson Junction as a Base for Low-Noise Superconducting Qubits

Maxim N. Chernodub, Julien Garaud, Dmitri Kharzeev

▶ To cite this version:

Maxim N. Chernodub, Julien Garaud, Dmitri Kharzeev. Chiral Magnetic Josephson Junction as a Base for Low-Noise Superconducting Qubits. Universe, 2022, 8 (12), pp.657. 10.3390/universe8120657. hal-02263645v1

HAL Id: hal-02263645 https://hal.science/hal-02263645v1

Submitted on 9 Nov 2020 (v1), last revised 5 Jan 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chiral Magnetic Josephson junction: a base for low-noise superconducting qubits?

M. N. Chernodub, 1,2,* J. Garaud, 1,† and D. E. Kharzeev 3,4,5,‡

¹Institut Denis Poisson CNRS/UMR 7013, Université de Tours, 37200 France

²Laboratory of Physics of Living Matter, Far Eastern Federal University, Sukhanova 8, Vladivostok, 690950, Russia

³Department of Physics and Astronomy, Stony Brook University, New York 11794-3800, USA

⁴Department of Physics and RIKEN-BNL Research Center,

Brookhaven National Laboratory, Upton, New York 11973, USA

⁵Le Studium, Loire Valley Institute for Advanced Studies, Tours and Orléans, France

(Dated: August 2, 2019)

Superconducting materials with non-centrosymmetric lattices lacking the space inversion symmetry are known to exhibit a variety of interesting parity-breaking phenomena, including the anomalous Josephson effect. Here we consider a Josephson junction consisting of two non-centrosymmetric superconductors (NCSs) connected by a uniaxial ferromagnet, and demonstrate that it exhibits a direct analog of the Chiral Magnetic Effect observed in Dirac and Weyl semimetals. We propose to use this "Chiral Magnetic Josephson junction" (CMJ junction) as an element of a qubit with a Hamiltonian tunable by the ferromagnet's magnetization. The CMJ junction allows to avoid the use of an offset magnetic flux in inductively shunted qubits, thus enabling a simpler and more robust architecture. The resulting "chiral magnetic qubit" is protected from the noise caused by fluctuations in magnetization when the easy axis of the uniaxial ferromagnet is directed across the junction.

The discovery of superconductors lacking the spatial inversion symmetry [1–5] has opened the possibility to study spontaneous breaking of a continuous symmetry in a parity-violating material. In particular, the superconducting order parameter in these non-centrosymmetric superconductors (NCSs) is a parity-odd quantity [4, 5], enabling a number of interesting magnetoelectric phenomena due to mixing of singlet and triplet superconducting parameters, correlations between supercurrents and spin polarization, appearance of helical states and peculiar structure of Abrikosov vortices (see [6, 7] for a review).

Parity breaking in NCSs also results in an unconventional Josephson effect, where the junction features a phase-shifted current relation [8, 9]:

$$J(\varphi, \varphi_g) = J_c \sin(\varphi - \varphi_g). \tag{1}$$

Here φ is the superconducting phase difference across the junction, J_c the critical Josephson current, and φ_g is the parity-breaking phase offset. Nonzero bias $\varphi_g \neq 0$ results in a nonvanishing current across the junction, even when the phase difference φ is zero. Since the current is a parity-odd quantity, this clearly signals parity violation.

Phase-biased junctions (often called " φ_0 -junctions") were suggested to appear in a wide range of systems including non-centrosymmetric [8], and multilayered [10] ferromagnetic links between conventional superconductors, topological insulators [11, 12], nanowires [13, 14], quantum point contacts [15], and quantum dots [16– 18]. The first experimental realization of Josephson φ_0 junctions has been reported in superconductor–quantum dot structures, where the phase offset φ_g can be controlled via electrostatic gating [19].



Figure 1. The Chiral Magnetic Josephson junction: two non-centrosymmetric superconductors (NCSs) weakly linked by a uniaxial ferromagnet (F). The exchange field h of the ferromagnet, oriented across the link, induces an inversion symmetry-breaking component of the supercurrent (represented here by the spiral) in the junction.

In this Letter, we introduce the Josephson junction made of two NCSs weakly linked by a uniaxial ferromagnet with an easy axis normal to the interface, i.e. parallel to the electric current (see Fig. 1). Unlike in previous proposals [8, 20, 21], the ferromagnetic exchange field \boldsymbol{h} here is directed normally to the NCS/F/NCS interfaces. Parity breaking in NCS couples the magnetization \boldsymbol{h} to the supercurrent \boldsymbol{j} , resulting in a term $\boldsymbol{j} \cdot \boldsymbol{h}$ in the Ginzburg-Landau free energy functional describing crystal structure with O point group symmetry. As derived below, it results in a nonzero current even in the absence of phase gradients across the junction. This current, directed along the magnetic field, stems from the breaking of parity in a non-equilibrium state and is thus a direct analog of the Chiral Magnetic Effect [22] predicted for systems of chiral fermions, and observed in Dirac and Weyl semimetals [23–26]. This analogy motivates our terminology "Chiral Magnetic Josephson junction" (CMJ junction) to describe the NCS/F/NCS junction displayed in Fig. 1. Below, we demonstrate that the current across the CMJ junction is still given by the expression (1), where the magnitude of the bias φ_g can be tuned by the ferromagnet's magnetization.

We propose to use the CMJ junction as a constituent of a superconducting qubit. The junction's energy associated with the current (1) is

$$E(\varphi, \varphi_g) = E_J[1 - \cos(\varphi - \varphi_g)], \qquad (2)$$

where E_J is the Josephson energy. The total energy of the qubit E_Q is the sum of the junction's energy (2) and a term quadratic in the phase difference φ . For example, in the case of an inductively shunted junction [27], this quadratic term results from the inductive energy E_L :

$$E_Q(\varphi,\varphi_g) = E_J[1 - \cos(\varphi - \varphi_g)] + E_L \varphi^2.$$
(3)

Here, the offset φ_g plays the role of an offset flux, and can be used to control the form of the qubit Hamiltonian. Using the magnetization of the ferromagnetic link should simplify the qubit architecture by avoiding the use of an offset flux, and the corresponding source.

Noise in the offset flux is an important component of qubit decoherence [28, 29]. As demonstrated below, the noise in the offset phase φ_g results from the fluctuations of the component of the magnetization normal to the interface. In the proposed setup, this direction corresponds to the easy axis of the uniaxial ferromagnet. Thus only longitudinal fluctuations of magnetization contribute to the noise, but these are suppressed by the ratio of the qubit temperature to Curie temperature of the ferromagnet, which is about $10^{-5} - 10^{-4}$. Moreover, the current in the CMJ junction is parallel to magnetization, and thus is "force-free", i.e. not subjected to a Lorentz force. This greatly reduces the coupling between the current and magnetization that contributes to the noise.

The offset phase φ_g of the CMJ junction can be estimated within the Ginzburg-Landau (GL) framework. The superconducting state in non-centrosymmetric superconductors is commonly believed to be a mixture of singlet and triplet pseudo-spin states [6, 7] due to the spin-orbital coupling in the presence of the broken inversion symmetry [30]. Using $\hbar = c = 1$, the Ginzburg-Landau free energy describing the superconducting state of non-centrosymmetric material reads as [7, 31]:

$$f = a|\psi|^2 + \gamma |\mathbf{D}\psi|^2 + \frac{b}{2}|\psi|^4 + \frac{K}{2}\mathbf{j}\cdot\mathbf{h}.$$
 (4)

The single-component superconducting order parameter $\psi = |\psi|e^{i\varphi}$ is coupled to the vector potential \boldsymbol{A} of the magnetic field $\boldsymbol{h} = \boldsymbol{\nabla} \times \boldsymbol{A}$, via the gauge derivative $\boldsymbol{D} = -i\boldsymbol{\nabla} - 2e\boldsymbol{A}$, while the coefficients b, γ and $a = \alpha(T-T_c)$ are standard phenomenological GL parameters. The parity-odd nature of the non-centrosymmetric superconductor is reflected by the last term (Lifshitz invariant) of the free energy (4), which describes the direct coupling of the magnetic field \boldsymbol{h} to the usual, parity-odd component of the supercurrent density:

$$\boldsymbol{j} \equiv \boldsymbol{j}^{\text{odd}} = 2e\gamma \left[\psi^* \boldsymbol{D} \psi + \psi(\boldsymbol{D} \psi)^* \right].$$
 (5)

Note that the exchange field h of the ferromagnet plays here the role of the background magnetic field B. The parity-odd, last term in (4) yields an additional, parityeven, contribution to the total supercurrent J:

$$\boldsymbol{J} = \boldsymbol{j}^{\text{odd}} + \boldsymbol{j}^{\text{even}}, \qquad \boldsymbol{j}^{\text{even}} = 4e^2 \gamma K |\psi|^2 \boldsymbol{h}.$$
 (6)

The GL functional (4) describes NCS materials with O point symmetry, such as Li₂Pt₃B [5, 32] and Mo₃Al₂C [33, 34], and the coupling constant K determines the magnitude of the superconducting magnetoelectric effects that follow from the broken inversion symmetry. Our derivation equally applies to non-centrosymmetric superconductors with other crystallographic groups with a generic Lifshitz invariant $K_{\alpha\beta}h_{\alpha}j_{\beta}$. In this case, when the *x*-axis is directed across the normal link, the diagonal element K_{xx} should be nonzero. Notice that Lifshitz invariants of the type $\mathbf{n} \cdot \mathbf{h} \times \mathbf{j}$ do not have such diagonal element and thus cannot satisfy this requirement (see footnote [35]).

As illustrated in Fig. 1, we consider a pair of identical non-centrosymmetric superconductors separated by a uniaxial ferromagnetic weak link whose internal exchange field $\mathbf{h} \equiv h_x \mathbf{e}_x$ points across the link. We neglect the term quartic in the condensate and disregard inhomogeneities of both the condensate ψ and the exchange field \mathbf{h} in the transverse yz plane. The minimization of the GL free energy (4) with respect to the superconducting order parameter in the background of the ferromagnetic exchange field \mathbf{h} then yields the equation

$$a\psi - \gamma \frac{\partial^2 \psi}{\partial x^2} - 2ie\gamma K h_x \frac{\partial \psi}{\partial x} = 0 \tag{7}$$

that describes the tunneling of the Cooper pairs across the weak link. For the time being we assume the absence of an external electromagnetic field at the link, $\mathbf{A} = 0$. Due to proximity effects, the tunneling of the Cooper pairs between the non-centrosymmetric superconductors through the centrosymmetric weak link will not respect the parity inversion $x \to -x$, as can be seen from Eq. (7).

The general solution of Eq. (7) for the superconducting gap inside the weak link reads as:

$$\psi(x) = C_+ e^{q_+ x} + C_- e^{q_- x},\tag{8}$$

where the wavevectors

$$q_{\pm} = \pm \sqrt{\frac{a}{\gamma} - \left(eh_x K\right)^2} - ieh_x K \tag{9}$$

should have a nonzero real part so that the weak link is in a normal state, thus requiring $a > a_c = \gamma (eh_x K)^2$. The coefficients C_{\pm} in Eq. (8) are determined by the boundary conditions at the interfaces of the ferromagnetic weak link with the superconductors at $x = \pm L/2$. It is customary to make a simplification using the rigid boundary conditions [8, 36] which assume the absence of a barrier at the interfaces, and imply continuity of the superconducting order parameter:

$$\psi(x = \pm L/2) = |\Delta| e^{\pm i\varphi/2}; \tag{10}$$

here $|\Delta|$ is the absolute value of the order parameter at the superconducting leads.

Using the relations Eqs. (8-10), together with the definition of the total current (6), yields the phase-shifted current relation:

$$J = J_0 \sin\left(\varphi - \varphi_g\right), \quad J_0 = \frac{4e\gamma|\Delta|^2 \sqrt{\frac{a}{\gamma} - (eh_x K)^2}}{\sinh L \sqrt{\frac{a}{\gamma} - (eh_x K)^2}}, (11)$$

which exhibits the offset of the phase difference given by

$$\varphi_g = eh_x KL. \tag{12}$$

This offset, corresponding to the broken inversion symmetry, is proportional to the strength of the magnetic interaction. In other words, the presence of the nonzero phase bias $\varphi_g \neq 0$ signals the breaking of the inversion symmetry between leftward and rightward tunneling of the Cooper pairs, and leads to a nonzero current in the "steady state" of the junction even if the phase difference between the superconducting leads is zero, $\varphi = 0$.

In a long junction, $L\sqrt{a/\gamma - (eh_x K)^2} \gg 1$, the current (11) is an exponentially small quantity due to suppression of the Cooper-pair tunneling between widely separated superconducting leads. The limit of a short junction in the presence of parity breaking should be taken with care. In the thermodynamic equilibrium $\varphi = \varphi_g(L)$, and the current through the junction is always zero. However, in the steady state with zero phase difference $\varphi = 0$, the electric current does not vanish, and is given by the parity–even term (6):

$$\boldsymbol{J}(\varphi = 0, L \to 0) = 4e^2 \gamma K |\Delta|^2 \boldsymbol{h}.$$
 (13)

This current plays a crucial role in the dynamics of the chiral magnetic qubit. As mentioned earlier, the current (13) shares a striking similarity with the Chiral Magnetic Effect [22], with h and K respectively playing the roles of the external magnetic field and the source of the parity breaking.

Some features of the phenomenon that we discuss appear also in usual centrosymmetric *s*-wave superconductors separated by the Josephson junction made of an NCS-type ferromagnet with *tangentially*-oriented field h. Even though the underlying dynamics is quite different, the latter system is described by an equation similar to (7) [8]. However, in our case the current flows along the magnetic field, is force-free, and thus is not subjected to the noise resulting from the transverse fluctuations of magnetization.

Similar types of Josephson junctions with the ferromagnetic exchange field h oriented transversally between two non-centrosymmetric superconductors with the C_{4v} point group (corresponding to interactions of the type $n \cdot h \times j$) have been proposed in Refs. [20, 21]. The Josephson current in a non-ferromagnetic junction between two non-centrosymmetric superconductors does not exhibit the offset in the phase [37]. A finite phase offset for the Josephson current of chiral charge appears, however, for the junction between two Weyl superconductors separated by a Weyl semimetal, also for magnetic field oriented transversally [38].

We estimate the phase bias (12) numerically as follows:

$$\varphi_g \simeq 1.5 \times 10^{-3} \, h(\text{T}) \, L(\text{nm}) \, K(\text{nm}).$$
 (14)

The length of the ferromagnetic Josephson junction is typically of the order of tens of nanometers $(L \sim 30 \text{ nm})$ in Ref. [39]). The exchange field h should not exceed the upper critical field H_{c2} which, for a number of NCS superconductors, may reach significant values $H_{c2} \sim 10 \,\mathrm{T}$. The fields of this order and higher are known to be created by usual ferromagnets [40]. Note that even for magnetic fields larger than H_{c1} , the vortex formation can be avoided by choosing a weak link with a sufficiently small cross section, such that a total magnetic flux entering the superconductor is smaller than the flux quantum Φ_0 . The main uncertainty in our estimate comes from the poorly known parity-odd coupling K. Its value was estimated to be $K \simeq (10^{-3} \dots 10^{-2}) \lambda$ [41, 42], where $\lambda \simeq (0.1 \dots 1) \mu m$ is the penetration depth. In spite of this uncertainty, it appears that the phase bias may be tuned to take values of order $\varphi_g \sim \pi$. For a given NCS superconductor, the phase bias can be manipulated by the magnetization of the weak link.

The chiral magnetic Josephson junction sketched in Fig. 1 can be inductively shunted, for example by a series of conventional Josephson junctions, to form a "chiral magnetic qubit" (see Fig. 2). Such circuits include, in addition, two mixed Josephson junctions between the conventional and NCS superconductors. These mixed junctions do not generate the electric current across them at zero phase difference $\varphi = 0$ [43].

The Coulomb interactions between the Cooper pairs is described by the kinetic term in the Hamiltonian of the



Figure 2. (a) Fluxonium-type qubit based on conventional Josephson junction inductively shunted by a series of Josephson junctions. The qubit is biased by an external magnetic flux $\varphi_{\text{ext}} \equiv 2\pi \Phi/\Phi_0$, where Φ_0 is the elementary flux quantum. The gate phase offset for a conventional Josephson junction is absent, $\varphi_g = 0$. (b) Chiral magnetic qubit based on the Chiral Magnetic Josephson (CMJ) junction inductively shunted by a series of Josephson junctions. The CMJ junction possesses an internal phase offset $\varphi_g \neq 0$ eliminating the need for an external magnetic flux φ_{ext} .

qubit:

$$\hat{H} = 4E_C \hat{n}^2 + E_J [1 - \cos(\varphi - \varphi_g)] + E_L \varphi^2, \quad (15)$$

where $\hat{n} = -i\hbar\partial_{\varphi}$ is the operator of the Cooper-pair number, and last two terms describe the Josephson tunneling and the induction (3). The Hamiltonian (15) is generic for a family of inductively shunted qubits including the fluxonium [44, 45], one-junction flux qubits, and fluxbiased phase qubits [46].

As illustrated in Fig. 2(a), fluxonium qubits relate the phase offset to the externally applied flux Φ as $\varphi_g = 2\pi \Phi/\Phi_0$, where $\Phi_0 = h/(2e)$ is the flux quantum. These are further characterized by a specific set of model parameters, such as the small inductive energy $(E_L/E_J \simeq 0.045)$ and a moderate charging energy $(E_C/E_J \lesssim 1)$ which give a unique combination of long coherence time and large anharmonicity of the energy levels [47]. Transmon qubits, on the other hand, are characterized by Coulomb charging energy which is much smaller than the Josephson tunneling energy, $E_C \ll E_J$, thus allowing the reduction of noise caused by the offset charge fluctuations [48].

A nonzero phase bias $\varphi_g \neq 0$ imposes a large anharmonicity on the energy-level structure [27] determined by the Schröringer equation:

$$\hat{H}\psi_n(\varphi) = \varepsilon_n \psi_n(\varphi).$$
 (16)

The regime $\varphi_g = \pi/2$ provides maximum level splitting and the absence of nearly-degenerate level pairs [27]. Fig. 3 displays the structure of the energy levels corresponding to this Hamiltonian. The transitions between the first excited state and the ground state, $|1\rangle \rightarrow |0\rangle$, can be substantially suppressed by the barrier separating them. This barrier is almost absent in the typical fluxonium regime ($E_L = 0.045E_J$ and $E_C = E_J$), as the first excited energy level ε_1 practically coincides with the height of the barrier, Fig. 3(a). This conclusion is valid,



Figure 3. The potential energy (3) of the chiral magnetic qubit with the chiral magnetic Josephson junction possessing the phase offset $\varphi_g = \pi/2$ for various Coulomb charging energies E_C and inductive energies E_L . Lowest eigenstates $|n\rangle$ with $n = 0, 1, \ldots$ are shown along with the numerically computed energy levels ε_n and the corresponding wavefunctions $\psi_n(\varphi)$.

to a good accuracy, for a wide range of values of the phase offset φ_q .

Decreasing the Coulomb energy towards the transmon regime leads to the appearance of the prohibitive barrier for transitions $|1\rangle \rightarrow |0\rangle$ between the different wells, Fig. 3(b). As displayed in Fig. 3(c), further decrease of the Coulomb energy reduces the energy difference between $|1\rangle$ and $|0\rangle$ states. The lifetime of the first excited level may be enhanced by lowering the inductive energy E_L . Figure 3(d) shows that at $E_L = 0.01E_J$ the barrier is sufficiently high to ensure a quasi-classical protection of the first excited level. Related discussions of the energy levels can be found in Refs. [47, 49] for fluxonium-type qubits.

The conventional way to induce the phase φ_g in fluxonium qubits is to apply a background magnetic flux Φ . The noise $\delta \Phi / \Phi_0$ in magnetic flux is typically of the order of $10^{-3} - 10^{-2}$; overcoming this noise is a central problem in quantum computer design. In our case, the noise in φ_g is due to the noise in magnetization. Indeed, Eq. (12) yields the noise relation

$$\left(\frac{\delta\varphi_g}{\varphi_g}\right) = eKL\left(\frac{\delta h_x}{h_x}\right). \tag{17}$$

To reduce the noise in magnetization, we propose to use a highly anisotropic uniaxial ferromagnet, e.g. of magnetoplumbite type. In the case of a uniaxial ferromagnet, the 3D rotational symmetry is explicitly broken by the symmetry of the crystalline lattice, and the only surviving symmetry is 2D rotations around the easy symmetry axis of the ferromagnet, in the basal plane perpendicular to this axis. The fluctuations of magnetization in this case are given by the simplified form of the LandauLifshitz-Gilbert equation with no gyroscopic term. They correspond to the rotation of magnetization around the easy axis (which points here along the x axis), with fluctuating components of magnetization h_y and h_z , but with a fixed h_x which is an integral of motion. Since the phase offset φ_g (12) depends only on h_x , the transverse fluctuations of magnetization will not induce noise in this quantity.

Unlike the transverse ones, the longitudinal fluctuations of magnetization (i.e. fluctuations of the magnitude of h) will induce a noise in the offset phase φ_q . However, longitudinal fluctuations of the magnetization are expected to be suppressed compared to the transverse ones by a factor of cT/T_C , where T is the temperature, T_C is Curie temperature of the ferromagnet, and c is a constant of order one. Indeed, the transverse fluctuations correspond to gapless Goldstone modes with kinetic energy $\sim T$, while the longitudinal one is massive with energy $\sim T_C$. Analysis [50, 51] of the Landau-Lifshitz-Bloch equation (including both transverse and longitudinal fluctuations of magnetization) indicates that the constant $c \simeq 2/3$. Therefore, at temperatures of the superconducting qubits that are on the order of tens of milli-Kelvin, with Curie temperatures on the order of 1,000 K, we expect the suppression of longitudinal fluctuations by a factor of $\sim 10^{-4} - 10^{-5}$. This allows to expect a suppression in the noise resulting from the offset flux of CMJ junction, as compared to external flux noise, by a significant factor of 10^{-2} .

The domain structure in uniaxial ferromagnets is known to crucially depend on the anisotropy [52]: in weakly anisotropic ferromagnets (Landau-Lifshitz type), there is a branching of domains close to the surface, and the magnetic flux does not leave the ferromagnet. This is not a desirable domain configuration, as the magnetic field has to penetrate the superconductor. On the other hand, in strongly anisotropic ferromagnets (Kittel type), such as magnetoplumbite, the domains do not branch, and thus the magnetic flux does escape the ferromagnet. The domain structure of a thin ferromagnetic film [53] may be affected by the superconducting interface; this question requires further investigation.

To summarize, we have introduced a Josephson junction consisting of two non-centrosymmetric superconductors connected by a uniaxial ferromagnet, and have demonstrated that it exhibits a direct analog of the Chiral Magnetic Effect. We have proposed this Chiral Magnetic Josephson junction (CMJ junction) for use as an element of a qubit with parameters tunable by the ferromagnet's magnetization. The resulting Chiral Magnetic Qubit is protected from noise caused by fluctuations in magnetization, and does not require an external magnetic flux, allowing for a simpler and more robust architecture. The main uncertainty stems from the poorly known parity-odd response of non-centrosymmetric superconductors, and we believe that these materials and the properties of their interfaces deserve further studies.

The work of D.K. was supported by the U.S. Department of Energy, Office of Nuclear Physics, under contracts DE-FG-88ER40388 and DE-AC02-98CH10886, and by the Office of Basic Energy Science under contract DE-SC-0017662.

- * maxim.chernodub@idpoisson.fr
- [†] garaud.phys@gmail.com
- [‡] dmitri.kharzeev@stonybrook.edu
- A.I. Rusinov L.N. Bulaevskii, A.A. Guseinov, "Superconductivity in crystals without symmetry centers," Soviet Journal of Experimental and Theoretical Physics 44, 1243 (1976), [Russian original - Zh. Eksp. Teor. Fiz., Vol. 71, No. 6, p. 2356, December 1976].
- [2] L. S. Levitov, Y. V. Nazarov, and G. M. Éliashberg, "Magnetostatics of superconductors without an inversion center," Soviet Journal of Experimental and Theoretical Physics Letters 41, 445 (1985), [Russian original - Pis'ma Zh. Eksp. Teor. Fiz., Vol. 41, No. 9, p. 365, May 1985].
- [3] E. Bauer, G. Hilscher, H. Michor, Ch. Paul, E. W. Scheidt, A. Gribanov, Yu. Seropegin, H. Noël, M. Sigrist, and P. Rogl, "Heavy Fermion Superconductivity and Magnetic Order in Noncentrosymmetric CePt₃Si," Phys. Rev. Lett. **92**, 027003 (2004).
- [4] K. V. Samokhin, E. S. Zijlstra, and S. K. Bose, "CePt₃Si : An unconventional superconductor without inversion center," Phys. Rev. B 69, 094514 (2004).
- [5] H. Q. Yuan, D. F. Agterberg, N. Hayashi, P. Badica, D. Vandervelde, K. Togano, M. Sigrist, and M. B. Salamon, "S-Wave Spin-Triplet Order in Superconductors without Inversion Symmetry: Li₂Pd₃B and Li₂Pt₃B," Phys. Rev. Lett. **97**, 017006 (2006).
- [6] E. Bauer and M. Sigrist, Non-Centrosymmetric Superconductors: Introduction and Overview, edited by E. Bauer and M. Sigrist, Lecture notes in physics (Springer, 2012).
- [7] M. Smidman, M. B. Salamon, H. Q. Yuan, and D. F. Agterberg, "Superconductivity and spin-orbit coupling in non-centrosymmetric materials: a review," Reports on Progress in Physics 80, 036501 (2017).
- [8] A. Buzdin, "Direct Coupling Between Magnetism and Superconducting Current in the Josephson φ_0 Junction," Phys. Rev. Lett. **101**, 107005 (2008).
- [9] F. Konschelle and A. Buzdin, "Magnetic Moment Manipulation by a Josephson Current," Phys. Rev. Lett. 102, 017001 (2009).
- [10] Jun-Feng Liu and K. S. Chan, "Anomalous Josephson current through a ferromagnetic trilayer junction," Phys. Rev. B 82, 184533 (2010).
- [11] Yukio Tanaka, Takehito Yokoyama, and Naoto Nagaosa, "Manipulation of the Majorana Fermion, Andreev Reflection, and Josephson Current on Topological Insulators," Phys. Rev. Lett. **103**, 107002 (2009).
- [12] Fabrizio Dolcini, Manuel Houzet, and Julia S. Meyer, "Topological Josephson ϕ_0 junctions," Phys. Rev. B **92**, 035428 (2015).

- [13] Tomohiro Yokoyama, Mikio Eto, and Yuli V. Nazarov, "Anomalous Josephson effect induced by spin-orbit interaction and Zeeman effect in semiconductor nanowires," Phys. Rev. B 89, 195407 (2014).
- [14] G Campagnano, P Lucignano, D Giuliano, and A Tagliacozzo, "Spin-orbit coupling and anomalous Josephson effect in nanowires," Journal of Physics: Condensed Matter 27, 205301 (2015).
- [15] A. A. Reynoso, Gonzalo Usaj, C. A. Balseiro, D. Feinberg, and M. Avignon, "Anomalous Josephson Current in Junctions with Spin Polarizing Quantum Point Contacts," Phys. Rev. Lett. **101**, 107001 (2008).
- [16] A. Zazunov, R. Egger, T. Jonckheere, and T. Martin, "Anomalous Josephson Current through a Spin-Orbit Coupled Quantum Dot," Phys. Rev. Lett. 103, 147004 (2009).
- [17] L. Dell'Anna, A. Zazunov, R. Egger, and T. Martin, "Josephson current through a quantum dot with spinorbit coupling," Phys. Rev. B 75, 085305 (2007).
- [18] Aldo Brunetti, Alex Zazunov, Arijit Kundu, and Reinhold Egger, "Anomalous Josephson current, incipient time-reversal symmetry breaking, and Majorana bound states in interacting multilevel dots," Phys. Rev. B 88, 144515 (2013).
- [19] D. B. Szombati, S. Nadj-Perge, D. Car, S. R. Plissard, E. P. A. M. Bakkers, and L. P. Kouwenhoven, "Josephson Φ_0 -junction in nanowire quantum dots," Nature Physics **12**, 568 (2016).
- [20] Huan Zhang, Jun Wang, and Jun-Feng Liu, "Anomalous Josephson effect in noncentrosymmetric superconductors," Applied Physics Letters 108, 102601 (2016).
- [21] Yousef Rahnavard, Dirk Manske, and Gaetano Annunziata, "Magnetic Josephson junctions with noncentrosymmetric superconductors," Phys. Rev. B 89, 214501 (2014).
- [22] Kenji Fukushima, Dmitri E. Kharzeev, and Harmen J. Warringa, "Chiral magnetic effect," Phys. Rev. D 78, 074033 (2008).
- [23] D. E. Kharzeev and H. J. Warringa, "Chiral Magnetic conductivity," Physical Review D 80, 034028 (2009).
- [24] Q. Li, D. E. Kharzeev, C. Zhang, Y. Huang, I. Pletikosić, A. V. Fedorov, R. D. Zhong, J. A. Schneeloch, G. D. Gu, and T. Valla, "Chiral magnetic effect in ZrTe₅," Nature Physics **12**, 550 (2016).
- [25] J. Xiong, S. K. Kushwaha, T. Liang, J. W. Krizan, M. Hirschberger, W. Wang, R. J. Cava, and N. P. Ong, "Evidence for the chiral anomaly in the Dirac semimetal Na₃Bi," Science **350**, 413 (2015).
- [26] X. Huang, L. Zhao, Y. Long, P. Wang, D. Chen, Z. Yang, H. Liang, M. Xue, H. Weng, Z. Fang, X. Dai, and G. Chen, "Observation of the chiral anomaly induced negative magneto-resistance in 3D Weyl semimetal TaAs," Physical Review X 5, 031023 (2015).
- [27] S. M. Girvin, "Quantum Machines: Measurement and Control of Engineered Quantum Systems," (Oxford Scholarship Online, Les Houches, France, 2014) Chap. Circuit QED: superconducting qubits coupled to microwave photons, pp. 113–256, Oxford University Press.
- [28] A. A. Clerk, M. H. Devoret, S. M. Girvin, Florian Marquardt, and R. J. Schoelkopf, "Introduction to quantum noise, measurement, and amplification," Rev. Mod. Phys. 82, 1155–1208 (2010).
- [29] M. H. Devoret, A. Wallraff, and J. M. Martinis, "Superconducting Qubits: A Short Review," arXiv e-prints

, cond-mat/0411174 (2004), arXiv:cond-mat/0411174.

- [30] Lev P. Gor'kov and Emmanuel I. Rashba, "Superconducting 2D System with Lifted Spin Degeneracy: Mixed Singlet-Triplet State," Phys. Rev. Lett. 87, 037004 (2001).
- [31] D. F. Agterberg, "Non-Centrosymmetric Superconductors: Introduction and Overview," in Non-Centrosymmetric Superconductors: Introduction and Overview, edited by Ernst Bauer and Manfred Sigrist (Springer Berlin Heidelberg, Berlin, Heidelberg, 2012) Chap. Magnetoelectric Effects, Helical Phases, and FFLO Phases, pp. 155–170.
- [32] Petre Badica, Takaaki Kondo, and Kazumasa Togano, "Superconductivity in a New Pseudo-Binary Li₂B(Pd_{1-x}Pt_x)₃ (x=0-1) Boride System," Journal of the Physical Society of Japan **74**, 1014–1019 (2005).
- [33] A. B. Karki, Y. M. Xiong, I. Vekhter, D. Browne, P. W. Adams, D. P. Young, K. R. Thomas, Julia Y. Chan, H. Kim, and R. Prozorov, "Structure and physical properties of the noncentrosymmetric superconductor Mo₃Al₂C," Phys. Rev. B 82, 064512 (2010).
- [34] E. Bauer, G. Rogl, Xing-Qiu Chen, R. T. Khan, H. Michor, G. Hilscher, E. Royanian, K. Kumagai, D. Z. Li, Y. Y. Li, R. Podloucky, and P. Rogl, "Unconventional superconducting phase in the weakly correlated noncentrosymmetric Mo₃Al₂C compound," Phys. Rev. B 82, 064511 (2010).
- [35] Possible candidates for the NCS superconductors should thus have a crystalline structure with either the point group O (Li₂Pt₃B, Mo₃Al₂C), T point group (e.g. LaRhSi, LaIrSi) or C_4 (La₅B₂C₆), C_2 (UIr) etc. On the other hand, the point groups $C_{n\nu}$ with n = 2, 3, 4, 6(possessed, for example, by the compounds MoS₂, MoN, GaN, CePt₃Si, CeRhSi₃, CeIrSi₃ [7]) correspond to the Lifshitz invariants of the type $\boldsymbol{n} \cdot \boldsymbol{h} \times \boldsymbol{j}$ that do not fit our proposal.
- [36] A. A. Golubov, M. Yu. Kupriyanov, and E. Il'ichev, "The current-phase relation in Josephson junctions," Rev. Mod. Phys. 76, 411–469 (2004).
- [37] Yasuhiro Asano and Satoshi Yamano, "Josephson effect in noncentrosymmetric superconductor junctions," Phys. Rev. B 84, 064526 (2011).
- [38] Song-Bo Zhang, Johanna Erdmenger, and Björn Trauzettel, "Chirality Josephson Current Due to a Novel Quantum Anomaly in Inversion-Asymmetric Weyl Semimetals," Phys. Rev. Lett. **121**, 226604 (2018).
- [39] V. V. Ryazanov, V. A. Oboznov, A. Yu. Rusanov, A. V. Veretennikov, A. A. Golubov, and J. Aarts, "Coupling of Two Superconductors through a Ferromagnet: Evidence for a π Junction," Phys. Rev. Lett. **86**, 2427–2430 (2001).
- [40] A. I. Buzdin, "Proximity effects in superconductorferromagnet heterostructures," Rev. Mod. Phys. 77, 935– 976 (2005).
- [41] Chi-Ken Lu and Sungkit Yip, "Transverse Magnetic Field Distribution in the Vortex State of Noncentrosymmetric Superconductor with OSymmetry," Journal of Low Temperature Physics 155, 160–168 (2009).
- [42] M. K. Kashyap and D. F. Agterberg, "Vortices in cubic noncentrosymmetric superconductors," Phys. Rev. B 88, 104515 (2013).
- [43] Nobuhiko Hayashi, Christian Iniotakis, Masahiko Machida, and Manfred Sigrist, "Josephson effect between conventional and non-centrosymmetric superconductors," Journal of Physics and Chemistry of Solids 69,

3225 - 3227 (2008).

- [44] Vladimir E. Manucharyan, Jens Koch, Leonid I. Glazman, and Michel H. Devoret, "Fluxonium: Single Cooper-Pair Circuit Free of Charge Offsets," Science **326**, 113–116 (2009).
- [45] Ioan M. Pop, Kurtis Geerlings, Gianluigi Catelani, Robert J. Schoelkopf, Leonid I. Glazman, and Michel H. Devoret, "Coherent suppression of electromagnetic dissipation due to superconducting quasiparticles," Nature 508, 369 (2014).
- [46] G Wendin, "Quantum information processing with superconducting circuits: a review," Reports on Progress in Physics 80, 106001 (2017).
- [47] Yen-Hsiang Lin, Long B. Nguyen, Nicholas Grabon, Jonathan San Miguel, Natalia Pankratova, and Vladimir E. Manucharyan, "Demonstration of Protection of a Superconducting Qubit from Energy Decay," Phys. Rev. Lett. **120**, 150503 (2018).
- [48] Jens Koch, Terri M. Yu, Jay Gambetta, A. A. Houck, D. I. Schuster, J. Majer, Alexandre Blais, M. H. Devoret, S. M. Girvin, and R. J. Schoelkopf, "Charge-insensitive qubit design derived from the Cooper pair box," Phys. Rev. A 76, 042319 (2007).

- [49] N. Earnest, S. Chakram, Y. Lu, N. Irons, R. K. Naik, N. Leung, L. Ocola, D. A. Czaplewski, B. Baker, Jay Lawrence, Jens Koch, and D. I. Schuster, "Realization of a Λ System with Metastable States of a Capacitively Shunted Fluxonium," Phys. Rev. Lett. **120**, 150504 (2018).
- [50] D. A. Garanin, V. V. Ishchenko, and L. V. Panina, "Dynamics of an ensemble of single-domain magnetic particles," Theoretical and Mathematical Physics 82, 169–179 (1990).
- [51] O. Chubykalo-Fesenko, U. Nowak, R. W. Chantrell, and D. Garanin, "Dynamic approach for micromagnetics close to the Curie temperature," Phys. Rev. B 74, 094436 (2006).
- [52] Jan Kaczér, "On the domain structure of uniaxial ferromagnets," Soviet Journal of Experimental and Theoretical Physics **19**, 1204 (1964), [Russian original - ZhETF, Vol. 46, No. 5, p. 1787, November 1964].
- [53] L. N. Bulaevskii and E. M. Chudnovsky, "Ferromagnetic film on a superconducting substrate," Phys. Rev. B 63, 012502 (2000).