



Extreme hydroclimate response gradients within the western Cape Floristic region of South Africa since the Last Glacial Maximum

Brian Chase, Arnoud Boom, Andrew Carr, Manuel Chevalier, Lynne J Quick, G. Anthony Verboom, Paula Reimer

► To cite this version:

Brian Chase, Arnoud Boom, Andrew Carr, Manuel Chevalier, Lynne J Quick, et al.. Extreme hydroclimate response gradients within the western Cape Floristic region of South Africa since the Last Glacial Maximum. *Quaternary Science Reviews*, 2019, 219, pp.297-307. 10.1016/j.quascirev.2019.07.006 . hal-02264496

HAL Id: hal-02264496

<https://hal.science/hal-02264496>

Submitted on 7 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Extreme hydroclimate response gradients within the western Cape Floristic region of South Africa since the Last Glacial Maximum

The authors do not recommend the distribution of this version of this article.

The article is freely available upon request.

To receive a copy, please send a request to Brian Chase at:
brian.chase@umontpellier.fr

Brian M. Chase^{*,a}, Arnoud Boom^b, Andrew S. Carr^b, Manuel Chevalier^c, Lynne J. Quick^d,
G. Anthony Verboom^e, Paula J. Reimer^f

^a*Institut des Sciences de l'Evolution-Montpellier (ISEM), University of Montpellier, Centre National de la Recherche Scientifique (CNRS), EPHE, IRD, Montpellier, France;*

^b*School of Geography, Geology and the Environment, University of Leicester, Leicester, LE1 7RH, UK;*

^c*Institute of Earth Surface Dynamics, Geopolis, University of Lausanne, Quartier UNIL-Mouline, Bâtiment Géopolis, CH-1015 Lausanne, Switzerland;*

^d*African Centre for Coastal Palaeoscience, Nelson Mandela University, 6031 Port Elizabeth, South Africa;*

^e*Department of Biological Sciences, University of Cape Town, 7700 Rondebosch, South Africa;*

^f*School of Natural and Built Environment, Queen's University Belfast, Belfast, BT7 1NN, Northern Ireland, UK;*

***Correspondence and requests should be addressed to:** Brian.Chase@umontpellier.fr, +33 (0)6 01 06 64 04

26 **Abstract**

27 The Cape Floristic Region (CFR) is one of the world's major biodiversity hotspots, and much
28 work has gone into identifying the drivers of this diversity. Considered regionally in the context
29 of Quaternary climate change, climate stability is generally accepted as being one of the major
30 factors promoting the abundance of species now present in the CFR. However, little direct
31 evidence is available from the region, and responses to changes in global boundary conditions
32 have been difficult to assess. In this paper, we present new high-resolution stable isotope data
33 from Pakhuis Pass, in the species-rich western CFR, and contextualise our findings through
34 comparison with other records from the region. Combined, they indicate clear, coherent
35 changes in regional hydroclimate, which we relate to broader forcing mechanisms. However,
36 while these climate change events share similar timings (indicating shared macro-scale drivers),
37 the responses are distinct between sites, in some cases expressing opposing trends over very
38 short spatial gradients (<50 km). We describe the evolution of these trends, and propose that
39 while long-term (10^5 yr) general climatic stability may have fostered high diversity in the region
40 through low extinction rates, the strong, abrupt changes in hydroclimate gradients observed in
41 our records may have driven a form of allopatric speciation pump, promoting the diversification
42 of plant lineages through the periodic isolation and recombination of plant populations.

43 **Keywords**

44 Palaeoclimate, climate dynamics, South Africa, Cape Floristic Region, biodiversity, rock hyrax
45 middens, stable isotopes

46 **Highlights**

- 47 • New high resolution $\delta^{15}\text{N}$ data from rock hyrax middens from the western Cape Floristic
48 Region.
- 49 • Position of the southern westerlies apparent as a strong determinant of regional
50 hydroclimates.
- 51 • High amplitude, rapid changes in hydroclimate are revealed in the region.
- 52 • Opposing responses to shared climate change events establish highly variable climate
53 gradients.

1 Introduction

Southern African hydroclimate is primarily defined by the advection and precipitation of moisture from tropical Atlantic and Indian Ocean sources during the austral summer (Figure 1). In contrast, southwestern South Africa experiences a distinct, inverse rainfall regime. Seasonal expansions of the circumpolar vortex and equatorward migration of the westerly storm track bring increased precipitation during winter months, while displacements of the South Atlantic Anticyclone during the summer intensify upwelling along the west coast, blocking the westward propagation of easterly waves, generating strong summer drought in the region (Tyson, 1986; Tyson and Preston-Whyte, 2000). During the Quaternary, southern Africa is thought to have been sensitive to long-term changes in these tropical and temperate systems. Driven by changes in global boundary conditions, tropical systems are invigorated during interglacial periods and temperate systems exert increased influence during glacial periods (Chase et al., 2015a; Chase et al., 2017; see Chase and Meadows, 2007; Cockcroft et al., 1988; Quick et al., 2011; van Zinderen Bakker, 1976).

This dynamic, which is thought to have become dominant in the Plio-Pleistocene (5.0 – 2.6 Ma) when trade wind-controlled upwelling was established as a dominant factor in the regional climate system (Diekmann et al., 2003), has created a Mediterranean climate zone in the southwestern Cape of South Africa (Figure 2). This climatic evolution has been a key factor in fostering the development of the vegetation of the Cape Floristic Region (CFR), which is remarkable for its high level of endemism and its species richness (Cowling, 1992; Goldblatt, 1978).

It has been recognised, however, that species diversity within the CFR, is not homogeneous, with the winter-rain dominated western CFR having more than twice the species per area of the eastern CFR, which currently experiences an aseasonal rainfall regime (Figure 2) (Cowling et al., 1992; Cowling and Lombard, 2002; Cowling et al., 1997). During the Quaternary, the western CFR has putatively experienced a continuous dominance of temperate systems, whereas the eastern CFR rainfall regimes may have oscillated between temperate and tropical rainfall dominance (see Chase and Meadows, 2007) with much more significant consequences on the distribution and nature of environmental niches (Cowling et al., 1999; Cowling et al., 1992). With a reliable climate regime, and relatively muted cycles of Quaternary climate change compared to the extratropical regions in the Northern Hemisphere (see Chase and Meadows, 2007; Dynesius and Jansson, 2000), it is believed that the CFR's great floral diversity – particularly in the west – is largely attributable to relative climatic stability and resulting low extinction rates (Cowling et al., 1992; Cowling and Lombard, 2002; Cowling et al., 2015; Cowling et al., 1997; Linder, 2005; Verboom et al., 2014). These hypotheses, and the conclusion that the western CFR represented a more durable climatic niche than the eastern CFR is supported by analyses of phylogenetic diversity, which indicate that the western CFR experienced higher levels of in situ radiation, while the eastern CFR indicates significant mixing of lineages from different biomes (Forest et al., 2007; Verboom et al., 2014).

At broad spatio-temporal scales, it thus seems apparent that the development of, and the dynamics within, the southwestern Cape's Mediterranean climate region were fundamental determinants for the evolution of the CFR. At finer spatial and temporal scales, these considerations become more complex. In the western CFR, it has been observed that species

richness is higher in the mountains than the lowlands, a situation related to topographic variability and the resulting diversity of edaphic and climate spaces (Linder, 1991; Verboom et al., 2015). Linder (1991) also found that species richness correlated most strongly with mean annual precipitation, which might further suggest that climatic stability is a key determinant of species diversity (as regions of currently high rainfall may be acting as interglacial refugia for many CFR taxa (Chase and Meadows, 2007; Forest et al., 2007)). But how stable were climates in these montane regions?

In this paper, we consider the climatic context of the high-diversity montane regions of the western CFR in the light of new, high resolution palaeoclimatic data obtain from the region. While neither spatially nor temporally comprehensive, it does enable the beginnings of a more detailed analysis and understanding of the nature of climate change dynamics in the region. While some studies have suggested long-term millennial-scale climate and vegetation stability (Meadows et al., 2010; Meadows and Sugden, 1991), a growing body of data indicates potentially more dynamic patterns of climate change, and further indicates that spatial climate gradients may be much more complex than previously supposed (Chase et al., 2015a; Chase et al., 2011). To address this, we present a new 19,300-year stable isotope composite record derived from rock hyrax middens from Pakhuis Pass in the Cederberg Mountains. Considered together with the records from other hyrax midden records from De Rif (Chase et al., 2015a; Chase et al., 2011) and Katbakkies Pass (Chase et al., 2015b; Meadows et al., 2010), we are able to assess the spatio-temporal dynamics of climate change along a modern climatic gradient in the Cape Fold Belt Mountains of the western CFR – across the eastern slope of the Cederberg , and determine to what extent regional climates can be considered to be stable.

1.1 Regional setting

Pakhuis Pass is located in the northern Cederberg Mountains, the dominant range of the north-south axis of the Cape Fold Mountains to the east and northeast of Cape Town (Figures 2, 3). Extending for ~200 km parallel to the Atlantic Ocean (50-100 km to the west), this western limb of the Cape Fold Belt is a significant divide between the relatively humid climates of the southwestern Cape and the arid Karoo, which dominates much of South Africa's western continental interior. The range also broadly marks the divide between southern Africa's two major climate regimes: the winter rainfall zone (WRZ) to the west and the summer rainfall zone (SRZ) to the northeast (cf. Chase and Meadows, 2007). The winter rainfall zone is defined by the seasonal intensification and northward expansion of the westerlies and associated frontal depressions that transport moisture to the region during the austral winter months. To the east, and across most of South Africa, tropical easterly flow transports moisture from the Indian Ocean during the austral summer. The Cederberg and adjacent ranges act as an orographic divide between these climate zones, with the mountains creating a distinct rainshadow for westerly derived rainfall (Figure 3; Tyson, 1986; Tyson and Preston-Whyte, 2000). The higher elevations receive five times the precipitation of the lowlands to the east, but perhaps more importantly, while the Cederberg Mountains receive more than 75% of their rainfall during the winter, the lowlands to the east currently receive 50% or more of their precipitation during the summer (Hijmans et al., 2005) (Figure 1).

Pakhuis Pass is located at ~485 m.a.s.l. on the eastern slopes of the Cederberg, in the rainshadow of the Pakhuisberge massif (~1000 m.a.s.l.) (Figure 3). Mean annual rainfall at the site is ~250-300 mm/yr, ~80% of which falls in the austral winter between April and September

(Hijmans et al., 2005). In comparison, the hyrax midden sites at De Rif (Chase et al., 2015a; Chase et al., 2011; Quick et al., 2011) and Katbakkies Pass (Chase et al., 2015b; Meadows et al., 2010) (Figure 2, 3) are both situated at ~1150 m.a.s.l., receive ~400 mm/yr and ~300 mm/yr of mean annual rainfall respectively, and by virtue of their elevation and position relative to the region's major topographic features they are more likely to receive precipitation during the summer months through orographic amplification of local or larger synoptic-scale systems (Tyson, 1986; Tyson and Preston-Whyte, 2000).

2 Material and methods

Middens for this study, PK08 and PK10-1 (Figure 4) were collected from the same ~400 m long, 15 m high cliff band (32.093°S, 19.065°E) as those described by Scott and Woodborne (2007a, b) and were selected for their high hyraceum (crystallised urine) relative to faecal pellet content. Apart from having greater structural integrity and less variable deposition rates, hyraceum represents environmental conditions more clearly than samples containing faecal pellets, which may potentially include a degree of dietary bias and often exhibit far more discontinuous/irregular deposition (Chase et al., 2012). Sections of each midden were cut perpendicular to the stratigraphy using an angle grinder and/or rotary impact hammer (Figure 4) and transported back to the laboratory for analysis. In addition to these newly collected middens, we analysed one of the Pakhuis Pass middens (PK1173) considered in the papers of Scott and Woodborne (2007a, b)

2.1 Chronology

Radiocarbon age determinations for the PK08 and PK10-1 middens (n=14) were processed at the ¹⁴CHRONO Centre, Queen's University Belfast using accelerator mass spectrometry (AMS) (Table 1; Figure 5). Samples were pre-treated with 2% HCl for one hour at room temperature to remove carbonates and dried at 60°C. They were then weighed into quartz tubes with an excess of CuO, sealed under vacuum and combusted to CO₂. The CO₂ was converted to graphite on an iron catalyst using the zinc reduction method (Slota et al., 1987). The radiocarbon ages were corrected for isotope fractionation using the AMS measured $\delta^{13}\text{C}$. These ages and those obtained from PK1173 (Scott and Woodborne, 2007a) were calibrated using the SHCal13 calibration data (Hogg et al., 2013). The Clam 2.2 software package (Blaauw, 2010) was used to generate all age-depth models (Figure 5). Clam was chosen over Bayesian techniques such as Bacon (Blaauw and Christen, 2011) because strong changes in accumulation rate may occur in hyrax middens (such as PK08 and PK10-1), and Clam – using linear models - is better suited to such sequences.

2.2 Stable nitrogen isotopes

Stable nitrogen isotope analysis of midden hyraceum samples were performed at the Department of Archaeology, University of Cape Town following Chase et al. (2010; 2009), with a contiguous/overlapping samples obtained two series of offset 1 mm holes. The standard deviation derived from replicate analyses of homogeneous material was better than 0.2 ‰. Results are expressed relative to atmospheric nitrogen.

3 Results

3.1 Chronology

Radiocarbon analyses indicate that the Pakhuis Pass hyrax middens accumulated during the late Pleistocene and Holocene, spanning the last ~19,300 years cal (calibrated) BP. The age-depth models for the two middens suggest continuous deposition, although accumulation rates do vary considerably (Table 1; Figure 5). Accumulation rates for the PK08 midden average ~28 $\mu\text{m yr}^{-1}$, with a period of more rapid accumulation centred on 6000 cal BP (~132 $\mu\text{m yr}^{-1}$). Each 1 mm isotope sample from PK08 therefore integrates between ~8 and 105 years of hyraceum accumulation (averaging 42 mm yr^{-1}). Accumulation rates for the PK10-1 midden increase with age/depth, from ~2.6 $\mu\text{m yr}^{-1}$ in the uppermost 30 mm to ~106 $\mu\text{m yr}^{-1}$ in the bottom 13 mm. Therein, each 1 mm isotope sample from PK10-1 integrates between ~380 years (upper section) to ~10 years (lower section) of hyraceum accumulation. Because of this significant change in accumulation rate, we show data from the extremely low resolution of the Holocene portion of PK10-1, but do not consider it our analyses, favouring? the much more highly resolved record from PK08. Accumulation of the PK1173 midden (Scott and Woodborne, 2007a) was more regular, averaging ~50 $\mu\text{m yr}^{-1}$, with each isotope sample integrating ~20 years of accumulation.

3.2 Stable nitrogen isotopes

The $\delta^{15}\text{N}$ values from the PK08, PK10-1 and PK1173 middens vary from 3.6 to 8.6‰ (Figure 6). Variations in midden $\delta^{15}\text{N}$ are interpreted to reflect changes in water availability (see more extensive discussion in Chase et al., 2012). At the global scale, a relationship has been recognised between aridity and foliar ^{15}N (e.g. Craine et al., 2009; Hartman and Danin, 2010),

and replication of this signal in plant and animal tissues and in faecal matter (e.g. Carr et al., 2016; Hartman, 2011; Murphy and Bowman, 2006; Newsome et al., 2011) has been demonstrated. This is thought to be generally a function of a more open nitrogen cycle in arid regions. Fractionating pathways in the soil (nitrification, denitrification, etc.) mean that nitrogen lost through transformation and the release of gaseous products is depleted in ^{15}N , and the remaining nitrogen is enriched. While in more humid regions N is cycled between live and dead organic pools, in drier regions more N flows to mineral pools where it is subject to gaseous loss (Amundson et al., 2003), and the $\delta^{15}\text{N}$ value of soils is thus higher with increasing aridity (Austin and Vitousek, 1998; Handley et al., 1999; Murphy and Bowman, 2009). The environmental processes relating to this recycling or loss of ^{15}N are not tied exclusively to rainfall amount, but in climatic terms are more accurately considered to relate to water availability (Murphy and Bowman, 2006). Studies of ^{15}N in hyrax middens from a wide range of environments indicate consistently strong correlations between midden ^{15}N , local vegetation/soil ^{15}N , as well as independent climate proxy records, supporting the conclusion that environmental moisture availability is a major driver of midden ^{15}N records (Carr et al., 2016; Chase et al., 2015a; Chase et al., 2013; Chase et al., 2017; Chase et al., 2015b; Chase et al., 2010; Chase et al., 2009; Chase et al., 2011).

This general relationship may also be influenced to some extent by other (e.g. microclimatic) factors, as is reflected in the variability observed in the relationship between modern foliar ^{15}N and aridity estimates (e.g. Hartman and Danin, 2010; Murphy and Bowman, 2006; Peri et al., 2012). This influence can be observed in the offset between the PK08 and PK1173 records and the record from PK10-1 (Figure 6). This is not uncommon when comparing

records from different middens, even when in relatively close proximity (<20 m) (Chase et al., 2013). These differences are believed to be driven by micro-topographic variations influencing water-availability in the primary feeding zones associated with each shelter (cf. Chase et al., 2013; Chase et al., 2017). In the case of the Pakhuis Pass middens, PK10-1 was retrieved from a narrow upper tier of the cliff complex, with the primary foraging range was likely to have been dominated by plants able to establish in the exposed landscape and thin soils found on the rock shelves above the cliff. PK08 and PK1173, in contrast, are located near the cliff base, and surrounded by deeper soils and higher groundwater recharge potential, which supports a more mesic, denser vegetation. The result is that while trends in $\delta^{15}\text{N}$ variability are similar between the PK08, PK1173 and PK10-1 middens, the PK10-1 values are approximately 0.95‰ higher (established by comparison of average values for the overlapping sections of midden sequences). For combination and consideration of the records, we have applied a 0.95‰ correction factor to the PK10-1 data.

To create a single composite record, we selected the PK08 record over the much lower resolution PK10-1 record for the mid- to late Holocene, as the latter is little informative in this context, with greater temporal averaging attenuating the variability evident in the higher resolution PK08 record. For the late Pleistocene, PK10-1 and PK1173 are of similar resolution and indicate similar trends. To incorporate information from both middens, we combined and smoothed the age-ranked data ensemble using Gaussian kernel-based interpolation (Rehfeld et al., 2011). This technique has been shown to be the most appropriate to interpolate irregularly sampled time series. We followed the recommendation of Rehfeld et al. (2011) and used the average temporal resolution of the data from these sections (31.97 years) to define the width

of our Gaussian kernel. To maintain the true temporal resolution of our composite record, the values were interpolated at the sample ages.

As a whole, the Pakhuis Pass $\delta^{15}\text{N}$ data indicate a general aridification at the site (change towards higher $\delta^{15}\text{N}$ values) from the late Pleistocene to the Holocene (Figure 6). The most humid periods recorded occur during the last glacial period, between 19,300 and 15,300 cal BP, and from 6100-5500 cal BP, following the early to mid-Holocene transition. Maximum aridity is reached during the early Holocene. The length and resolution of the records do not allow for definitive comparisons, but variability during the Pleistocene portion of the record – even considering its lower resolution - appears relatively muted compared to the strong, abrupt changes apparent in the Holocene.

Using change point analysis (Killick et al., 2012; Trauth et al., 2018), major transitions in the mean climate state (not accounting for changes in variability) in the record (where the root-mean-square level of the signal changes most significantly) were identified. Most notably, the period following the termination of early Holocene aridity at ~6800 cal BP is marked by a series of abrupt changes reflecting a rapid increase in water availability between ~6800 and 6140 cal BP. This period of relatively humid conditions ends equally abruptly, with distinctly drier conditions persisting from 5520 - 4480 cal BP. A subsequent sharp increase in water availability is followed by a gradual aridification between 4480 and 2830 cal BP, and apparently more stable conditions during the late Holocene. It is worth noting that the amplitude of change registered across these transitions (as much as 4.5‰) is similar to the amplitude of change between the Last Glacial Maximum and Holocene (5‰). Using the data derived from modern

hyrax faecal pellets presented by Carr et al. (2016), this may speculatively be translated to a ~70% increase in water availability (Aridity Index from 0.25 to 0.42) for the period between ~6800 and 6100 cal BP (Figure 6). Unfortunately, the midden records obtained do not provide detailed information regarding modern/sub-modern conditions, and a full contextualisation of these results in comparison with modern climates and recent change is not yet possible.

4 Discussion

4.1 Glacial-interglacial scale variability

Considering the combined influences of lower Pleistocene temperatures (Chevalier and Chase, 2016; Lim et al., 2016; Stute and Talma, 1998; Talma and Vogel, 1992), and the position of Pakhuis Pass in the winter rainfall zone, which is considered to have received increased precipitation during glacial periods (Chase et al., 2017; Chase and Meadows, 2007; Cockcroft et al., 1987; van Zinderen Bakker, 1976), it is not surprising that the late Pleistocene is characterised by more humid conditions (Figure 7). What is most remarkable about the Pakhuis Pass $\delta^{15}\text{N}$ record is that it indicates dramatically different patterns of change than those recorded at the De Rif hyrax midden site, located only 42 km to the south (Figure 2,3; Chase et al. 2011, 2015). This implies past climate change dynamics characterised by marked levels of spatio-temporal heterogeneity. Previously, inter-regional differences have been indicated by $\delta^{15}\text{N}$ records from Seweweekspoort (Figure 2; 230 km east-southeast of De Rif). In isolation, the differences between the De Rif and Seweweekspoort might otherwise have been explained by the distinct natures of eastern and western CFR climate histories (Chase and Meadows, 2007; Cowling et al., 1992). The new data from Pakhuis Pass imply a more complex dynamic.

Interpretations of the Seweweekspoort and De Rif records, which extend into the Last Glacial Maximum (LGM; 19-26.5 ka (Clark et al., 2009)) have highlighted the influence of different elements of the regional climate system as they were impacted by changing global boundary conditions. At glacial-interglacial timescales, the $\delta^{15}\text{N}$ record from Seweweekspoort (Chase et al., 2017) has been identified as primarily reflecting changes in the climate system related to global temperature and Antarctic sea-ice, which has been hypothesised as being a significant control on the position of the southern westerlies storm track (Chase and Meadows, 2007; Cockcroft et al., 1987; van Zinderen Bakker, 1976). In contrast, at De Rif, despite being closer to the core of the WRZ, where temperate systems define the modern regional rainfall regime, the primary control on hydrologic balance at millennial and multi-millennial timescales appears to be variations in the length and intensity of the summer drought season, potentially modulated by variations in the intensity of the South Atlantic Anticyclone, which blocks the incursion of tropical air masses and limits convection (Chase et al., 2015a). In the higher elevations of the Cederberg mountains, while winter rainfall is consistent and abundant, changes in summer rainfall may have had a more significant influence on driving drought-stress at the site. Including the new Pakhuis Pass $\delta^{15}\text{N}$ record in this regional consideration, it seems likely (based on similarities in first-order trends with SWP, as well as EPICA Dronning Maud Land ice core sea-salt sodium flux record, considered as a general proxy for sea-ice extent (Fischer et al., 2007)) that at glacial-interglacial timescales conditions at Pakhuis Pass is predominantly impacted by changes in temperate system influence, with summer rainfall playing a relatively limited role in defining water availability.

4.2 The Holocene

Focussing on the Holocene, more, and more precise data can be brought to bear on the question of CFR climate dynamics. While data coverage remains far from comprehensive, high resolution stable isotope data from rock hyrax middens have provided important clues as to some of the drivers and spatial dynamics that have defined climate change in the region during the Holocene (Figure 8). Building from the basis of the conceptual models developed by van Zinderen Bakker (1976) and Cockcroft et al. (1987), which indicate that periods of global cooling (warming) will generally result in wetter (drier) conditions in the WRZ (SRZ), and that a coeval inverse relationship in terms of precipitation amount exists between the two regions, more nuanced scenarios regarding observed variability are being proposed.

To constrain the relationship between CFR climate dynamics and the underlying drivers of the observed variability, we have considered the diatom-based Atlantic sector sea-ice extent and Southern Ocean summer sea-surface temperature (SSST) reconstructions of Nielsen et al. (2004) from marine core TN057-17. Similarities with these records can be observed at sites across the CFR, but it appears that the relationship between humidity and mid-latitude SSSTs changes from positive to negative moving from the eastern to western CFR. At the coastal eastern CFR site of Eilandvlei (Quick et al., 2018), for example, higher percentages of afrotemperate forest pollen (considered to indicate increased humidity and reduced drought stress) correlate well with higher SSSTs at TN057-17 (Figure 8). In contrast, higher humidity at Pakhuis and Katbakkies passes are more clearly linked to lower SSSTs. Climatically, this most likely indicates that increased westerly influence results in more humid conditions in the lee of

the Cederberg mountains, perhaps related to a greater frequency of more powerful frontal systems.

Considering the relationship between the Pakhuis and Katbakkies pass sites, the impact of changes in westerly influence is variable even between these two sites, despite their proximity and their shared position in the lee of the Cederberg. Katbakkies Pass $\delta^{15}\text{N}$ record exhibits changes in both the timing and amplitude that are similar to changes in TN057-17 SSTs, perhaps indicating that variations in humidity are more tightly coupled to temperate frontal systems than at Pakhuis Pass, where the timing and direction of the anomalies is shared, but the local response is less consistent (Figure 8). As has been indicated (Chase et al., 2017), increased frontal activity during the Holocene is not necessarily inconsistent with the increases in summer rain that have been previously suggested to drive moisture availability at Katbakkies Pass (Chase et al., 2015b), but it is clear that further work is required to adequately understand how temperate and tropical systems interact to create precipitation events in the region on the timescales considered here.

Underscoring the regional heterogeneity and the variable influence of the region's moisture bearing systems, the Holocene record from De Rif once again is markedly different from the Pakhuis and Katbakkies pass sites (Figure 8). Whereas the Pakhuis and Katbakkies Pass records exhibit a generally negative relationship between humidity and TN057-17 SSTs, at De Rif the relationship is generally positive, with more humid conditions occurring during periods of relatively elevated SSTs. In the early Holocene, while sea ice presence is low in the Atlantic sector of the Southern Ocean, conditions at De Rif are at their most humid for the last 19,400

years. With cooling SSTs and increased sea ice presence after 7000 cal BP, conditions become significantly drier. This extreme change during the early to mid-Holocene transition exemplifies the apparently sharply contrasting trends across the region, with the records from both Pakhuis and Katbakkies pass showing a strong increase in humidity evident from ~7000 – 6000 cal BP. While more muted, this coeval inverse relationship between the Pakhuis and Katbakkies Pass passes? sites and the De Rif persists throughout the Holocene, indicating that conditions at all of the sites are controlled by a shared dynamic, but that opposing factors determine conditions in the mountains (De Rif) and the rainshadow (Pakhuis and Katbakkies passes).

Of particular significance is how the opposing responses affect climatic gradients across the region. During the early Holocene, based on the $\delta^{15}\text{N}$ data, a very strong hydroclimatic gradient existed along the eastern slope of the Cederberg Mountains (Figure 9). Following the early to mid-Holocene transition, this gradient became much weaker, with more similar hydroclimatic conditions existing between the western CFR sites considered here. While only the records from De Rif and Pakhuis Pass extend to the Last Glacial Maximum, indications are that these gradients were weakened even further during the late Pleistocene, with similar $\delta^{15}\text{N}$ values at both sites suggesting nearly equivalent hydroclimatic conditions.

We hypothesise that these changes in environmental gradients may have had a significant influence on the diversity and distinction of environmental niches in the region, and that changes in these gradients may have acted as locks for gene flow, serving to alternatively isolate or enable the migration and mixing of plant species across the region. Considering the observation that species richness in the western CFR is higher in montane regions, the data

presented here provide palaeoclimatic evidence that highly variable spatial environmental gradients may have acted as a kind of allopatric speciation pump (e.g. Goldblatt and Manning, 2002; Haffer, 1969; Linder, 1985; Linder, 2003; Verboom et al., 2015), with the repeated isolation and mixing of promoting the exceptionally high species diversity of the CFR.

Further work will be required to establish where and to what extent variability on the scale discussed in this paper has been biologically significant. Has the degree or rate of change exceeded the tolerance of specific plant species or their ability to migrate? Can the spatial configuration of climate response anomalies be reliably linked to spatial patterns of species richness? Or, are the changes observed in the records discussed largely insignificant, and play no consequential role in determining species richness. Studies of fossil pollen records from the region provide a mixed response, with some indicating only minor changes in vegetation composition during the late Pleistocene and Holocene (Meadows et al., 2010; Meadows and Sugden, 1991), while other indicate substantial changes in vegetation have occurred (Scott and Woodborne, 2007a; Valsecchi et al., 2013). To adequately evaluate and compare these findings, more sites from transects across the region will need to be studied using a consistent methodology that employs a range of proxies capable of differentiating climate and vegetation change, and assessing spatial patterns of genetic diversity.

5 Conclusions

- We present a new, high resolution $\delta^{15}\text{N}$ record from Pakhuis Pass in the Cederberg Mountains of the western Cape Floristic Region.

- Like recently published records from the region (Chase et al., 2015a; Chase et al., 2015b), the data from Pakhuis Pass indicate substantial, rapid changes in hydroclimatic conditions, contrasting with earlier suggestions that the Cederberg experienced relatively little environmental change during since the LGM (Meadows et al., 2010; Meadows and Sugden, 1991).
- The record from Pakhuis Pass indicates similarities with patterns of change observed in comparable records from Katbakkies Pass ((Chase et al., 2015b), also in the rainshadow of the Cederberg) but contrasts sharply with conditions at De Rif (Chase et al., 2015a), where a coeval inverse pattern of variability is observed. Commonalities in the *timing* of change suggest a shared driver of regional climate dynamics, but distinct local responses, perhaps as a function of orographic influences.
- The opposing regional responses indicate significant changes in environmental gradients across the western CFR.
- While hydroclimatic conditions at Pakhuis Pass and De Rif may have been similar immediately after the LGM, subsequent responses to changes in global boundary conditions during deglaciation resulted in the establishment of a steep environmental gradient between the sites.
- We hypothesise that changes in the slope of this gradient likely reflect similar patterns across the region, and that these changes may have driven an allopatric speciation pump, contributing to the elevated species diversity observed in the montane regions of the western CFR.

- The spatial heterogeneity of hydroclimatic conditions suggested by the data included in this study indicates the need for more sites and comparable data from across the CFR in order to resolve the complexity of response to long-term climate change.

6 Acknowledgements

The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013), ERC Starting Grant “HYRAX”, grant agreement no. 258657. We are very grateful to Louis Scott for providing a portion of the PK1173 midden for analysis.

7 References

- Amundson, R., Austin, A.T., Schuur, E.A.G., Yoo, K., Matzek, V., Kendall, C., Uebersax, A., Brenner, D., Baisden, W.T., 2003. Global patterns of the isotopic composition of soil and plant nitrogen. *Global Biogeochemical Cycles* 17.
- Austin, A.T., Vitousek, P.M., 1998. Nutrient dynamics on a precipitation gradient in Hawai'i. *Oecologia* 113, 519-529.
- Blaauw, M., 2010. Methods and code for 'classical' age-modelling of radiocarbon sequences. *Quaternary Geochronology* 5, 512-518.
- Blaauw, M., Christen, J.A., 2011. Flexible paleoclimate age-depth models using an autoregressive gamma process. *Bayesian Analysis* 6, 457-474.
- Carr, A.S., Chase, B.M., Boom, A., Medina-Sanchez, J., 2016. Stable isotope analyses of rock hyrax faecal pellets, hyraceum and associated vegetation in southern Africa: Implications for dietary ecology and palaeoenvironmental reconstructions. *Journal of Arid Environments* 134, 33-48.
- Chase, B.M., Boom, A., Carr, A.S., Carré, M., Chevalier, M., Meadows, M.E., Pedro, J.B., Stager, J.C., Reimer, P.J., 2015a. Evolving southwest African response to abrupt deglacial North Atlantic climate change events. *Quaternary Science Reviews* 121, 132-136.
- Chase, B.M., Boom, A., Carr, A.S., Meadows, M.E., Reimer, P.J., 2013. Holocene climate change in southernmost South Africa: rock hyrax middens record shifts in the southern westerlies. *Quaternary Science Reviews* 82, 199-205.
- Chase, B.M., Chevalier, M., Boom, A., Carr, A.S., 2017. The dynamic relationship between temperate and tropical circulation systems across South Africa since the last glacial maximum. *Quaternary Science Reviews* 174, 54-62.

443 Chase, B.M., Lim, S., Chevalier, M., Boom, A., Carr, A.S., Meadows, M.E., Reimer, P.J., 2015b.
 444 Influence of tropical easterlies in southern Africa's winter rainfall zone during the Holocene.
 445 Quaternary Science Reviews 107, 138-148.

446 Chase, B.M., Meadows, M.E., 2007. Late Quaternary dynamics of southern Africa's winter
 447 rainfall zone. Earth-Science Reviews 84, 103-138.

448 Chase, B.M., Meadows, M.E., Carr, A.S., Reimer, P.J., 2010. Evidence for progressive Holocene
 449 aridification in southern Africa recorded in Namibian hyrax middens: implications for African
 450 Monsoon dynamics and the "African Humid Period". Quaternary Research 74, 36-45.

451 Chase, B.M., Meadows, M.E., Scott, L., Thomas, D.S.G., Marais, E., Sealy, J., Reimer, P.J.,
 452 2009. A record of rapid Holocene climate change preserved in hyrax middens from southwestern
 453 Africa. Geology 37, 703-706.

454 Chase, B.M., Quick, L.J., Meadows, M.E., Scott, L., Thomas, D.S.G., Reimer, P.J., 2011. Late
 455 glacial interhemispheric climate dynamics revealed in South African hyrax middens. Geology
 456 39, 19-22.

457 Chase, B.M., Scott, L., Meadows, M.E., Gil-Romera, G., Boom, A., Carr, A.S., Reimer, P.J.,
 458 Truc, L., Valsecchi, V., Quick, L.J., 2012. Rock hyrax middens: a palaeoenvironmental archive
 459 for southern African drylands. Quaternary Science Reviews 56, 107-125.

460 Chevalier, M., 2019. Enabling possibilities to quantify past climate from fossil assemblages at a
 461 global scale. Global and Planetary Change 175, 27-35.

462 Chevalier, M., Chase, B.M., 2016. Determining the drivers of long-term aridity variability: a
 463 southern African case study. Journal of Quaternary Science 31, 143-151.

464 Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B., Mitrovica, J.X.,
 465 Hostetler, S.W., McCabe, A.M., 2009. The Last Glacial Maximum. Science 325, 710-714.

466 Cockcroft, M.J., Wilkinson, M.J., Tyson, P.D., 1987. The application of a present-day climatic
 467 model to the late Quaternary in southern Africa. Climatic Change 10, 161-181.

468 Cockcroft, M.J., Wilkinson, M.J., Tyson, P.D., 1988. A palaeoclimatic model for the late
 469 Quaternary in southern Africa. Palaeoecology of Africa 19, 279-282.

470 Cowling, R.M., 1992. The Ecology of Fynbos: Nutrients, Fire, and Diversity. Oxford University
 471 Press, Cape Town.

472 Cowling, R.M., Cartwright, C.R., Parkington, J.E., Allsopp, J.C., 1999. Fossil wood charcoal
 473 assemblages from Elands Bay Cave, South Africa: implications for Late Quaternary vegetation
 474 and climates in the winter-rainfall fynbos biome. Journal of Biogeography 26, 367-378.

475 Cowling, R.M., Holmes, P.M., Rebelo, A.G., 1992. Plant diversity and endemism, in: Cowling,
 476 R.M. (Ed.), Ecology of fynbos. Nutrients, Fire and Diversity. Oxford University Press, Cape
 477 Town, pp. 62-112.

478 Cowling, R.M., Lombard, A.T., 2002. Heterogeneity, speciation/extinction history and climate:
 479 explaining regional plant diversity patterns in the Cape Floristic Region. Diversity and
 480 Distributions 8, 163-179.

481 Cowling, R.M., Potts, A.J., Bradshaw, P.L., Colville, J., Arianoutsou, M., Ferrier, S., Forest, F.,
 482 Fyllas, N.M., Hopper, S.D., Ojeda, F., Procheş, Ş., Smith, R.J., Rundel, P.W., Vassilakis, E.,
 483 Zutta, B.R., 2015. Variation in plant diversity in mediterranean-climate ecosystems: the role of
 484 climatic and topographical stability. Journal of Biogeography 42, 552-564.

485 Cowling, R.M., Richardson, D.M., Schulze, R.J., Hoffman, M.T., Midgley, J.J., Hilton-Taylor,
 486 C., 1997. Species diversity at the regional scale, in: Cowling, R.M., Richardson, D.M., Pierce,
 487 S.M. (Eds.), Vegetation of Southern Africa. Cambridge University Press, Cambridge, pp. 447-
 488 473.

489 Craine, J.M., Elmore, A.J., Aida, M.P.M., Bustamante, M., Dawson, T.E., Hobbie, E.A.,
 490 Kahmen, A., Mack, M.C., McLauchlan, K.K., Michelsen, A., Nardoto, G.B., Pardo, L.H.,
 491 Peñuelas, J., Reich, P.B., Schuur, E.A.G., Stock, W.D., Templer, P.H., Virginia, R.A., Welker,
 492 J.M., Wright, I.J., 2009. Global patterns of foliar nitrogen isotopes and their relationships with
 493 climate, mycorrhizal fungi, foliar nutrient concentrations, and nitrogen availability. *New*
 494 *Phytologist* 183, 980-992.
 495 Diekmann, B., Falker, M., Kuhn, G., 2003. Environmental history of the south-eastern South
 496 Atlantic since the Middle Miocene: evidence from the sedimentological records of ODP Sites
 497 1088 and 1092. *Sedimentology* 50, 511-529.
 498 Dynesius, M., Jansson, R., 2000. Evolutionary consequences of changes in species' geographical
 499 distributions driven by Milankovitch climate oscillations. *Proceedings of the National Academy*
 500 *of Sciences* 97, 9115-9120.
 501 Fischer, H., Fundel, F., Ruth, U., Twarloh, B., Wegner, A., Udisti, R., Becagli, S., Castellano, E.,
 502 Morganti, A., Severi, M., Wolff, E., Littot, G., Röthlisberger, R., Mulvaney, R., Hutterli, M.A.,
 503 Kaufmann, P., Federer, U., Lambert, F., Bigler, M., Hansson, M., Jonsell, U., de Angelis, M.,
 504 Boutron, C., Siggaard-Andersen, M.-L., Steffensen, J.P., Barbante, C., Gaspari, V., Gabrielli, P.,
 505 Wagenbach, D., 2007. Reconstruction of millennial changes in dust emission, transport and
 506 regional sea ice coverage using the deep EPICA ice cores from the Atlantic and Indian Ocean
 507 sector of Antarctica. *Earth and Planetary Science Letters* 260, 340-354.
 508 Forest, F., Grenyer, R., Rouget, M., Davies, T.J., Cowling, R.M., Faith, D.P., Balmford, A.,
 509 Manning, J.C., Procheş, Ş., van der Bank, M., Reeves, G., Hedderson, T.A.J., Savolainen, V.,
 510 2007. Preserving the evolutionary potential of floras in biodiversity hotspots. *Nature* 445, 757.
 511 GBIF.org, 2018a. Anthocerotopsida occurrence data downloaded on September 13th. GBIF.org.
 512 GBIF.org, 2018b. Cycadopsidae occurrence data downloaded on March 30th. GBIF.org.
 513 GBIF.org, 2018c. Ginkgoopsidae occurrence data downloaded on March 30th. GBIF.org.
 514 GBIF.org, 2018d. Gnetopsidae occurrence data downloaded on March 30th. GBIF.org.
 515 GBIF.org, 2018e. Liliopsida occurrence data downloaded on September 13th. GBIF.org.
 516 GBIF.org, 2018f. Lycopodiopsida occurrence data downloaded on September 13th. GBIF.org.
 517 GBIF.org, 2018g. Magnoliopsida occurrence data downloaded on March 30th. GBIF.org.
 518 GBIF.org, 2018h. Pinopsidae occurrence data downloaded on March 30th. GBIF.org.
 519 GBIF.org, 2018i. Polypodiopsida occurrence data downloaded on September 13th. GBIF.org.
 520 Goldblatt, P., 1978. An analysis of the flora of southern Africa: its characteristics, relationships,
 521 and origins. *Annals of the Missouri Botanical Garden* 65, 369-436.
 522 Goldblatt, P., Manning, J.C., 2002. Plant diversity of the Cape Region of southern Africa. *Annals*
 523 *of the Missouri Botanical Garden* 89, 281-302.
 524 Haffer, J., 1969. Speciation in Amazonian Forest Birds. *Science* 165, 131-137.
 525 Handley, L.L., Austin, A.T., Stewart, G.R., Robinson, D., Scrimgeour, C.M., Raven, J.A.,
 526 Heaton, T.H.E., Schmidt, S., 1999. The ^{15}N natural abundance ($\delta^{15}\text{N}$) of ecosystem samples
 527 reflects measures of water availability. *Functional Plant Biology* 26, 185-199.
 528 Hartman, G., 2011. Are elevated $\delta^{15}\text{N}$ values in herbivores in hot and arid environments caused
 529 by diet or animal physiology? *Functional Ecology* 25, 122-131.
 530 Hartman, G., Danin, A., 2010. Isotopic values of plants in relation to water availability in the
 531 Eastern Mediterranean region. *Oecologia* 162, 837-852.
 532 Hijmans, R., Cameron, S.E., Parra, J.L., Jones, P.G., Jarvis, A., 2005. Very high resolution
 533 interpolated climate surfaces for global land areas. *International Journal of Climatology* 25,
 534 1965-1978.

535 Hogg, A.G., Hua, Q., Blackwell, P.G., Niu, M., Buck, C.E., Guilderson, T.P., Heaton, T.J.,
 536 Palmer, J.G., Reimer, P.J., Reimer, R.W., Turney, C.S.M., Zimmerman, S.R.H., 2013. SHCal13
 537 Southern Hemisphere Calibration, 0–50,000 Years cal BP. *Radiocarbon* 55, 1889-1903.
 538 Killick, R., Fearnhead, P., Eckley, I.A., 2012. Optimal Detection of Changepoints With a Linear
 539 Computational Cost. *Journal of the American Statistical Association* 107, 1590-1598.
 540 Lim, S., Chase, B.M., Chevalier, M., Reimer, P.J., 2016. 50,000 years of vegetation and climate
 541 change in the southern Namib Desert, Pella, South Africa. *Palaeogeography, Palaeoclimatology,*
 542 *Palaeoecology* 451, 197-209.
 543 Linder, H.P., 1985. Gene flow, speciation, and species diversity patterns in a species-rich area:
 544 the Cape Flora. *Species and speciation* 4, 53-57.
 545 Linder, H.P., 1991. Environmental correlates of patterns of species richness in the south-western
 546 Cape Province of South Africa. *Journal of Biogeography* 18, 509-518.
 547 Linder, H.P., 2003. The radiation of the Cape flora, southern Africa. *Biological Reviews of the*
 548 *Cambridge Philosophical Society* 78, 597-638.
 549 Linder, H.P., 2005. Evolution of diversity: the Cape flora. *Trends in Plant Science* 10, 536-541.
 550 Meadows, M.E., Chase, B.M., Selanie, M., 2010. Holocene palaeoenvironments of the
 551 Cederberg and Swartruggens mountains, Western Cape, South Africa: Pollen and stable isotope
 552 evidence from hyrax dung middens. *Journal of Arid Environments* 74, 786-793.
 553 Meadows, M.E., Sugden, J.M., 1991. A vegetation history of the last 14,000 years on the
 554 Cederberg, southwestern Cape Province. *South African Journal of Science* 87, 34-43.
 555 Mucina, L., Rutherford, M.C., 2006. The vegetation of South Africa, Lesotho and Swaziland,
 556 *Strelitzia*. South African National Biodiversity Institute, Pretoria.
 557 Murphy, B.P., Bowman, D.M.J.S., 2006. Kangaroo metabolism does not cause the relationship
 558 between bone collagen $\delta^{15}\text{N}$ and water availability. *Functional Ecology* 20, 1062-1069.
 559 Murphy, B.P., Bowman, D.M.J.S., 2009. The carbon and nitrogen isotope composition of
 560 Australian grasses in relation to climate. *Functional Ecology* 23, 1040-1049.
 561 Newsome, S.D., Miller, G.H., Magee, J.W., Fogel, M.L., 2011. Quaternary record of aridity and
 562 mean annual precipitation based on $\delta^{15}\text{N}$ in ratite and dromornithid eggshells from Lake Eyre,
 563 Australia. *Oecologia* 167, 1151-1162.
 564 Nielsen, S.H.H., Koç, N., Crosta, X., 2004. Holocene climate in the Atlantic sector of the
 565 Southern Ocean: Controlled by insolation or oceanic circulation? *Geology* 32, 317-320.
 566 Peri, P.L., Ladd, B., Pepper, D.A., Bonser, S.P., Laffan, S.W., Amelung, W., 2012. Carbon
 567 ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) stable isotope composition in plant and soil in Southern Patagonia's
 568 native forests. *Global Change Biology* 18, 311-321.
 569 Quick, L.J., Chase, B.M., Meadows, M.E., Scott, L., Reimer, P.J., 2011. A 19.5 kyr vegetation
 570 history from the central Cederberg Mountains, South Africa: Palynological evidence from rock
 571 hyrax middens. *Palaeogeography, Palaeoclimatology, Palaeoecology* 309, 253-270.
 572 Quick, L.J., Chase, B.M., Wündsche, M., Kirsten, K.L., Chevalier, M., Mäusbacher, R.,
 573 Meadows, M.E., Haberzettl, T., 2018. A high-resolution record of Holocene climate and
 574 vegetation dynamics from the southern Cape coast of South Africa: pollen and microcharcoal
 575 evidence from Eilandvlei. *Journal of Quaternary Science* 33, 487-500.
 576 Rehfeld, K., Marwan, N., Heitzig, J., Kurths, J., 2011. Comparison of correlation analysis
 577 techniques for irregularly sampled time series. *Nonlin. Processes Geophys.* 18, 389-404.
 578 Roberts, D.L., Neumann, F.H., Cawthra, H.C., Carr, A.S., Scott, L., Durugbo, E.U., Humphries,
 579 M.S., Cowling, R.M., Bamford, M.K., Musekiwa, C., MacHutchon, M., 2017.

Palaeoenvironments during a terminal Oligocene or early Miocene transgression in a fluvial system at the southwestern tip of Africa. *Global and Planetary Change* 150, 1-23.

Scott, L., Woodborne, S., 2007a. Pollen analysis and dating of Late Quaternary faecal deposits (hyraceum) in the Cederberg, Western Cape, South Africa. *Review of Palaeobotany and Palynology* 144, 123-134.

Scott, L., Woodborne, S., 2007b. Vegetation history inferred from pollen in Late Quaternary faecal deposits (hyraceum) in the Cape winter-rain region and its bearing on past climates in South Africa. *Quaternary Science Reviews* 26, 941-953.

Slota, P.J., Jull, A.J.T., Linick, T.W., Toolin, L.J., 1987. Preparation of small samples for ^{14}C accelerator targets by catalytic reduction of CO. *Radiocarbon* 29, 303-306.

Stute, M., Talma, A.S., 1998. Glacial temperatures and moisture transport regimes reconstructed from noble gas and $\delta^{18}\text{O}$, Stampriet aquifer, Namibia, *Isotope Techniques in the Study of Past and Current Environmental Changes in the Hydrosphere and the Atmosphere*. IAEA Vienna Symposium 1997, Vienna, Austria, pp. 307-318.

Talma, A.S., Vogel, J.C., 1992. Late Quaternary paleotemperatures derived from a speleothem from Cango Caves, Cape Province, South Africa. *Quaternary Research* 37, 203-213.

Trabucco, A., Zomer, R.J., 2009. Global Aridity Index (Global-Aridity) and Global Potential Evapo-Transpiration (Global-PET) Geospatial Database. , in: Information, C.C.f.S. (Ed.). CGIAR-CSI GeoPortal, <http://www.csi.cgiar.org>.

Trauth, M.H., Foerster, V., Junginger, A., Asrat, A., Lamb, H.F., Schaebitz, F., 2018. Abrupt or gradual? Change point analysis of the late Pleistocene–Holocene climate record from Chew Bahir, southern Ethiopia. *Quaternary Research*, 1-10.

Tyson, P.D., 1986. *Climatic Change and Variability in Southern Africa*. Oxford University Press, Cape Town.

Tyson, P.D., Preston-Whyte, R.A., 2000. *The Weather and Climate of Southern Africa*. Oxford University Press, Cape Town.

UNEP, 1997. *World atlas of desertification*, 2nd ed. United Nations Environment Programme, London.

Valsecchi, V., Chase, B.M., Slingsby, J.A., Carr, A.S., Quick, L.J., Meadows, M.E., Cheddadi, R., Reimer, P.J., 2013. A high resolution 15,600-year pollen and microcharcoal record from the Cederberg Mountains, South Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* 387, 6-16.

van Zinderen Bakker, E.M., 1976. The evolution of late Quaternary paleoclimates of Southern Africa. *Palaeoecology of Africa* 9, 160-202.

Verboom, G.A., Linder, H.P., Forest, F., Hoffman, V., Bergh, N.G., Cowling, R.M., 2014. Cenozoic assembly of the greater Cape flora, in: Allsopp, N., Colville, J.F., Verboom, G.A. (Eds.), *Fynbos: Ecology, Evolution and Conservation of a Megadiverse Region*. Oxford University Press, Oxford, pp. 93-118.

Verboom, G.A., Nicola, G.B., Sarah, A.H., Vera, H., Matthew, N.B., 2015. Topography as a driver of diversification in the Cape Floristic Region of South Africa. *The New Phytologist* 207, 368-376.

Yamazaki, D., Ikeshima, D., Tawatari, R., Yamaguchi, T., O'Loughlin, F., Neal, J.C., Sampson, C.C., Kanae, S., Bates, P.D., 2017. A high-accuracy map of global terrain elevations. *Geophysical Research Letters* 44, 5844-5853.

Table captions

Table 1: Radiocarbon ages and calibration information for the PK08, PK10-1 and PK1173 (Scott and Woodborne, 2007a, b) rock hyrax middens.

Figure captions

Figure 1: Map of southern Africa showing seasonality of rainfall and climatic gradients dictated by the zones of summer/tropical (red) and winter/temperate (blue) rainfall dominance. Major atmospheric (white arrows) and oceanic (blue arrows) circulation systems are indicated.

Figure 2: Map of southernmost Africa showing **A)** showing seasonality of rainfall and climatic gradients defined in terms of percentage of winter rainfall, **B)** the extent of the vegetation types comprising the Cape Floristic Region (fynbos, renosterveld and Albany Thicket (data from Mucina and Rutherford, 2006)), **C)** a map of mean Aridity Index values for the region (Trabucco and Zomer, 2009), and **D)** a map of species richness using data derived from the GBIF database (Chevalier, 2019; 2018; GBIF.org, 2018, b, c, d, e, f, g, h, i); number of species per quarter-degree grid cell were interpolated to create the contour map. Classification of aridity values follows that of the United Nations Environment Programme (1997). Sites considered in this study are indicated: Pakhuis Pass (PK: this study; Scott and Woodborne, 2007a; Scott and Woodborne, 2007b), De Rif (DR: Chase et al., 2015a; Chase et al., 2011; Quick et al., 2011), Katbakkies Pass (KB: Chase et al., 2015b; Meadows et al., 2010), Seweweekspoort (SWP: Chase et al., 2015a; Chase et al., 2013; Chase et al., 2017) and Eilandvlei (Quick et al., 2018).

Figure 3: Map of the N-S axis of the Cape Fold Mountains in the southwestern Cape, including the Cederberg Mountains and adjacent ranges. Topographic variability (MERIT digital elevation model; Yamazaki et al., 2017) is overlain by a map of mean Aridity Index values (Trabucco and Zomer, 2009) for the region, as in Figure 1. The dominant vector of temperate moisture-bearing systems related to the westerly storm track is shown, as are the sites considered in this study: Pakhuis Pass (this study; Scott and Woodborne, 2007a; Scott and Woodborne, 2007b), De Rif (Chase et al., 2015a; Chase et al., 2011; Quick et al., 2011), Katbakkies Pass (Chase et al., 2015b; Meadows et al., 2010). The relative aridity of each site is shown on the legend. Inset indicates position of the region within the southwestern Cape.

Figure 4: Pakhuis Pass rock hyrax middens sampled for this study: **a)** PK08, 7 cm tool atop midden for scale; **b)** collecting a portion of the PK08 midden; **c)** PK10-1 midden, 16 cm GPS unit for scale.

Figure 5: Age-depth models for the PK08, PK10-1 and PK1173 rock hyrax middens.

Figure 6: Pane (a): The $\delta^{15}\text{N}$ data from the PK08 (red), PK10-1 (blue, sample location provided for lower resolution Holocene portion) and PK1173 (green) rock hyrax middens. Pane (b): The same records following corrections made to establish a single composite record for the site. The PK10-1 midden is associated with a more exposed foraging range and thinner soils, and a 0.95‰ correction factor was applied based on average values from PK08 and PK1173 (recovered from more mesic locations) for overlapping time periods. For the >15,500 cal BP portion of the sequence the similarly resolved, but irregularly sampled overlapping portions of PK10-1 and PK1173 (dashed lines) have been combined (shown in black) using Gaussian kernel-

based interpolation (Rehfeld et al., 2011). The timing of major changes as identified from the composite record using change point analysis (Killick et al., 2012) are indicated by vertical dashed lines.

Figure 7: The $\delta^{15}\text{N}$ data from the Pakhuis Pass (this paper), De Rif (Chase et al., 2011) and Seweweekspoort (Chase et al., 2017) rock hyrax middens, and the sea-salt sodium flux data from the EPICA Dronning Maud Land ice core from Antarctica, a proxy for sea-ice extent (Fischer et al., 2007).

Figure 8: Holocene $\delta^{15}\text{N}$ records from the Pakhuis Pass (this paper), Katbakkies Pass (Chase et al., 2015b), De Rif (Chase et al., 2015a) and Seweweekspoort (Chase et al., 2017) rock hyrax middens, as well as the afrotemperate forest pollen record from Eilandvlei (Quick et al., 2018), and proxies records relating to the position of the southern westerlies, including reconstructions of Southern Ocean summer sea-surface temperatures and sea ice presence (Nielsen et al., 2004) and the sea-salt sodium flux data from the EPICA Dronning Maud Land ice core from Antarctica, a proxy for sea-ice extent (Fischer et al., 2007; Roberts et al., 2017).

Figure 9: Holocene western Cape Floristic Region $\delta^{15}\text{N}$ records from the Pakhuis Pass (this paper), Katbakkies Pass (Chase et al., 2015b) and De Rif (Chase et al., 2015a) rock hyrax middens, resampled to a common 100-year resolution (linear interpolation). Three time slices have been selected across the mid-Holocene (4600, 5600 and 7000 cal BP), and the $\delta^{15}\text{N}$ values from the sites, as proxies for water availability, have been used to interpolate the steepness of hydroclimatic ($\delta^{15}\text{N}$) gradients between the sites. White isolines are described at 0.2‰ $\delta^{15}\text{N}$ intervals. *Note, as only three sites were used in this calculation, extrapolated gradients not*

686 *along the vector described by these three sites should not be considered as reliable*
687 *reconstructions.*