



# Modification of cell walls of apple, red beet and kiwifruit by heating in acid conditions: common and specific responses

Xuwei Liu, Catherine M.G.C. Renard, Agnès Rolland-Sabaté, Sylvie Bureau, Carine Le Bourvellec

## ► To cite this version:

Xuwei Liu, Catherine M.G.C. Renard, Agnès Rolland-Sabaté, Sylvie Bureau, Carine Le Bourvellec. Modification of cell walls of apple, red beet and kiwifruit by heating in acid conditions: common and specific responses. 15th Cell wall meeting, Jul 2019, Cambridge, United Kingdom. hal-02264926

**HAL Id: hal-02264926**

**<https://hal.science/hal-02264926>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Modification of cell walls of apple, red beet and kiwifruit by heating in acid conditions: common and specific responses.

Xuwei Liu<sup>1</sup>, Catherine M.G.C. Renard<sup>2</sup>, Agnès Rolland-Sabaté<sup>1</sup>, Sylvie Bureau<sup>1</sup>, Carine Le Bourvellec<sup>1\*</sup>

1 : UMR408 SQPOV « Sécurité et Qualité des Produits d'Origine Végétale », INRA, Avignon University, F-84000 Avignon, France

2 : CEPIA Division, INRA, F-44000 Nantes, France

## Abstract

Most fruit- and vegetable-based products are commonly processed before being consumed. It is known that cooking and processing induce significant changes in chemical composition and physical properties of cell wall polysaccharides. In this study, cell walls were isolated from three species: apple, red beet and kiwifruit using the alcohol-insoluble solids (AIS) procedure. The composition of the cell wall polysaccharides were compared before and after modification by boiling (100 °C, 20 minutes) at different pH values (2.0, 3.5 and 6.0) to explore the impact of acid hydrolysis and  $\beta$ -elimination. Native apple cell walls were mainly composed of cellulose, highly methylated pectins relatively rich in arabinan and galactan side chains and fucogalactoxyloglucan; red beet cell walls were rich in cellulose, highly methylated pectins relatively rich in arabinan and ferulic acid, kiwi fruit cell walls were mostly composed of cellulose, highly methylated pectins with low side chain contents and xyloglucans. After modification, the galacturonic acid contents of whole cell walls decreased the most at pH 6.0 indicating prevalence of  $\beta$ -elimination. Galacturonic acid contents decreased for all the cell walls whereas the degree of methylation (DM) decreased mostly in apple and red beet cell walls (from 82% to 60% and from 65% to 42%, respectively). At pH 2.0, acid hydrolysis decreased the content of neutral sugar side chains, especially arabinan from apple and red beet cell walls. Due to the much lower proportion of arabinose in kiwifruit cell walls, no obvious changes were recorded for kiwifruits at pH 2. Heating at pH 3.5 was as expected the least disruptive condition. Infrared spectra differentiated the samples by species and treatment as the composition done, the work is under progress in order to identified related peaks to the principal cell wall components.

Keywords: Polysaccharide, FT-IR, Processing, Side chain, Pectin, Methylation