



**HAL**  
open science

## Trial for the long neutron counter TETRA using $^{96,97}\text{Rb}$ radioactive sources

D. Testov, D. Verney, Y. Penionzhkevich, A. Boso, C. Delafosse, F. Didierjean, I. Deloncle, F. Ibrahim, F. Le Blanc, C. Gaulard, et al.

### ► To cite this version:

D. Testov, D. Verney, Y. Penionzhkevich, A. Boso, C. Delafosse, et al.. Trial for the long neutron counter TETRA using  $^{96,97}\text{Rb}$  radioactive sources. *Journal of Instrumentation*, 2019, 14 (8), pp.P08002. 10.1088/1748-0221/14/08/P08002 . hal-02266320

**HAL Id: hal-02266320**

**<https://hal.science/hal-02266320v1>**

Submitted on 27 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Test of long $^3\text{He}$ neutron counter TETRA using $^{96,97}\text{Rb}$ radioactive sources.

---

**D. Testov<sup>a,b,1</sup> D. Verney<sup>c</sup> Yu. Penionzhkevich<sup>a,d</sup> A. Boso<sup>e</sup> C. Delafosse<sup>f</sup> F. Didierjean<sup>g</sup>  
I. Deloncle<sup>h</sup> F. Ibrahim<sup>c</sup> F. Le Blanc<sup>c</sup> C. Gaulard<sup>h</sup> J. Guillot<sup>c</sup> S. Lukyanov<sup>a</sup> V. Maslov<sup>a</sup>  
I. Matea<sup>c</sup> B. Roussière<sup>c</sup> A. Severyukhin<sup>a</sup> Yu. Sobolev<sup>a</sup> E. Sokol<sup>a</sup> V. Smirnov<sup>a</sup> R. Thoër<sup>c</sup>  
D. Thisse<sup>c</sup> G. Tocabens<sup>c</sup> and ALTO people**

<sup>a</sup>Joint Institute for Nuclear Research, Dubna, Moscow region, Russia

<sup>b</sup>Dipartimento di Fisica e Astronomia and INFN, Sezione di Padova, Padova, Italy.

<sup>c</sup>Institut de Physique Nucléaire, IN2P3/CNRS and University Paris Sud, Orsay, France

<sup>d</sup>National Research Nuclear University, Kashirskoye Shosse 31, Moscow 115409, Russia

<sup>e</sup>National Physical Laboratory, Teddington, Middlesex, UK

<sup>f</sup>University of Jyväskylä, Jyväskylä, Finland

<sup>g</sup>Université de Strasbourg, CNRS, Strasbourg, France

<sup>h</sup>Centre de Sciences Nucléaires et de Sciences de la Matière, IN2P3/CNRS, Orsay, France

E-mail: [dmitry.testov@lnl.infn.it](mailto:dmitry.testov@lnl.infn.it)

**ABSTRACT:** TETRA long neutron counter is operated at ALTO ISOL facility behind mass separator PARNNe. TETRA has been proven to be a unique instrument to measure  $\beta$ -decay properties of short-lived neutron-rich nuclei in the interest of nuclear structure and astrophysical r-process calculations. A proper calibration of TETRA and validation of experimental procedure of measurements of a probability of  $\beta$ -delayed neutron emission ( $P_{1n}$ ) is only possible using a well-known  $\beta$ -neutron decaying radioactive source which can be only produced and measured on-line due to a short half-life. Thus, the present papers reports on measurements of  $P_{1n}$  and  $T_{\frac{1}{2}}$  properties of  $^{96,97}\text{Rb}$  nuclei using TETRA to be in a good agreement with the literature values. This proves that the developed techniques can be extended to disclose unknown  $P_{1n}$  and  $T_{\frac{1}{2}}$  of neutron-rich species.

**KEYWORDS:** Neutron detectors, Gaseous detectors

---

<sup>1</sup>Corresponding author.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Experiment Details</b>	<b>1</b>
<b>3</b>	<b>Conclusions</b>	<b>7</b>

---

## 1 Introduction

Nowadays the interest to the phenomenon of  $\beta$ -delayed (multi-neutron) neutron emission is arising. The mechanism which drives this process in neutron-rich nuclei is still not understood. Whereas,  $\beta$ -delayed neutrons play a significant role in the astrophysical calculations and the nuclear reactor physics. Therefore, many nuclei have been revisited for detailed examination of this phenomenon. Modern detection systems such as TETRA [1, 2] or BEDO [3] installed at ALTO ISOL facility [4] allow to perform measurements of  $\beta$ -decay properties of nuclei far away from the line of nuclei  $\beta$ -stability.

Beta-decay decay ground state properties have been the aim for many experiments performed at TETRA. The recent results obtained using TETRA/BEDO lead to observation of unusual population resonance structures in  $\beta$ -decay of  $^{83}\text{Ga}$  [5] and high energy  $\gamma$ -ray emission [6] which can be interpreted as a competition between  $\gamma$ -ray and neutron emission beyond the neutron separation energy threshold. Thus, measurements of half-life ( $T_{1/2}$ ) and probability of  $\beta$ -delayed (multi) neutron emission ( $P_{xn}$ ) are very sensitive to the parameters of the experimental setup. Performance of this detector can be characterized using standard neutron calibration (spontaneous-fission) sources, for an example  $^{252}\text{Cf}$ . The limitation of this source is the different energy ranger of prompt fission neutrons emitted by the source and  $\beta$ -delayed neutrons. Nevertheless, response of TETRA to prompt neutrons from this kind of a source is well understood by extensive simulations validated by measurements [7]. However,  $^{252}\text{Cf}$  source, obviously, does not allow verification of analysis procedure to proceed data from a  $\beta$ -decay experiment. Therefore, test the setup and the analysis approaches we performed two experiments using short-lived radioactive sources created at the centre of TETRA:  $^{96}\text{Rb}$  in the first and  $^{97}\text{Rb}$  runs correspondingly. The ground state properties of these nuclei are know. We report  $T_{\frac{1}{2}}$  and  $P_{1n}$  measured in the present experiment using previously adopted methods [1, 5] to be in a fair agreement with the literature values.

## 2 Experiment Details

The  $\text{UC}_x$  target placed in a Ta oven heated up to  $>2000^\circ\text{C}$  was exposed to the primary 50 MeV electron beam delivered bt the ALTO linear accelerator [4]. The average beam current on the tareget was  $10\ \mu\text{A}$ . For this experiment, the oven was connected to a W tube used to surface ionized fission products. The beam was then accelated at 30 keV. TETRA was operated behind the mass separator

PARNNe whose resolution ( $\delta m/m=1300$ ) was high enough to provide isobaric selection  $A = 96, 97$ . The other members of the isobaric chain, (Sr, Y) have higher (than Rb) ionization potential and thus their production rate can be neglected in the experiment. This fact makes  $^{96,97}\text{Rb}$  good references nuclei to verify experimental methods.


The  $\beta$ -decay station installed at the ALTO experimental hall was used in the neutron detection mode [3, 7]. The detailed description of the setup is provided in Ref. [7]: the surface-ionized mass-separated beams were collected on a Al-coated mylar tape at the centre of the long-neutron counter TETRA creating a radioactive source. The collection point was surrounded by a plastic scintillator for  $\beta$  detection. A High Purity Ge detector was used for  $\gamma$  detection. One cycle of data taking consisted of a short background measurement ( $T_{bg}$ ) followed by irradiation ( $T_{beam}$ ) when the beam was impinging on the tape; and the decay time ( $T_{decay}$ ) when the beam was deviated allowed decay of the accumulated isotopes. Therefore, the data acquisition system was recording neutron,  $\beta$  and  $\gamma$  activity from the source during  $T_m=T_{bg}+T_{beam}+T_{decay}$ . Then the tape was moved for 2 m to transport the source outside the detection system. The time settings and number of repeated cycles for  $A = 96, 97$  is listed in Table 1.

**Table 1:** Tape-cycle parameters used for the  $A = 96, 97$  settings:  $T_{bg}$  is the background counting time before beam collection,  $T_{coll}$  the duration of the beam collection and  $T_{dec}$  the beam-off source decay counting time, in ms.  $N_{cycles}$  is the total number of tape cycles for each mass setting. Summary of the results obtained from the analysis of the cumulated  $\beta$ - and neutron-activity curves for the  $A = 96, 97$  settings.

Beam	$T_{bg}$ ms	$T_{coll}$ ms	$T_{dec}$ ms	$N_{cycles}$	$T_{\frac{1}{2}}$ ms	$P_{1n}$ %	$\Phi$ pps	$\epsilon_{eff}^{\beta}$	$\epsilon_{eff}^n$
$^{96}\text{Rb}$	500	100	1900	191	200(1)	12.0(1.1)	$\sim 3.4 \cdot 10^4$	29(4)	58(4)
$^{97}\text{Rb}$	500	500	1500	500	169(1)	27.8(2.4)	$\sim 0.9 \cdot 10^4$	29(4)	56(4)

The resulting  $\beta$ - and neutron activity curves for  $A=96, 97$  mass separator settings accumulated over  $N_{cycles}$ , see Table 1, are plotted in Fig. 1. All neutrons detected for a selected mass of rubidium were attributed either to the background or to  $\beta$ -n decay of rubidium isotopes. Even if,  $\beta$ -delayed neutron emission is energetically allowed, there is no experimental evidence of  $\beta$ -delayed neutron emission of  $^{96,97}\text{Sr}$  so far. The simultaneous fit of the grow-in and decay patterns of accumulated neutron curves leads to half-lives of  $^{96,97}\text{Rb}$  as reported in Table 1, where the uncertainty is the uncertainty from the fit.

The effective efficiency of the  $\beta$  detector ( $\epsilon_{\beta}$ ) and the TETRA array ( $\epsilon_n$ ) was derived from coincidence  $\gamma$ -ray spectra recorded for  $A=96$  and  $A=97$  mass separator settings plotted in Fig. 2. The efficiencies were obtained from the observed ratios of an area of  $i^{th}$  peak in singles ( $S_{\gamma}^i$ ),  $\beta$  gated ( $S_{\gamma\beta}^i$ ), and  $\beta$ -n gated ( $S_{\gamma\beta n}^i$ )  $\gamma$ -spectra. The relative intensities of the observed transitions are summarized in Tables 2 and 3. The  $\epsilon_{\beta}$  was derived as a weighted average of  $S_{\gamma\beta}^i/S_{\gamma}^i$  ratios measured for the  $i$ -th transition. Due to lack of statistic in the  $\beta$ -n  $\gamma$ -gated spectrum  $\epsilon_n$  was found using only the most strong transitions at 352 keV and 815 keV in  $^{95}\text{Sr}$  and  $^{96}\text{Sr}$  respectively as is reported in Table 1. All the  $\gamma$  activities recorded could be identified and no isobaric contaminants were observed within our detection limits. Moreover, no evidence for contamination from surface ionized




**Figure 1:** Left:  $\beta$  (top) and neutron (bottom) activity curves recorded for the  $A = 96$  setting of the mass separator. These curves were accumulated over 191 tape cycles; the different  $\beta$ -activity components are singled out with coloured curves for  $A = 96$ :  $^{96}\text{Rb}$  in green,  $^{96}\text{Sr}$  in orange,  $^{96}\text{Y}$  in purple and the background level is represented by the dotted line; one component of the  $\beta$ -delayed neutron activity curve originating from  $^{96}\text{Rb}$  (green) decays is singled out. Right: the same for the  $A = 97$  setting of the mass separator:  $\beta$ -activity components are singled out with coloured curves for  $A = 97$ :  $^{97}\text{Rb}$  in green,  $^{97}\text{Sr}$  in orange,  $^{97}\text{Y}$  in purple; the  $\beta$ -delayed neutron activity curve originating from  $^{97}\text{Rb}$  (green) decays is singled out.

$^{96,97}\text{Sr}$  isotopes was found in the analysis of the activity curves. Indeed, such a contamination is unlikely due to the higher ionization potential of Sr.

To extract  $P_{1n}$  values and the production rates  $\phi$  we used the method reported previously [7]. The method is based on the system of Bateman equations describing decay of a given radioactive source. In this method the  $P_{1n}$  and  $\phi$  values are determined as roots of a corresponding system of Bateman equations 2.1.

$$\left\{ \begin{array}{l} (N_{\beta}^{exp} - N_{\beta}^{bg}) \frac{1}{\epsilon_{\beta}} = \int_{T_{bg}}^{T_m} \left( A_{(A,Z)}(t) + A_{(A,Z+1)}(t) + A_{(A,Z+2)}(t) + \right. \\ \left. + A_{(A-1,Z+1)}(t) + A_{(A-1,Z+2)}(t) \right) dt, \\ (N_n^{exp} - N_n^{bg}) \frac{1}{\epsilon_n} = P_{1n} \cdot \int_{T_{bg}}^{T_{meas}} A_{(A,Z)}(t) dt, \end{array} \right. \quad (2.1)$$

where  $N_{\beta}^{exp}$  and  $N_n^{exp}$  are total numbers of  $\beta$  and neutrons detected;  $N_{\beta}^{bg}$  and  $N_n^{bg}$  are the total numbers of background events;  $\epsilon_{\beta}$  and  $\epsilon_n$  the measured, as explained above, the effective



**Figure 2:**  $\beta$  gated (blue) and  $\beta$ -n gated (red)  $\gamma$ -spectra recorded for the  $A = 96$  (top) and  $A = 97$  (bottom) settings of the mass separator [spectra to be re-formatted].

efficiencies of  $\beta$  detector and the TETRA array;  $A_{(A,Z+1)}$ , ...,  $A_{(A-1,Z+2)}$  are the activities of daughter, grand-daughter nuclei characterized by their decay constants  $\lambda_{(A,Z+1)}$ , ...,  $\lambda_{(A-1,Z+2)}$  respectively. Meantime the activity of the mother nuclei  $A_{(A,Z)}(t)$  at a  $t$ -moment populated with the beam depends on both the decay constant  $\lambda_{(A,Z)}$  and the intensity of the beam,  $\phi$ :

$$A_{(A,Z)}(t) = \begin{cases} \frac{1}{\lambda_{(A,Z)}} \phi - e^{-t \cdot \lambda_{(A,Z)}}, & T_{bg} \leq t \leq T_{coll} \\ \frac{1}{\lambda_{(A,Z)}} (e^{-(T_{coll}+t) \cdot \lambda_{(A,Z)}} (e^{T_{coll} \cdot \lambda_{(A,Z)}} - 1) \cdot \phi, & T_{coll} \leq t \leq T_m \end{cases} \quad (2.2)$$

The system of the Equations 2.1 was solved for  $A=96$  and  $A=97$  collected datasets. Whereas

**Table 2:** Photopeaks attributed to the  $\beta$  and  $\beta$ -n decays of  $^{96}\text{Rb}$  identified in the spectrum recorded for the  $A = 96$  setting of the mass-separator.  $\gamma$ -intensities are given relative to the transition at 815.5 keV or 122.5 keV observed in  $^{96}\text{Sr}$  or  $^{96}\text{Y}$  respectively.

$E_\gamma$ [keV]	Spectrum	$I_\gamma^{rel}$ [%]	$S_\gamma^i/S_{\gamma\beta}^i$ [%]	assignment
122.5(2)	$\gamma$	100(2)	29(1)	$^{96}\text{Sr}(\beta)$
	$\gamma\beta$	100(3)		
204.5(2)	$\gamma$	28(4)	-	$^{96}\text{Rb}(\beta n)$
	$\gamma n$	25(4)		
279.5(2)	$\gamma$	13.0(5)	27(2)	$^{96}\text{Sr}(\beta)$
	$\gamma\beta$	12.0(9)		
353.2(5)	$\gamma$	100(8)	29(3)	$^{96}\text{Rb}(\beta n)$
	$\gamma\beta$	100(14)		
	$\gamma\beta n$	100(15)		
	$\gamma n$	100(8)		
530.1(2)	$\gamma$	11.3(6)	30(3)	$^{96}\text{Sr}(\beta)$
	$\gamma\beta$	11.0(9)		
692.2(2)	$\gamma$	9.0(1.5)	36(6)	$^{96}\text{Rb}(\beta)$
	$\gamma\beta$	10.5(1.5)		
809.6(2)	$\gamma$	110(3)	28(1)	$^{96}\text{Sr}(\beta)$
	$\gamma\beta$	108(3)		
815.2(2)	$\gamma$	100(3)	31(1)	$^{96}\text{Rb}(\beta)$
	$\gamma\beta$	100(4)		
931.9(2)	$\gamma$	23(1)	32(3)	$^{96}\text{Sr}(\beta)$
	$\gamma\beta$	25(2)		
977.9(2)	$\gamma$	23(1)	30(5)	$^{96}\text{Rb}(\beta)$
	$\gamma\beta$	25(2)		
1037.5(2)	$\gamma$	7.5(5)	27(4)	$^{96}\text{Rb}(\beta)$
	$\gamma\beta$	6.6(8)		
1180.5(2)	$\gamma$	4.0(4)	31(7)	$^{96}\text{Rb}(\beta)$
	$\gamma\beta$	4.1(6)		


$\lambda_{(A,Z)}$  was fixed to the value obtained from the fit of neutron activities, the  $\lambda_{(A,Z+1)}, \dots, \lambda_{(A-1,Z+2)}$  were fixed to their table values. The obtained  $P_{1n}$  and  $\phi$  for  $^{96}\text{Rb}$  and  $^{97}\text{Rb}$  are reported in Table 1. A constant average production yield was assumed. The errors on  $P_{1n}$  and  $\phi$  are mostly dominated by associated statistical errors but also by uncertainty on  $\beta$  and neutron efficiencies and uncertainties on half-lives of Sr and Y daughters known from the literature. Once  $P_{1n}$  and  $\phi$  were determined contributions to the  $\beta$ -activity curve accumulated after  $N_{\text{cycles}}$  from  $\beta$ -decay of the parent nucleus and its daughters in two experiments ( $^{96}\text{Rb}$  or  $^{97}\text{Rb}$ ) were derived and are shown by different colour code in Fig. 1 .

In Fig. 3, it is plotted  $P_{1n}$  values versus  $T_{\frac{1}{2}}$  for  $^{96}\text{Rb}$  (left panel) and  $^{97}\text{Rb}$  (right panel) from the experiments cited in the literature where these two properties of  $\beta$ -decay were measured simul-

**Table 3:** Photopeaks attributed to the  $\beta$  and  $\beta$ -n decays of  $^{97}\text{Rb}$  identified in the spectrum recorded for the  $A = 97$  setting of the mass-separator.  $\gamma$ -intensities are given relatively to the transition at 167.0 keV or 1905.2 keV observed in  $^{97}\text{Sr}$  or  $^{97}\text{Y}$  respectively.

$E_\gamma$ [keV]	Spectrum	$I_\gamma^{rel}$ [%]	$S_\gamma^i/S_{\gamma\beta}^i$ [%]	assignment
167.0(4)	$\gamma$	100(2)		$^{97}\text{Rb}(\beta)$
	$\gamma\beta$	100(4)	27(1)	
418.2(2)	$\gamma$	23(1)	33(2)	$^{97}\text{Rb}(\beta n)$
	$\gamma\beta$	27(2)		
585.1.5(2)	$\gamma$	13.0(5)	28(1)	$^{97}\text{Rb}(\beta)$
	$\gamma\beta$	12.0(9)		
600.1(2)	$\gamma$	37(1)	32(2)	$^{97}\text{Rb}(\beta)$
	$\gamma\beta$	43(3)		
644.8(2)	$\gamma$	11.9(6)	28(3)	$^{97}\text{Rb}(\beta)$
	$\gamma\beta$	12.5(5)		
815.4(2)	$\gamma$	100(2)	27(1)	$^{97}\text{Rb}(\beta n)$
	$\gamma\beta$	100(6)		
	$\gamma\beta n$	100(11)		
953.9(2)	$\gamma$	103(4)	28(1)	$^{97}\text{Sr}(\beta)$
	$\gamma\beta$	103(7)		
1905.2(2)	$\gamma$	100(4)	29(1)	$^{97}\text{Sr}(\beta)$
	$\gamma\beta$	100(8)		
2212.3(2)	$\gamma$	39(2)	28(3)	$^{97}\text{Sr}(\beta)$
	$\gamma\beta$	38(4)		

taneously. As seen, our  $P_{1n}$  and  $T_{\frac{1}{2}}$  are in a remarkable agreement with the existing systematics.



**Figure 3:** Comparison of  $P_{1n}$  (x-axis) versus  $T_{\frac{1}{2}}$  for  $^{96}\text{Rb}$  and  $^{97}\text{Rb}$  known from the literature [8–14] to the values measured in the present work.

### 3 Conclusions

TETRA long neutron counter is a unique device which is currently used at ALTO ISOL facility to measure gross-properties of  $\beta$ -decay of neutron-rich nuclei. The configuration of  $\beta$ -decay station configured to the neutron detection mode allows possibility of simultaneous detection of  $\beta$ ,  $\gamma$  and neutron radioactivity followed  $\beta$ -decay of the radioactive source accumulated with the beam at the centre of the detection system. The performance of the setup was characterized using radioactive sources of  $^{96,97}\text{Rb}$  produced from the photo-fission of  $^{238}\text{U}$ . The sources were carefully chosen because their  $\beta$ -decay properties are known and thus represent well-known reference cases to test neutron detectors and validate the experimental procedures. The half-lives and the probabilities of  $\beta$ -delayed neutron emission for  $^{96,97}\text{Rb}$  measured in the experiment are in good agreement with the previously reported values. This experimental procedure was necessary to validate the method exploited to extract un-known values of  $P_{1n}$  of different nuclei with large N/Z ratio (to be) produced at ALTO ISOL facility.

### Acknowledgments

We would like to thank the technical staff of the Tandem/ALTO facility for their assistance to provide excellent operation of the electron accelerator and mass separator PARNNe. The authors wish to acknowledge support from the bilateral agreement JINR- CNRS/IN2P3 No. XX-XX. This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No. 771036 (ERC CoG MAIDEN). The research leading to this results has received funding from the European Union's HORIZON2020 Program under the grant agreement No. 654002.

### References

- [1] *D. Testov*, D. Verney, B. Roussiere et al., Nucl. Inst. Meth. A **815** (2016) 96
- [2] *D. Testov*, Ch.Briançon, S.Dmitriev et al., Physics of Atomic Nuclei, **72** (2009) 1
- [3] *A. Etile*, D. Verney, N. N. Arsenyev et al., Phys. Rev. C, **91** (2015) 064317
- [4] *F. Azaiez*, S. Essabaa , F. Ibrahim, and D. Verney, Nucl. Phys. News. **23**, (2013)
- [5] *D. Verney*, D. Testov, B. Roussiere et al, Phys. Rev. C **95** (2017) 054320.
- [6] *A. Gottardo*, D. Verney, I. Deloncle, et al, Phys. Lett. B **772** (2017) 359
- [7] *D. Testov*, E. Kuznetsova, J.N. Wilson J. Instrum. **10** (2015) P09011
- [8] *I. Amarel*, H. Gauvin, A. Johnson, J. Nucl. Chem., **31** (1969) 577
- [9] *E. Roeckl*, P.F. Dittner, R. Klapischh et al., Nucl. Phys A **222** (1974) 621
- [10] *C. Ristori*, J. Crancon, K.D. Wunsch et al., Z.Phys. A **290** (1979) 311
- [11] *E. Lund*, P. Hoff, K. Aleklett et al., Z.Phys. A **294** (1980) 233
- [12] *G. Engler*, and E. Ne'Eman Nucl. Phys. A **367** (1981) 29-40
- [13] *B. Pfeiffer*, K.Kratz, P.Moller, Prog.Nucl.Energy **41** (2002) 39
- [14] *Ray A. Warner* and Paul L. Reeder Rad. Eff. **94** (1986) 27