



HAL
open science

Erratum: Comet 67P outbursts and quiescent coma at 1.3 AU from the Sun: dust properties from Rosetta/VIRTIS-H observations

Dominique Bockelée-Morvan, G. Rinaldi, S. Erard, C. Leyrat, S Capaccioni,
P. Drossart, G. Filacchione, S Migliorini, G Quirico, G. Mottola, et al.

► **To cite this version:**

Dominique Bockelée-Morvan, G. Rinaldi, S. Erard, C. Leyrat, S Capaccioni, et al.. Erratum: Comet 67P outbursts and quiescent coma at 1.3 AU from the Sun: dust properties from Rosetta/VIRTIS-H observations. *Monthly Notices of the Royal Astronomical Society*, 2017, 469 (Suppl_2), pp.S842-S843. 10.1093/mnras/sty533 . hal-02266787

HAL Id: hal-02266787

<https://hal.science/hal-02266787>

Submitted on 24 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Erratum: Comet 67P outbursts and quiescent coma at 1.3 AU from the Sun: dust properties from Rosetta/VIRTIS-H observations

by Dominique Bockelée-Morvan,¹★ G. Rinaldi,² S. Erard,¹ C. Leyrat,¹ F. Capaccioni,² P. Drossart,¹ G. Filacchione,² A. Migliorini,² E. Quirico,³ S. Mottola,⁴ G. Tozzi,⁵ G. Arnold,⁴ N. Biver,¹ M. Combes,¹† J. Crovisier,¹ A. Longobardo,² M. Blecka⁶ and M.-T. Capria²

¹LESIA, Observatoire de Paris, PSL Research University, CNRS, Sorbonne Universités, UPMC Univ. Paris 06, Univ. Paris-Diderot, Sorbonne Paris Cité, 5 place Jules Janssen, F-92195 Meudon, France

²INAF-IAPS, Istituto di Astrofisica e Planetologia Spaziali, via del fosso del Cavaliere, 100, I-00133, Rome, Italy

³CNRS, Institut de Planétologie et d'Astrophysique de Grenoble, Université Grenoble Alpes, 414 rue de la Piscine, BP53, F-38041 Grenoble, France

⁴Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institute for Planetary Research, Rutherfordstrasse 2, D-12489 Berlin, Germany

⁵INAF, Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, I-50125 Firenze, Italy

⁶Space Research Centre, Polish Academy of Sciences, Bartycka 18a, PL-00716, Warsaw, Poland

Key words: errata, addenda – comets: general – comets: individual: 67P/Churyumov-Gerasimenko – infrared: planetary systems.




Figure 1. Replaces Fig. 8 of Bockelée-Morvan et al. (2017). Results of Mie-scattering calculations for moderately porous ($P_{\max} = 0.5$) carbon/olivine mixtures. The grey area (not included in the original figure) shows the range of observed values on 13 and 14 September 2015 (pre-outburst).

The paper ‘Comet 67P outbursts and quiescent coma at 1.3 AU from the Sun: dust properties from Rosetta/VIRTIS-H observations’ was published in MNRAS 469, S443 (2017). While performing a follow-up investigation, we discovered a numerical error in the algorithms that were developed to model the infrared continuum emission from a population of dust particles. The results of the scattering models for compact or moderately porous grains (Mie theory) and fluffy grains (Rayleigh-Gans-Debye theory, RGD) are both affected. Though the general conclusions of the paper are




Figure 2. Replaces Fig. 9 of Bockelée-Morvan et al. (2017). Results of Rayleigh-Gans-Debye calculations for fractal aggregates made of carbon/olivine mixtures.

unchanged, the quantitative constraints obtained on the dust size distribution in the quiescent coma are slightly different.

Figs 8, 9, 10 and 11 should be replaced by, respectively, Figs 1, 2, 3 and 4 of this erratum. The new results introduce the following changes :

(i) Section 4.2, second paragraph (quiescent coma, Mie results): for moderately porous grains ($P_{\max} = 0.5$), the sets of size distribution parameters which fit satisfactorily the measurements are now (a_{\min}, β) = (0.3 μm , 2.5) and (0.9 μm , 3.0) (see Fig. 1). For non-porous grains, the updated parameters fitting the colour temperature and colour (whereas overestimating the bolometric albedo) are (a_{\min}, β) = (0.1 μm , 2.5) and (0.5 μm , 3.0).

(ii) Section 4.2, last paragraph (quiescent coma, RGD results): we find now that, for a relative number of fluffy aggregates of

* E-mail: dominique.bockelee@obspm.fr

† deceased.




Figure 3. Replaces Fig. 10 of Bockelée-Morvan et al. (2017). Results from Mie-scattering calculations: correlation between spectral properties.




Figure 4. Replaces Fig. 11 of Bockelée-Morvan et al. (2017). Results for relative fractions of 25 per cent of fractal aggregates and 75 per cent of moderately porous carbon/olivine mixtures ($P_{\max} = 0.5$). The grey area (not included in the original figure) shows the range of observed values on 13 and 14 September 2015 (pre-outburst).

25 per cent, the VIRTIS-H data can be explained providing the minimum radius of the particles exceeds $10 \mu\text{m}$ (Fig. 4). Good matches of colour, colour temperature and bolometric albedo are obtained for $(a_{\min}, \beta) = (30 \mu\text{m}, 3.0)$ and $(20 \mu\text{m}, 3.5)$. Size indexes smaller than 2.5 are excluded (Fig. 4).

(iii) Section 6 (Summary and conclusions), fourth paragraph (Size distribution in quiescent coma): the parameters (a_{\min}, β) fitting the data assuming moderately porous grains should be updated following the corrections provided above. It was previously derived that, when considering mixtures of compact particles and fractal aggregates, a size index in the range 2–2.5 is favoured. We now find that the size index should be ≥ 3 , which is consistent with the constraint obtained by Fulle et al. (2016) from GIADA data.

REFERENCES

- Bockelée-Morvan D. et al., 2017, MNRAS, 469, S443
 Fulle M. et al., 2016, ApJ, 821, 19

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.