



HAL
open science

DIRICHLET PRODUCT OF DERIVATIVE ARITHMETIC WITH AN ARITHMETIC FUNCTION MULTIPLICATIVE

Es-Said En-Naoui

► **To cite this version:**

Es-Said En-Naoui. DIRICHLET PRODUCT OF DERIVATIVE ARITHMETIC WITH AN ARITHMETIC FUNCTION MULTIPLICATIVE. 2019. hal-02267882

HAL Id: hal-02267882

<https://hal.science/hal-02267882>

Preprint submitted on 22 Aug 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

DIRICHLET PRODUCT OF DERIVATIVE ARITHMETIC WITH AN ARITHMETIC FUNCTION MULTIPLICATIVE

A PREPRINT

Es-said En-naoui
essaidennaoui1@gmail.com

August 21, 2019

ABSTRACT

We define the derivative of an integer to be the map sending every prime to 1 and satisfying the Leibniz rule. The aim of this article is to calculate the Dirichlet product of this map with a function arithmetic multiplicative.

1 Introduction

Barbeau [1] defined the arithmetic derivative as the function $\delta : \mathbb{N} \rightarrow \mathbb{N}$, defined by the rules :

1. $\delta(p) = 1$ for any prime $p \in \mathbb{P} := \{2, 3, 5, 7, \dots, p_i, \dots\}$.
2. $\delta(ab) = \delta(a)b + a\delta(b)$ for any $a, b \in \mathbb{N}$ (the Leibnitz rule).

Let n a positive integer, if $n = \prod_{i=1}^s p_i^{\alpha_i}$ is the prime factorization of n , then the formula for computing the arithmetic derivative of n is (see, e.g., [1, 3]) giving by :

$$\delta(n) = n \sum_{i=1}^s \frac{\alpha_i}{p_i} = n \sum_{p^\alpha || n} \frac{\alpha}{p} \quad (1)$$

A brief summary on the history of arithmetic derivative and its generalizations to other number sets can be found, e.g., in [4].

First of all, to cultivate analytic number theory one must acquire a considerable skill for operating with arithmetic functions. We begin with a few elementary considerations.

Definition 1 (arithmetic function). An **arithmetic function** is a function $f : \mathbb{N} \rightarrow \mathbb{C}$ with domain of definition the set of natural numbers \mathbb{N} and range a subset of the set of complex numbers \mathbb{C} .

Definition 2 (multiplicative function). A function f is called an **multiplicative function** if and only if :

$$f(nm) = f(n)f(m) \quad (2)$$

for every pair of coprime integers n, m . In case (2) is satisfied for every pair of integers n and m , which are not necessarily coprime, then the function f is called **completely multiplicative**.

Clearly, if f are a multiplicative function, then $f(n) = f(p_1^{\alpha_1}) \dots f(p_s^{\alpha_s})$, for any positive integer n such that $n = p_1^{\alpha_1} \dots p_s^{\alpha_s}$, and if f is completely multiplicative, so we have : $f(n) = f(p_1)^{\alpha_1} \dots f(p_s)^{\alpha_s}$.

Example 3. Let $n \in \mathbb{N}^*$, This is the same classical arithmetic functions used in this article :

1. **Identity function** : The function defined by $Id(n) = n$ for all positive integer n .

$$2. \text{ The Euler phi function : } \varphi(n) = \sum_{\substack{k=1 \\ \gcd(k,n)=1}}^n 1.$$

$$3. \text{ The number of distinct prime divisors of } n : \omega(n) = \sum_{p|n} 1.$$

$$4. \text{ The Mobius function : } \mu(n) = \begin{cases} 1 & \text{if } n = 1 \\ 0 & \text{if } p^2|n \text{ for some prime } p \\ (-1)^{\omega(n)} & \text{otherwise} \end{cases}$$

$$5. \text{ number of positive divisors of } n \text{ defined by : } \tau(n) = \sum_{d|n} 1.$$

$$6. \text{ sum of divisors function of } n \text{ defined by : } \sigma(n) = \sum_{d|n} d.$$

Now ,if $f, g : \mathbb{N} \rightarrow \mathbb{C}$ are two arithmetic functions from the positive integers to the complex numbers, the Dirichlet convolution $f * g$ is a new arithmetic function defined by:

$$(f * g)(n) = \sum_{d|n} f(d)g\left(\frac{n}{d}\right) = \sum_{ab=n} f(a)g(b) \quad (3)$$

where the sum extends over all positive divisors d of n , or equivalently over all distinct pairs (a, b) of positive integers whose product is n .

In particular, we have $(f * g)(1) = f(1)g(1)$, $(f * g)(p) = f(1)g(p) + f(p)g(1)$ for any prime p and for any power prime p^m we have :

$$(f * g)(p^m) = \sum_{j=0}^m f(p^j)g(p^{m-j}) \quad (4)$$

This product occurs naturally in the study of Dirichlet series such as the Riemann zeta function. It describes the multiplication of two Dirichlet series in terms of their coefficients:

$$\left(\sum_{n \geq 1} \frac{(f * g)(n)}{n^s} \right) = \left(\sum_{n \geq 1} \frac{f(n)}{n^s} \right) \left(\sum_{n \geq 1} \frac{g(n)}{n^s} \right) \quad (5)$$

with Riemann zeta function or is defined by :

$$\zeta(s) = \sum_{n \geq 1} \frac{1}{n^s}$$

These functions are widely studied in the literature (see, e.g., [5, 6, 7]).

Now before to past to main result we need this propriety , if f and g are multiplicative function , then $f * g$ is multiplicative.

2 Main results

In this section we give the new result of Dirichlet product between derivative arithmetic and an arithmetic function multiplicative f , and we will give the relation between τ and the derivative arithmetic .

Theorem 4. *Given a multiplicative function f , and lets n and m two positive integers such that $\gcd(n, m) = 1$, Then we have :*

$$(f * \delta)(nm) = (Id * f)(n).(f * \delta)(m) + (Id * f)(m).(f * \delta)(n) \quad (6)$$

Proof. Lets n and m two positive integers such that $\gcd(n, m) = 1$, and let f an arithmetic function multiplicative , then we have :

$$\begin{aligned}
(f * \delta)(nm) &= \sum_{d|nm} f\left(\frac{nm}{d}\right)\delta(d) = \sum_{\substack{d_1|n \\ d_2|m}} f\left(\frac{nm}{d_1 d_2}\right)\delta(d_1 d_2) = \sum_{\substack{d_1|n \\ d_2|m}} f\left(\frac{n}{d_1}\right)f\left(\frac{m}{d_2}\right)\left(d_1\delta(d_2) + d_2\delta(d_1)\right) \\
&= \sum_{\substack{d_1|n \\ d_2|m}} \left(d_1 f\left(\frac{n}{d_1}\right)f\left(\frac{m}{d_2}\right)\delta(d_2) + d_2 f\left(\frac{m}{d_2}\right)f\left(\frac{n}{d_1}\right)\delta(d_1)\right) \\
&= \left(\sum_{d_1|n} d_1 f\left(\frac{n}{d_1}\right)\right)\left(\sum_{d_2|m} f\left(\frac{m}{d_2}\right)\delta(d_2)\right) + \left(\sum_{d_2|m} d_2 f\left(\frac{m}{d_2}\right)\right)\left(\sum_{d_1|n} f\left(\frac{n}{d_1}\right)\delta(d_1)\right) \\
&= (Id * f)(n) \cdot (f * \delta)(m) + (Id * f)(m) \cdot (f * \delta)(n)
\end{aligned}$$

□

Lemma 5. For any natural number n , if $n = \prod_{i=1}^s p_i^{\alpha_i}$ is the prime factorization of n , then :

$$(f * \delta)(n) = (Id * f)(n) \sum_{i=1}^s \frac{(f * \delta)(p_i^{\alpha_i})}{(Id * f)(p_i^{\alpha_i})} \quad (7)$$

Proof. Let n a positive integer such that $n = p_1^{\alpha_1} \dots p_s^{\alpha_s}$ and let f an arithmetic function , Then :

$$\begin{aligned}
(f * \delta)(n) &= (f * \delta)(p_1^{\alpha_1} \dots p_s^{\alpha_s}) \\
&= (Id * f)(p_2^{\alpha_2} \dots p_s^{\alpha_s}) \cdot (f * \delta)(p_1^{\alpha_1}) + (Id * f)(p_1^{\alpha_1}) \cdot (f * \delta)(p_2^{\alpha_2} \dots p_s^{\alpha_s}) \\
&= (Id * f)(n) \cdot \frac{(f * \delta)(p_1^{\alpha_1})}{(Id * f)(p_1^{\alpha_1})} + (Id * f)(p_1^{\alpha_1}) \cdot \left[(Id * f)(p_3^{\alpha_3} \dots p_s^{\alpha_s}) \cdot (f * \delta)(p_2^{\alpha_2}) + \right. \\
&\quad \left. + (Id * f)(p_2^{\alpha_2}) \cdot (f * \delta)(p_3^{\alpha_3} \dots p_s^{\alpha_s}) \right] \\
&= (Id * f)(n) \frac{(f * \delta)(p_1^{\alpha_1})}{(Id * f)(p_1^{\alpha_1})} + (Id * f)(n) \frac{(f * \delta)(p_2^{\alpha_2})}{(Id * f)(p_2^{\alpha_2})} + \\
&\quad + (Id * f)(p_1^{\alpha_1}) (Id * f)(p_2^{\alpha_2}) (f * \delta)(p_3^{\alpha_3} \dots p_s^{\alpha_s}) \\
&\vdots \\
&= (Id * f)(n) \frac{(f * \delta)(p_1^{\alpha_1})}{(Id * f)(p_1^{\alpha_1})} + (Id * f)(n) \frac{(f * \delta)(p_2^{\alpha_2})}{(Id * f)(p_2^{\alpha_2})} + \dots + (Id * f)(n) \frac{(f * \delta)(p_s^{\alpha_s})}{(Id * f)(p_s^{\alpha_s})} \\
&= (Id * f)(n) \left[\frac{(f * \delta)(p_1^{\alpha_1})}{(Id * f)(p_1^{\alpha_1})} + \dots + \frac{(f * \delta)(p_s^{\alpha_s})}{(Id * f)(p_s^{\alpha_s})} \right] \\
&= (Id * f)(n) \sum_{i=1}^s \frac{(f * \delta)(p_i^{\alpha_i})}{(Id * f)(p_i^{\alpha_i})}
\end{aligned}$$

□

an other prof by induction on s that if $n = \prod_{i=1}^s p_i^{\alpha_i}$ then $(f * \delta)(n) = (Id * f)(n) \sum_{i=1}^s \frac{(f * \delta)(p_i^{\alpha_i})}{(Id * f)(p_i^{\alpha_i})}$.

Proof. Consider $n \in \mathbb{N}$ and express $n = \prod_{i=1}^s p_i^{\alpha_i}$ where all p_i are distinct .

where $s = 1$, it is clear that $(f * \delta)(n) = (Id * f)(n) \sum_{i=1}^1 \frac{(f * \delta)(p_i^{\alpha_i})}{(Id * f)(p_i^{\alpha_i})} = (Id * f)(p_1^{\alpha_1}) \frac{(f * \delta)(p_1^{\alpha_1})}{(Id * f)(p_1^{\alpha_1})} = (f * \delta)(p_1^{\alpha_1})$.

Assume that $n = \prod_{i=1}^s p_i^{\alpha_i}$, then we have :

$$\begin{aligned}
(id * \delta)(n.p_{s+1}^{\alpha_{s+1}}) &= (Id * f)(p_{s+1}^{\alpha_{s+1}}) \cdot (f * \delta)(n) + (Id * f)(n) \cdot (f * \delta)(p_{s+1}^{\alpha_{s+1}}) \\
&= (Id * f)(p_{s+1}^{\alpha_{s+1}}) \cdot (Id * f)(n) \sum_{i=1}^s \frac{(f * \delta)(p_i^{\alpha_i})}{(Id * f)(p_i^{\alpha_i})} + (Id * f)(p_{s+1}^{\alpha_{s+1}}) \cdot (Id * f)(n) \frac{(f * \delta)(p_{s+1}^{\alpha_{s+1}})}{(Id * f)(p_{s+1}^{\alpha_{s+1}})} \\
&= (Id * f)(n.p_{s+1}^{\alpha_{s+1}}) \sum_{i=1}^s \frac{(f * \delta)(p_i^{\alpha_i})}{(Id * f)(p_i^{\alpha_i})} + (Id * f)(n.p_{s+1}^{\alpha_{s+1}}) \frac{(f * \delta)(p_{s+1}^{\alpha_{s+1}})}{(Id * f)(p_{s+1}^{\alpha_{s+1}})} \\
&= (Id * f)(n.p_{s+1}^{\alpha_{s+1}}) \left[\sum_{i=1}^s \frac{(f * \delta)(p_i^{\alpha_i})}{(Id * f)(p_i^{\alpha_i})} + \frac{(f * \delta)(p_{s+1}^{\alpha_{s+1}})}{(Id * f)(p_{s+1}^{\alpha_{s+1}})} \right] \\
&= (Id * f)(n.p_{s+1}^{\alpha_{s+1}}) \sum_{i=1}^{s+1} \frac{(f * \delta)(p_i^{\alpha_i})}{(Id * f)(p_i^{\alpha_i})}
\end{aligned}$$

□

Proposition 6. Let f a function arithmetic multiplicative, and δ the derivative arithmetic, then we have :

$$(Id * \delta)(n) = \frac{1}{2} \tau(n) \delta(n) \quad (8)$$

Proof. Since $(Id * Id)(n) = \sum_{d|n} \frac{n}{d} = n \sum_{d|n} 1 = n\tau(n)$.

and : $(Id * \delta)(p^\alpha) = \sum_{j=1}^{\alpha} \delta(p^j) Id(p^{\alpha-j}) = \sum_{j=1}^{\alpha} j p^{j-1} p^{\alpha-j} = \frac{1}{2} \alpha(\alpha+1) p^{\alpha-1}$.

Then for every a positive integer n such that $n = p_1^{\alpha_1} \dots p_s^{\alpha_s}$, we have :

$$\begin{aligned}
(Id * \delta)(n) &= (Id * Id)(n) \sum_{i=1}^s \frac{(Id * \delta)(p_i^{\alpha_i})}{(Id * Id)(p_i^{\alpha_i})} \\
&= n\tau(n) \sum_{i=1}^s \frac{\frac{1}{2} \alpha_i (\alpha_i + 1) p_i^{\alpha_i - 1}}{p_i^{\alpha_i} \tau(p_i^{\alpha_i})} \\
&= n\tau(n) \sum_{i=1}^s \frac{\frac{1}{2} \alpha_i (\alpha_i + 1) p_i^{\alpha_i - 1}}{p_i^{\alpha_i} (\alpha_i + 1)} \\
&= \frac{1}{2} n\tau(n) \sum_{i=1}^s \frac{\alpha_i}{p_i} = \frac{1}{2} \tau(n) \delta(n)
\end{aligned}$$

□

So by the proposition 6, and the equality 5 we have this relation between arithmetic derivative and the function τ :

$$2\zeta(s-1) \sum_{n \geq 1} \frac{\delta(n)}{n^s} = \sum_{n \geq 1} \frac{\delta(n) \tau(n)}{n^s} \quad (9)$$

Let defined the new function arithmetic called **En-naoui** function, by :

$$\Phi_\varphi(n) = n \sum_{p|n} \left(1 - \frac{1}{p} \right) \quad (10)$$

Then we have this equality related between δ arithmetic function :

$$(\mu * \delta)(n) = \varphi(n) \left(\delta(n) - 2\omega(n) + B(n) + \frac{\Phi_\varphi(n)}{n} + \frac{(B * Id)(n)}{\sigma(n)} \right). \quad (11)$$

with B is the arithmetic function defined by : $B(n) = \sum_{p^\alpha || n} \alpha p$. In next article i will prove this equality, just I need to submit an article about this new function .

3 Acknowledgements

I would like to thank S.Allo for the opportunity to collaborate on my research and for his continued mentorship, and friendship, and I'm indebted to the Professor Zouhair Mouayn to endorse me to submit my article here. I also extend my deepest appreciation to my family for opening their minds, hearts, and homes to me and my work.

References

- [1] E. J. Barbeau. Remark on an arithmetic derivative, *Canad. Math. Bull.* 4 (1961), 117–122.
- [2] A. Buium. Arithmetic analogues of derivations, *J. Algebra*, 198 (1997), 290–299.
- [3] V. Ufnarovski and B. Åhlander. How to differentiate a number, *J. Integer Seq.*, 6 , (2003).
- [4] M. Stay. Generalized number derivatives, *J. Integer Seq.*, 8 , (2005).
- [5] T. M. Apostol. Introduction to Analytic Number Theory, *Springer-Verlag, New York* , (1976).
- [6] M.Lewinter and J.Meyer. Elementary number theory with programming , *Wiley* , (2016).
- [7] Heng Huat Chan, Analytic number theory for undergraduates , *World Scientific Publishing Co. Pte. Ltd.* , (2009).