

Telomere dynamic in humans and animals: Review and perspectives in environmental toxicology

Maxime Louzon, Michael Coeurdassier, Frédéric Gimbert, Benjamin Pauget,

Annette de Vaufleury

► To cite this version:

Maxime Louzon, Michael Coeurdassier, Frédéric Gimbert, Benjamin Pauget, Annette de Vaufleury. Telomere dynamic in humans and animals: Review and perspectives in environmental toxicology. Environment International, 2019, 131, 10.1016/j.envint.2019.105025 . hal-02269457

HAL Id: hal-02269457 https://hal.science/hal-02269457

Submitted on 25 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Telomere dynamic in humans and animals: review and perspectives in 1 environmental toxicology 2 3 4 Maxime Louzon¹, Michael Coeurdassier¹, Frédéric Gimbert¹, Benjamin Pauget², Annette de 5 Vaufleury1* ¹ Department Chrono-Environnement, UMR UFC/CNRS 6249 USC INRA University of Bourgogne 6 7 Franche-Comté, 16 route de Gray, 25000 Besançon, France. 8 ² TESORA, Le Visium, 22 avenue Aristide Briand, 94110 Arcueil, France. 9 *Corresponding author: annette.devaufleury@univ-fcomte.fr 10 **Keywords** 11 telomeres, attrition, vertebrates, invertebrates, ageing, pollutants 12 13 14 15 16 17 18 19

20

21 Abstract

22 Telomeres (TLs) play major roles in stabilizing the genome and are usually shortened 23 with ageing. The maintenance of TLs is ensured by two mechanisms involving telomerase 24 (TA) enzyme and alternative lengthening telomeres (ALT). TL shortening and/or TA 25 inhibition have been related to health effects on organisms (leading to reduced reproductive 26 lifespan and survival), suggesting that they could be key processes in toxicity mechanisms (at 27 molecular and cellular levels) and relevant as an early warning of exposure and effect of 28 chemicals on human health and animal population dynamics. Consequently, a critical analysis 29 of knowledge about relationships between TL dynamic and environmental pollution is 30 essential to highlight the relevance of TL measurement in environmental toxicology.

31 The first objective of this review is to provide a survey on the basic knowledge about 32 TL structure, roles, maintenance mechanisms and causes of shortening in both vertebrates 33 (including humans) and invertebrates. Overall, TL length decreases with ageing but some 34 unexpected exceptions are reported (e.g., in species with different lifespans, such as the 35 nematode *Caenorhabditis elegans* or the crustacean *Homarus americanus*). Inconsistent 36 results reported in various biological groups or even between species of the same genus (e.g., 37 the microcrustacean Daphnia sp.) indicate that the relation usually proposed between TL 38 shortening and a decrease in TA activity cannot be generalized and depends on the species, 39 stage of development or lifespan. Although the scientific literature provides evidence of the 40 effect of ageing on TL shortening, much less information on the relationships between 41 shortening, maintenance of TLs, influence of other endogenous and environmental drivers, 42 including exposure to chemical pollutants, is available, especially in invertebrates.

43 The second objective of this review is to connect knowledge on TL dynamic and 44 exposure to contaminants. Most of the studies published on humans rely on correlative epidemiological approaches and few *in vitro* experiments. They have shown TL attrition when exposed to contaminants, such as polycyclic aromatic hydrocarbons (PAH), polychlorinated biphenyls (PCB), pesticides and metallic elements (ME). In other vertebrates, the studies we found deals mainly with birds and, overall, report a disturbance of TL dynamic consecutively to exposure to chemicals, including metals and organic compounds. In invertebrates, no data are available and the potential of TL dynamic in environmental risk assessment remains to be explored.

52 On the basis of the main gaps identified some research perspectives (*e.g.*, impact of 53 endogenous and environmental drivers, dose response effects, link between TL length, TA 54 activity, longevity and ageing) are proposed to better understand the potential of TL and TA 55 measurements in humans and animals in environmental toxicology.

56

57 **1.** <u>Introduction</u>

58 Telomeres (TLs) are repetitive, non-coding DNA sequences located at the ends of 59 chromosomes. TLs were identified in the 1930s by McClintock (1941) and Müller (1938), 60 who described their potential functions in maintaining chromosome stability by observing the 61 chromosomes of corn and drosophila. It was postulated that cells were not able to divide themselves into "infinity" (Weismann, 1889). This theory was confirmed in the 1960s 62 63 (Hayflick and Moorhead, 1961; Hayflick, 1965) on a human foetal cell culture (division 64 ability between 40 and 60 times before entrance in a senescence phase). Olovnikov theorized 65 the *Hayflick limit*, which postulates that somatic cells have a limited number of divisions in connection with the loss of DNA sequences at the extremities of chromosomes, *i.e.*, TLs. 66 67 When this limit is exceeded, cells enter senescence (Olovnikov, 1971). The popularity of TLs grown over the last four decades since the discovery of a repeated sequence of six telomeric 68

nucleotides of a ciliated protozoan, Tetrahymena thermophila (CCCCAA), and of the 69 70 protective role of this sequence against chromosome degradation (Yao et al., 1981; Szostak 71 and Blackburn, 1982; Shampay et al., 1984). TLs are mainly known to protect and stabilize 72 DNA by protecting chromosomes from degradation (end-to-end fusion and erosion), 73 stabilizing broken chromosome ends, modulating the DNA repair machinery of double strand 74 break (DSB) damage with non-homologous end-joining (NHEJ), counting the number of cell 75 divisions (replicative potential of cells), participating in mitotic and meiotic chromosome 76 segregation and the attachment of chromosomes to the nuclear envelope (Haber, 1999; de Lange, 2005; Oeseburg et al., 2010; Grach, 2013). For humans, TL length decreases with 77 78 ageing (Aubert and Lansdorp, 2008) and TL attrition is recognized as a biomarker of the nine 79 hallmarks of ageing (Lopez-Otin et al., 2013). Mechanisms of TL maintenance limit fast TL 80 attrition and are closely associated with the activities of some enzymes, the most well known 81 being telomerase, or TA (Greider and Blackburn, 1985; Greider and Blackburn, 1989).

82 If TL attrition occurs naturally, some studies report that it can be accelerated by various 83 biological (e.g., infections) and environmental (e.g., chemical substances) stressors known for their toxicological effects. However, there is a lack of knowledge of various stressors (e.g., 84 85 contaminants, biotic and abiotic factors) and species, including invertebrates. Furthermore, data regarding the actions of these stressors on TL maintenance pathways that are key to 86 87 protecting and stabilizing the genome against various stressors, including pollutants, are 88 lacking. Based on data showing that chemicals are able to induce human premature ageing 89 and decrease lifespan (Chen et al., 2013; Puri et al., 2017), it appears relevant to investigate 90 on TL dynamic to search for bridges between molecular and cellular effects (impact on TL length and/or TL maintenance), health effects (e.g., survival, reproductive lifespan) and 91 92 consequences on populations (e.g., density, growth).

93

94 The aim of this review is to identify research questions on TL dynamic in vertebrates and 95 invertebrates in relation to ageing, lifespan and environmental toxicology. The main question 96 concerns the potential use of TL measurement as a relevant biomarker, especially in the 97 framework of the One Health concept, to assess the effects of multiple stressors at the human-98 animal-ecosystem interface (Destournieux-Garzón et al., 2018). Thus, the first objective of 99 this review is to report basic knowledge about the structure, roles and maintenance 100 mechanisms of TLs and causes of TL shortening in the animal kingdom. On the basis of this 101 premise, the second objective is to connect knowledge on TL dynamic and exposure to 102 contaminants. Finally, based on the lack of knowledge of TL dynamic in environmental 103 toxicology, some perspectives are proposed.

104

2. <u>Structure variability of telomeres in the animal kingdom</u>

105 In animals, TL sequences (TTAGGG)_n are highly conserved (Meyne et al., 1989; Traut et 106 al., 2007), with slight variations in phyla such as Arthropoda (Gomes et al., 2010). Sequence 107 conservation results, for example, in similar TL sequences for a heterothermic invertebrate 108 (the snail *Cantareus aspersus*) and a homeothermic vertebrate (*Homo sapiens*), while most 109 insects possess a pentameric motif (TTAGG), with the exception of Coleoptera (Frydrichovà 110 et al., 2004; Vitkovà et al., 2005; Gomes et al., 2011). TL sequences have been determined in 111 various species (*Table 1*), among which some are used as bioindicators in environmental 112 toxicology (e.g., Daphnia pulex, Eisenia foetida, C. aspersus). The repetition number of TL 113 sequences is variable according to species or age and can frequently reach several thousands 114 of nucleotides in Metazoa. For instance, in human lymphocytes, the size ranges between 2.5 115 to 15 kilo base pairs (kbp) depending on the patient (O'Callaghan and Fenech, 2011).

The end of the TL is commonly called a loop because it looks like a hairpin structure (Griffith et al., 1999). Loops are formed by double-stranded TL repeats, the T-loop and the Dloop (Greider, 1999). The terminal region of the TL is a single-stranded overhang that is 3' G-

rich and is known as the G-tail (guanine-rich hairpin tail). This region forms G-quadruplexes, 119 120 where each guanine base participates in hydrogen bond formation (Lipps and Rhodes, 2009) 121 in a square planar arrangement that is stabilized by central cations (Williamson et al., 1989; 122 Move et al., 2015). TL functions are dependent on G-quadruplexes that are associated with 123 proteins (Paeschke et al., 2005). In humans, TLs bear a protein complex, the shelterin 124 complex, which is mainly composed of six different proteins: TRF1, TRF2, RAP1, TIN2, 125 TPP1 and POT1 (Palm and de Lange, 2008; Oeseburg et al., 2010) (Figure 1). This DNA-126 protein complex is essential to protect TLs by the implication of TL maintenance machinery, 127 as it avoids the inhibition of various DNA damage response (DDR) and DNA repair 128 machinery (de Lange, 2009; Lu et al., 2013). Some proteins in humans associated with the 129 shelterin complex have also been found in several animal species, such as POT1 in 130 Caenorhabditis elegans (Ferreira et al., 2013) or in Daphnia sp (Schumpert et al., 2015). All 131 these proteins play various roles in TL maintenance (Martinez and Blasco, 2010) and are 132 summarized in *Table 2*.

133 Although the structure presented in *Figure 1* is accepted as the most commonly found in 134 animal kingdom, some studies have shown that it is not ubiquitous, notably for insects, such 135 as, Chironomus sp., Drosophila melanogaster and Bombyx mori (Louis, 2002; Fujiwara et al., 136 2005; George et al., 2006; Mason et al., 2008; Pardue and DeBaryshe, 2008). For D. 137 *melanogaster*, the shelterin complex is replaced by a same architecture, the terminin complex 138 (Raffa et al., 2013), composed by five proteins (HOAP, HIPHOP, MOI, VER and TEA) and 139 the function of terminin for diptera is mainly to protect telomere from fusion events or DNA 140 damage response (Raffa et al., 2013; Zhang et al., 2016; Cicconi et al., 2017).

141

3. <u>Maintenance of telomeres (TLs)</u>

142 Various mechanisms operate to limit TL attrition (*i.e.*, TA, ALT) and/or to repair TLs
143 when they are damaged.

144

145 **3.1.** Roles of telomerase (TA)

146 Telomerases are a group of enzymes coupled to RNA that synthesize de novo the 147 extremities of the TL to limit its fast attrition in proliferative cell types (Greider and 148 Blackburn, 1985; Greider and Blackburn, 1989). For example, a single TA enzyme able to 149 synthesize 74 telomeric nucleotides per minute in vitro has been described in the ciliated 150 protozoan Tetrahymena thermophila (Greider, 1991). To synthetize TLs, TA uses a catalytic 151 component, TERT (telomerase reverse transcriptase; Nakamura et al., 1997), that has been 152 characterized in many animals (e.g., Xenopus laevis and Bombyx mori) (Kuramoto et al., 153 2001; Osanai et al., 2006). The activation of TERT requires its binding with two proteins of 154 the shelterin complex, TPP1 and POT1 (Sandin and Rhodes, 2014). Then, TERT fosters the 155 de novo synthesis of TLs by using the second subunit of TA, an RNA template (TERC or 156 telomerase RNA component). In addition, TA can interact with G-quadruplexes to allow TL 157 extension (Moye et al., 2015).

158 TA stability and activity can be reduced by the actions of hormone (e.g., corticosterone; 159 Gopalakrishnan et al., 2013; Angelier et al., 2018) and/or altered genes (e.g., DKC1, dyskerin 160 pseudouridine synthase 1; Bertuch, 2016). The instability and inactivity of TA are a cause of 161 diseases such as dyskeratosis congenita (Zinsser-Cole-Engman syndrome), a telomeropathy 162 leading to premature ageing (Drachtman and Alter, 1992; Vulliamy et al., 2001; Armanios 163 and Blackburn, 2012; Armanios et al., 2013). If TA limits the shortening of TLs by adding 164 small repeated segments of DNA at the ends of chromosomes at each cell division, their 165 overexpression can be detrimental to individuals by causing diseases, such as cancer. In 80% 166 to 90% of all cancers, cancer cells often have high TA activity (Jafri et al., 2016) due to mutations in the core promoter region of TERT (Horn et al., 2013; Grüner et al., 2018; Liu et 167 168 al., 2018) which is considered a key factor for tumour development.

169

170

3.2. Roles of alternative lengthening telomeres (ALT)

171 The ALT pathway, an alternative pathway used to synthetize TLs, has been reported in 172 immortal human cell lines (Bryan et al., 1995; Bryan et al., 1997). The ALT pathway can be 173 activated after epigenetic aberrations and mutations in genes encoding proteins involved in 174 chromatin remodelling at TLs and other genomic sites or tumour suppressors, such as p53 175 (Chung et al., 2012; Cesare and Reddel, 2013; Conomos et al., 2013; Naderlinger and 176 Holzmann, 2017). There are various mechanisms involved in the ALT pathway (e.g., TL 177 elongation performed by homologous recombination) (Natarajan and McEachern, 2002; 178 Natarajan et al., 2003; Topcu et al., 2005; Cesare and Reddel, 2010; Cesare and Reddel, 2013; 179 Conomos et al., 2013). Some proteins are associated with the ALT pathway and are involved 180 in the restart of telomeric DNA replication, such as HR protein complexes (MRN, MRE11, 181 RAD50, NSB1, NBN), proteins located on chromosome 5 for the structural maintenance of 182 TLs (SMC5, SMC6), Fanconi anaemia proteins (FANDCD2, FANCA) and endonucleases 183 (FEN1) (Cesare and Reddel, 2010). The ALT pathway was evidenced in human somatic cells 184 (especially those that lose tumour protein 53 (TP53), which are genetically unstable) and 185 promotes proliferative cancer cells in humans (15% of cancers) (Henson et al., 2002; Cesare 186 et al., 2009; Lovejoy et al., 2012; Conomos et al., 2013; Neumann et al., 2013; Roake and 187 Artendi, 2016; Chen et al., 2017; Min et al., 2017).

188

3.3. DNA repair of telomeres

The maintenance of functional TLs is critical for genome stability. When damages affect TL extremities, TA or the ALT pathway is recruited for maintenance. However, damage (*e.g.*, breaks, oxidation of nucleotides, alkylations) can also take place at various locations in the TL sequence. In this case, other mechanisms, *i.e.*, DNA repair pathways are mobilized. These

193 pathways are the same as those that watch over the rest of the genome, such as base excision 194 repair (BER), nucleotide excision repair (NER), homologous recombination (HR), non-195 homologous end-joining (NHEJ) and mismatch repair (MMR) (Webb et al., 2013; Jia et al., 196 2015). The DNA BER pathway involves endonucleases (such as APE1) and is a major way to 197 protect genomes from oxidative stress in association with the shelterin complex through, for 198 instance, interaction with TRF2 to stabilize telomeric DNA (Madlner et al., 2013). Moreover, 199 DNA repair mechanisms are also essential when damage occurs in genes encoding proteins 200 involved in TL dynamic (e.g., TERT, POT1, MRN).

201

202 4. <u>Causes and consequences of telomere attrition</u>

203 <u>4.1. Ageing</u>

204 Several causes of TL shortening have been identified (*Figure 2*). The main one, ageing is 205 natural and is due to cell division (Blackburn, 2005), because at each division, chromosomes 206 lose telomeric patterns. TLs have a critical minimal size that, when reached, induces an 207 inhibition of cell division cycles and cellular senescence processes (Harley et al., 1990; 208 Blackburn, 1991; Amanions and Blackburn, 2012; Reichert et al., 2017). In various animal 209 groups, it has been reported that TLs shorten with ageing (Table 3). The relation between 210 ageing and shortened TLs has been evidenced mainly for vertebrates (i.e., mammals, fish and 211 birds), for example in cats (McKevitt et al., 2003) and rodents (Coviello-McLaughlin et al., 212 1997). However, for some vertebrates (e.g., long-lived species like the bird, Oceanodroma 213 *leucorhoa*, or the water python, *Liasis fuscus*), no relationship between the age of individuals 214 and TL length has been highlighted, probably due to reduced oxidative damage (Haussman et 215 al., 2003) or high TA activity stimulated by hormones notably estrogens (Ujvari and Masden, 216 2009). Table 3 shows a lack of knowledge on TL dynamic as a function of age in 217 invertebrates and the mechanism of ageing. Based on the few data available with a sufficient 218 number of individuals by age group, three patterns can be identified regarding relationships 219 between TLs, TA and ageing: increase, decrease and no variation. In the Baikal gastropod 220 Kobeltocochlea martensiana, an enlargement of TLs with ageing has been demonstrated 221 (Maximova et al., 2017). Studies on the planaria Schmidtea mediterranea, which has an 222 apparently limitless regenerative capacity, showed that TLs decrease with age in sexual 223 animals, while asexual planaria can maintain TL length during fission or regeneration (Tan et 224 al., 2012). In the aquatic arthropod *Daphnia sp.*, no relationship between age and TL length 225 was found in D. pulex while D. pulicaria, whose life cycle is twice as long as that of D. pulex, 226 showed TL attrition with ageing (Schumpert et al., 2015). In both species, TL dynamic 227 appears to be correlated with TA activity. These different patterns underline that phylogenetic 228 proximity does not mean similar patterns between TLs or TA responses and ageing. For the 229 nematode Caenorhabditis elegans (which can be described as a short-lived species with a 230 temperature-dependent lifespan between 5 and 100 days), no correlation between ageing and 231 TL length was detected (Raices et al., 2005). The same observation was made for the long-232 lived arthropod, Homarus americanus (lifespan from 50 and 100 years), which was 233 unexpected regarding its special longevity in an optimal environment. Ageing phenomena are 234 absent in this species, and mortality is often caused by the increased weight of individuals that 235 preclude moult (Klapper et al., 1998; Krupp et al., 2000).

In humans (Aubert and Lansdorp, 2008) and mice (Rudolph et al., 1999; Tomas-Loba et al., 2008), TL attrition is considered one of the nine hallmarks of ageing (Lopez-Otin et al., 2013) and is a marker of cellular senescence (Bernadotte et al., 2016). A relationship between the shortening of TLs in human leukocytes and all causes of death has been demonstrated (Boonekamp et al., 2013; Opresko et al., 2017). However, according to Farzaneh-Far et al. (2008), death cannot be predicted with a simple measure of TL length, and depends also on disease advancements or genotype. For example, it has been found that in humans, the
heterozygous genotype for some genes, including SERPINE, is associated with a longer TL
length, a longer lifespan (7 or more years on average), better cardiovascular capabilities,
lower fasting insulin levels, and a lower incidence of diabetes mellitus (Khan et al., 2017).

Hence, although a correlation between TL attrition, decrease in TA activity and ageing hasbeen often reported, it does not seem to be generalizable throughout the animal kingdom.

248 **4.2. Other causes**

Various mechanisms (e.g., genotoxicity, oxidative stress) can impact TL length (Figure 249 250 2). DNA damage such as a single strand break (SSB) or a double strand break (DSB) can be a 251 source of TL attrition (Pottier et al., 2013; Grach, 2013), notably when the DNA repair 252 machinery is inefficient (Webb et al., 2013; Jia et al., 2015). Deficiencies in enzymes 253 supported by the BER pathway (e.g., OGG1 and UNG) can contribute to TL shortening by 254 altered TL sister chromatid exchanges, an increased number of SSBs and DSBs, preferential 255 TL G-strand losses, accumulation of uracil on TLs, increased TL fragility and aberrant TL 256 recombination (Oikawa and Kawanishi, 1999; Wang et al., 2010; Vallabhaneni et al., 2013; 257 Webb et al., 2013; An et al., 2015; Jia et al., 2015; Vallabhaneni et al., 2015). Telomeric DNA 258 is highly susceptible to oxidative-induced DNA damage because of its high guanine content 259 (Kawanishi and Oikawa, 2004). Among oxidative lesion bases at TLs, 8-oxo-dG is the most 260 frequent (Zhou et al., 2017). Therefore, TLs can also be indirectly altered by mutagenicity 261 when reactive oxygen species (ROS) induce mutations (e.g., 8-oxo-dG induces a transversion 262 G:C to A:T) in genes coding for TL structure and maintenance or the DNA repair of TLs 263 (Wood et al., 1990; Henle et al., 1999; von Zglinicki, 2002; Chang et al., 2003; Vasseur and 264 Cossu-Leguille, 2003; Houben et al., 2008; Coluzzi et al., 2017; Stauffer et al., 2018). A disorder in protein recruitment in the TL complex shelterin can also be the cause of TL 265

attrition because the T-loop cannot be easily formed, making the end of the chromosome unprotected. This phenomenon can also be at the origin of apoptosis or senescence via activation of the p53 or p16link4a pathways (Oeseburg et al., 2010). In addition, the absence or deficiency of TA may also be a cause of premature TL shortening (Ziegler et al., 2017).

270 TL shortening in human populations is related to psychological stress, diseases, infections, 271 diets and lifestyles including smoking, as reported for example by Valdes et al. (2005), 272 Lansdorp (2009), Mirabello et al. (2009), and Gomes et al. (2012). Similar studies have been performed on mammals (Illmonen et al., 2008), showing that repeated infection by 273 274 Salmonella enterica can cause TL attrition in house mice (Mus musculus). Many biotic and 275 abiotic factors, such as nutrient (e.g., folate and selenium) deprivation, can accelerate TL 276 attrition. Indeed, the antioxidant activity of these nutrients has a protective effect against oxidative DNA telomeric damage (Liu et al., 2004; Boccardi et al., 2013). Among other 277 278 factors that can influence TL dynamic, food availability, number of breeding events and 279 antioxidant capabilities have been described in birds (Hall et al., 2004; Foote et al., 2011; 280 Heidinger et al., 2012; Bebbington et al., 2016; Sletten et al., 2016; Angelier et al., 2018).

281

4.3. Consequences of TL attrition on organism and population

282 Telomere shortening and short TL length is associated with various diseases including 283 diabetes (Sampson et al., 2006; Uziel et al., 2007), cardiovascular diseases (Fitzpatrick et al., 284 2007), obesity (Valdes et al., 2005), cancers (Srinivas et al., 2019) and pathologies of the 285 female reproductive tract (Kalmbach et al., 2013). These studies focusing on humans support that TL length alteration can affect individual fitness in relation to a reduced lifespan reduced 286 287 for various animal species (Monaghan and Haussman, 2006). Interspecies comparisons 288 provide additional information to understand relations between TL dynamic and lifespan. Higher rates of TL loss are associated with a shorter lifespan when comparing different bird 289

(Tricola et al., 2018) or mammal species. For instance, in humans, the rate of TL loss is 20-40
bp/year for a lifespan of 75 years while, it is 360 bp/year in dogs with a lifespan of 12 years
(Unryn et al., 2015). Generally, in vertebrates, TL length is inversely correlated with the
lifespan of a species (Gomes et al., 2011).

294 If TLs length is predictive of the longevity of individuals of species as demonstrated for 295 Acrocephalus arundinaceus or Daphnia pulicaria (Asghar et al., 2015; Schumpert et al., 296 2015), an accelerated attrition of TLs within a population may cause a decrease of the mean 297 reproductive lifespan and of individual fitness, so reduce fertility rate, which should lead to 298 changes in population dynamics (Reichert, 2013). However, if theoretical links between TL 299 dynamic, individual life-history traits and population dynamic seems obvious, the published 300 literature dealing with the ecological repercussion of accelerated TL shortening remains very 301 limited.

302 5. Contaminants and effects on telomere dynamic

303 Besides ageing and the other factors influencing TL dynamic we listed above, effects of 304 toxic contaminants released by anthropogenic activities on individual health, life-history traits 305 and population dynamic have been widely reported in the scientific literature. The interactions 306 between a toxic compound and a biological system occurred first at the molecular and cellular 307 levels. Then, ecological processes may also act, for instance when a predator suffers a 308 decrease in prey availability because of poisoning. According to the mechanism(s) of action 309 and depending on the sensitivity of the exposed species, this usually involves that the 310 compound reaches one (or several) molecular target(s) and/or induces oxidative stress to 311 express its toxicity at higher levels of biological organization. As reported above, TL dynamic 312 is related to life-history traits and potentially to population dynamic. Thus, it can be assume 313 that changes in TL dynamic is among the key processes linking toxic compounds,

314 molecular/cellular disturbances and consequences on individual health, fitness and315 population.

316

5.1. In humans

Currently, most of the studies questioning the impact of contaminants on TL length and TA activity concern humans, especially leukocytes. The general trend is that TL length and TA activity decrease concomitantly when cells or humans are exposed to contaminants (*Table 4*). This premise fits well with the commonly proposed hypothesis that exposure to pollutants causes cell damage together with a decrease in TL length and TA activity.

323 Exposure to metallic elements (ME) has been related to TL length changes, two patterns 324 can be distinguished: decrease or increase (Table 4). For example, cadmium (Cd) can cause 325 TL attrition, as shown in placenta from newborns (Lin et al., 2013), in saliva from adolescents 326 (Fillman et al., 2016) and in leukocytes from adults (Zota et al., 2015; Patel et al., 2017). 327 Similar trends were reported for lead (Pb) (Pottier et al., 2013; Pawlas et al., 2015; Pawlas et al., 2016), arsenic (As) (Ferrario et al., 2009; Srinivas et al., 2019) and the combination of As, 328 329 Cd and/or tungsten (W) (Wai et al., 2018; Grau-Perez et al., 2019). Organic chemicals, such 330 aromatic hydrocarbons (PAH), polychlorinated biphenyls (PCB), polycyclic as polybrominated diphenyl ethers (PBDE) and air pollutants (e.g., benzene, toluene and fuel 331 332 smoke) can lead to TL attrition (Table 4). Some PCBs and PBDEs induced a decrease in TL 333 length, TA activity and the expression of fundamental genes implicated in TL maintenance 334 and stability (Xin et al., 2016; Ziegler et al., 2017; Yuan et al., 2018).

Toxicologists have also paid special attention to carcinogenic compounds such as ME (*e.g.*, As), some persistent organic pollutant (POP) or air contaminants (NO_x, NO₂, PM_{2.5} and 337 PM₁₀) (Dioni et al., 2011; Gao et al., 2015; Hou et al., 2015; Joyce and Hou, 2015; Mitro et al., 2016; Jimenez Villarreal et al., 2018; Zhao et al., 2018). A study conducted by Gao et al. 338 339 (2015) on leukocytes from 167 participants demonstrated that a higher As urinary 340 concentration was associated with increased TL length, overexpression of WRN (a gene 341 involved in the ALT pathway), and decreased expression of TERF2, TERF2IP (shelterin 342 complex), DKC1 (upregulation of TA), and OBFC1 (upregulation of TA). A similar trend 343 was reported on leukocytes that presented an elongation of TL and an increase of the level of 344 two proteins of the shelterin complex (TRF1 and TRF2) following As exposure (Chatterjee et 345 al. 2015). As previously stated, abnormal TL elongation should not necessarily be considered 346 as positive, since strong activities of TA or ALT can be related to tumour development. In the 347 study of Gao et al. (2015), the WRN gene, implicated in ALT pathway, was up-regulated in 348 leukocytes and TL were lengthened after exposure to As. Moreover, it was demonstrated that 349 methylation of CHK1 or PML genes may activate ALT pathway in human or mouse 350 (Naderlinger and Holzmann, 2017), especially after exposure by PCBs and PCDEs (Yuan et 351 al., 2018). Therefore, for carcinogenic contaminants, a possible mechanism of action could be 352 an increase in TL length caused by TA and/or ALT upregulation rather than TL attrition 353 (Finnon et al., 2000; Jafri et al., 2016; Dagg et al., 2017). A pathway explaining an 354 overactivation of the ALT pathway could be epigenetic alterations, such as methylations, 355 induced by some contaminants (Reichard and Puga, 2010; Nica et al., 2017).

356

5.2. In animals

There is a growing interest in ecology to measure TLs in various organisms, including vertebrates and invertebrates, with approximately five publications in 1995 vs one hundred in 2015 (Angelier et al., 2018). However, in ecotoxicology, studies are limited to vertebrates and mainly birds (*Table 5*), but no studies on invertebrates has been published. In some of the few 361 studies that assessed with the impact of contaminants on TLs or TA in animals, TL attrition 362 was related to high exposure. Stauffer et al. (2017) showed inconsistent trends when 363 investigating the interaction between exposure to ME, oxidative stress and TL length in wild 364 populations of two birds (Parus major and Ficedula hypoleuca). They demonstrated that great 365 tit nestlings exposed to ME had shorter TLs than those living in unpolluted areas. However, 366 this correlation was not found when investigating pied flycatcher nestlings or adult females of 367 two species. The authors investigated whether the redox status that varied among species and 368 age groups may explain these differences but no link with TL length was found. For Rissa 369 tridactyla, an arctic seabird species, Blévin et al. (2016) showed that TL attrition was higher 370 in wild adult females exposed to the major metabolite of chlordane mixture (oxychlordane) 371 than in males. More expectedly, oxidative stress could accelerate TL attrition, as shown 372 experimentally in wild-caught house mice (*Mus musculus musculus*), for which an inhibition 373 of superoxide dismutase (SOD) activity was related to TL attrition (Stauffer et al., 2018). A 374 study on the fish Epinephelus coiodes showed a concomitant decrease in TA activity with 375 high exposure to benzo(a)pyrene (a PAH) (Peterson et al., 2015). As reported in Figure 2, 376 potential relationships between TL attrition and increases of DNA damages has been shown 377 on Ctenomys torquatus exposed to PAH and ME from coal (Matzenbacher et al., 2019). All 378 these studies show that TL length and TA activity can be altered by contaminants in various 379 vertebrate species, including birds, fishes and mammals.

However, other studies have not revealed that TL attrition is associated with chemical exposure. White-tailed eagle (*Haliaeetus albicilla*) nestlings sampled from a site contaminated by organohalogenated compounds did not exhibit TL shortening or reduced SOD activity (Sletten et al., 2015). Blévin et al. (2017) measured an increase in TL length in both sexes of *Rissa tridactyla* (Black-legged kittiwake) exposed to perfluoroalkyl substances (PFAs). 386 The impact of pollutants on TLs and TA is still a new field of research in environmental 387 toxicology; on the basis of the recent data published, it remains difficult to identify general 388 trends. This could be partly due to a lack of knowledge about the influence of various 389 individual (e.g., gender, age, species) and/or environmental (e.g., weather conditions, food 390 availability, pathogens, pollutions) factors on TL dynamic. In the following section, we 391 propose several lines of research to improve knowledge on the impact of pollutants on TLs 392 and TA in environmental toxicology for humans and animals (mainly those used as 393 bioindicators for ecotoxicological purposes).

394 6. <u>Perspectives for the assessment of telomere dynamic in environmental toxicology</u>

395 Environmental toxicology concerns the toxic effects of environmental contaminants on 396 living organisms in their environments. Gaining more knowledge on the impact of 397 contaminants on TL dynamic could be relevant to understand these effects and possibly to 398 propose adverse outcome pathways (AOP), which would impact life traits and population, as 399 recommended by the OECD (2018). This review showed that the level of knowledge on TLs, 400 in general and particularly in environmental toxicology, is quite different for humans, 401 vertebrates and invertebrates. Thus, perspectives for the possible use of TL length and TA 402 activity as biomarkers in environmental toxicology depend on the considered group; they will 403 be briefly developed for each of these groups of organisms, before moving to more specific 404 consideration (either technical or biological).

For humans, although TLs are already used as biomarkers (*Table 4*), further studies are necessary to better fulfil some specifications (*e.g.*, the link between TL variation and biological consequence or between TL lengthening and exposure to compounds with carcinogenic properties). For other vertebrates (*Table 5*), with the possible exception of birds, the use of TL length and TA activity as biomarkers appears far from ready in routine use for 410 ecotoxicological purposes. This review highlights a lack of knowledge on TL dynamic in 411 almost all bioindicator species: for the majority, the relation between ageing and TL length 412 remains to be characterized, as well as the influence of many endogenous and environmental 413 factors (acting possibly in interactions) on TL dynamic. This is also true for invertebrates, for 414 which this area of research is virtually untouched.

The basic question here is whether early TL shortening could be a biomarker to reveal the 415 416 impact of stress (physical, chemical and biological) on premature ageing and thus organism 417 health and longevity. For this purpose, TL length and TA activity must satisfy some 418 specifications: good sensitivity, knowledge about biological consequences for interpreting the 419 responses of the biomarker, feasibility, field application, knowledge about variations in the 420 amplitudes of responses, and the identification of confounding factors (i.e., capacities to discriminate variations that could be caused by pollutants or environmental factors) (Garric et 421 422 al., 2010). According to these specifications, many technical and biological issues must be 423 better considered for reliable TL length and TA activity assessments in environmental 424 toxicology:

The choice of analytical technique to assess TL length and TA activity. Currently, 425 426 many methods exist to measure TLs (qPCR, flow-fluorescence in situ 427 hybridization (FISH), qFISH, single TL length analysis (STELA), terminal 428 restriction fragment by southern blot or in-gel hybridization, dot blot) and detect 429 TA activity (ddPCR). The choice of methods depends on scientific questions and 430 technical considerations (see, for example, the reviews of Montpetit et al., 2014 431 and Nussey et al., 2014). For example, for the same sample, qPCR provides an 432 estimate of the average TL length in a DNA extract but the diversity of the TL size 433 for each cells cannot be determined. Molecular biology kits exist and have often used (*e.g.*, by Schumpert et al. (2015) to measure TA activity on two *Daphnia sp*).
For reasons related to quickness, simplicity, robustness and cost, the qPCR TL
measurement technique seems well appropriate and is widely used (Cawthon et al.,
2002; Shay and Wright, 2019), even if the results are semiquantitative (but
sufficient to identify variations between various exposures to contaminants).

- Influence of the quality of DNA extraction. In terms of measurement techniques
 and TL size, different kits can be preferred. It is important to use the same
 extraction method during all the studies (Nussey et al., 2014; Raschenberger et al.,
 2016).
- Influence of storage conditions of the sample that can impact measurements. For
 example, FTA cards (*Flinders Technology Associates cards*) are not recommended
 for long-term storage (Reichert et al., 2017) and repeated cycles of
 freezing/thawing can cause DNA alterations.
- 447 Influence of the choice of biological tissue or cell type (Gardner et al., 2007; ٠ 448 Kesäniemi et al., 2019) according to physiological characteristics (*i.e.*: TA 449 activity, ALT, cell division properties) and toxicological characteristics (i.e.: 450 target, storage or metabolism of xenobiotics) is important to be considered. For 451 example, Yip et al. (2017) chose to measure TLs on Medaka fish liver cells 452 because of the regulatory functions of metabolism and homeostasis. For small organisms (e.g., arthropods) the recourse of pools and whole body may be 453 454 relevant.
- Experiments must use of a sufficient number of individuals by age and/or
 environmental condition class to make reliable statistical comparisons to account
 for interindividual variations.

458 The link between TL length, TA activity, longevity and ageing needs to be ٠ 459 investigated in a wide range of animals, including invertebrates (notably those 460 commonly used in ecotoxicological risk assessment, e.g., crustacean, earthworms and mollusks, for which no data are available). Indeed, interspecific comparison to 461 462 estimate variation of TL length or TA activity with ageing for species for which 463 the dynamic of TLs is unknown is difficult because various patterns have been 464 demonstrated for phylogenetically close species (e.g., between various bird 465 species). For example, a relation between TL length and ageing for Daphnia pulicaria has been evidenced (Schumpert et al., 2015) and suggests that it could be 466 467 interesting to assess the sublethal effects of contaminants or other stressors on this 468 taxon with TL length measurements. However, this relation is not the same for 469 Daphnia pulex where TL length does not evolve with ageing. Understanding the 470 reason for the differences observed in phylogenetically close species, offers an 471 exciting challenge.

Many studies concerning the impact of contaminants on TL dynamic are
correlative. To better evaluate the relevance of TL dynamic studies in
environmental toxicology and to highlight by which mechanisms TL changes are
related to exposure to chemicals, it is essential to develop more laboratory
approaches. Some experiments conducted under controlled conditions should
address the following questions:

What is the influence of environmental and individual factors on TL
dynamic such as, gender, hormones, food source, crowding, disturbance,
noise, infection, pathogens, parasites, and climatic factors, as described by
Angelier et al. (2018)? For example, in female Medaka fish, oestradiol may
be involved in TL maintenance by i) increasing TA activity (TERT mRNA

483 upregulation, TERT phosphorylation) and ii) limiting oxidative DNA
484 damage at the level of TLs (antioxidant gene upregulation) (Yip et al.,
485 2017).

- 486 What are the consequences of TL attrition on survival or reproductive
 487 abilities and consequently on population dynamic?
- What is the pattern(s) of the relationship between TL length and
 contaminant exposure? This question is supported by the fact that some
 carcinogenic compounds could lead to an elongation of TLs and because
 dose-reponse relationships have not been established even for the most
 prevalent pollutants either single or in mixture. This should be done
 considering acute but also subchronic and chronic exposure scenario.
- What are the main mechanisms, including direct (DNA damages) and
 indirect ones (modulation of maintenance processes, oxidative stress...),
 linking chemical exposure and the alteration of the natural TL dynamic?
- 497

498 **7.** <u>Conclusion</u>

499 TLs are highly conserved in the animal kingdom, ubiquitous and appear "universal" in 500 living organisms. However, many studies are still needed, as noted in the present review, 501 before TLs can be used as a relevant biomarker (in species belonging to many biological 502 groups) to predict the effects of various stressors (e.g., chemical contaminants) on health for 503 humans and animal populations in the wild. More data on TL dynamic could help build new 504 AOP for chemical risk assessment based on mechanistic reasoning. TL dynamic studies could 505 improve environmental monitoring and our understanding of the impact of disrupted 506 anthropic ecosystems (on the health status, lifespans and dynamic of populations) and could reduce disciplinary borders in the One Health concept, because epidemiologists, 507

508 ecotoxicologists, veterinarians and economists look for universal information concerning an509 organism's health.

510 Acknowledgements

511 The authors thank the organizations that support them in their work: the University of

512 Bourgogne-Franche-Comté (UBFC) and the company TESORA. The authors warmly thank

513 the reviewers for their constructive remarks.

514 **Disclosures of interest**

515 The authors declare no disclosures of interest.

516 Authors

517 All authors reviewed and approved the final version of the manuscript.

518 <u>References</u>519

- An, N., Fleming, A.M., White, H.S., Burrows, C.J., 2015. Nanopore detection of 8oxoguanine in the human telomere repeat sequence. ACS Nano 9, 4296–4307.
 https://doi.org/10.1021/acsnano.5b00722
- Andreotti, G., Hoppin, J.A., Hou, L., Koutros, S., Gadalla, S.M., Savage, S.A., Lubin, J.,
 Blair, A., Hoxha, M., Baccarelli, A., Sandler, D., Alavanja, M., Freeman, L.E.B., 2015.
 Pesticide use and relative leukocyte telomere length in the agricultural health study. PLoS
 ONE 10, e0133382. <u>https://doi.org/10.1371/journal.pone.0133382</u>
- Angelier, F., Costantini, D., Blévin, P., Chastel, O., 2018. Do glucocorticoids mediate the link
 between environmental conditions and telomere dynamics in wild vertebrates? A review.
 Gen. Comp. Endocrinol. 256, 99–111. https://doi.org/10.1016/j.ygcen.2017.07.007
- Armanios, M., 2013. Telomeres and age-related disease: how telomere biology informs
 clinical paradigms. J Clin Invest 123, 996–1002. <u>https://doi.org/10.1172/JCI66370</u>
- Armanios, M., Blackburn, E.H., 2012. The telomere syndromes. Nat Rev Genet 13, 693–704.
 <u>https://doi.org/10.1038/nrg3246</u>
- Asghar, M., Bensch, S., Tarka, M., Hannsson, B., Hasselquit, D., 2015. Maternal and genetic
 factors determine early life telomere length. Proc. R. Soc. B. 282.
 https://doi.org/10.1098/rspb.2014.2263
- 537 Aubert, G., Lansdorp, P.M., 2008. Telomeres and aging. Physiol. Rev. 88, 557–579.
 538 <u>https://doi.org/10.1152/physrev.00026.2007</u>
- Bassig, B.A., Zhang, L., Cawthon, R.M., Smith, M.T., Yin, S., Li, G., Hu, W., Shen, M.,
 Rappaport, S., Barone-Adesi, F., Rothman, N., Vermeulen, R., Lan, Q., 2014. Alterations
 in leukocyte telomere length in workers occupationally exposed to benzene. Environ Mol
 Mutagen 55, 673-678. https://doi.org/10.1002/em.21880

- Bebbington, K., Spurgin, L.G., Fairfiled, E.A., Dugdale, H.L., Komdeur, J., Burke, T.,
 Richardon, D.S., 2016. Telomere length reveals cumulative individual and
 transgenerational inbreeding effects in a passerine bird. Mol. Ecol. 25, 2949-2960.
 https://doi.org/10.1111/mec.13670
- 547 Benetos, A., Okuda, K., Lajemi, M., Kimura, M., Thomas, F., Skurnick, J., Labat, C., Bean,
 548 K., Aviv, A., 2001. Telomere length as an indicator of biological aging: the gender effect
 549 and relation with pulse pressure and pulse wave velocity. Hypertension 37, 381–385.
- Bernadotte, A., Mikhelson, M.V., Spivak, M.I., 2016. Markers of cellular senescence.
 Telomere shortening as a marker of cellular senescence. Aging 8(1), 3-11.
 <u>https://doi.org/10.18632/aging.100871</u>
- Bertuch, A.A., 2016. The molecular genetics of the telomere biology disorders. RNA Biol 13,
 696–706. <u>https://doi.org/10.1080/15476286.2015.1094596</u>
- Bijnens, E.M., Zeegers, M.P., Derom, C., Martens, D.S., Gielen, M., Hageman, G.J., 555 Plusquin, M., Thiery, E., Vlietinck, R., Nawrot, T.S., 2017. Telomere tracking from birth 556 557 adulthood and residential traffic exposure. Med 15. BMC to 558 https://doi.org/10.1186/s12916-017-0964-8
- 559 Blackburn, E.H., 1991. Structure and function of telomeres. Nature 350, 569–573.
 <u>https://doi.org/10.1038/350569a0</u>
- Blackburn, E.H., 2005. Telomeres and telomerase: their mechanisms of action and the effects
 of altering their functions. FEBS Lett. 579, 859–862.
 https://doi.org/10.1016/j.febslet.2004.11.036
- Blasco, M.A., 2005. Telomeres and human disease: ageing, cancer and beyond. Nat. Rev.
 Genet. 6, 611–622. <u>https://doi.org/10.1038/nrg1656</u>
- Blévin, P., Angelier, F., Tartu, S., Bustamante, P., Herzke, D., Moe, B., Bech, C., Gabrielsen,
 G.W., Bustnes, J.O., Chastel, O., 2017. Perfluorinated substances and telomeres in an
 Arctic seabird: Cross-sectional and longitudinal approaches. Environ. Pollut. 230, 360–
 367. <u>https://doi.org/10.1016/j.envpol.2017.06.060</u>
- Blévin, P., Angelier, F., Tartu, S., Ruault, S., Bustamante, P., Herzke, D., Moe, B., Bech, C.,
 Gabrielsen, G.W., Bustnes, J.O., Chastel, O., 2016. Exposure to oxychlordane is
 associated with shorter telomeres in arctic breeding kittiwakes. Sci. Total Environ. 563–
 564, 125–130. <u>https://doi.org/10.1016/j.scitotenv.2016.04.096</u>
- Boccardi, V., Esposito, A., Rizzo, M.R., Marfella, R., Barbieri, M., Paolisso, G., 2013.
 Mediterranean diet, telomere maintenance and health status among elderly. PLoS ONE 8, e62781. <u>https://doi.org/10.1371/journal.pone.0062781</u>
- Boonekamp, J.J., Simons, M.J.P., Hemerik, L., Verhulst, S., 2013. Telomere length behaves
 as biomarker of somatic redundancy rather than biological age. Aging Cell 12, 330–332.
 <u>https://doi.org/10.1111/acel.12050</u>
- Bryan, T.M., Englezou, A., Dalla-Pozza, L., Dunham, M.A., Reddel, R.R., 1997. Evidence
 for an alternative mechanism for maintaining telomere length in human tumors and
 tumor-derived cell lines. Nat. Med. 3, 1271–1274.
- Bryan, T.M., Englezou, A., Gupta, J., Bacchetti, S., Reddel, R.R., 1995. Telomere elongation
 in immortal human cells without detectable telomerase activity. EMBO J. 14, 4240–4248.
- 585 Cawthon, R.M., 2002. Telomere measurement by quantitative PCR. Nucleic Acids Res 30,
 586 e47.
- 587 Cesare, A.J., Kaul, Z., Cohen, S.B., Napier, C.E., Pickett, H.A., Neumann, A.A., Reddel, 588 R.R., 2009. Spontaneous occurrence of telomeric DNA damage response in the absence 589 of chromosome fusions. Nat. Struct. Mol. Biol. 16, 1244-1251. 590 https://doi.org/10.1038/nsmb.1725
- 591 Cesare, A.J., Reddel, R.R., 2010. Alternative lengthening of telomeres: models, mechanisms
 592 and implications. Nat. Rev. Genet. 11, 319–330. <u>https://doi.org/10.1038/nrg2763</u>

- 593 Cesare, A.J., Reddel, R.R., 2013. Alternative Lengthening of Telomeres in Mammalian Cells.
 594 Landes Bioscience.
- Chang, S., Khoo, C.M., Naylor, M.L., Maser, R.S., DePinho, R.A., 2003. Telomere-based
 crisis: functional differences between telomerase activation and ALT in tumor
 progression. Genes Dev 17, 88–100. https://doi.org/10.1101/gad.1029903
- Chatterjee, D., Bhattacharjee, P., Sau, T.J., Das, J.K., Sarma, N., Bandyopadhyay, A.K., Roy,
 S.S., Giri, A.K., 2015. Arsenic exposure through drinking water leads to senescence and
 alteration of telomere length in humans: A case-control study in West Bengal, India. Mol.
 Carcinog. 54(9), 800-809. https://doi.org/10.1002/mc.22150
- 602 Chatterjee, D., Adak, S., Banerjee, N., Bhattacharjee, P., Bandyopadhyay, A.K., Giri, A.K.,
 603 2018. Evaluation of health effects, genetic damage and telomere length in children
 604 exposed to arsenic in West Bengal, India. Mut Res Genet Toxicol Environ Mutagen 836,
 605 Part A, 82-88. https://doi.org/10.1016/j.mrgentox.2018.06.012
- 606 Chen, Y., Ebenstein, A., Greenstone, M., Li, H., 2013. Evidence on the impact of sustained
 607 exposure to air pollution on life expectancy from China's Huai River policy. Proc. Natl.
 608 Acad. Sci. U.S.A., 110(32), 12936-12941. https://doi.org/10.1073/pnas.1300018110
- Chen, Y.-A., Shen, Y.-L., Hsia, H.-Y., Tiang, Y.-P., Sung, T.-L., Chen, L.-Y., 2017.
 Extrachromosomal telomere repeat DNA is linked to ALT development via cGASSTING DNA sensing pathway. Nat. Struct. Mol. Biol. 24, 1124–1131.
 https://doi.org/10.1038/nsmb.3498
- 613 Chung, I., Osterwald, S., Deeg, K.-I., Rippe, K., 2012. PML body meets telomere. The
 614 beginning of an ALTernate ending? Nucleus 3:3, 263-275.
 615 https://doi.org/10.4161/nucl.20326
- 616 Cicconi, A., Micheli, E., Verni, F., Jackson, A., Gradilla, A.C., Cipressa, F., Raimondo, D.,
 617 Bosso, G., Wakefield, J.G., Ciapponi, L., Cenci, G., Gatti, M., Cacchione, S., Raffa,
 618 G.D., 2017. The *Drosophila* telomere-capping protein Verrocchio binds single-stranded
 619 DNA and protects telomeres from DNA damage response. Nucleic Acids Res. 45(6),
 620 1068-3085. https://doi.org/10.1093/nar/gkw1244
- 621 Coluzzi, E., Buonsante, R., Leone, S., Asmar, A.J., Miller, K.L., Cimini, D., Sgura, A., 2017. 622 Transient ALT activation protects human primary cells from chromosome instability 623 induced low chronic oxidative stress. Sci Rep. 43309. by 7, https://doi.org/10.1038/srep43309 624
- Conomos, D., Pickett, H.A., Reddel, R.R., 2013. Alternative lengthening of telomeres:
 remodeling the telomere architecture. Front Oncol 3, 27.
 <u>https://doi.org/10.3389/fonc.2013.00027</u>
- 628 Coviello-McLaughlin, G.M., Prowse, K.R., 1997. Telomere length regulation during postnatal
 629 development and ageing in *Mus spretus*. Nucleic Acids Res. 25, 3051–3058.
- Dagg, R.A., Pickett, H.A., Neumann, A.A., Napier, C.E., Henson, J.D., Teber, E.T., Arthur,
 J.W., Reynolds, C.P., Murray, J., Haber, M., Sobinoff, A.P., Lau, L.M.S., Reddel, R.R.,
 2017. Extensive proliferation of human cancer cells with ever-shorter telomeres. Cell
 Rep. 19(12), 2544-2556. https://doi.org/10.1016/j.celrep.2017.05.087
- de Felice, B., Nappi, C., Zizolfi, B., Guida, Marco, Di Spiezio Sardo, A., Bifulco, G., Guida,
 Maurizio, 2012. Telomere shortening in women resident close to waste landfill sites.
 Gene 500, 101–106. <u>https://doi.org/10.1016/j.gene.2012.03.040</u>
- de Lange, T., 2005. Telomere-related genome instability in cancer. Cold Spring Harb. Symp.
 Quant. Biol. 70, 197–204. <u>https://doi.org/10.1101/sqb.2005.70.032</u>
- de Lange, T., 2009. How telomeres solve the end-protection problem. Science 326, 948–952.
 <u>https://doi.org/10.1126/science.1170633</u>
- 641 Destoumieux-Garzón, D., Mavingui, P., Boetsch, G., Boissier, J., Darriet, F., Duboz, P.,
 642 Fritsch, C., Giraudoux, P., Le Roux, F., Morand, S., Paillard, C., Pontier, D., Sueur, C.,

- 643 Voituron, Y., 2018. The One Health concept: 10 years old and a long road ahead. Front.
 644 Vet. Sci. 5. <u>https://doi.org/10.3389/fvets.2018.00014</u>
- Dioni, L., Hoxha, M., Nordio, F., Bonzini, M., Tarantini, L., Albetti, B., Savarese, A.,
 Schwartz, J., Bertazzi, P.A., Apostoli, P., Hou, L., Baccarelli, A., 2011. Effects of shortterm exposure to inhalable particular matter on telomere length, telomerase expression,
 and telomerase methylation in steel workers. Environ Health Perspect 119(5).
 https://doi.org/10.1289/ehp.1002486
- Drachtman, R.A., Alter, B.P., 1992. Dyskeratosis congenita: clinical and genetic
 heterogeneity. Report of a new case and review of the literature. Am J Pediatr Hematol
 Oncol 14, 297–304.
- Farzaneh-Far, R., Cawthon, R.M., Na, B., Browner, W.S., Schiller, N.B., Whooley, M.A.,
 2008. Prognostic value of leukocyte telomere length in patients with stable coronary
 artery disease: data from the Heart and Soul Study. Arterioscler. Thromb. Vasc. Biol. 28,
 1379–1384. https://doi.org/10.1161/ATVBAHA.108.167049
- Ferreira, H.C., Towbin, B.D., Jegou, T., Gasser, S.M., 2013. The shelterin protein POT-1
 anchors *Caenorhabditis elegans* telomeres through SUN-1 at the nuclear periphery. J.
 Cell Biol 203 (5), 727-735. https://doi.org/10.1083/jcb.201307181
- Ferrario, D., Collotta, A., Carfi, M., Bowe, G., Vahter, M., Hartung, T., Gribaldo, L., 2009.
 Arsenic induces telomerase expression and maintains telomere length in human cord
 blood cells. Toxicology 260, 132–141. https://doi.org/10.1016/j.tox.2009.03.019
- Fillman, T., Shimizu-Furusawa, H., Ng, C.F.S., Parajuli, R.P., Watanabe, C., 2016.
 Association of cadmium and arsenic exposure with salivary telomere length in adolescents in Terai, Nepal. Environ. Res. 149, 8–14.
 https://doi.org/10.1016/j.envres.2016.04.037
- Finnon, P., Silver, A.R.J., Bouffler, S.D., 2000. Upregulation of telomerase activity by Xirradiation in mouse leukaemia cell is independent of Tert, Terc, Tnks and Myc
 transcription. Carcinogenesis 21(4), 573-578. https://doi.org/10.1093/carcin/21.4.573
- Fitzpatrick, A.L., Kronmal, R.A., Gardner, J.P., Psaty, B.M., Jenny, N.S., Tracy, R.P.,
 Walston, J., Kimura, M., Aviv, A., 2007. Leukocyte telomere length and cardiovascular
 disease in the cardiovascular health study. Am. J. Epidemiol. 165(1), 14-21.
 https://doi.org/10.1093/aje/kwj346
- Foote, C.G., Daunt, F., González-Solís, J., Nasir, L., Phillips, R.A., Monaghan, P., 2011.
 Individual state and survival prospects: age, sex, and telomere length in a long-lived
 seabird. Behav Ecol 22, 156–161. <u>https://doi.org/10.1093/beheco/arq178</u>
- Francis, N., Gregg, T., Owen, R., Ebert, T., Bodnar, A., 2006. Lack of age-associated
 telomere shortening in long- and short-lived species of sea urchins. FEBS Letters 580,
 4713–4717. https://doi.org/10.1016/j.febslet.2006.07.049
- Frenck, R.W., Blackburn, E.H., Shannon, K.M., 1998. The rate of telomere sequence loss in
 human leukocytes varies with age. Proc Natl Acad Sci U S A 95, 5607–5610.
- Frydrychová, R., Grossmann, P., Trubac, P., Vítková, M., Marec, F., 2004. Phylogenetic
 distribution of TTAGG telomeric repeats in insects. Genome 47, 163–178.
 <u>https://doi.org/10.1139/g03-100</u>
- Fujiwara, H., Osanai, M., Matsumoto, T., Kojima, K.K., 2005. Telomere-specific non-LTR
 retrotransposons and telomere maintenance in the silkworm, *Bombyx mori*. Chromosome
 Res 13, 455-467. https://doi.org/10.1007/s10577-005-0990-9
- Gardner, J.P., Kimura, M., Chai, W., Durrani, J.F., Tchakmakjian, L., Cao, X., Lu, X., Li, G.,
 Peppas, A.P., Skurnick, J., Wright, W.E., Shay, J.W., Aviv, A., 2007. Telomere
 Dynamics in Macaques and Humans. J Gerontol A Biol Sci Med Sci 62, 367–374.
 <u>https://doi.org/10.1093/gerona/62.4.367</u>

- George, J.A., DeBaryshe, P.G., Traverse, K.L., Celniker, S.E., Pardue, M.L., 2006. Genomic
 organization of the *Drosophila* telomere retrotransposable elements. Genome Res. 16,
 1231-1240. https://doi.org/10.1101/gr.5348806
- Gomes, N.M.V., Ryder, O.A., Houck, M.L., Charter, S.J., Walker, W., Forsyth, N.R., Austad,
 S.N., Venditt, C., Pagel, M., Shay, J.W., Wright, W.E., 2011. Comparative biology of
 mammalian telomeres: hypotheses on ancestral states and the roles of telomeres in
 longevity determination. Aging Cell 10, 761–768. <u>https://doi.org/10.1111/j.1474-</u>
 <u>9726.2011.00718.x</u>
- Gomes, N.M.V., Shay, J.W., Wright, W.E., 2010. Telomere biology in Metazoa. FEBS Lett.
 584, 3741–3751. <u>https://doi.org/10.1016/j.febslet.2010.07.031</u>
- Gopalakrishnan, S., Cheung, N.K., Yip, B.W., Au, D.W., 2013. Medaka fish exhibits
 longevity gender gap, a natural drop in estrogen and telomere shortening during aging: a
 unique model for studying sex-dependent longevity. Front. Zool. 10, 78.
 https://doi.org/10.1186/1742-9994-10-78
- Grach, A., 2013. Telomere Shortening Mechanisms. The Mechanisms of DNA Replication.
 <u>https://doi.org/10.5772/55244</u>
- Grau-Perez, M., Zhao, J., Pierce, B., Francesconi, K.A., Goessler, W., Zhu, Y., Umans, J., 708 709 Best, L., Cole, S.A., Navas-Acien, A., Tellez-Plaza, M., 2019. Urinary metals and 710 leukocyte telomere length in American Indian communities: The strong heart and the 711 strong heart family study. Environ Pollut. 246, 311-318. 712 https://doi.org/10.1016/j.envpol.2018.12.010
- Gao, J., Roy, S., Tong, L., Argos, M., Jasmine, F., Rahaman, R., Rakibuz-Zaman, M., Parvez,
 F., Ahmed, A., Hore, S.K., Sarwar, G., Slavkovich, V., Yunus, M., Rahman, M., Baron,
 J.A., Graziano, J.H., Ahsan, H., Pierce, B.L., 2015. Arsenic exposure, telomere length,
 and expression of telomere-related genes among Bangladeshi individuals. Env. Research.
 136:462-469. https://doi.org/10.1016/j.envres.2014.09.040
- Garric, J., Morin, S., Vincent-Hubert, 2010. Les biomarqueurs en écotoxicologie : définition,
 intérêt, limite, usage. Revue Sciences Eaux & Territoires, 1, 12-19.
 https://doi.org/10.14758/SET-REVUE.2010.1.03
- Greider, C.W., Blackburn, E.H., 1989. A telomeric sequence in the RNA of Tetrahymena
 telomerase required for telomere repeat synthesis. Nature 337, 331–337.
 <u>https://doi.org/10.1038/337331a0</u>
- 724 Greider, C.W., 1991. Chromosome first aid. Cell 67, 645–647.
- 725 Greider, C.W., 1999. Telomeres do D-loop-T-loop. Cell 97, 419–422.
- Greider, C.W., Blackburn, E.H., 1985. Identification of a specific telomere terminal
 transferase activity in Tetrahymena extracts. Cell 43, 405–413.
- Griffith, J.D., Comeau, L., Rosenfield, S., Stansel, R.M., Bianchi, A., Moss, H., de Lange, T.,
 1999. Mammalian telomeres end in a large duplex loop. Cell 97, 503–514.
 https://doi.org/10.1016/S0092-8674(00)80760-6
- Gruber, H., Schaible, R., Ridgway, I.D., Chow, T.T., Held, C., Philipp, E.E.R., 2014.
 Telomere-independent ageing in the longest-lived non-colonial animal, *Arctica islandica*.
 Exp. Gerontol. 51, 38–45. https://doi.org/10.1016/j.exger.2013.12.014
- Günes, C., Wezel, F., Southgate, J., Bolenz, C., 2018. Implications of TERT promoter
 mutations and telomerase activity in urothelial carcinogenesis. Nat Rev Urol 15, 386-393.
 https://doi.org/10.1038/s41585-018-0001-5
- Haber, J.E., 1999. Sir-Ku-itous routes to make ends meet. Cell 97, 829–832.
 https://doi.org/10.1016/S0092-8674(00)80795-3
- Hall, M.E., Nasir, L., Daunt, F., Gault, E.A., Croxall, J.P., Wanless, S., Monaghan, P., 2004.
 Telomere loss in relation to age and early environment in long-lived birds. Proc R Soc
 Lond [Biol] 271, 1571–1576. https://doi.org/10.1098/rspb.2004.2768

- Harley, C.B., Futcher, A.B., Greider, C.W., 1990. Telomeres shorten during ageing of human
 fibroblasts. Nature 345, 458–460. <u>https://doi.org/10.1038/345458a0</u>
- 744 Haussmann, M.F., Winkler, D.W., O'Reilly, K.M., Huntington, C.E., Nisbet, I.C.T., Vleck, 745 C.M., 2003. Telomeres shorten more slowly in long-lived birds and mammals than in 746 Soc Lond [Biol] short-lived ones. Proc R 270. 1387–1392. 747 https://doi.org/10.1098/rspb.2003.2385
- Hayflick, L., Moorhead, P.S., 1961. The serial cultivation of human diploid cell strains.
 Experimental Cell Res. 25, 585–621. <u>https://doi.org/10.1016/0014-4827(61)90192-6</u>
- Hayflick, L., 1965. The limited in vitro lifetime of human diploid cell strains. Experimental
 Cell Res. 37, 614–636. <u>https://doi.org/10.1016/0014-4827(65)90211-9</u>
- Heidinger, B.J., Blount, J.D., Boner, W., Griffiths, K., Metcalfe, N.B., Monaghan, P., 2012.
 Telomere length in early life predicts lifespan. Proc. Natl. Acad. Sci. U.S.A. 109, 1743–
 1748. <u>https://doi.org/10.1073/pnas.1113306109</u>
- Henle, E.S., Han, Z., Tang, N., Rai, P., Luo, Y., Linn, S., 1999. Sequence-specific DNA
 cleavage by Fe2+-mediated fenton reactions has possible biological implications. J. Biol.
 Chem. 274, 962–971.
- Henriques, C.M., Carneiro, M.C., Tenente, I.M., Jacinto, A., Ferreira, M.G., 2017.
 Correction: telomerase is required for zebrafish lifespan. PLoS Genet. 13, e1006652.
 https://doi.org/10.1371/journal.pgen.1006652
- Henriques, C.M., Carneiro, M.C., Tenente, I.M., Jacinto, A., Ferreira, M.G., 2013.
 Telomerase is required for zebrafish lifespan. PLoS Genet. 9, e1003214.
 <u>https://doi.org/10.1371/journal.pgen.1003214</u>
- Henson, J.D., Neumann, A.A., Yeager, T.R., Reddel, R.R., 2002. Alternative lengthening of
 telomeres in mammalian cells. Oncogene 21, 598–610.
 <u>https://doi.org/10.1038/sj.onc.1205058</u>
- 767 Horn, S., Figl, A., Rachakonda, P.S., Fischer, C., Sucker, A., Gast, A., Kadel, S., Moll, I., 768 Nagore, E., Hemminki, K., Schadendorf, D., Kumar, R., 2013. TERT promoter mutations 769 familial and sporadic melanoma. Science, 22, 339(6122), 959-961. in 770 https://doi.org/10.1126/science.1230062
- Hou, L., Wang, S., Dou, C., Zhang, X., Yu, Y., Zheng, Y., Avula, U., Hoxha, M., Díaz, A., 771 772 McCracken, J., Barretta, F., Marinelli, B., Bertazzi, P.A., Schwartz, J., Baccarelli, A.A., 773 2012. Air pollution exposure and telomere length in highly exposed subjects in Beijing, 774 repeated-measure China: А study. Environ Int 48. 775 https://doi.org/10.1016/j.envint.2012.06.020
- Houben, J.M.J., Moonen, H.J.J., van Schooten, F.J., Hageman, G.J., 2008. Telomere length
 assessment: biomarker of chronic oxidative stress? Free Radic. Biol. Med. 44, 235–246.
 https://doi.org/10.1016/j.freeradbiomed.2007.10.001
- Hoxha, M., Dioni, L., Bonzini, M., Pesatori, A.C., Fustinoni, S., Cavallo, D., Carugno, M.,
 Albetti, B., Marinelli, B., Schwartz, J., Bertazzi, P.A., Baccarelli, A., 2009. Association
 between leukocyte telomere shortening and exposure to traffic pollution: a cross-sectional
 study on traffic officers and indoor office workers. Environ Health 8, 41.
 <u>https://doi.org/10.1186/1476-069X-8-41</u>
- Huppert, J.L., Balasubramanian, S., 2005. Prevalence of quadruplexes in the human genome.
 Nucleic Acids Res. 33, 2908–2916. <u>https://doi.org/10.1093/nar/gki609</u>
- Ilmonen, P., Kotrschal, A., Penn, D.J., 2008. Telomere attrition due to infection. PLoS One 3,
 e2143. <u>https://doi.org/10.1371/journal.pone.0002143</u>
- Iwano, T., Tachibana, M., Reth, M., Shinkai, Y., 2004. Importance of TRF1 for functional telomere structure. J. Biol. Chem. 279, 1442–1448.
 https://doi.org/10.1074/jbc.M309138200

- Jafri, M.A., Ansari, S.A., Alqahtani, M.H., Shay, J.W., 2016. Roles of telomeres and telomerase in cancer, and advances in telomerase-targeted therapies. Genome Med 8, 69.
 <u>https://doi.org/10.1186/s13073-016-0324-x</u>
- Jia, P., Her, C., Chai, W., 2015. DNA excision repair at telomeres. DNA Repair (Amst.) 36,
 137–145. <u>https://doi.org/10.1016/j.dnarep.2015.09.017</u>
- Jimenez Villarreal, J., Ortiz, B.M., Garza, S.M., Armendariz, D.I.R., Villa, V.D.B., Rosales,
 P.C., Martinez, N.D.B., Aguirre, H.D., Martinez, J.M., 2018. Telomere length analysis in
 residents of a community exposed to arsenic. J Biochem Mol Toxicol 2018 Oct 6:e22230.
 https://doi.org/10.1002/jbt.22230
- 800Joyce, B.T., Hou, L., 2015. Organic pollutants and telomere length: A new facet of
EBioMedicine2,1854–1855.802https://doi.org/10.1016/j.ebiom.2015.11.0462,1854–1855.
- Kawanishi, S., Oikawa, S., 2004. Mechanism of telomere shortening by oxidative stress. Ann.
 N. Y. Acad. Sci. 1019, 278–284. <u>https://doi.org/10.1196/annals.1297.047</u>
- Kesäniemi, J., Lavrinienko, A., Tukalenko, E., Boratynski, Z., Kivisaari, K., Mappes, T.,
 Milinevsky, G., Moller, A.P., Mousseau, T.A., Watts, P.C., 2019. Exposure to
 environmental radionuclides associates with tissue-specific impacts on telomerase
 expression and telomere length. Sci Rep. 9, 850. https://doi.org/10.1038/s41598-01837164-8
- Khan, S.S., Shah, S.J., Klyachko, E., Baldridge, A.S., Eren, M., Place, A.T., Aviv, A.,
 Puterman, E., Lloyd-Jones, D.M., Heiman, M., Miyata, T., Gupta, S., Shapiro, A.D.,
 Vaughan, D.E., 2017. A null mutation in SERPINE1 protects against biological aging in
 humans. Sci Adv 3. https://doi.org/10.1126/sciadv.aao1617
- Klapper, W., Kühne, K., Singh, K.K., Heidorn, K., Parwaresch, R., Krupp, G., 1998.
 Longevity of lobsters is linked to ubiquitous telomerase expression. FEBS Lett. 439, 143–146.
- Koroleva, A.G., Evtushenko, E.V., Maximova, N.V., Vershinin, A.V., Sitnikova, T.Y.,
 Kirilchik, S.V., 2015. Length and structure of telomeric DNA in three species of Baikal
 gastropods (Caenogastropoda: Hydrobioidea: Benedictiidae). Russ J Genet 51, 300–307.
 https://doi.org/10.1134/S1022795415030060
- Krupp, G., Bonatz, G., Parwaresch, R., 2000. Telomerase, immortality and cancer. Biotechnol
 Annu Rev 6, 103–140.
- Kuramoto, M., Ohsumi, K., Kishimoto, T., Ishikawa, F., 2001. Identification and analyses of
 the *Xenopus* TERT gene that encodes the catalytic subunit of telomerase. Gene 277 (1-2),
 101-110. https://doi.org/10.1016/S0378-1119(01)00684-9
- 826 Lansdorp, P.M., 2009. Telomeres and disease. EMBO J 28, 2532–2540.
 827 <u>https://doi.org/10.1038/emboj.2009.172</u>
- Lee, E.Y., Lin, J., Noth, E.M., Hammond, S.K., Nadeau, K.C., Eisen, E.A., Balmes, J.R.,
 2017. Traffic-related air pollution and telomere length in children and adolescents living
 in Fresno, CA: A pilot study. Int J Occup Environ Med. 59, 446.
 https://doi.org/10.1097/JOM.0000000000996
- Li, H., Jönsson, B.A.G., Lindh, C.H., Albin, M., Broberg, K., 2011. N-nitrosamines are
 associated with shorter telomere length. Scand J Work Environ Health 37 (4), 316-324.
 https://doi.org/10.5271/sjweh.3150
- Li, H., Hedmer, M., Wojdacz, T., Hossain, M.B., Lindh, C.H., Tinnerberg, H., Albin, M.,
 Broberg, K., 2015. Oxidative stress, telomere shortening, and DNA methylation in
 relation to low-to-moderate occupational exposure to welding fumes. Environ. Mol.
 Mutagen. 56, 684–693. <u>https://doi.org/10.1002/em.21958</u>

- Lin, N., Mu, X., Wang, G., Ren, Y., Su, S., Li, Z., Wang, B., Tao, S., 2017. Accumulative
 effects of indoor air pollution exposure on leukocyte telomere length among nonsmokers. Environ. Pollut. 227, 1–7. <u>https://doi.org/10.1016/j.envpol.2017.04.054</u>
- 842 Lin, S., Huo, X., Zhang, Q., Fan, X., Du, L., Xu, X., Qiu, S., Zhang, Y., Wang, Y., Gu, J., 843 2013. Short placental telomere was associated with cadmium pollution in an electronic 844 waste recycling town in China. PLoS One 8, e60815. 845 https://doi.org/10.1371/journal.pone.0060815
- Lipps, H.J., Rhodes, D., 2009. G-quadruplex structures: in vivo evidence and function. Trends
 Cell Biol. 19, 414–422. <u>https://doi.org/10.1016/j.tcb.2009.05.002</u>
- Liu, D., O'Connor, M.S., Qin, J., Songyang, Z., 2004. Telosome, a mammalian telomereassociated complex formed by multiple telomeric proteins. J. Biol. Chem. 279, 51338– 51342. <u>https://doi.org/10.1074/jbc.M409293200</u>
- Liu, R., Zhang, T., Zhu, G., Xing, M., 2018. Regulation of mutant TERT by BRAF
 V600E/MAP kinase pathway through FOS/GABP in human cancer. Nat Commun 9, 579.
 https://doi.org/10.1038/s41467-018-03033-1
- López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., Kroemer, G., 2013. The hallmarks
 of aging. Cell 153, 1194–1217. <u>https://doi.org/10.1016/j.cell.2013.05.039</u>
- Louis, E.J., 2002. Are *Drosophila* telomeres an exception or the rule? Genome Biol. 3(10),
 reviews0007.1-6. https://doi.org/10.1186/gb-2002-3-10-reviews0007
- Lu, W., Zhang, Y., Liu, D., Songyang, Z., Wan, M., 2013. Telomeres-structure, function, and
 regulation. Exp. Cell Res. 319, 133–141. <u>https://doi.org/10.1016/j.yexcr.2012.09.005</u>
- Madlener, S., Ströbel, T., Vose, S., Saydam, O., Price, B.D., Demple, B., Saydam, N., 2013.
 Essential role for mammalian apurinic/apyrimidinic (AP) endonuclease Ape1/Ref-1 in
 telomere maintenance. Proc. Natl. Acad. Sci. U.S.A. 110, 17844–17849.
 https://doi.org/10.1073/pnas.1304784110
- Martínez, P., Blasco, M.A., 2010. Role of shelterin in cancer and aging. Aging Cell 9, 653–
 666. <u>https://doi.org/10.1111/j.1474-9726.2010.00596.x</u>
- Matzenbacher, C.A., Da Silva, J., Garcia, A.L.H., Cappetta, M., de Freitas, T.R.O., 2019.
 Anthropogenic effects on natural mammalian populations: correlation between telomere
 length and coal exposure. Sci Rep. 9, 6325. https://doi.org/10.1038/s41598-019-42894-8
- Maximova, N., Koroleva, A., Sitnikova, T., Khanaev, I., Bukin, Y, Kirilchik, S., 2017. Age
 dynamics of telomere length of Baikal gastropods is sex specific and multidirectional.
 Folia Biol, 65(4), 187-197. https://doi.org/10.3409/fb65_4.187
- McClintock, B., 1941. The stability of broken ends of chromosomes in zea mays. Genetics 26,
 234–282.
- McKevitt, T.P., Nasir, L., Wallis, C.V., Argyle, D.J., 2003. A cohort study of telomere and
 telomerase biology in cats. Am. J. Vet. Res. 64, 1496–1499.
- Meyne, J., Ratliff, R.L., Moyzis, R.K., 1989. Conservation of the human telomere sequence
 (TTAGGG)n among vertebrates. Proc. Natl. Acad. Sci. U.S.A. 86, 7049–7053.
- Min, J., Wright, W.E., Shay, J.W., 2017. Alternative lengthening of telomeres mediated by
 mitotic DNA synthesis engages break-induced replication processes. Mol. Cell. Biol. 37.
 <u>https://doi.org/10.1128/MCB.00226-17</u>
- Mirabello, L., Huang, W.-Y., Wong, J.Y.Y., Chatterjee, N., Reding, D., Crawford, E.D., De
 Vivo, I., Hayes, R.B., Savage, S.A., 2009. The association between leukocyte telomere
 length and cigarette smoking, dietary and physical variables, and risk of prostate cancer.
 Aging Cell 8, 405–413. https://doi.org/10.1111/j.1474-9726.2009.00485.x
- Mitro, S.D., Birnbaum, L.S., Needham, B.L., Zota, A.R., 2016. Cross-sectional associations
 between exposure to persistent organic pollutants and leukocyte telomere length among
 U.S. adults in NHANES, 2001-2002. Environ Health Perspect. 124(5), 651-658.
 https://doi.org/10.1289/ehp.1510187

- Mizuno, Y., Konishi, S., Hideki, I., Fujimori, E., Kojima, N., Yoshinaga, J., 2019. Cadmium
 exposure and blood telomere length in female university students in Japan. Biol Trace
 Elem Res. 2019, 1-8, https://doi.org/10.1007/s12011-019-1656-3
- Monaghan, P., Haussmann, M.F., 2006. Do telomere dynamics link lifestyle and lifespan ?
 TRENDS Ecol Evol. 21(1), 47-53. https://doi.org/10.1016/j.tree.2005.11.007
- Montpetit, A.J., Alhareeri, A.A., Montpetit, M., Starkweather, A.R., Elmore, L.W., Filler, K.,
 Mohanraj, L., Burton, C.W., Menzies, V.S., Lyon, D.E., Collins, J.B., Teefey, J.M.,
 Jackson-Cook, C.K., 2014. Telomere length: A review of methods for measurement.
 Nurs Res 63, 289–299. https://doi.org/10.1097/NNR.00000000000037
- Moye, A.L., Porter, K.C., Cohen, S.B., Phan, T., Zyner, K.G., Sasaki, N., Lovrecz, G.O.,
 Beck, J.L., Bryan, T.M., 2015. Telomeric G-quadruplexes are a substrate and site of
 localization for human telomerase. Nat Commun 6, 7643.
 https://doi.org/10.1038/ncomms8643
- 902 Muller, H.J., 1938. The remaking of chromosomes. Collecting Net. 13, 181-198, 18.
- Naderlinger, E., Holzmann, K., 2017. Epigenetic regulation of telomere maintenance for
 therapeutic interventions in gliomas. Genes (Basel) 8(5), 145.
 https://doi.org/10.3390/genes8050145
- Nakamura, T.M., Morin, G.B., Chapman, K.B., Weinrich, S.L., Andrews, W.H., Lingner, J.,
 Harley, C.B., Cech, T.R., 1997. Telomerase catalytic subunit homologs from fission yeast
 and human. Science 277, 955–959.
- Natarajan, S., McEachern, M.J., 2002. Recombinational telomere elongation promoted by
 DNA circles. Mol. Cell. Biol. 22:4512-4521.
- 911 Natarajan, S., Groff-Vindman, C., McEachern, M.J., 2003. Factors influencing the
 912 recombinational expansion and spread of Telomeric tandem arrays in *Kluyveromices*913 *lactis*. Eurkaryot Cell, 2:1115-1127.
- Neumann, A.A., Watson, C.M., Noble, J.R., Pickett, H.A., Tam, P.P.L., Reddel, R.R., 2013.
 Alternative lengthening of telomeres in normal mammalian somatic cells. Genes Dev. 27, 18–23. https://doi.org/10.1101/gad.205062.112
- Nica, D.V., Popescu, C., Draghici, G.A., Andrica, F.M., Privistirescu, I.A., Gergen, I.I.,
 Stöger, R., 2017. High-level dietary cadmium exposure is associated with global DNA
 hypermethylation in the gastropod hepatopancreas. PLoS One 12(9), e0184221,
 https://doi.org/10.1371/journal.pone.0184221
- Nussey, D.H., Baird, D., Barrett, E., Boner, W., Fairlie, J., Gemmell, N., Hartmann, N., Horn,
 T., Haussmann, M., Olsson, M., Turbill, C., Verhulst, S., Zahn, S., Monaghan, P., 2014.
 Measuring telomere length and telomere dynamics in evolutionary biology and ecology.
 Methods Ecol Evol 5, 299–310. <u>https://doi.org/10.1111/2041-210X.12161</u>
- 925 O'Callaghan, N.J., Fenech, M., 2011. A quantitative PCR method for measuring absolute
 926 telomere length. Biol Proced Online 13, 3. <u>https://doi.org/10.1186/1480-9222-13-3</u>
- 927 OECD, 2018. OECD Series on Adverse Outcome Pathways. ISSN: 2415170X.
 928 https://doi.org/10.1787/2415170X
- Oeseburg, H., Boer, R.A. de, Gilst, W.H. van, Harst, P. van der, 2010. Telomere biology in
 healthy aging and disease. Pflugers Arch Eur J Physiol 459, 259–268.
 <u>https://doi.org/10.1007/s00424-009-0728-1</u>
- Oikawa, S., Kawanishi, S., 1999. Site-specific DNA damage at GGG sequence by oxidative
 stress may accelerate telomere shortening. FEBS Lett. 453, 365–368.
- Okamoto, K., Iwano, T., Tachibana, M., Shinkai, Y., 2008. Distinct roles of TRF1 in the
 regulation of telomere structure and lengthening. J. Biol. Chem. 283, 23981–23988.
 https://doi.org/10.1074/jbc.M802395200

- Okazaki, S., Tsuchida, K., Maekawa, H., Ishikawa, H., Fujiwara, H., 1993. Identification of a
 pentanucleotide telomeric sequence, (TTAGG)n, in the silkworm *Bombyx mori* and in
 other insects. Mol. Cell. Biol. 13, 1424–1432.
- 940 Olovnikov, A.M., 1971. Principles of marginotomy in template synthesis of polynucleotides.
 941 Doklady Akad Nauk SSSR. 201, 1496-1499.
- 942 Opresko, P.L., Shay, J.W., 2017. Telomere-associated aging disorders. Ageing Res. Rev. 33,
 943 52–66. <u>https://doi.org/10.1016/j.arr.2016.05.009</u>
- Osanai, M., Kojima, K.K., Futahashi, R., Yaguchi, S., Fujiwara, H., 2006. Identification and characterization of the telomerase reverse transcriptase of *Bombyx mori* (silkworm) and Tribolium castaneum (flour beetle). Gene 376, 281–289.
 <u>https://doi.org/10.1016/j.gene.2006.04.022</u>
- Paeschke, K., Simonsson, T., Postberg, J., Rhodes, D., Lipps, H.J., 2005. Telomere endbinding proteins control the formation of G-quadruplex DNA structures in vivo. Nat.
 Struct. Mol. Biol. 12, 847. https://doi.org/10.1038/nsmb982
- Palm, W., de Lange, T., 2008. How shelterin protects mammalian telomeres. Annu. Rev.
 Genet. 42, 301–334. <u>https://doi.org/10.1146/annurev.genet.41.110306.130350</u>
- Pardue, M.L., DeBaryshe, P.G., 2008. Drosophila telomeres: A variation on the telomerase
 theme. Fly (Austin) 2(3), 101-110. https://doi.org/10.4161/fly.6393
- Patel, A., Joshi, G., Fu, Y., Shaju, S., Messina, A., 2017. Telomere elongation and telomerase
 activity in normal and cancer cell lines: HEK-293, HeLa and A549. bioRxiv 100446.
 https://doi.org/10.1101/100446
- Pauliny, A., Larsson, K., Blomqvist, D., 2012. Telomere dynamics in a long-lived bird, the
 barnacle goose. BMC Evol Biol. 12, 257. <u>https://doi.org/10.1186/1471-2148-12-257</u>
- Pavanello, S., Hoxha, M., Dioni, L., Bertazzi, P.A., Snenghi, R., Nalesso, A., Ferrara, S.D.,
 Montisci, M., Baccarelli, A., 2011. Shortened telomeres in individuals with abuse in
 alcohol consumption. Int. J. Cancer 129, 983–992. <u>https://doi.org/10.1002/ijc.25999</u>
- 963 Paviolo, N.S., Santiñaque, F.F., Castrogiovanni, D.C., Folle, G.A., Bolzán, A.D., 2015. The methylating agent streptozotocin induces persistent telomere dysfunction in mammalian 964 Environ 965 cells. Mutat Res Genet Toxicol Mutagen 794, 17-24. 966 https://doi.org/10.1016/j.mrgentox.2015.09.007
- Pawlas, N., Płachetka, A., Kozłowska, A., Broberg, K., Kasperczyk, S., 2015. Telomere
 length in children environmentally exposed to low-to-moderate levels of lead. Toxicol.
 Appl. Pharmacol. 287, 111–118. <u>https://doi.org/10.1016/j.taap.2015.05.005</u>
- Pawlas, N., Płachetka, A., Kozłowska, A., Mikołajczyk, A., Kasperczyk, A., Dobrakowski,
 M., Kasperczyk, S., 2016. Telomere length, telomerase expression, and oxidative stress
 in lead smelters. Toxicol Ind Health 32, 1961–1970.
 https://doi.org/10.1177/0748233715601758
- Perera, F., Lin, C.-J., Qu, L., Tang, D., 2018. Shorter telomere length in cord blood associated
 with prenatal air pollution exposure: Benefits of intervention. Environ Int 113, 335–340.
 https://doi.org/10.1016/j.envint.2018.01.005
- Peterson, D.R., Mok, H.O.L., Au, D.W.T., 2015. Modulation of telomerase activity in fish
 muscle by biological and environmental factors. Comp. Biochem. Physiol. C Toxicol.
 Pharmacol. 178, 51–59. <u>https://doi.org/10.1016/j.cbpc.2015.09.004</u>
- Pottier, G., Viau, M., Ricoul, M., Shim, G., Bellamy, M., Cuceu, C., Hempel, W.M., Sabatier,
 L., 2013. Lead exposure induces telomere instability in human cells. PLoS One 8,
 e67501. <u>https://doi.org/10.1371/journal.pone.0067501</u>
- Puri, P., Nandar, S.K., Kathuria, S., Ramesh, V., 2017. Effects of air pollution on the skin: A
 review. Indian J. Dermatol. Venereol. Leprol. 83(4), 415-423.
 <u>https://doi.org/10.4103/0378-6323.199579</u>

- Raffa, G.D., Ciapponi, L., Cenci, G., Gatti, M., 2011. Terminin: A protein complex that
 mediates epigenetic maintenance of *Drosophila* telomeres. Nucleus 2(5), 383-391.
 https://doi.org/10.4161/nucl.2.5.17873
- Raffa, G.D., Cenci, G., Ciapponi, L., Gatti, M., 2013. Organization and evolution of
 Drosophila terminin: similarities and differences between *Drosophila* and human
 telomeres. Front Oncol. 3:112. https://doi.org/10.3389/fonc.2013.00112
- Raices, M., Maruyama, H., Dillin, A., Karlseder, J., 2005. Uncoupling of longevity and
 telomere length in *C. elegans*. PLoS Genet. 1, e30.
 <u>https://doi.org/10.1371/journal.pgen.0010030</u>
- Raschenberger, J., Lamina, C., Haun, M., Kollerits, B., Coassin, S., Boes, E., Kedenko, L.,
 Köttgen, A., Kronenberg, F., 2016. Influence of DNA extraction methods on relative
 telomere length measurements and its impact on epidemiological studies. Sci. Rep.
 6:25398. https://doi.org/10.1038/srep25398
- Reichard, J.F., Puga, A., 2010. Effects of arsenic exposure on DNA methylation and
 epigenetic gene regulation. Epigenomics 2(1), 87-104. https://doi.org/10.2217/epi.09.45
- 1001 Reichert, S., 2013. Determinants of telomere length and implications in life history trade-offs.
 1002 Thèse de doctorat de l'université de Strasbourg.
- Reichert, S., Froy, H., Boner, W., Burg, T.M., Daunt, F., Gillespie, R., Griffiths, K., Lewis,
 S., Phillips, R.A., Nussey, D.H., Monaghan, P., 2017. Telomere length measurement by
 qPCR in birds is affected by storage method of blood samples. Oecologia 184, 341–350.
 <u>https://doi.org/10.1007/s00442-017-3887-3</u>
- Roake, C.M., Artandi, S.E., 2016. Telomere-lengthening mechanism revealed. Nature 539,
 35-36. https://doi.org/10.1038/nature19483
- Rosén M., Edström J.E., 2001. DNA structures common for chironomid telomeres
 terminating with complex repeats. Insect Mol Biol 9, 341–347.
 https://doi.org/10.1046/j.1365-2583.2000.00193.x
- Rudolph, K.L., Chang, S., Lee, H.W., Blasco, M., Gottlieb, G.J., Greider, C., DePinho, R.A.,
 1013 1999. Longevity, stress response, and cancer in aging telomerase-deficient mice. Cell 96,
 1014 701–712.
- Sampson, M.J., Winterbone, M.S., Hughes, J.C., Dozio, N., Hughes, D.A., 2006. Monocyte
 telomere shortening and oxidative DNA damage in type 2 diabetes. Diabetes Care 29(2),
 283-289. https://doi.org/10.2337/diacare.29.02.06.dc05-1715
- Sandin, S., Rhodes, D., 2014. Telomerase structure. Curr. Opin. Struct. Biol. 25, 104–110.
 <u>https://doi.org/10.1016/j.sbi.2014.02.003</u>
- 1020Schumpert, C., Nelson, J., Kim, E., Dudycha, J.L., Patel, R.C., 2015. Telomerase activity and1021telomerelengthindaphnia.PLoSOne10,e0127196.1022https://doi.org/10.1371/journal.pone.0127196
- Scott, N.M., Haussmann, M.F., Elsey, R.M., Trosclair, P.L., Vleck, C.M., 2006. Telomere
 length shortens with body length in *Alligator mississippiensis*. Southeastern Naturalist 5,
 685–692. <u>https://doi.org/10.1656/1528-7092(2006)5[685:TLSWBL]2.0.CO;2</u>
- Shampay, J., Szostak, J.W., Blackburn, E.H., 1984. DNA sequences of telomeres maintained
 in yeast. Nature 310, 154–157.
- Shay, J.W., Wright, W.E., 2019. Telomeres and telomerases: three decades of progress. Nat
 Rev Genet. 20, 299-309. https://doi.org/10.1038/s41576-019-0099-1
- Sletten, S., Bourgeon, S., Bårdsen, B.-J., Herzke, D., Criscuolo, F., Massemin, S., Zahn, S.,
 Johnsen, T.V., Bustnes, J.O., 2016. Organohalogenated contaminants in white-tailed
 eagle (*Haliaeetus albicilla*) nestlings: An assessment of relationships to immunoglobulin
 levels, telomeres and oxidative stress. Sci. Total Environ. 539, 337–349.
 https://doi.org/10.1016/j.scitotenv.2015.08.123

- Srinivas, N., Rachakonda, S., Hielscher, T., Calderazzo, S., Rudnai, P., Gurzau, E., Koppova,
 K., Fletcher, T., Kumar, R., 2019. Telomere length, arsenic exposure and risk of basal
 cell carcinoma of skin. Carcinogenesis, bgz059, https://doi.org/10.1093/carcin/bgz059
- Stauffer, J., Panda, B., Eeva, T., Rainio, M., Ilmonen, P., 2017. Telomere damage and redox
 status alterations in free-living passerines exposed to metals. Sci Total Environ 575, 841–
 848. https://doi.org/10.1016/j.scitotenv.2016.09.131
- Stauffer, J., Panda, B., Ilmonen, P., 2018. Telomere length, sibling competition and development of antioxidant defense in wild house mice. Mech. Ageing Dev. 169, 45–52.
 <u>https://doi.org/10.1016/j.mad.2017.10.002</u>
- Szostak, J.W., Blackburn, E.H., 1982. Cloning yeast telomeres on linear plasmid vectors. Cell
 29, 245–255.
- Tan, T.C.J., Rahman, R., Jaber-Hijazi, F., Felix, D.A., Chen, C., Louis, E.J., Aboobaker, A.,
 2012. Telomere maintenance and telomerase activity are differentially regulated in
 asexual and sexual worms. Proc Natl Acad Sci U S A 109, 4209–4214.
 https://doi.org/10.1073/pnas.1118885109
- Tasaka, K., Yokoyama, N., Nodono, H., Hoshi, M., Matsumoto, M., 2013. Innate sexuality
 determines the mechanisms of telomere maintenance. Int. J. Dev. Biol. 57, 69–72.
 <u>https://doi.org/10.1387/ijdb.120114mm</u>
- Tomás-Loba, A., Flores, I., Fernández-Marcos, P.J., Cayuela, M.L., Maraver, A., Tejera, A.,
 Borrás, C., Matheu, A., Klatt, P., Flores, J.M., Viña, J., Serrano, M., Blasco, M.A., 2008.
 Telomerase reverse transcriptase delays aging in cancer-resistant mice. Cell 135, 609–
 622. <u>https://doi.org/10.1016/j.cell.2008.09.034</u>
- Topcu, Z., Nickles, K., Davis, C., McEachern, M.J., 2005. Abrupt disruption of capping and a single source for recombinationally elongated telomeres in *Kluyveromyces lactis*. Proc. Natl. Acad. Sci. U.S.A. 102, 3348–3353. <u>https://doi.org/10.1073/pnas.0408770102</u>
- Traut, W., Szczepanowski, M., Vítková, M., Opitz, C., Marec, F., Zrzavý, J., 2007. The
 telomere repeat motif of basal Metazoa. Chromosome Res. 15, 371–382.
 <u>https://doi.org/10.1007/s10577-007-1132-3</u>
- 1063 Tricola, G.M., Simons, M.J.P., Atema, E., Boughton, R.K., Brown, J.L., Dearborn, D.C., 1064 Divoky, G., Eimes, J.A., Huntington, C.E., Kitaysky, A.S., Juola, F.A., Lank, D.B., 1065 Litwa, H.P., Mulder, E.G.A., Nisbet, I.C.T., Okanoya, K., Safran, R.J., Schoech, S.J., Schreiber, E.A., Thompson, P.M., Verhulst, S., Wheelwright, N.T., Winkler, D.W., 1066 1067 Young, R., Vleck, C.M., Haussmann, M.F., 2018. The rate of telomere loss is related to 1068 maximum lifespan in birds. Phil. Trans. R. Soc. В 373. 20160445. https://doi.org/10.1098/rstb.2016.0445 1069
- 1070 Ujvari, B., Madsen, T., 2009. Short telomeres in hatchling snakes: erythrocyte telomere
 1071 dynamics and longevity in tropical pythons. PLoS One 4, e7493.
 1072 <u>https://doi.org/10.1371/journal.pone.0007493</u>
- 1073 Unryn, B.M., Cook, L.S., Riabowol, K.T., 2005. Paternal age is positively linked to telomere
 1074 length of children. Aging Cell 4, 97-101. https://doi.org/10.1111/j.14741075 9728.2005.00144.x
- 1076 Uziel, O., Singer, J.A., Danicek, V., Sahar, G., Berkov, E., Luchansky, M., Fraser, A., Ram,
 1077 R., Lahav, M., 2007. Telomere dynamics in arteries and mononuclear cells of diabetic
 1078 patients: effect of diabetes and of glycemic control. Exp Gerontol. 42(10), 971-978.
 1079 https://doi.org/10.1016/j.exger/2007.07.005
- 1080 Valdes, A.M., Andrew, T., Gardner, J.P., Kimura, M., Oelsner, E., Cherkas, L.F., Aviv, A.,
- 1081Spector, T.D., 2005. Obesity, cigarette smoking, and telomere length in women. Lancet1082366, 662–664. https://doi.org/10.1016/S0140-6736(05)66630-5

- Vallabhaneni, H., O'Callaghan, N., Sidorova, J., Liu, Y., 2013. Defective repair of oxidative
 base lesions by the DNA glycosylase Nth1 associates with multiple telomere defects.
 PLoS Genet. 9, e1003639. <u>https://doi.org/10.1371/journal.pgen.1003639</u>
- Vallabhaneni, H., Zhou, F., Maul, R.W., Sarkar, J., Yin, J., Lei, M., Harrington, L., Gearhart,
 P.J., Liu, Y., 2015. Defective repair of uracil causes telomere defects in mouse
 hematopoietic cells. J. Biol. Chem. 290, 5502–5511.
 https://doi.org/10.1074/jbc.M114.607101
- 1090 Vasseur, P., Cossu-Leguille, C., 2003. Biomarkers and community indices as complementary
 1091 tools for environmental safety. Env Int. 28(8), 711-717.
- 1092 Vítková, M., Král, J., Traut, W., Zrzavý, J., Marec, F., 2005. The evolutionary origin of insect
 1093 telomeric repeats, (TTAGG)n. Chromosome Res. 13, 145–156.
 1094 <u>https://doi.org/10.1007/s10577-005-7721-0</u>
- 1095 Vitturi, R., Colomba, M.S., Pirrone, A., Libertini, A., 2000. Physical mapping of rDNA genes, (TTAGGG)n telomeric sequence and other karyological features in two
 1097 earthworms of the family Lumbricidae (Annelida : Oligochaeta). Heredity, Sep;85, Pt3, 203-207.
- 1099 Vitturi, R., Libertini, A., Sineo, L., Sparacio, I., Lannino, A., Gregorini, A., Colomba, M.,
 1100 2005. Cytogenetics of the land snails *Cantareus aspersus* and *C. mazzullii* (Mollusca:
 1101 Gastropoda: Pulmonata). Micron 36, 351–357.
 1102 https://doi.org/10.1016/j.micron.2004.12.010
- von Zglinicki, T., 2002. Oxidative stress shortens telomeres. Trends Biochem. Sci. 27, 339–
 344.
- Vulliamy, T.J., Knight, S.W., Mason, P.J., Dokal, I., 2001. Very short telomeres in the
 peripheral blood of patients with X-linked and autosomal dyskeratosis congenita. Blood
 Cells Mol. Dis. 27, 353–357. <u>https://doi.org/10.1006/bcmd.2001.0389</u>
- Wai, K.M., Umezaki, M., Kosaka, S., Mar, O., Umemura, M., Fillman, T., Watanabe, C.,
 2018. Impact of prenatal heavy metal exposure on newborn leucocyte telomere length: A
 birth-cohort study. Environ Pollut 243(B), 1414-1421.
 https://doi.org/10.1016/j.envpol.2018.09.090
- Walter, M.F., Biessmann, M.R., Benitez, C., Török, T., Mason, J.M., Biessmann, H., 2007.
 Effects of telomere length *in Drosophila melanogaster* on life span, fecundity and fertility. Chromosoma 116, 41–51. <u>https://doi.org/10.1007/s00412-006-0081-5</u>
- Walton, R.T., Mudway, I.S., Dundas, I., Marlin, N., Koh, L.C., Aitlhadj, L., Vulliamy, T.,
 Jamaludin, J.B., Wood, H.E., Barratt, B.M., Beevers, S., Dajnak, D., Sheikh, A., Kelly,
 F.J., Griffiths, C.J., Grigg, J., 2016. Air pollution, ethnicity and telomere length in east
 London schoolchildren: An observational study. Environ Int 96, 41–47.
 <u>https://doi.org/10.1016/j.envint.2016.08.021</u>
- Wang, Z., Rhee, D.B., Lu, J., Bohr, C.T., Zhou, F., Vallabhaneni, H., de Souza-Pinto, N.C.,
 Liu, Y., 2010. Characterization of oxidative guanine damage and repair in mammalian
 telomeres. PLoS Genet. 6, e1000951. <u>https://doi.org/10.1371/journal.pgen.1000951</u>
- Webb, C.J., Wu, Y., Zakian, V.A., 2013. DNA repair at telomeres: keeping the ends intact.
 Cold Spring Harb Perspect Biol 5. <u>https://doi.org/10.1101/cshperspect.a012666</u>
- Weismann, A., 1889. Collected essays upon heredity and kindred biological problems.
 Poulton, E.B., Clarendon, Oxford.
- Wicky, C., Villeneuve, A.M., Lauper, N., Codourey, L., Tobler, H., Müller, F., 1996.
 Telomeric repeats (TTAGGC)n are sufficient for chromosome capping function in *Caenorhabditis elegans*. Proc Natl Acad Sci U S A 93, 8983–8988.
- Williamson, J.R., Raghuraman, M.K., Cech, T.R., 1989. Monovalent cation-induced structure
 of telomeric DNA: the G-quartet model. Cell 59, 871–880.

- Wood, M.L., Dizdaroglu, M., Gajewski, E., Essigmann, J.M., 1990. Mechanistic studies of
 ionizing radiation and oxidative mutagenesis: genetic effects of a single 8hydroxyguanine (7-hydro-8-oxoguanine) residue inserted at a unique site in a viral
 genome. Biochemistry 29, 7024–7032.
- Xin, X., Senthilkumar, P.K., Schnoor, J.L., Ludewig, G., 2016. Effects of PCB126 and
 PCB153 on telomerase activity and telomere length in undifferentiated and differentiated
 HL-60 cells. Environ Sci Pollut Res Int 23, 2173–2185. <u>https://doi.org/10.1007/s11356-</u>
 015-5187-y
- Xu, M., Wu, X.-B., Yan, P., Zhu, H., 2009. Telomere length shortens with age in chinese
 alligators (*Alligator sinensis*). J. Appl. Anim. Res. 36, 109–112.
 <u>https://doi.org/10.1080/09712119.2009.9707042</u>
- Xu, Y., Tokar, E.J., Waalkes, M.P., 2014, Arsenic-induced cancer cell phenotype in human
 breast epithelia is estrogen receptor-independent but involves aromatase activation. Arch.
 Toxicol. 88(2), 263-274. https://doi.org/10.1007/s00204-013-1131-4
- Yao, M.C., Blackburn, E., Gall, J., 1981. Tandemly repeated C-C-C-C-A-A hexanucleotide of
 Tetrahymena rDNA is present elsewhere in the genome and may be related to the
 alteration of the somatic genome. J. Cell Biol. 90:515-520.
- Yip, B.W., Mok, H.O., Peterson, D.R., Wan, M.T., Taniguchi, Y., Ge, W., Au, D.W., 2017.
 Sex-dependent telomere shortening, telomerase activity and oxidative damage in marine
 medaka *Oryzias melastigma* during aging. Mar. Pollut. Bull. 124, 701–709.
 <u>https://doi.org/10.1016/j.marpolbul.2017.01.021</u>
- Yuan, J., Liu, Y., Wang, J., Zhao, Y., Li, K., Jing, Y., Zhang, X., Liu, Q., Geng, X., Li, G.,
 Wang, F., 2018. Long-term persistent organic pollutants exposure induced telomere
 dysfunction and senescence-associated secretary phenotype. J Gerontol A Biol Sci Med
 Sci 73(8), 1027-1035. https://doi.org/10.1093/gerona/gly002.
- Zeljezic, D., Bjelis, M., Mladinic, M., 2015. Evaluation of the mechanism of nucleoplasmic
 bridge formation due to premature telomere shortening in agricultural workers exposed to
 mixed pesticides: indication for further studies. Chemosphere 120, 45–51.
 <u>https://doi.org/10.1016/j.chemosphere.2014.05.085</u>
- Zhang, X., Lin, S., Funk, W.E., Hou, L., 2013. Environmental and occupational exposure to chemicals and telomere length in human studies. Occup Environ Med 70, 743–749.
 <u>https://doi.org/10.1136/oemed-2012-101350</u>
- Zhang, Y., Zhang, L., Tang, X., Bhardwaj, S.R., Ji, J., Rong, Y.S., 2016. MTV, an ssDNA
 protecting complex essential for transposon-based telomere maintenance in *Drosophila*.
 PLoS Genet 12(11), e1006435. https://doi.org/10.1371/journal.pgen.1006435
- 1167 Zhang, C., Kibriya, M.G., Jasmine, F., Roy, S., Gao, J., Sabarinathan, M., Shinkle, J., 1168 Delgado, D., Ahmed, A., Islam, T., Eunus, M., Islam, T., Hasan, R., Graziano, J.H., 1169 Ahsan, H., Pierce, B.L., 2018. A study of telomere length, arsenic exposure, and arsenic 1170 toxicity a Bangladeshi cohort. Environ Res. 164, 346-355. in 1171 https://doi.org/10.1016/j.envres.2018.03.005
- Zhang, L., Song, L., Liu, B., Wu, M., Wang, L., Zhang, B., Xiong, C., Xia, W., Li, Y., Cao,
 Z., Wang, Y., Xu, S., 2019. Prenatal cadmium exposure is associated with shorter
 leukocyte telomere length in Chinese newborns. BMC Med 17:27, 1-8.
 https://doi.org/10.1186/s12916-019-1262-4
- Zhao, J., Chan, J., Lambelé, M., Yusufzai, T., Stumpff, J., Opresko, P.L., Thali, M., Wallace,
 S.S., 2018. Air pollution and telomere length: a systematic review of 12,058 subjects.
 Cardiovasc Diagn Ther 8(4), 480-492. https://doi/org/10.21037/cdt.2018.06.05
- Zhou, J., Chan, J., Lambelé, M., Yusufzai, T., Stumpff, J., Opresko, P.L., Thali, M., Wallace,
 S.S., 2017. NEIL3 repairs telomere damage during S phase to secure chromosome

- 1181
 segregation
 at
 mitosis.
 Cell
 Rep
 20,
 2044–2056.

 1182
 https://doi.org/10.1016/j.celrep.2017.08.020
 20
 2044–2056.
 20
- Ziegler, S., Schettgen, T., Beier, F., Wilop, S., Quinete, N., Esser, A., Masouleh, B.K.,
 Ferreira, M.S.V., Vankann, L., Uciechowski, P., Rink, L., Kraus, T., Brümmendorf, T.H.,
 Ziegler, P., 2017. Accelerated telomere shortening in peripheral blood lymphocytes after
 occupational polychlorinated biphenyls exposure. Arch Toxicol 91, 289–300.
 https://doi.org/10.1007/s00204-016-1725-8
- 1188 Zota, A.R., Needham, B.L., Blackburn, E.H., Lin, J., Park, S.K., Rehkopf, D.H., Epel, E.S.,
- 1189 2015. Associations of cadmium and lead exposure with leukocyte telomere length: 1190 findings from national health and nutrition examination survey, 1999–2002. Am J
- 1191 Epidemiol 181, 127–136. <u>https://doi.org/10.1093/aje/kwu293</u>

1192





1 2	Telomere dynamic in humans and animals: review and perspectives in environmental toxicology
3	
4	List of the figures
5	
6 7 8	<u>File fig1.tif:</u> <i>Figure 1: Human TL structure and six TL binding proteins: TRF1, TRF2, RAP1, TPP1, TIN2, POT1 (modified from Blasco, 2005, Martinez and Blasco, 2010, Lu et al., 2013, Sandin and Rhodes, 2014).</i>
9	File fig2.tif: Figure 2: Some causes of TL attrition.
10	
11	List of the Tables
12	Table 1: No-exhaustiv list of telomere sequences of bioindicator taxons in environmental
13	toxicology.
14	Table 2: Roles of the six telomere binding proteins: TRF1, TRF2, RAP1, TPP1, TIN2, POT1
15	(adapted from Iwano et al., 2004, Okamoto et al., 2008, Martinez and Blasco, 2010).
16	Table 3: Links between telomere shortening and ageing with others biologicals markers in
17	various biological groups of animals.
18	Table 4: Relationship between telomere dynamics and environmental contamination by
19	contaminants in humans.
20	Table 5: Relationship between telomere dynamics and environmental contamination by
21	contaminants in animal.
22	
23	
24	
25	
26	
27	

- **Table 1:** No-exhaustiv list of telomere sequences of bioindicator taxons in environmental
- *toxicology*

	SEQUENCES	SPECIES	SOURCES
	(TTAGGG) _N	Vertebrates (Fishs, Birds, Mammals, Amphibians), Invertebrates (Snails as <i>Cantareus aspersus</i> , Earthworms as <i>Eisenia foetida</i> , Planaria)	Meyne et al., 1989, Vitturi et al., 2000, Vitturi et al., 2005, Koroleva et al., 2013, Tasaka et al., 2013.
	(TTAGG) _N	Invertebrates (Daphnia sp., Chironomus sp., Bombyx mori)	Okazaki et al., 1993, Rosen et Edstrom, 2000, Schumpert et al., 2015
30	(TTAGGC) _N	Invertebrates (Nematod, Caenorhabditis elegans)	Wicky et al., 1996
31			
32			
33			
34			
35			
36			
37			
38			
39			
40			
41			
42			
43			
44			
45			
46			
47			
48	Table 2: Rol	es of the six telomere binding proteins: TRF1, 7	TRF2, RAP1, TPP1, TIN2, POT1
49	(adapted from	ı Iwano et al., 2004, Okamoto et al., 2008, Mai	rtinez and Blasco, 2010)

PROTEINS ROLES	TIN2	RAP1	TPP1	TRF2	TRF1	POT1
BRIDGING SHELTERIN	x		x			
CAPPING SHELTERIN			x	x	Х	
G-STRAND PROTECTION						x
INTERACTIONS WITH DNA DAMAGE RESPONSE (DDR)		x		X	Х	x
INTERACTIONS WITH MITOSIS REGULATIONS					Х	
PREVENTION OF TELOMERE FRAGILITY		x	x		х	
PREVENTION OF TELOMERE RECOMBINATION AND END-TO-END FUSION		х		х		х
STIMULATING SUBTELOMERIC SILENCING		х				
ALLOWS TA RECRUITMENT			X			
TL ACCESSIBILITY FOR TA INHIBITION						X
TELOMERIC LOCALIZATION OF OTHER SHELTERIN COMPONENTS REGULATION	X				Х	

Table 3: Links between telomere shortening and ageing with others biologicals markers in various biological groups of animals.

BIOLOGICAL SPECIES	STUDIED AGES	Ν	BIOLOGICAL TISSUES	EFFECTS OF AGEING ON TELOMERE DYNAMICS INVERTEBRATES	OF AGEING ON EFFECTS OF AGEING ON OTHERS RE DYNAMICS BIOLOGICALS MARKERS ATES	
Strongylocentrotus franciscanus (Sea urchin)	4 to > 100 years	?	Some tissues (Aristotle's lantern muscle, ampullae,	↓ telomere lengths, O telomerase activity	↑ body size	Francis et al., 2006
Lytechnius variegatus (Sea urchin)	> 3 years	?	esophagus, intestine, gonad)	↓ telomere lengths, O telomerase activity	↑ body size	Francis et al., 2006
<i>Schmidtea mediterranea</i> (Planaria)	7 days to 3 years	73 (min. 7 per ages)	Somatic cells	↓ telomere lengths	-	Tan et al., 2012
Daphnia pulex (Daphnia)	Embryos, 1, 2 and 3 weeks	25 to 30 individuals in pools/age	Body	O telomere lengths, ↑ telomerase activity	↑ mortality	Schumpert et al., 2015
Daphnia pulicaria (Daphnia)	Embryos, 1, 4 and 8 weeks	25 to 30 individuals in pools/age	Body	\downarrow telomere lengths, \downarrow telomerase activity	↑ mortality	Schumpert et al., 2015
Drosophila melanogaster (Vinegar fly)	0 to 100 days	2000 (100 per 5 days)	Body	O telomere lengths	\downarrow fecundity, \downarrow fertility	Walter et al., 2007
Benedictia fragilis (Baikal gastropods)	0 to 8 years	95	Foot	↓ telomere lengths	-	Maximova et al., 2017
Benedictia baicalensis (Baikal gastropods)	0 to 8 years	121	Foot	↑ telomere lengths	-	Maximova et al., 2017
Kobeltocochlea martensiana (Baikal gastropods)	0 to 8 years	111	Foot	↑ telomere lengths	-	Maximova et al., 2017
				FISHS		
Oryzias melastigma (Medaka)	4, 8, 12 months	15 (5/age)	Liver	, ↓ telomere lengths, ∆ telomerase activity, ∆ TERT mRNA expression in liver,	In liver: \uparrow 8-OH-dG level, \uparrow protein carbonyl level, Δ omER α mRNA expression, in plasma: Δ testosterone (T) and Δ estrogen (E2) levels	Yip et al., 2017
Danio rerio (Zebrafish)	8 days to 24 months	140 (~5/age, ~100 at 8 days)	Some tissues (testis, gut, head kidney, muscle, skin)	↓ telomere lengths, †/↓ telomerase activity (∆ f(tissues))	-	Henriques et al., 2013, Henriques et al., 2017
				REPTILES		
Alligator sinensis (Chinese alligator)	3 to 10 years	30	Erythrocytes	↓ telomere lengths	-	Xu et al., 2009
<i>Alligator</i> <i>mississippiensis</i> (American Alligator)	0 to 7 years	34	Erythrocytes	↓ telomere lengths	↑ body size	Scott et al., 2006
<i>Liasis fuscus</i> (Water python)	0 to 20 years	70	Erythrocytes	↑ telomere lengths	-	Ujvari et Masden, 2009
	. –			BIRDS		
<i>Taeniopygia guttata</i> (Zebra finch)	1 to 7 years	99	Erythrocytes	\downarrow telomere lengths	-	Heidinger et al., 2012
Taeniopygia guttata	0 to 8 years	25	Erythrocytes	↓ telomere lengths	-	Haussman et al., 2003
Tachycineta bicolor (Tree swallow)	0 to 7 years	42	Erythrocytes	\downarrow telomere lengths	-	Haussman et al., 2003

<i>Pygoscelis adeliae</i> (Adélie penguin)	1 to 14 years	22	Erythrocytes	↓ telomere lengths	-	Haussman et al., 2003
Sterna hirundo (Common tern)	0 to 25 years	41	Erythrocytes	\downarrow telomere lengths	-	Haussman et al., 2003
Oceanodroma leucorhoa (Leach's petrel)	0 to 35 years	33	Erythrocytes	↑ telomere lengths	-	Haussman et al., 2003
Phalacrocorax aristolelis (European shag)	0 to 20 years	144	Whole blood (chicks), erythrocytes (adults)	↓ telomere lengths	-	Hall et al., 2004
<i>Diomedea exulans</i> (Wandering albatross)	0 to 42 years	92	Whole blood (chicks), erythrocytes (adults)	↓ telomere lengths	-	Hall et al., 2004
Branta leucopsis (Barnacle goose)	2 to 26 years	30	Whole blood	↓ telomere lengths	-	Pauliny et al., 2012
Acrocephalus arundinaceus (Great reed warbler)	0 to 6 years	174	Whole blood	↓ telomere lengths	-	Asghar et al., 2015
(,				MAMMALS		
Canis lupus familiaris (Dogs)	0 to 15 years	175	Whole blood	\downarrow telomere lengths	↑ probability of death from cardiovascular disease	Fick et al., 2012
Felis catus (Cats)	cohort	32	Whole blood	\downarrow telomere lengths	-	McKevitt et al., 2003
<i>Mus spretus</i> (Western Mediterranean mouse)	0 to 2,5 years	20	Some tissues (testes, kidney, liver, spleen, brain)	↓ telomere lengths	-	Coviello- McLaughlin et al., 1997
Homo sapiens	0 to 82	72	Leukocytes	↓ telomere lengths	-	Frenck et al., 1998
(Humans) Homo sapiens	cohort	193	Leukocytes	↓ telomere lengths	\downarrow pulse wave velocity, pulse pression	Benetos et al., 2001

Legends: Δ = variations, \uparrow = increasing, \downarrow = decreasing, O = no effects

 Table 4: Relationship between telomere dynamics and environmental contamination by contaminants in humans

BIOLOGICAL MATERIALS	DLOGICAL EXPERIMENTALS ATERIALS CONDITIONS CONTAMINANTS EFFECTS ON TELOMERE DYNAMICS		OTHERS BIOLOGICALS EFFECTS AND ASSOCIATIONS WITH INTERNAL CONCENTRATIONS	SOURCES			
			IN VITRO STUDIES				
HL-60 cells	6 and 30 days in vitro	PCB126 and PCB153 (5 μM)	↓ Telomere length, ↓ Telomerase activity, ↓ hTERT, ↓ hTR, Δ POT1, Δ TRF1, Δ TRF2 expressions	↑ phases G0/G1 cells	Xin et al., 2016		
Primary cultured lymphocytes	red2, 7, 42 and 70 days in vitro3-OH-CB28 (metabolite of PCB28) (1-50 μ M) \downarrow Telomere length, \downarrow Telomerase activity, \downarrow Telomerase genes expressionsO lymphocytes numbers, O TF receptor excision circle)		O lymphocytes numbers, O TREC (T-cell receptor excision circle) levels	Ziegler et al., 2017			
Cells of umbilical cord blood (huUCB)	30 minutes – 14 days	As inorganic (0,1 nM, 1 and 10 µM)	\downarrow Telomere length, \downarrow hTERT expression	\downarrow Telomere length, \downarrow hTERT expression \uparrow ROS			
Cells of carcinoma cell line (Clone B3, EJ30 bladder cell)	ls of carcinoma cell te (Clone B3, EJ30 1 day Pb(NO ₃) ₂ (0-1000 μM) ↓ Telomere length bladder cell)		↑ chromosomal abnormalities	Pottier et al., 2013			
	STUDIES ON COHORTS OR ON PATIENTS						
			METALLIC ELEMENTS				
Placenta from 320 newborns	patients	Placental Cd and Pb concentration	↓ Telomere length	Association with [Cd] placental	Lin et al., 2013		
Leukocytes from 6796 adults	cohort	Cd and Pb in blood and urines	↓ Telomere length	Association with [Cd] _{blood}	Zota et al., 2015, Patel et al., 2017		
Leukocytes from 334 smealters	cohort	Cd and Pb in blood (occupational exposure by air)	↓ Telomere length	Association with [Pb]blood	Pawlas et al., 2016		
Leukocytes from 365 children	cohort	Pb in blood	↓ Telomere length	Association with [Pb]blood	Pawlas et al., 2015		
Salivary cells from adolescents (12-16 years olds, n=351)	patients	As and Cd in urines	↓ Telomere length	Association with [Cd] _{urinary}	Fillman et al., 2016		
Leukocytes from 73 female university students	cohort	Cd in urines	O Telomere length	-	Mizuno et al., 2019		

Umbilical cord blood leukocytes from 410 mother-newborn pairs	cohort	Cd in urines	↓ Telomere length	Association with [Cd] _{urinary}	Zhang et al., 2019
Leukocytes from 167 participants	patients	As in urines	↑ Telomere length, ↑ Expression of WRN (ALT), ↓ Expressions of genes including TERF2 (shelterin complex), DKC1 (up-regulation of telomerase), TERF2IP (sheltering complex), OBFC1 (up- regulation of telomerase)	Association with [As]urinary	Gao et al., 2015
Leukocytes from 188 male participants (76 exposed in drinking water > 10 µg.L ⁻¹ , 112 control group	cohort	As in urines	↑ Telomere length	Association with [As]drinking water and [As]urinary	Jimenez Villarreal et al., 2018
Leukocytes from 1061 patiens (528 with skin cancers, 533 hospital- based controls aged 30- 79 years)	patients	As in urines	↓ Telomere length	Association with [As]urinary and skin cancer	Srinivas et al., 2019
Leukocytes from 180 participants (60 unexposed, 120 exposed)	cohort	As in urines	↑ Telomere length, O telomerase activity, ↑ levels of TRF1 and TRF2 (shelterin complex)	Association with [As] _{urinary} , ↑ percentage of senescent cells, ↑ expression of cytokines (IL-6 and IL-8), ↑ expression of metalloproteases (MMP-1 and MMP-3), ↑ levels of p21 and p53, ↓ p16 level	Chatterjee et al., 2015
Leukocytes from 120 children (5 to 15 years) participants (68 exposed in drinking water > 10 µg.L ⁻¹ , 52 control group)	cohort	As in drinking water and in urines	O Telomere length	Association with [As] _{urinary} , ↑ micronucleus in buccal, urothelial cells and lymphocytes,	Chatterjee et al., 2018
Leukocytes from 1380 participants	cohort	As in urines	O/↓ Telomere length	Association with [As] _{urinary} /greater risk of subsequent skin lesion	Zhang et al., 2018
Leukocytes from 460 newborn	cohort	As, Cd and Pb in maternal urines	↓ Telomere length	Association with [As] or [Cd] _{urinary}	Wai et al., 2018
Leukocytes from 1702 participants	cohort	As, Cd and W in urines	↓ Telomere length	Association [As], [Cd] and [W]urinary	Grau-Perez et al., 2019
			ALCOHOL		
Leukocytes from 404	patients	Alcohol	↓ Telomere length	Association with alcohol abusers	Pavanello et al., 2011

participants

ATMOSPHERIC POLLUTION

Leukocytes from 333 schoolchildren	cohort	NO _x , NO ₂ , PM _{2.5} , PM ₁₀	↑ Telomere length	Association with annual air pollution exposure	Walton et al., 2016
Leukocytes from 120 truck drivers and 120 office workers	cohort	NO _x , NO ₂ , PM _{2.5} , PM ₁₀	↑ Telomere length	Association with PM _{2.5} , PM ₁₀	Hou et al., 2012
Leukocytes from 63 steel workers (3 days postexposure during 1 day)	cohort	PM ₁ , PM ₁₀	↑ Telomere length, ↓ hTERT mRNA	Association with PM_1 and PM_{10}	Dioni et al., 2011
Leukocytes from 264 patients	patients	Solid fuel combustion	↓ Telomere length	Association with solid fuel combustion for three decades	Lin et al., 2017
Leukocytes from 101 welders and 127 controls	patients	Respirable dust (between 1,2 mg/m ³ to 3,3 mg/m ³ , control: 0,1 mg/m ³)	O/↓ telomere length	O 8-oxodG levels, O/↓ DNA methylation	Li et al., 2015
Leukocytes from 50 prengnant women living in polluted area vs 50 in a non-polluted area	cohort	Atmospheric pollution and intense environmental pressure due to wastes	↓ Telomere length, ↓ hTERT mRNA	Association with polluted area	De Felice et al., 2012
Buccal cells and placenta cells from 233 patients	cohort	Residential traffic exposure	↓ Telomere length	Association with residential traffic exposure	Bijnens et al., 2017
Leukocytes from workers (77 street traffic officers and 57 office workers (referents))	cohort	Personal airborne benzene exposure (0 to 300 µg/m ³) and toluene exposure (12.5 to 1000 µg/m ³)	↓ Telomere length	Association with benzene and toluene exposure	Hoxha et al., 2009
Leukocytes from workers (43 and 43 in controls)	cohort	Workers exposed to benzene (31 ppm in 5 days)	↑ Telomere length	Association with benzene exposure (> 31 ppm in 5 days)	Bassig et al., 2014
Leukocytes from 157 rubber workers	cohort	Workers exposed to pyrene and nitrosamines	↓ Telomere length	Association with nitrosamines exposure	Li et al., 2011

Cells of cord blood on 255 newborns	patients	РАН	↓ Telomere length	Association with PAH-DNA adducts	Perera et al., 2018
Leukocytes from 14 children	patients	РАН	↓ Telomere length	Association with PAH exposure	Lee et al., 2017
Leukocytes from 1330 adults	cohort	18 PCBs, 7 dioxins, 9 furans	↑ Telomere length	Association with TEQ (Toxic Equivalency)	Mitro et al., 2015
Leukocytes from 452 adults (136 in controls, 316 exposed)	cohort	25 PCBs and 11 PBDEs	↓ Telomere length, ↑ frequences of telomere loss and multiple separate telomere at the ends of chromosomes, ↑ ALT activities,	↑ [PCBs] _{lipids} , ↑ [PBDEs] _{lipids} , ↑ abortion, stillborn, hypertension, diabetes and autoimmune diseases, ↑ relative expression of p53 and Rb in plasma, ↑ p16ink4a and p14-arf level in plasma, ↑ interleukines levels in plasma (IL-2, IL-6, IL-10),	Yuan et al., 2018
			PESTICIDES		
Leukocytes from 30 farmers and 30 townsfolk	patients	17 pesticides	O Telomere length	↑ micronuclei, ↑ nucleoplasmic bridges (correlation with [pesticides])	Zeljezic et al., 2015
Leukocytes from 568 cancer-free male farmers	cohort	57 pesticides	↓ Telomere length	Association with exposure of 2,4-D, diazinon, butuylate, malathion (except for alachlor it's opposite)	Andreotti et al., 2015

Legends: Δ = variations, \uparrow = increasing, \downarrow = decreasing, O = no effects

Table 5: Relationship between telomere dynamics and environmental contamination by contaminants in animal

BIOLOGICAL MATERIALS EXPERIMENTALS CONDITIONS

CONTAMINANTS

EFFECTS ON TELOMERE DYNAMICS

OTHERS BIOLOGICALS EFFECTS

SOURCES

FISHS

Epinephelus coioides (Orange-spotted grouper) Muscle (n=5)	Bio-assays	B[a]P : 12 μg/g de poissons/j since 1, 2 and 4 weeks	↓ telomerase activity	Association with B[a]p exposure	Peterson et al., 2015
		BIRDS			
Haliaeetus albicilla (White-tailed eagle) nestlings Red blood cells (n=35)	Biomonitoring <i>in situ</i> (Troms County, Northern Norway)	9 PCBs, p,p'DDE, HCB, oxy-chlordane, cis-chlordane, trans-nonachlor, cis- nonachlor	O Telomere length	\downarrow SOD activity, Δ IgY levels	Sletten et al., 2016
Parus major (Great tit) nestlings and adults females Livers cells (n=24 and n=21)	Biomonitoring <i>in situ</i> (Cu/Ni smelter in Harjavalta, Finland)	ME (Cu, Ni, As, Pb, Cd maj.)	↓ Telomere length (nestlings), O Telomere length (adults females)	↑ [As, Pb], ↓ SOD activity (nestlings), ↑ GPx activity, ↑ [Ca], ↓ lipoperoxidation	Stauffer et al., 2017
<i>Ficedula hypoleuca</i> (European pied flycatcher) nestlings and adults females Livers cells (n=22 and n=23)	Biomonitoring <i>in situ</i> (Cu/Ni smelter in Harjavalta, Finland)	ME (Cu, Ni, As, Pb, Cd maj.)	O Telomere length	↑ [As, Pb], ↓ Gr and GST activities, ↓ lipoperoxidation (nestlings),	Stauffer et al., 2017
<i>Rissa tridactyla</i> (Black-legged kittiwake) Red blood cells (n=38)	Biomonitoring <i>in situ</i> (Krvkkiefiellet colony of Kongsfjorden, Svalbard)	PCBs, p,p'DDE, HCB, cis-nonachlor, trans-nonachlor,[Oxychlordane] between 2000 to 7000 pg/g ww	↓ Telomere length (adults females), O Telomere length (adults males)	Association with [oxychlordane]	Blévin et al., 2016
<i>Rissa tridactyla</i> Red blood cells (n=55)	Biomonitoring <i>in situ</i> (Krvkkiefiellet colony of Kongsfjorden, Svalbard)	PFAs	↑ Telomere length (in both sexes)	Association with [PFAs]	Blévin et al., 2017
		MAMMAL	S		
Rattus norvergicus (Sprague-Dawley rats) Fibroblast-type cell line (ADIPO-P2)	18h to 15 days <i>in vitro</i>	Sodium citrate (20 mM) and Streptozotocin (2 mM)	↓ Telomere length, ↑ Telomere duplication, ↑ Translocation of telomeric sequences, ↑ Additional telomeric FISH signals, ↑ extra- chromosomal telomeric FISH signals	Association with chromosomic aberrations	Paviolo et al., 2015
Ctenomys torquatus (Collared tuco-tuco) kidney, muscle and skin tissues (n=30 control, n=36 exposed)	Biomonitoring <i>in situ</i> (Butia, Candiota and Pelotas in Brazil)	Coal (PAH, ME)	↓ Telomere length	↑ DNA damage, O global DNA methylation	Matzenbacher et al., 2019

Legends: Δ = variations, \uparrow = increasing, \downarrow = decreasing, O = no effects