



**HAL**  
open science

# The triggering factors of the Móafellshyrna debris slide in northern Iceland: Intense precipitation, earthquake activity and thawing of mountain permafrost

Þorsteinn Sæmundsson, Costanza Morino, Jón Kristinn Helgason, Susan J. Conway, Halldór G Pétursson

## ► To cite this version:

Þorsteinn Sæmundsson, Costanza Morino, Jón Kristinn Helgason, Susan J. Conway, Halldór G Pétursson. The triggering factors of the Móafellshyrna debris slide in northern Iceland: Intense precipitation, earthquake activity and thawing of mountain permafrost. *Science of the Total Environment*, 2018, 621, pp.1163-1175. 10.1016/j.scitotenv.2017.10.111 . hal-02270621

**HAL Id: hal-02270621**

**<https://hal.science/hal-02270621>**

Submitted on 26 Aug 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The triggering factors of the Móafellshyrna debris slide in northern Iceland: intense precipitation, earthquake activity and thawing of mountain permafrost.

Þorsteinn Sæmundsson<sup>(a)</sup>, Costanza Morino<sup>(b)</sup>, Jón Kristinn Helgason<sup>(c)</sup>, Susan J. Conway<sup>(d)</sup>, Halldór G. Pétursson<sup>(e)</sup>.

(a) Department of Geography and Tourism, University of Iceland, Askja, Sturlugata 7, 101 Reykjavík, Iceland

(b) School of Environment, Earth & Ecosystems, The Open University, Milton Keynes MK7 6AA, UK

(c) Icelandic Meteorological Office, Avalanche Centre, Suðurgata 12, 400 Ísafjörður, Iceland

(d) Laboratoire de Planétologie et Géodynamique de Nantes UMR-CNRS 6112, 2, rue de la Houssinière, BP 92208, 44322 NANTES Cedex 3, France

(e) Icelandic Institute of Natural History, Borgum við Norðurslóð, 602 Akureyri, Iceland

## **Abstract**

*On the 20<sup>th</sup> September 2012, a large debris slide occurred in the Móafellshyrna Mountain in the Tröllaskagi peninsula, central north Iceland. Our work describes and discusses the relative importance of the three factors that may have contributed to the failure of the slope: intense precipitation, earthquake activity and thawing of ground ice. We use data from weather stations, seismometers, witness reports and field observations to examine these factors. The slide initiated after an unusually warm and dry summer followed by a month of heavy precipitation. Furthermore, the slide occurred after three seismic episodes, whose epicentres were located ~60 km NNE of Móafellshyrna Mountain. The main source of material for the slide was ice-rich colluvium perched on a topographic bench. Blocks of ice-cemented colluvium slid and then broke off the frontal part of the talus slope, and the landslide also involved a component of debris slide, which mobilised around 312 000 - 480 000 m<sup>3</sup> (as estimated from field data and aerial images of erosional morphologies). From our analysis we infer that intense precipitation and seismic activity prior to the slide are the main preparatory factors for the slide. The presence of ice-cemented blocks in the slide's deposits leads us to infer that deep thawing of ground ice was likely the final triggering factor. Ice-cemented blocks of debris have been observed in the deposits of two other recent landslides in northern Iceland, in the Torfufell Mountain and the Árnesfjall Mountain. This suggests that discontinuous mountain permafrost is degrading in Iceland, consistent with the decadal trend of increasing atmospheric temperature in Iceland. This study highlights a newly identified hazard in Iceland: landslides as a result of ground ice thaw. Knowledge of the detailed distribution of mountain permafrost in colluvium on the island is poorly constrained and should be a priority for future research in order to identify zones at risk from this hazard.*


## 37 **1. Introduction**

38 On the 20<sup>th</sup> September 2012, the local residents of the Þrasastaðir farm in the Fljótin  
39 area (Tröllaskagi peninsula, central north Iceland; Figs. 1,3), heard a loud rumbling noise  
40 coming from the Móafellshyrna Mountain, as a large debris slide descended the north-  
41 western side of the mountain (Fig. 3). Nine days after the failure, we observed large  
42 blocks of ice-cemented deposits within the debris slide deposits.

43 Rapid mass movements, including rock falls, rock avalanches, debris flows and debris  
44 slides, are common geomorphological processes in Iceland and present a significant and  
45 direct threat to many towns, villages and farmhouses (Decaulne 2005; 2007).

46 Precipitation, snow melt, temperature variations and earthquake activity are the most  
47 common triggering factors for landslides in Iceland (Sæmundsson *et al.*, 2003;  
48 Sæmundsson and Decaulne, 2007, Decaulne & Sæmundsson 2007). However, during  
49 the last decade, three, somewhat unusual, rapid mass movements have occurred in  
50 northern Iceland: on the Torfufell Mountain (Eyjafjörður valley, central north Iceland) on  
51 14<sup>th</sup> October 2011, on the Móafellshyrna Mountain on 20<sup>th</sup> September 2012 (case study  
52 of this paper), and on the Árnesfjall Mountain (Westfjords) on 10<sup>th</sup> July 2014 (Sæmunds-  
53 son *et al.*, 2013; Sæmundsson *et al.*, 2014 a/b). In all these landslides ice-cemented  
54 debris was found within the deposits, a phenomenon that has never been reported  
55 previously in Iceland. The source areas of these slides are all located on steep (~45-60°)  
56 NW to NE facing-slopes, where discontinuous permafrost might be expected (e.g.  
57 Gorbunov, 1988; King, 1986). The source areas at Torfufell and Móafellshyrna are  
58 located at elevations of 750-870 m a.s.l., within the zone of discontinuous permafrost in  
59 Iceland as calculated by Etzelmüller *et al.* (2007a), whereas at Árnesfjall the source area  
60 is located at about 400 m a.s.l., which is much lower altitude than ever observed for  
61 mountain permafrost in Iceland (Etzelmüller *et al.*, 2007a). Intense precipitation was  
62 recorded prior to all of these slides (Sæmundsson *et al.*, 2013; Sæmundsson *et al.*, 2014  
63 a/b). In this paper, we present the case study of the Móafellshyrna debris slide, where we  
64 examine three factors could have contributed to this failure: intense precipitation,  
65 earthquake activity and ground-ice degradation (via increased annual temperatures). We  
66 emphasise the field evidence of ground ice-thaw, because permafrost degradation has  
67 never previously been considered as a triggering factor for gravitational mass  
68 movements in Iceland, although permafrost degradation is considered to be an  
69 preparatory factor for paraglacial slope failure (McColl 2012).

70 In the following sections we describe a) the state of knowledge of permafrost in Iceland,  
71 b) seismic conditions in Iceland and their role in previous mass wasting events, c)  
72 general meteorological conditions in Iceland. We then report our results, reconstructing  
73 the conditions that favoured the occurrence of the landslide. We then discuss our results,  
74 dividing the factors that induced the landslide into, i) preparatory and ii) triggering (as per  
75 McColl, 2012). We finally posit that Móafellshyrna landslide was preconditioned by  
76 combination of intense precipitation in the weeks prior to the slide and the seismic activity  
77 on 18<sup>th</sup> and 19<sup>th</sup> September and that the degradation of ground-ice was the final trigger.



Station name	Station ID	Altitude	Years analysed	Precipitation	Temperature
Siglufjörður	SIFJO	6.0 m a.s.l.	2012	X	X
Öxnadalshéiði	OXNAD	540.0 m a.s.l.	2005 - 2012	-	X
Ólafsfjörður	OLFJO	5.0 m a.s.l.	2000 - 2012	X	X

79 Figure 1 – The Móafellshyrna site, located in the Tröllaskagi peninsula, northern Iceland (see Fig. 3 for  
80 detailed location), and the location of the weather stations used for this study. The hillshaded digital  
81 elevation model used as a basis of this map is from the Digital Elevation Model over Europe (EU-DEM)  
82 from the Global Monitoring for Environment and Security service for geospatial reference data access  
83 project (GMES RDA). The table provides details on the Icelandic Meteorological Office weather stations,  
84 whose datasets have been used for this study. Symbols “X” and “-“ mean that data are or are not available  
85 at the stations, respectively.

## 86 **1.1 Permafrost distribution in Iceland**

87 During the last decade our knowledge of the regional distribution of permafrost in Iceland  
88 has increased considerably (e.g. Etzelmüller *et al.*, 2007a/b; Farbrot *et al.*, 2007a/b;  
89 Kellerer-Pirklbauer, 2007; Lilleören *et al.*, 2013; Kneisel *et al.*, 2007; Kneisel, 2010).  
90 There are several published works on permafrost in the central highlands of Iceland  
91 dating back to the 1950s, and these studies focused on geomorphological features, such  
92 as palsas and patterned ground (e.g., Thorarinsson, 1951; 1964; Friedman *et al.*, 1971;  
93 Schunke, 1974; Stingl and Herrmann, 1976; Priesnitz and Schunke 1978; Hirakawa,  
94 1986; Thorhallsdottir, 1994; 1996; 1997; Sæmundsson *et al.*, 2012). These works both  
95 mapped permafrost conditions and a recent study at the Orravatnsrústir palsa site, NE of  
96 the Hofsjökull ice cap, showed clear indications of declining permafrost conditions from  
97 2000-2010 in the highlands (Sæmundsson *et al.*, 2012). Studies have also sought to  
98 better define the spatial distribution of mountain permafrost in Iceland. Such studies used  
99 inventories of rock glaciers and stable ice-corded moraines combined with meteorological  
100 data analyses (e.g., Etzelmüller *et al.*, 2007a/b; Farbrot *et al.*, 2007a/b; Kellerer-  
101 Pirklbauer, 2007; Lilleören *et al.*, 2013) and Electrical Resistivity Tomography (e.g.,  
102 Kneisel *et al.*, 2007; Kneisel, 2010; Sæmundsson *et al.*, 2012). These works did not  
103 comment on the dynamics of the permafrost.


104 Etzelmüller *et al.* (2007a) published the first regional distribution map of mountain perma-  
105 frost in Iceland. This was the first attempt to understand the overall extent of mountain  
106 permafrost on the island. The map was based on a model which used meteorological  
107 data calibrated with ground surface temperature data and validated using ground  
108 temperature data from four shallow boreholes and an inventory of rock glaciers.  
109 According to Etzelmüller *et al.* (2007a), mountain permafrost is widespread in the  
110 northern and eastern Iceland above 800-900 m, covering 8% of the island, or around  
111 8.000 km<sup>2</sup>.

112 A warming trend in atmospheric temperature has been observed in Iceland over the last  
113 two centuries. In Stykkishólmur in western Iceland records show an increase of about  
114 0.7°C per century (Nawri & Björnsson, 2010; Jónsson, 2008). From 1975 to 2008 the  
115 warming rate was much higher or about 0.35°C per decade (Ministry for the Environment  
116 of Iceland, 2010). However, so far relatively little attention has been paid to the  
117 consequences of the recent climate change on the possible degradation of mountain  
118 permafrost in Iceland.

## 119 **1.2 Seismic activity in northern Iceland**

120 The seismic activity in Iceland is related to its position on the Mid-Atlantic plate boundary,  
121 which crosses the island and its location over the Icelandic Hotspot (e.g. Tryggvason *et*  
122 *al.*, 1983; Wolfe *et al.*, 1997; Allen *et al.*, 2002; Bjarnason, 2008; Einarsson, 2008;  
123 Thordarson & Höskuldsson 2002). The seismic activity in the northernmost region of the  
124 island is related to the Tjörnes Fracture Zone (Einarsson and Björnsson, 1979;  
125 Sæmundsson, 1974), which is defined by three seismically active lineaments - the  
126 Grímsey Oblique Rift, the Húsavík Flatey Fault and the Dalvík lineament; (e.g.,  
127 Sæmundsson, 1974; Gudmundsson, 2000; 2006) - and includes the Eyjafjarðaráll N-S  
128 extensional graben, located north offshore the Tröllaskagi peninsula (Fig. 2). It was  
129 activity on this fault system which caused the earthquakes prior to the Móafellshyrna  
130 slide.

131 Earthquakes are known to have triggered landslide and rockfall activity in Iceland in the  
132 past (e.g. Jónsson, 1957; Jónsson and Pétursson, 1992; Thorarinsson, 1937, 1959;  
133 Halldórsson, 1984; Jensen, 2000; Sæmundsson *et al.*, 2003; Ágústsson and Pétursson,  
134 2013), but no study has explored in detail the influence of earthquake activity as a  
135 preparatory and/or triggering factor on rockfall or landslides in Iceland, as has been done  
136 elsewhere in the world (e.g., Harp and Jibson, 1996; Yin *et al.*, 2009). The above  
137 mentioned Icelandic studies relate mass movements to larger earthquakes than those  
138 prior to the slide in Móafellshyrna, e.g. in June 2000, when two earthquakes of magnitude  
139 6.4 occurred in Iceland, with the epicentre in the middle of the southern lowlands.  
140 Rockfall activity was reported as a result of this event as far as 75 km from the epicentre  
141 (Sæmundsson *et al.*, 2003).



142  
 143 Figure 2 - The structural elements of the Tjörnes Fracture Zone marked in black (Grímsey lineament  
 144 (GOR), Húsavík-Flatey fault (HFF) and Dalvík lineament (DL); from Stefánsson *et al.*, 2008) and the  
 145 Eyjafjarðaráll graben marked in red (EY); the position of the epicenter zone for the earthquakes preceding  
 146 the Móafellshyrna slide is marked with a yellow star, the Icelandic Meteorological Office (IMO)  
 147 seismometers in the area marked with the green dots and the labels refer to their abbreviated names, as  
 148 given in full in the key. On the right are reported the timing and magnitude of the earthquake sequence in  
 149 the Eyjafjarðaráll graben from 19<sup>th</sup> to 20<sup>th</sup> September 2012 (from Gudmundsson *et al.*, 2014).

150 **1.3 General weather conditions in Iceland**

151 Weather patterns in Iceland are highly variable, with frequent and strong variation in  
 152 precipitation and temperature; this is mainly because Iceland is located on the main path  
 153 taken by North Atlantic low-pressure systems (Einarsson 1984). The mean annual air  
 154 temperature for the period 1971–2000 was 4–5°C in the south, 3–4°C in the east and  
 155 west and 2–3°C in northern coastal parts of the country (Tveito *et al.*, 2000). Hence  
 156 precipitation can fall as both snow and rain. The two main dominant precipitation wind  
 157 direction in the Tröllaskagi area are NE and SW (Brynjólfsson and Ólafsson 2008,  
 158 Arnalds *et al.*, 2001). The precipitation is heaviest during strong NE winds.  
 159 Consequently, mean annual precipitation increases from about 500 – 1000 mm per year  
 160 in the central and northern parts of the country to more than 3000 mm/yr in the southeast  
 161 (Crochet *et al.*, 2007). During the winter months from October to April the precipitation in  
 162 the outer part of the Tröllaskagi peninsula is almost exclusive snow or sleet and the main  
 163 part of the snow avalanche activity is associated with strong north-easterly wind. The  
 164 northern part of Tröllaskagi peninsula is generally a heavy snow prone area and  
 165 Siglufjörður and Fijótin area are generally considered to be one of the heaviest snow  
 166 prone areas in Iceland (Arnalds et al. 2001).



## 167 **2. Geographic and Geologic Setting of Móafellshyrna**

168 The Tröllaskagi peninsula is a mountain massif located between the Eyjafjörður fjord in  
169 the east and the Skagafjörður fjord in the west (Fig. 1). The peninsula is topped by flat  
170 summits reaching up to 1000 – 1500 m a.s.l. and sculptured by glacial erosion with  
171 glacially carved fjords, valleys and cirques. Such over-steepened landscapes are  
172 recognised as one of the key pre-conditioning factors for failures (McColl 2012). Over  
173 150 alpine glaciers, mainly north facing, have been mapped in this massif (Sigurðsson  
174 and Williams, 2008).


175 The bedrock in the outermost part of the Tröllaskagi peninsula falls within the Tertiary  
176 basalt series (16-3.3 M years; e.g., Moorbath *et al.*, 1968; McDougall *et al.*, 1984;  
177 Watkins and Walker, 1977). This series is mostly jointed basaltic lava flows, composed of  
178 2 to 30 m thick individual flows separated by lithified sedimentary horizons (from few  
179 centimetres up to tenths of meters thick) (Sæmundsson 1979; Sæmundsson *et al.*, 1980).  
180 The bedrock is heavily jointed and intersected by dikes, and the general dip angle of the  
181 lava beds is towards the southwest (Sæmundsson *et al.*, 1980; Johannesson and  
182 Sæmundsson, 2009; Hjartarson & Sæmundsson 2014).

183 The Tröllaskagi peninsula is located on the main path of the North Atlantic low pressure  
184 system, with a variable and turbulent weather. Winters are long and mild and summers  
185 brief and cool (Einarsson, 1984). The Mean Annual Air Temperature (MAAT) for the  
186 period 1971-2000 was 2-3°C in the northern coastal areas (Tveito *et al.*, 2000; Crochet *et*  
187 *al.*, 2007). The data series for the Tröllaskagi area between 1940-1970 show MAAT of 2-  
188 4°C in the coastal areas and -2 to -4°C at the summits (Einarsson, 1964). According to  
189 Crochet *et al.* (2007), the annual precipitation from 1971-2000 in the Tröllaskagi area  
190 varies from 1000-1500 mm in the coastal lowlands up to 2000-2500 mm on the summits.  
191 Localised orographic effects mean that in the Tröllaskagi peninsula precipitation is higher  
192 near the coastline when there are northerly winds (Brynjólfsson and Ólafsson 2008).  
193 Conversely, the precipitation is likely to be higher in the lowlands in the interior of the  
194 Tröllaskagi peninsula area than at the coast during periods with southerly winds. In the  
195 mountains of the peninsular the orographic effect would also play a role during southerly  
196 winds precipitation may in fact be even higher at a higher elevation. According to Arnalds  
197 *et al.* (2001), more than half of the recorded precipitation in Siglufjörður occurs  
198 throughout the winter months (Oct-Apr), where the precipitation is almost exclusively  
199 snow or sleet. This can also be said about the rest of Tröllaskagi peninsula. Avalanches

200 during winter months in Tröllaskagi peninsula can be associated with strong north-  
201 easterly winds with snowfall or snowfall during south-westerly winds associated with  
202 lower wind speed. Slush flows are not common in Tröllaskagi area.

203 The Móafellshyrna Mountain is located in the Fljótin area in the outermost part of the  
204 Tröllaskagi peninsula in central north Iceland (Figs.1, 3). The mountain is 1044 m high,  
205 with narrow alpine-mountain ridge orientated in the north-south direction. It is located  
206 between the Móafellsdalur tributary valley in the west and the Hvarfdalur tributary valley  
207 in the east (Fig. 3). In the outermost part of the western valley side of Móafellsdalur are  
208 old remnants of a landslide that originated from the western mountain side. The age of  
209 these old landslide deposits is not known. Jónsson (1957, 1976) stated that the landslide  
210 activity in Iceland was most intense shortly after the last deglaciation. His statement has  
211 been confirmed by later studies (e.g. Jonson *et al.* 2004, Sigurgeirsson and Hjartarson  
212 2011, Mercier *et al.* 2012 and 2013, Coquin *et al.* 2015 and 2016, Decaulne *et al.* 2016).  
213 The farm Þrasastaðir, the innermost inhabited farm in the Stífludalur valley, is situated on  
214 the northern side of the Stífludalur on the northern side of the Fljótaá River facing the  
215 Móafellshyrna mountain. The residents of the farm witnessed and observed the slide  
216 (Fig. 3) and their report is given in Section 4.1.

217 The landscape in the Fljótin area is predominantly carved by glacial erosion. The area  
218 was heavily glaciated during the maximum extent of the last glaciation, but the  
219 deglaciation history of the area is not well documented. Researchers suggest that the  
220 area was deglaciated in Early Preboreal times (e.g., Norðdahl and Pétursson, 2005;  
221 Norðdahl *et al.*, 2008; Ingólfsson *et al.*, 2010; Coquin *et al.* 2016). The upper part of the  
222 valley sides (up to 600-900 m a.s.l.) are often very steep, with fractured and loose  
223 bedrock produced by, erosion during deglaciation, unloading by the glacier, frost  
224 shattering and freeze-thaw processes, while the lower parts are more gentle and  
225 generally covered by glacial deposits or talus.



226  
 227 Figure 3 – Geographical setting of MÓafellshyrna mountain (see Fig. 1 for location). (A) Hillshaded digital  
 228 elevation model and contours (in green, metres above sea level) of the MÓafellshyrna region. Elevation  
 229 data are from EU-DEM from GMES RDA. (B) Aerial photograph of the MÓafellshyrna site taken before the  
 230 slide in 2012 from the website Samsyn. (C) Oblique photo of the MÓafellshyrna debris slide in July 2015  
 231 taken by Costanza Morino. In panels B and C the perimeter of the slide is marked with red dashed line  
 232 taken from the trimble data for the deposits and reconstructed from photographs for the upper part, in C  
 233 this line has been manually traced onto the oblique image.

234

### 235 **3. Methods**

236 The triggering factors of the MÓafellshyrna debris slide event in September 2012 were  
 237 analysed using meteorological data from the Icelandic Meteorological Office (IMO) for the  
 238 three months prior to the slide and for the period 2010-2012. Data for the seismic activity  
 239 of the north coast were also obtained from the IMO. We also interviewed the inhabitants  
 240 of the valley and performed our own field investigations.

#### 241 **3.1 Meteorological data**

242 Only two weather stations in the northern part of the Tröllaskagi peninsula measure  
 243 precipitation, one located in the town of Siglufjörður at 6 m a.s.l. (35 km north of the site)

244 and the other one in the town of Ólafsfjörður at 5 m a.s.l. (21 km northeast of the site).  
245 Three weather-stations were used for this study that collect temperature records: two  
246 operated by IMO - Siglufjörður (WMO (World Meteorological Organization) ID: 4157),  
247 Ólafsfjörður (WMO ID: 4155) and one station located at the Öxnadalsheiði highlands  
248 pass (WMO ID: 4859) at 540 m a.s.l. operated by the Icelandic Road and Coastal  
249 Administration (Fig. 1). A problem in our approach is that the majority of the stations used  
250 to establish mean atmospheric temperatures are located near the coast and therefore at  
251 low altitudes. This leads to a potential bias when evaluating trends in temperature,  
252 because such stations may not be representative of the atmospheric temperatures  
253 experienced in the highlands. To overcome this, we applied the environmental lapse rate  
254 of 0.649°C per 100 m (Sheridan *et al.* 2010) to the mean temperatures recorded at all  
255 three stations as an estimate of the temperature at the source zone of the Móafellshyrna  
256 debris slide. We do not attempt to correct the precipitation data collected near the coast  
257 for the inland conditions, because this would not only require a temperature correction,  
258 but we would need to take into account variations in wind speed, wind direction and  
259 pressure in a meteorological model, which is beyond the scope of this paper.

### 260 **3.2 Direct report from witnesses**

261 The local residents of the farm Prasastaðir witnessed the release of the landslide, and  
262 were interviewed on the day of the event regarding the earthquake activity prior to the  
263 slide, timing of the slide and the events that occurred during the first few hours of the  
264 slide. The slide was photographed by the first author only a few hours after it had  
265 occurred. Ongoing rockfall activity on the slope prohibited more detailed observations  
266 immediately after the event. It was not considered safe to perform field analysis until 29<sup>th</sup>  
267 September 2012, when direct observations and measurements were made. The site was  
268 revisited in summer 2015 to track changes in the slide.

### 269 **3.3 Field measurements and survey**

270 The boundary of the landslide body was mapped on the 29<sup>th</sup> June 2012 with Trimble  
271 GEOXT from the GeoExplorer CE series with an accuracy of 1-2 m. Thickness  
272 measurements of the slide were performed both with direct measurements in the field on  
273 29<sup>th</sup> September 2012 and from aerial photographs and erosional and depositional  
274 features mapped. For geomorphological mapping both ground photographs and aerial  
275 photographs from the National Land Survey of Iceland ([www.lmi.is](http://www.lmi.is)) and Loftmyndir ehf  
276 ([www.map.is](http://www.map.is)) and the UK's Natural Environment Research Council Airborne Research

277 and Survey Facility (NERC ARSF) flown in 2015 were used in concert with field  
278 observations. Landslide volume estimates were derived from morphometric properties of  
279 the deposits measured directly in the field and from aerial photographs.

## 280 **4. Results**

281

### 282 **4.1 Witness report**

283 The residents of the Þrasastaðir farm, located at the junction between the Móafellsdalur  
284 and the Stífludalur valleys and 1.7 km from the terminal deposits of the landslide (Fig. 3),  
285 were interviewed only few hours after the slide. They recounted that on the 20<sup>th</sup> of  
286 September 2012 at around 12:30 they heard a rumbling noise, which originated from the  
287 Móafellshyrna mountain. They also recounted that a black tension crack, in the snow  
288 covered mountain, progressively formed above the colluvial cone at around 850 m a.s.l.  
289 They saw large blocks of debris that broke off the frontal part of the cone and fell onto the  
290 talus slope below. This activity was most intense in the first 1-1.5 hours, but they reported  
291 that there were intermittent noises and rock fall activity throughout the day.

292 The residents of the Þrasastaðir farm felt all the three earthquakes that occurred on the  
293 19<sup>th</sup> and 20<sup>th</sup> September, with the last one only three hours before the debris slide event.  
294 It was estimated that less than 1 m of snow was on the ground at the time of the slide.


### 295 **4.2. The debris slide morphology**

296 The debris slide detached from a main scarp at an elevation of 880 m a.s.l. (Fig. 4A). The  
297 source material of the landslide was a colluvial cone composed of talus deposits perched  
298 on a topographic bench at 790 m a.s.l. (Fig. 3C), and lying against a rockwall composed  
299 of lava and tephra layers of the Tertiary Basalt formation dipping less than 10° towards  
300 SSW. The colluvial material comprises a mixture of materials derived from the above  
301 rockwall (basalt with intercalated sediment layers) and under nominal conditions (with or  
302 without ice) should lie stable against the rockwall under the angle of repose.

303 The colluvial material partially slid off the steep rockwall ledge at the edge of the bench,  
304 as described by the local residents. From field observations and aerial photographs we  
305 estimate that the horizontal displacement in the upper part of the cone was around 40 m  
306 (Fig. 4A). Later observations revealed that the bedrock was exposed from beneath the  
307 colluvium as it slid (at the time of the slide it was hidden by a thin veneer of mud, which  
308 was later cleaned by rainfall). Part of the colluvial deposits remained perched on the

309 topographic bench after the event (Fig. 3C, 4B). Based on field observation, we  
310 estimated that the thickness of the frontal part of the still ice-cemented colluvial cone that  
311 was preserved at the edge of the bench after the slide was around 20-30 m thick (Fig.  
312 4B). As the colluvial cone slid off the rockwall edge, large blocks of ice-cemented debris  
313 broke off and fell onto the talus slope located below the topographic bench (Fig. 3C). Fig.  
314 3B shows decametre-scale blocks of ice cemented colluvium that have remained on the  
315 bench. A 300 m long, 80-100 m wide and 8-10 m deep channel was carved into the talus  
316 slope deposits (Fig. 3C, 4C), entraining further material to the landslide. This channel  
317 shows that the landslide materials must already have been fluidised (saturated) in order  
318 to scour out this channel.

319 At 495-505 m a.s.l. at the foot of the talus slope, we observed that both loose muddy  
320 debris and blocks of ice-cemented deposits comprised the landslide deposits (Figs.  
321 4C,D,E). We infer that these ice cemented blocks had fallen from the topographic bench,  
322 as they had the same aspect and size of those still perched in the source area (Fig. 4B).  
323 These blocks ranged from less than 1 m up to 12 m wide and from less than 1 m up to 10  
324 m tall. The ice-cemented blocks were composed of layers of sand to boulder-size (up to  
325 50 cm in long-axis) angular rock fragments (Fig. 4F), interbedded with layers composed  
326 of clay to sand-size materials. The red colour of the horizons of fine material (Fig. 4E)  
327 derives from the original source material, namely paleo-soils interbedded in the lava  
328 layers composing the Tertiary Basalt Formation. Nine days after the failure, from visual  
329 inspections we estimated a content of ground ice that was cementing the blocks of  
330 around 15-20% of the total volume (Fig. 4F), i.e. pore-filling ice, rather than excess ice.  
331 No massive ice deposits were observed.



333 Figure 4 – (A) An overview of the western side of the Móafellshyrna Mountain after the debris slide. The  
334 uppermost and lowermost elevation limits of the debris slide path are labelled in m a.s.l. as are the  
335 locations of panels B to G. (B) The ice-cemented blocks perched, up to 20 m in thickness on the 100 m  
336 high rockwall. (C) The erosional area in the lower talus slope below the rockwall. Note the large block of  
337 ice-cemented deposits on the right. (D) Landslide debris and an ice-cemented block resting at 495-505 m  
338 a.s.l. (E) The same large block of ice-cemented deposits as in D, around 12 m wide and 4 m high. (F)  
339 Close up of the block in E showing stratified deposits of coarse angular clasts with an icy matrix. (G)  
340 Looking downwards onto the deposits in the terminal part of the slide. Photos: Þ. Sæmundsson, A taken on  
341 the 20<sup>th</sup> September and B-G taken on the 29<sup>th</sup> September 2012.

342 Some of the landslide deposits came to rest below the talus slope at 495-505 m a.s.l.  
343 (Fig. 4D), where the average slope angle is less than 10°, but the rest traversed down the  
344 mountain slope, finally stopping at 330 m elevation (Fig. 4G), a few meters from the river  
345 Galtará draining the bottom of the valley (Fig. 3). The part of the material that continued  
346 down the mountain side is composed of clay to boulder-size material which followed a  
347 well-defined path with distinct lateral boundaries. The deposit boundaries are upstanding  
348 with 1-2 m of relief. The lack of sorting in the deposits, their lobate planform and their  
349 upstanding boundaries are consistent with this landslide being classed as a debris side  
350 transforming to a debris avalanche from the Hungr *et al.* (2001, 2014) classification, i.e.  
351 loose debris fluidised by the inclusion of water exceeding the pore-space (Iverson 1997)  
352 without a well-defined central channel and generally shallow motion. This shows that  
353 water volumes of 20-40% of the final deposits were required to mobilise this flow and that  
354 it was not a dry rock avalanche.

355 Based on aerial photographs the dimensions of the upper colluvial cone perched on the  
356 bench was around 80 m in height and around 150 m wide at the lower end prior to the  
357 slide. Based on photographs taken after the slide the total movement of the cone was  
358 estimated around 40 m and the frontal part of the cone perched on the bench was 150 m  
359 wide and about 20-30 m thick (Fig. 4B). Based on these dimensions we estimated that  
360 the volume of the deposits that broke off the frontal part of the cone and fell down onto  
361 the lower talus had a volume up to ~120 000 – 180 000 m<sup>3</sup>. From the dimensions of the  
362 channel carved into the lower talus on the southern side of the slide (with 80-100 m,  
363 depth 8-10 m, length 300 m) (Fig. 4C) we calculated that an additional volume of ~192  
364 000 - 300 000 m<sup>3</sup> of material was mobilized. Combining these volumes gives an estimate  
365 of the total volume of material mobilized in the event of 312 000 - 480 000 m<sup>3</sup>. From GPS  
366 measurements the areal extent of the landslide is ~0.3 km<sup>2</sup>, giving an average depth of



367 deposits of 1-2 m, consistent with our direct observations of thickness at the deposit-  
368 boundaries.

### 369 **4.3 Antecedent conditions**

#### 370 *4.3.1 Precipitation*



371 The spring and summer months preceding the Móafellshyrna event were dry, and the  
372 autumn was unusually wet (Figs. 5, 6) (Jónsson 2013). From April until 28<sup>th</sup> August 2012,  
373 dry conditions prevailed in the outer part of the Tröllaskagi peninsula, with only one day  
374 with precipitation greater than 10 mm: 23<sup>rd</sup> to 24<sup>th</sup> July, when 70 to 90 mm of rain was  
375 recorded at Siglufjörður and Ólafsfjörður weather stations (Fig. 7).

376 From 20<sup>th</sup> August to 20<sup>th</sup> September around 1/3 of the precipitation for 2012 fell in the  
377 area (~400-550 mm). For comparison, the average annual precipitation in the town of  
378 Ólafsfjörður is ~400 mm for the period 2000-2012 (Figs. 5 & 6). In detail, from 28<sup>th</sup> August  
379 to 8<sup>th</sup> September the cumulative precipitation in Siglufjörður was 190 mm and 120 mm in  
380 Ólafsfjörður, with an additional 30 to 40 mm precipitation at these stations on the 3<sup>rd</sup>, 6<sup>th</sup>  
381 and 7<sup>th</sup> September. From 9<sup>th</sup> to 11<sup>th</sup> September an unseasonal and severe snowstorm hit  
382 the north eastern and northern parts of the country (Jónsson 2013, Hermannsdóttir  
383 2012). Following this snowfall, around 100 mm rain was measured in only two days at  
384 Siglufjörður and almost 150 mm at Ólafsfjörður. The precipitation continued from 11<sup>th</sup> to  
385 17<sup>th</sup> September either as snow, sleet or rain at these weather stations. From September  
386 17<sup>th</sup> to 19<sup>th</sup> less than 10 mm of precipitation was recorded, but at the time of the event  
387 540 mm precipitation had been recorded in Siglufjörður and 490 mm at Ólafsfjörður  
388 weather stations since 23<sup>rd</sup> July, which corresponds to 40-45% of the mean annual  
389 precipitation from 2000-2012 (Fig. 5). The monthly precipitation data from Ólafsfjörður  
390 station from 2000 to 2012 show that September is the month with maximum precipitation  
391 for any given year, with a range between 70 and 250 mm (Figs. 5 & 6). The year of 2012,  
392 however, had precipitation exceeding the average for this month (Fig. 5)


393 Unfortunately, there is no weather station located in the mountainous region of  
394 Tröllaskagi area and as previously mentioned there is no weather station in Fljótin area.  
395 This means that we cannot report absolute precipitation data for the Móafellshyrna site,  
396 but it is reasonable to assume that on a month by month basis the trends should be  
397 similar to those of surrounding weather stations. Predicting whether precipitation falls as  
398 snow or rain at Móafellshyrna is outside the scope of this study and is complicated by a

399 number of factors including snow drifting, wind-dependant snowmelt and variable  
400 orographic effects dependant on wind direction (Brynjólfsson and Ólafsson 2008) – see  
401 Section 1.3. Snow was visible on the ground on the day of the slide and was less than  
402 1m thick from eye witness accounts (equivalent to ~100 mm of precipitation, depending  
403 on snow density) Therefore, the majority of the precipitation received up to 20<sup>th</sup>  
404 September had been absorbed by the ground. The fact that snow was present on the  
405 ground argues against a sudden influx of water into the ground via snowmelt, known to  
406 trigger other mass wasting phenomena in Iceland (e.g. Decaulne et al. 2005). Hence, we  
407 did not pursue an analysis of the wind data from the weather stations, because this would  
408 only be important if melt or precipitation were the primary triggers for the slide.




409



410  
 411  
 412 Figure 5 – Matrix plots of the difference between the average monthly temperature (top)  
 413 and precipitation (bottom) and the average value for that month for the period 2000-2012  
 414 for the Ólafsfjörður station (Data supplied by the IMO in 2016).



415  
 416 Figure 6 – A boxplot of the precipitation data from the Ólafsfjörður station for each month between the  
 417 years 2000 to 2012 (Data from IMO 2016). The end of the dotted lines is where the max and min values of  
 418 precipitation were measured for each month, excluding the outliers that are displayed as dots. The blue  
 419 boxes is where the 50% of accumulated measured precipitation falls and the black lines are the medians of  
 420 each month.





423 Figure 7- Top: daily precipitation (black bars), cumulative precipitation (blue lines) for temperature  
424 measurements from the Siglufjörður (SIFJO) and Ólafsfjörður (OLFJO) weather stations. Middle: daily  
425 temperature data from the Siglufjörður (SIFJO), Ólafsfjörður (OLFJO) and Öxnadalsheiði (OXAN) weather  
426 stations. Bottom: mean temperature data for all three stations adjusted to 880 m altitude of Móafellshyrna  
427 (red= SIFJO, green= OXAN and blue= OLFJO). All data from 1<sup>st</sup> July to 30<sup>th</sup> September 2012 (Data  
428 obtained from IMO in 2016).

429

430

#### 431 4.3.2 Temperature

432 The temperature patterns in 2012 were also unusual, the summer and spring were  
433 unusually warm on the whole, but the autumn was particularly cold (Fig. 5; Jónsson  
434 2013). The average temperature measured in the town of Ólafsfjörður was in 10.4°C July  
435 and 10.9°C in August in 2012. The average temperatures for these months for the period  
436 2000-2012 are 9.8°C and 9.6°C respectively. The average temperature for September  
437 2012 was on the other hand 5.7°C compared to the average temperature of 7.2°C for  
438 2000-2012.

439 The average daily air temperature from 6<sup>th</sup> to 20<sup>th</sup> September at the Siglufjörður and  
440 Ólafsfjörður stations ranged from 3 to 6°C and at the Öxnadalsheiði weather station  
441 fluctuated around zero, but lowered to approximately -3°C the night before the slide (Fig.  
442 7). Our corrected temperature data indicate average daily temperatures of at the altitude  
443 of the Móafellshyrna slide of around -1 to -2°C in the days preceding the slide and hence  
444 night-time temperatures would have been even lower. During the evening of 19<sup>th</sup>  
445 September, a drop below 0°C in the atmospheric temperature in the mountains was  
446 measured in the Öxnadalsheiði weather station (Fig. 7). The snow on the ground at the  
447 time of the Móafellshyrna debris slide shows that similar sub-freezing conditions also  
448 prevailed at this altitude prior to the slide. These low temperatures combined with the  
449 snow cover are strong evidence that sudden influx of water from precipitation was not the  
450 trigger for the Móafellshyrna slide.

#### 451 4.3.3 Earthquake sequence

452 In the Eyjafjarðaráll graben, three earthquakes with magnitudes M 4.2 to M 4.5 were  
453 registered on 19<sup>th</sup> and 20<sup>th</sup> September (Gudmundsson *et al.*, 2014). Their epicentres are  
454 located 25-27 km north-northeast of the town of Siglufjörður, and 60 km north-northeast  
455 of the Móafellshyrna site (Fig. 4). On the morning of 19<sup>th</sup> September, one day prior to the

456 slide, two earthquakes with magnitudes M 4.5 and M 4.3 occurred at 07:57 and 08:28  
457 respectively. A number of smaller aftershocks continued to occur throughout the day  
458 following these earthquakes. Another earthquake of magnitude M 4.2, occurred at 9:27  
459 on the 20<sup>th</sup> September, approximately three hours prior to the first observations of the  
460 Móafellshyrna slide (Fig. 4).

461

## 462 **5. Discussion**

463 The Móafellshyrna debris slide is a rare example of a gravitational mass movement  
464 where three triggering factors may have contributed to the failure of the slope: heavy  
465 precipitation, earthquake activity and ground ice degradation (via rising average surface  
466 temperatures). We infer that precipitation was the main preparatory factor for this debris  
467 slide for the following reasons. Heavy prolonged precipitation was recorded across the  
468 area, where nearly half of the usual annual precipitation fell in less than one month (Fig.  
469 5). Many case studies have shown that high magnitude water input, either by rainfall  
470 (Rapp and Nyberg 1981) or snowmelt (Decaulne *et al.* 2007), leads to oversaturation of  
471 soil directly triggering debris flows and shallow landslides. These studies also point out  
472 the role of intense rainfall as a preparatory (rather than direct trigger) factor to failure (e.g.  
473 Rapp, 1964, 1985 and 1995; Johnson and Rahn, 1970; Johnson and Rodine, 1984;  
474 Rapp and Nyberg, 1981; Addisson, 1987; Innes, 1989; Luckman, 1992; Becht, 1995;  
475 Cannon and Reneau, 2000; Sæmundsson *et al.*, 2003, Decaulne *et al.* 2007,  
476 Sæmundsson and Decaulne 2007, Guzzetti *et al.* 2007). However, the role of water  
477 infiltration in triggering of shallow landslides and debris flow in permafrost areas is rarely  
478 well documented (Harris and Gustafsson, 1993). We do not favour sudden water input as  
479 a direct trigger of the Móafellshyrna debris slide, because snow was present on the  
480 ground at the time of the failure. However, we think this is a necessary pre-condition for  
481 the failure to occur.

482 The direct influence of the seismic activity and associated ground acceleration on the  
483 motion of the debris in the source area seems unlikely, as the event did not occur  
484 immediately following any seismic event. Earthquakes are a common triggering factor for  
485 landslide activity and are considered as a major cause for landslides worldwide (e.g.  
486 Keefer, 1984, 1994, 2002, Malamund *et al.* 2004). Yet, no other debris flow or rockfall  
487 activity was reported on the northern part of the Tröllaskagi area on 19<sup>th</sup> or 20<sup>th</sup>



488 September, which might be expected if ground acceleration were sufficient to trigger  
489 mass movement. However, the short time interval (3hrs) between the last earthquake  
490 and the failure indicates a possible connection between the slide and the seismic events.

491 Selby (1993) argued that “stability of the slope, orientation of the earthquake in relation to  
492 the slide mass, earthquake magnitude, focal depth, seismic attenuation and after-shock  
493 distribution” are factors that determine whether earthquakes trigger landslides. According  
494 to Keefer (1984), the maximum area likely to be affected by landslides in a seismic event  
495 increases from approximately 0 km<sup>2</sup> at M=4.0 up to 500.000 km<sup>2</sup> at M=9.2. According to  
496 Malamund *et al.* (2004) the lowest earthquake-magnitude is M 4.3 +/-0.4. for triggering  
497 gravitational mass movements. Tatard *et al.* (2010) state that earthquakes of M4 and  
498 lower have little or no influence on landslide triggering. Nevertheless, several studies  
499 (e.g., Sassa *et al.*, 2007, Walter and Joswig, 2008) mention that small earthquakes  
500 (maximum M 3.6 in southern Italy according to Del Gaudio *et al.*, 2000) and repeated  
501 shocks can influence hydrogeological settings and can possibly cause landslides,  
502 sometimes with delay between the earthquake and the mass movement. Jibson *et al.*  
503 (1994) also discuss delayed landslide movements, from larger earthquakes (M 7.0), and  
504 state that the simplest explanation for the delay is a change in the ground-water  
505 conditions. Based on the above mentioned studies, it is unlikely that an earthquake of M  
506 4.3 was the only triggering factor for the Móafellshyrna debris slide, having taken place  
507 60 km away from the epicentre. On the other hand, since the Móafellshyrna slide  
508 occurred only three hours after a seismic event, the seismic sequence is likely playing  
509 some indirect role.

510 The ground water flow system of the colluvial cone composing the source material of the  
511 Móafellshyrna landslide is expected to be very limited. This is due to several factors: i)  
512 the catchment area above the source area is not very large (around 350 m long); ii) the  
513 colluvial cone is confined uphill by a vertical rockwall, and downhill by the edge of the  
514 topographic bench; iii) the presence of ground ice cementing the deposits; iv) the sub-  
515 horizontal dipping of the bedrock layers where the deposits are perched. However, it has  
516 been shown that talus slopes can contain multiple and distinct groundwater flow systems  
517 beneath or within them, and that they have a rapid and localised response to precipitation  
518 and melt inputs (Roy and Hayashi, 2009; McClymont *et al.*, 2010). One component of the  
519 groundwater flow in the colluvial cone of Móafellshyrna may originate in the pervasive  
520 system of sub-horizontal and sub-vertical discontinuities affecting the bedrock. If a

521 groundwater system was present before the failure, the response of the water table  
522 should be rapid. Since seismic activity can release water by coseismic liquefaction or  
523 consolidation of loose sediments (e.g., Manga *et al.*, 2003; Montgomery *et al.*, 2003), a  
524 change in the hydrogeological equilibrium of the colluvial cone caused by seismic activity  
525 could have contributed to the occurrence of the failure.

526 Field evidence from the Móafellshyrna debris slide strongly suggests the involvement of  
527 ground ice thaw in triggering the event. Because the month of the event had  
528 temperatures lower than average and the days prior to the event were mostly below zero  
529 Celsius (as evidenced by snow on the ground), we do not think thaw water from the  
530 ground ice in the perched talus contributed significantly to the event. However, longer  
531 term, deeper thawing caused by an annual rise in temperature and therefore a shift in the  
532 permafrost table, including an anomalously warm preceding summer, we believe is a  
533 more likely contributor. In recent years, there has been an increasing interest worldwide  
534 in the influence of climate warming and associated decline of mountain permafrost on the  
535 occurrence of mass wasting phenomena (e.g., Rebetz *et al.*, 1997; Gruber *et al.* 2004;  
536 Gruber and Haeberli, 2007; Fischer *et al.*, 2006; Sattler *et al.*, 2011; Stoffel and Huggel,  
537 2012; Damm and Felderer, 2013; Stoffel *et al.*, 2014). The increasing frequency of rapid  
538 mass movements, such as debris flows, debris slides, rock falls and rock avalanches, in  
539 mountainous areas have been linked in several cases with mountain permafrost  
540 degradation (e.g., Clague *et al.*, 2012; Wirz *et al.*, 2013; Barboux *et al.*, 2015; Darrow *et*  
541 *al.*, 2015; Haeberli *et al.*, 2016). Loss of ice-cementation, the presence of segregated ice,  
542 increased hydrostatic pressure and the associated reduction of shear strength can all  
543 lead to reduction of stability with increasing atmospheric temperature via permafrost  
544 degradation (e.g., Gruber and Haeberli, 2007; Krautblatter *et al.*, 2013; Pogliotti *et al.*,  
545 2015). Although, these previous studies have focused on the stability of massive rock  
546 masses, a similar (perhaps exaggerated) effect might be expected in ice cemented talus.

547 The increase of mean annual temperature, which has been observed in Iceland over the  
548 last few decades (Björnsson *et al.* 2008), should be leading to degradation of  
549 discontinuous permafrost in Iceland, which is thought to be present in the Tröllaskagi  
550 peninsula (Etzelmüller *et al.*, 2007a). Our observations of the ice-cemented deposits  
551 shows that the Móafellshyrna slide originated from deposits containing pore-filling ground  
552 ice and equally that these deposits were still frozen at the time of the slide. Together  
553 these argue for a permafrost origin for this ground ice. The increasing average

554 temperatures over the last decades (Björnsson *et al.* 2008, Jónsson 2013) before the  
555 event may have initiated the degradation of ground ice in the talus cone where the slide  
556 initiated, but not from the top-down, but from the base-up. This thawing may have: i)  
557 lubricated of the base of the cemented colluvial cone, ii) lowered the effective friction  
558 angle (cohesion), and hence iii) caused the slow movement of the colluvial cone perched  
559 on the bench and the sliding along a detachment surface of the whole ice-cemented  
560 mass. The warming of the rock mass onto which the colluvial deposit was previously  
561 cemented could have been brought about by a combination of propagation of the thermal  
562 wave through the rock mass from the warmer southeast-facing side (e.g., Noetzli *et al.*  
563 2007) and the delivery of warmer liquid water (derived from the intense precipitation) to  
564 the talus-rock interface from the south-westward dipping strata. Hence, the rupture  
565 occurred beneath the permafrost table. Perhaps the ice-cemented colluvium was in effect  
566 forming an underground “ice dam” that was holding back water saturated debris until its  
567 own weight and the seismic shaking caused it to fail. However, we cannot substantiate  
568 this link with certainty as we lack direct temperature measurements in the talus cone. Our  
569 hypothesis is supported by the slow widening of the tension crack at the top of the source  
570 area as observed by the eye-witnesses and the fact that the landslide was fluidised (a  
571 water content higher than expected for such a small catchment) even as it fell down the  
572 talus slope (causing the channelized erosion).

573 Ice-cemented deposits have been observed in two other landslides in Iceland, e.g. on the  
574 Torfufell mountain (source area at ~750 m a.s.l.) and the Árnesfjall Mountain (source  
575 area at ~350 m a.s.l.). These further events provide additional evidence to support our  
576 hypothesis that the lower limit of permafrost degradation extends to lower altitudes. The  
577 source zones for the Móafellshyrna and the Torfufell slides are at the lower elevation limit  
578 of discontinuous mountain permafrost in northern Iceland (i.e., 840 m a.s.l.; Etzelmüller *et*  
579 *al.*, 2007a). On the other hand, the source zone of the slide in the Árnesfjall Mountain is  
580 at an unexpected much lower elevation, which shows that the knowledge of mountain  
581 permafrost in Iceland is incomplete. The setting of talus perched on benches is not a rare  
582 situation in Iceland because of the sub-horizontal basalt layers create topographic  
583 benches on which loose material can accumulate. Hence investigating whether those  
584 with permafrost conditions, particularly above inhabited areas, contain ground ice and  
585 establishing its condition, should be a priority.

586 **6. Conclusions**

587 The debris slide in the Móafellshyrna Mountain began with a slow movement of a  
588 perched colluvial cone, as described by the local residents of Þrasastaðir. This colluvium  
589 is composed of stratified ice-cemented deposits at 840 m elevation and large blocks and  
590 boulders broke off the frontal part of the cone and fell onto the talus slope below. The  
591 mass movement transformed into a rapid debris slide, travelling down the mountainside  
592 with the final deposits coming to rest at 330 m a.s.l. The total volume of the slide is  
593 estimated to be around 312 000 – 480 000 m<sup>3</sup> (including the initial mass and mass added  
594 via bulking), covering an area of 0.3 km<sup>2</sup>.

595 We suggest that heavy precipitation prior to the slide was the main preparatory factor,  
596 with over 400 mm of precipitation recorded in one month prior to the event after an  
597 unusually dry summer season. The influence of seismic activity is unclear, but the close  
598 temporal association between the last earthquake series and the failure suggests that the  
599 shaking could have pre-conditioned the landslide weakening the cohesion between the  
600 ice-cemented colluvium and the bedrock and/or changing of the hydrology. The presence  
601 of ice-cementing the source colluvium at 880 m confirms the presence of discontinuous  
602 mountain permafrost at that elevation. We suggest that the partial thaw of these deposits  
603 was a trigger for the failure for three reasons: i) the landslide followed an usually warm  
604 spring and summer, ii) mean annual air temperatures are generally increasing in Iceland  
605 and iii) the colluvial cone initially slid as a single cohesive mass suggesting basal  
606 lubrication/melting. The fact that two other recent landslides contain similar ice cemented  
607 deposits suggests that mountain permafrost degradation could be more prevalent in  
608 triggering landslides in Iceland than has previously been thought.

609 The ice-cemented deposits within the slides of the Móafellshyrna, Torfufell and Árnesfjall  
610 Mountains have highlighted the need for a more detailed understanding of the distribution  
611 and condition of mountain permafrost within perched talus deposits in Iceland. Future  
612 studies should focus on the relationship between rapid mass wasting processes and the  
613 degradation of mountain permafrost in such deposits in Iceland. These three landslides  
614 occurred in uninhabited areas, but future similar landslides might not, and therefore they  
615 could pose a potential hazard to society and infrastructure in the island.

616

617 **Acknowledgements**

618 This work would not have been possible without a postgraduate studentship grant from  
619 the CENTA Doctoral Training Partnership funded by the U.K. Natural Environment  
620 Research Council (NERC) and the British Geological Survey University Funding Initiative.  
621 Aerial photos used in this research are courtesy of NERC Airborne Research and Survey  
622 Facility via the European Facility for Airborne Research (EuFAR), projects HidHaz and  
623 EUFAR15\_48. Especially we would like to thank Íris Jónsdóttir and Jón Elvar Númasson  
624 residents at the Þrasastaðir farm for their valuable information and detailed description of  
625 the behaviour of the slide and all their hospitality and help at the site. Thanks to Gestur  
626 Hansson, a snow observer in Siglufjörður, for providing us with photos and assisting us at  
627 the site.

628

## 629 **References**

- 630 Addisson, K. 1987: Debris flow during intense rainfall in Snowdonia, North Wales: a  
631 preliminary survey. *Earth Surface Processes and Landforms* 12, 561-566.
- 632 Ágústsson, K. & Pétursson, H.G. 2013: Skriður og grjóthrun. In: Sólnes, J. (ed).  
633 Náttúruvá á Íslandi, eldgos og jarðskjálftar. Háskólaútgáfan. pp. 639-645. (in  
634 Icelandic).
- 635 Allen, R.G. Nolet, Morgan, W.J., Vogfjörð, K.S., Netles, M., Ekström, G., Bergsson, B.H.,  
636 Erlendsson, P., Foulger, G., Jakobsdóttir, S.S., Julian, B., Pritchard, M., Ragnarsson,  
637 S. & Stefánsson, R. 2002: Plume driven plumbing and crustal formation in Iceland. *J.*  
638 *Geophys. Res.* 107(B8), doi:10.1029/2001JB000584
- 639 Arnalds, Þ., Sauermoser, S., Jóhannesson, T., Grímsdóttir, T. 2001: Hazard zoning for  
640 Siglufjörður – Technical report. Icelandic Meteorological Office, Report 01020. VÍ-  
641 ÚR11. 68 pp. Reykjavík.
- 642 Barboux, C., Strozzi, T., Delaloye, R., Wegmüller, U., & Collet, C. 2015: Mapping slope  
643 movements in Alpine environments using TerraSAR-X interferometric methods. *ISPRS*  
644 *Journal of Photogrammetry and Remote Sensing*, 109, 178-192.
- 645 Becht, M. 1995: Slope erosion processes in the Alps. In: O: Slaymaker (ed.): *Steepland*  
646 *Geomorphology*, 54-61.
- 647 Bjarnason, I. Th. 2008: An Iceland hotspot saga. *Jökull* 58, 3-16.
- 648 Björnsson, H., Sveinbjörnsdóttir, Á.E., Daníelsdóttir, A.K., Snorrason, Á., Sigurðsson,  
649 B.D., Sveinbjörnsson, E., Viggósson, G., Sigurjónsson, J., Baldursson, S.,  
650 Þorvaldsdóttir, S. & Jónsson, T. 2008: Hnattrænar loftslagsbreytingar og áhrif þeirra á  
651 Íslandi – Skýrsla vísindanefndar um loftslagsbreytingar. (e. Global warming and it's  
652 effect on Iceland). Umhverfissráðuneytið. (in Icelandic).
- 653 Brynjólfsson, S. & Ólafsson, H. 2008: Precipitation in the Svarfadardalur region, North-  
654 Iceland. *Meteorology and Atmospheric Physics* 103: 57-66.
- 655 Clague, J.J., Huggel, C., Korup, O. & McGuire, B. 2012: Climate change and hazardous  
656 processes in high mountains. *Revista de la Asociación Geológica Argentina*, 69(3), pp.  
657 328-338.

- 658 Coquin, J., Mercier, D., Bourgeois, O., Cossart, E. & Decaulne, A., 2015: Gravitational  
659 spreading of mountain ridges coeval with Late Weichselian deglaciation: impact on  
660 glacial landscapes in Tröllaskagi, northern Iceland. *Quaternary Science Reviews* 107.  
661 197-213.
- 662 Coquin, J., Mercier, D., Bourgeois, O., Feuillet, T. & Decaulne, A. 2016: Is gravitational  
663 spreading a precursor for the Stífluhólar landslide (Skagafjörður, Northern Iceland)?  
664 *Géomorphologie: Relief, Processus, Environment*. Vol. 22, no 1, p. 9-24.
- 665 Crochet, P., Jóhannesson, T., Jónsson, T., Sigurdsson, O., Björnsson, H., Pálsson, F. &  
666 Barstad, I. 2007: Estimating the spatial distribution of precipitation in Iceland using a  
667 linear model of orographic precipitation. *Journal of Hydrometeorology*, 8(6): 18 1285-  
668 1306.
- 669 Damm, B. & Felderer, A. 2013: Impact of atmospheric warming on periglacial degradation  
670 and debris flow initiation – a case study from the eastern European Alps. *Quaternary  
671 Science Journal*. Vol. 62, (2), 136-148.
- 672 Darrow, M.M., Gyswyt, N.L., Simpson, J.M., Daanen, R.P. & Hubbard, T.D. 2016: Frozen  
673 debris lobe morphology and movement: an overview of eight dynamic features,  
674 southern Brooks Range, Alaska. *The Cryosphere*, 10(3), p.977.
- 675 Decaulne, A. 2005: Slope processes and related risk appearance within the Icelandic  
676 Westfjords during the twentieth century. *Nat. Hazards Earth Syst. Sci.*, 5, 309-318.
- 677 Decaulne, A. 2007: Snow-avalanche and debris-flow hazards in the fjords of north-  
678 western Iceland, mitigation and prevention. *Natural Hazards*, 41(1), 81-98.
- 679 Decaulne, A., Sæmundsson, Þ. & Pétursson, O. 2005: Debris flow triggered by rapid  
680 snowmelt: A case study in the Gleiðarhjalli area, Northwestern Iceland. *Geografiska  
681 Annaler*. 87 A(4): 487-500.
- 682 Decaulne, A. & Sæmundsson, Þ. 2007: Spatial and temporal diversity for debris-flow  
683 meteorological control in subarctic oceanic periglacial environments in Iceland. *Earth  
684 Surface Processes and Landforms*. 32(13), 1971–1983.
- 685 Decaulne, A., Cossart E., Mercier, D., Feuillet, T., Coquin, J. & Jónsson, H.P. 2016: An  
686 early Holocene age for the Vatn landslide (Skagafjörður, central northern Iceland):  
687 Insights into the role of postglacial landsliding on slope development. *The Holocene*.  
688 Vol. 26(8) 1304–1318.
- 689 Del Gaudio, V., Trizzino, R., Calcagnile, G., Calvaruso, A. & Pierri, P. 2000: Landsliding  
690 in seismic areas: The case of the Acquara-Vadoncello landslide (southern Italy), *Bull.  
691 Eng. Geol. Environ.*, 59(1), 23–37.
- 692 Einarsson, M. A. 1984: Climate in Iceland. In: H.van.Loos (ed), *World survey of  
693 climatology*, 15, climate of the oceans: 673-697. Amsterdam, Elsevier.
- 694 Einarsson, P. 2008: Plate boundaries, rifts and transforms in Iceland. *Jökull* 58. 35-58.
- 695 Einarsson, P. and Björnsson, S. 1979: Earthquakes in Iceland. *Jökull*, 29. 37-46.
- 696 Etzelmüller, B., Farbrot, H., Guðmundsson, Á., Humlum, O., Tveito, O.E. & Björnsson, H.  
697 2007a: The regional distribution of mountain permafrost in Iceland. *Permafrost and  
698 periglacial processes*, 18: 185-1999.
- 699 Etzelmüller, B., Schuler, T.V., Farbrot, H. & Gudmundsson, A. 2007b: Permafrost in  
700 Iceland: Thermal state and climate change impact. *American Geophysical Union, Fall  
701 Meeting 2007*, abstract #C31A-02.

- 702 Farbrót, H., Etzelmüller, B., Gudmundsson, A., Humlum, O., Kellerer-Pirklbauer, A.,  
703 Eiken, T. & Wangensteen, B. 2007a: Rock glaciers and permafrost in Trollaskagi,  
704 northern Iceland. *Zeitschrift Fur Geomorphologie* 51: 1–16.
- 705 Farbrót, H., Etzelmüller, B., Schuler, T.V., Gudmundsson, A., Eiken, T., Humlum, O. &  
706 Björnsson, H. 2007b: Thermal characteristics and impact of climate change on  
707 mountain permafrost in Iceland. *Journal of Geophysical Research* 112:
- 708 Fischer, L., Kaab, A., Huggel, C. & Noetzel, J. 2006: Geology, glacier retreat and  
709 permafrost degradation as controlling factors of slope instabilities in a high mountain  
710 rock wall: the Monte Rosa east face. *Natural Hazards and Earth System Science*, 6  
711 (5), pp. 761-772.
- 712 Friedman, J.D., Johansson, C.E., Oskarsson, H., Svensson, H., Thorarinsson, S. &  
713 Williams Jr., R.S. 1971: Observations on Icelandic polygon surfaces and palsa areas.  
714 Photo interpretation and field studies. *Geografiska Annaler* A53, 115–145.
- 715 Gorbunov, A. P. 1988: The alpine permafrost zone of the USSR. In *Proceedings, Fifth*  
716 *International Conference on Permafrost*, Vol. 1.
- 717 Gudmundsson, A. 2000: Dynamics of Volcanic Systems in Iceland: Example of  
718 Tectonism and Volcanism at Juxtaposed Hot Spot and Mid-Ocean Ridge Systems.  
719 *Annual Review of Earth and Planetary Sciences*. Vol. 28, 107–140.
- 720 Gudmundsson, A. 2006: Infrastructure and evolution of ocean-ridge discontinuities in  
721 Iceland. *Journal of Geodynamics*, 43, 6-29.
- 722 Gudmundsson, G.B., Martin Hensch, Matthew Roberts and the SIL monitoring group  
723 2014: The fall 2012 earthquake sequence in Eyjafjarðaráll, western Húsavík-Flatey  
724 Fault. *International Workshop on Earthquakes in North Iceland, Húsavík, North*  
725 *Iceland*, 6-8 June 2013, pp.71-73.
- 726 Guzzetti, F., Peruccacci, S., Rossi, M. & Stark, C.P. 2007: Rainfall thresholds for the  
727 initiation of landslides in central and southern Europe. *Meteorology and Atmospheric*  
728 *Physic*, 98, 239–267. DOI 10.1007/s00703-007-0262-7
- 729 Gruber, S., Hoelzle, M. & Haeberli, W. 2004: Permafrost thaw and destabilization of  
730 Alpine rock walls in the hot summer of 2003. *Geophysical Research Letters*, Vol. 31,  
731 L13504, doi:10.1029/2004GL020051.
- 732 Gruber, S. & Haeberli, W. 2007: Permafrost in steep bedrock slopes and its temperature-  
733 related destabilization following climate change. *Journal of Geophysical Research*.  
734 Vol. 112, F02S18, doi:10.1029/2006JF000547.
- 735 Haeberli, W., Schaub, Y. & Huggel, C. 2016: Increasing risks related to landslides from  
736 degrading permafrost into new lakes in de-glaciating mountain ranges.  
737 *Geomorphology*, Vol.293 (B), 405-417.
- 738 Halldórsson, P. 1984: Skagafjarðarskjálftinn 1963. *Veðurstofa Íslands, Reykjavík* (in  
739 Icelandic).
- 740 Harris, S.A. & Gustafsson, C.A. 1993: Debris flow characteristics in an area of continuous  
741 permafrost, St Elias Range, Yukon Territory. *Zeitschrift fur Geomorphologie* NF 37, 41-  
742 56.
- 743 Harp, E.L. & Jibson, R.W. 1996: Landslides triggered by the 1994 Northridge, California,  
744 earthquake. *Bulletin of the Seismological Society of America* 86.1B: S319-S332.

- 745 Hermannsdóttir, K. 2012: Veðrið sem gekk yfir landið 9. – 11. September 2012.  
 746 www.vedur.is 19.2.2012. (<http://www.vedur.is/vedur/frodleikur/greinar/nr/2533>) (in  
 747 Icelandic).
- 748 Hirakawa, K. 1986: Development of palsa bog in central highland, Iceland. Geographical  
 749 Reports of Tokyo Metropolitan University 21, 111–122.
- 750 Hjartarson, Á., & Sæmundsson, K., 2014: Geological map of Iceland, 1:600.000,  
 751 Bedrock. ÍSOR, Icelandic Geosurvey. Reykjavík.
- 752 Hungr, O., Evans, S.G., Bovis, M.J., Hutchinson, J.N., 2001. A review of the classification  
 753 of landslides of the flow type. Environmental & Engineering Geoscience 7, 221–238.  
 754 doi:10.2113/gseegeosci.7.3.221
- 755 Hungr, O., Leroueil, S., Picarelli, L., 2014. The Varnes classification of landslide types, an  
 756 update. Landslides 11, 167–194. doi:10.1007/s10346-013-0436-y
- 757 Icelandic Met Office 2016: Icelandic weather record archives on [www.vedur.is](http://www.vedur.is).
- 758 Ingólfsson, Ó., Norðdahl, H. & Schomacker, A. 2010: Deglaciation and Holocene glacial  
 759 history of Iceland. Developments in Quaternary Sciences, vol. 13, 51-68.
- 760 Innes, J.L. 1989: Rapid mass-movements in Upland Britain: a review with particular  
 761 reference to debris flows. Studia Geomorphologica Carpatho-Baltanica 23, 53-67.
- 762 Iverson, R.M., 1997. The physics of debris flows. Rev. Geophys. 35, 245–296.
- 763 Jensen, E.H. 2000: Úttekt á jarðfræðilegum hættum eftir jarðskjálftana 17. og 21. Júní  
 764 2000. Veðurstofa Íslands, Reykjavík (in Icelandic).
- 765 Jibson, R.W., Prentice, C.S., Borissoff, B.A., Rogozhin, E.A. & Langer, C.J. 1994: Some  
 766 Observations of Landslides Triggered by the 29 April 1991 Racha Earthquake,  
 767 Republic of Georgia. Bulletin of the Seismological Society of America, Vol. 84, No. 4,  
 768 pp. 963-973.
- 769 Johannesson, H. & Sæmundsson, K. 2009: Geological Map of Iceland., 1:600000,  
 770 Tectonics. Museum of Natural History, Reykjavik.
- 771 Johnson, A.M. & Rahn, P.H. 1970: Mobilization of debris flows. Zeitschrift fur  
 772 Geomorphologie 9, 168-185.
- 773 Johnson, A.M. & Rodine, J.R. 1984: Debris flows. In D. Brunnsden and D.B. Prior (eds.):  
 774 Slope instability, Wiley and sons, 257-361.
- 775 Jónsson, Ó. 1957: Skriðuföll og snjóflóð, I bindi. Bókaútgáfan Norðri hf, Akureyri (in  
 776 Icelandic).
- 777 Jónsson, Ó. 1976: Berghlaup. Ræktunarfélag Norðurlands, Akureyri. 622 pp. (in  
 778 Icelandic).
- 779 Jónsson, Ó. & Pétursson, H.G. 1992: Skriðuföll og snjóflóð, II bindi. Skriðuannáll (2.  
 780 útgáfa). Skjaldborg, Reykjavík. (in Icelandic).
- 781 Jónsson, H.B., Norðdahl, H. & Pétursson, H.G. 2004: Myndaði Berghlaup Vatnsdalshóla?  
 782 Náttúrufræðingurinn 72 (3-4), 129-138. (in Icelandic).
- 783 Jónsson, T. 2008: Past temperature conditions in Iceland from 1798 to 2007.  
 784 ([www.vedur.is](http://www.vedur.is) article 26.2.2008 (<http://en.vedur.is/climatology/articles/nr/1213>)).
- 785 Jónsson, T. 2013: The weather in Iceland 2012. Climate summary. [www.vedur.is](http://www.vedur.is)  
 786 9.1.2013 (<http://en.vedur.is/weather/articles/nr/2614>).



- 787 Keefer, D.K. 1984: Landslides caused by earthquakes, *Geol. Soc. Am. Bull.* 95, 406–421.
- 788 Keefer, D.K. 1994: The importance of earthquake-induced landslides to long-term slope  
789 erosion and slope-failure hazards in seismically active regions, *Geomorphology* 10.  
790 265–284.
- 791 Keefer, D.K. 2002: Investigating landslides caused by earthquakes – A historical review.  
792 *Surveys in Geophysics* 23: 473–510, Kluwer.
- 793 Kellere-Pirklbauer, A., Wangesteen, B., Farbrøt, H. & Etzelmüller, B. 2007: Relative  
794 surface age-dating of rock glacier systems near Hólar in Hjaltadalur, northern Iceland.  
795 *J. Quaternary Sci.*, Vol. 23 pp. 137–151. ISSN 0267-8179.
- 796 King, L. 1986: Zonation and ecology of high mountain permafrost in Scandinavia.  
797 *Geografiska Annaler*, 68 A, no 3, pp. 131-139.
- 798 Kneisel, C. 2010: The nature and dynamics of frozen ground in alpine and subarctic  
799 periglacial environments. *The Holocene*, 20 (3), 423-445.
- 800 Kneisel, C., Sæmundsson, Þ. & Beylich, A.A. 2007: Reconnaissance surveys of  
801 contemporary permafrost environments in central Iceland using geoelectrical methods:  
802 Implications for permafrost degradation and sediment fluxes. *Geogr. Ann.* 89A (1): 41-  
803 50.
- 804 Krautblatter, M., Funk, D. & Günzel, F.K. 2013. Why permafrost rocks become unstable:  
805 a rock–ice-mechanical model in time and space. *Earth Surface Processes and*  
806 *Landforms*, 38(8), pp.876-887.
- 807 Lilleören, K.S., Etzelmüller, B., Gratner-Roer, I., Kaab, A., Sestermann, S. &  
808 Guðmundsson, Á. 2013: The distribution, thermal characteristics and dynamics of  
809 permafrost in Tröllaskagi, Northern Iceland, as inferred from the distribution of rock  
810 glaciers and ice-cored moraines. *Permafrost and Periglacial Processes* doi:  
811 10.1002/ppp.1792.
- 812 Luckman, B.H. 1992: Debris flows and snow avalanches landforms in the Lairig Ghru,  
813 Cairngorm Mountains, Scotland. *Geografiska Annaler* 74A (2-3), 109-121.
- 814 Malamud, B.D., Turcotte, D.L., Guzzetti, F. & Reichenbach, P. 2004: Landslides,  
815 earthquakes, and erosion. *Earth and Planetary Science Letters* 229, 45– 59.
- 816 Manga, M., Brodsky, E.E., and Boone, M., 2003, Response of streamflow to multiple  
817 earthquakes and implications for the origin of postseismic discharge changes:  
818 *Geophysical Research Letters*, v. 30, no. 5, 1214, doi: 10.1029/2002GL016618.
- 819 McClymont, A. F., et al. "Groundwater flow and storage within an alpine meadow-talus  
820 complex." *Hydrology and Earth System Sciences* 14.6 (2010): 859-872.
- 821 McColl, S.T. 2012: Paraglacial rock-slope stability. *Geomorphology* 153-154, 1-16. i
- 822 McDougall, I., Kristjánsson, L. & Sæmundsson, K. 1984: Magnetostratigraphy and  
823 geochronology of NW-Iceland. *Journal of Geophysical Research*, 89: 7029–7060.
- 824 Mercier, D., Cossart, E., Decaulne, A., Feuillet, T., Jónsson H.P. & Sæmundsson, Þ.  
825 2012: The Höfðahólar rock avalanche (sturzström): Chronological constraint of  
826 paraglacial landsliding on an Icelandic hillslope. *The Holocene* 23(3): 432– 446.
- 827 Mercier, D., Cossart, E., Decaulne, A., Feuillet, T., Jónsson, H.P. & Sæmundsson, Þ.  
828 2013: The Höfðahólar rock avalanche (sturzström): Chronological constraint of  
829 paraglacial landsliding on an Icelandic hillslope. *The Holocene*. 23(3) 432–446.

- 830 Ministry for the Environment of Iceland 2010: Iceland's Fifth National Communication on  
831 Climate Change Under the United Nations Framework Convention on Climate  
832 Change, 102 pp.
- 833 Montgomery, D.R., and Manga, M., 2003, Streamflow and water well responses to  
834 earthquakes: *Science*, v. 300p. 2047-2049.
- 835 Moorbath, S., Sigurdsson, H. & Goodwin, R. 1968: K-Ar ages of oldest exposed rocks in  
836 Iceland. *Earth and Planetary Science Letter*, 26 4: 197-205.
- 837 Nawri, N. & Björnsson, H. 2010. Surface Air Temperature and Precipitation Trends for  
838 Iceland in the 21st Century. Report VÍ 2010-005.
- 839 Norðdahl, H. & Pétursson, H.G. 2005: Relative sea level changes in Iceland. New aspect  
840 of the Weichselian deglaciation of Iceland. In: Caseldine, C., Russell, A., Harðardóttir,  
841 J., Knudsen, Ó., (Eds), *Iceland – modern processes and past environments.*  
842 *Developments in Quaternary Science* 5, 25–78.
- 843 Norðdahl, H., Ingólfsson, Ó., Pétursson, H.G., & Hallsdóttir, M. 2008: Late Weichselian  
844 and Holocene environmental history of Iceland. *Jökull* 58, 343–364.
- 845 Noetzi, J., Gruber, S., Kohl, T., Salzmann, N., Haerberli, W., 2007. Three-dimensional  
846 distribution and evolution of permafrost temperatures in idealized high-mountain  
847 topography. *Journal of Geophysical Research* 112. doi:10.1029/2006JF000545
- 848 Pogliotti, P., Guglielmin, M., Cremonese, E., Morra di Cella, U., Filippa, G., Pellet, C. &  
849 Hauck, C. 2015: Warming permafrost and active layer variability at Cime Bianche,  
850 Western European Alps. *The Cryosphere*, 9(2), pp.647-661.
- 851 Priesnitz K, & Schunke E. 1978: An approach to the ecology of permafrost in Central  
852 Iceland. Third International Conference on Permafrost, Edmonton, Canada. National  
853 Research Council of Canada, Ottawa; 474–479.
- 854 Rapp, A. 1964: Recordings of mass wasting in the Scandinavian Mountains. *Zeitschrift*  
855 *fur Geomorphologie* 5, 204-205.
- 856 Rapp, A. 1985: Extreme rainfall and rapid snowmelt as causes of mass movements in  
857 high latitude mountains. In M. Church and O. Slaymaker (eds): *Field and theory:*  
858 *lectures in geocryology*, University of British Columbia Press, Vancouver, 36-56.
- 859 Rapp, A. 1995: Case studies of geoprocesses and environmental changes in mountains  
860 of northern Sweden. *Geografiska Annaler* 77A (4), 189-198.
- 861 Rapp, A. & Nyberg, R. 1981: Alpine debris flows in northern Scandinavia, morphology  
862 and dating by lichenometry. *Geografiska Annaler* 58A (3), 193-200.
- 863 Rebetz, M.; Lugon, R. & Baeriswyl, P-A. 1997: Climate changes and debris flow in high  
864 mountain regions: The case study of the Ritigraben Torrent (Swiss Alps). *Climatic*  
865 *Change* 36: 371-389.
- 866 Roy, James W., and Masaki Hayashi. "Multiple, distinct groundwater flow systems of a  
867 single moraine–talus feature in an alpine watershed." *Journal of Hydrology* 373.1  
868 (2009): 139-150.
- 869 Sassa, K., H. Fukuoka, F. Wang, & G. Wang (Eds.) 2007: Landslides induced by a  
870 combined effect of earthquake and rainfall, in *Progress in Landslide Science*, pp. 193–  
871 207, Springer, Berlin.

- 872 Sattler, K., Keiler, M., Zischg, A. & Schrott, L. 2011: On the connection between debris  
873 flow activity and permafrost degradation: A case study from the Schnalstal, South  
874 Tyrolean Alps, Italy. *Permafrost and periglacial processes*, 22, 254-265.
- 875 Schunke, E. 1974: Frostspaltenmakropolygone im westlichen Zentral-Island, ihre  
876 klimatischen und edaphischen Bedingungen. *Eiszeitalter und Gegenwart* 25: 157–165.
- 877 Selby, M. J. 1993: *Hillslope Materials and Processes*. Oxford University Press, New York.
- 878 Sheridan, P., Smith, S., Brown, A., Vosper, S., 2010. A simple height-based correction for  
879 temperature downscaling in complex terrain. *Meteorological Applications*  
880 doi:10.1002/met.177
- 881 Sigurgeirsson, M.Á. & Hjartarson, Á. 2011: Gjóskulög og fjörumór á berghlaupi við  
882 Sjárvarhóla á Kjalarnesi Náttúrufræðingurinn 81 (3–4), pp. 123–130. (in Icelandic).
- 883 Sigurðsson, O. & Williams, R. S. 2008: Geographic names of Iceland's Glaciers: Historic  
884 and Modern. U.S. Geological Survey Professional Paper 1746, 225 p.
- 885 Stefánsson, R., Gunnar B. Guðmundsson, G.B., Þorbjarnardóttir, B. & Halldórsson, P.  
886 2008: Tjörnes fracture zone. New and old seismic evidences for the link between the  
887 North Iceland rift zone and the Mid-Atlantic ridge. *Tectonophysics*, 447, 117–126.
- 888 Stingl H. & Herrmann R. 1976: Untersuchungen zum Strukturbodenproblem auf Island;  
889 Gelaendebeobachtungen und statistische Auswertung. *Zeitschrift für Geomorphologie*  
890 20: 205–226.
- 891 Stoffel, M. & Huggel, C. 2012: Effects of climate change on mass movements in  
892 mountain environments. *Process in Physical Geography*. 36 (3), 421-439.
- 893 Stoffel, M., Mendlik, T., Schneuwly-Bollschweiler, M. & Gobiet, A. 2014: Possible impacts  
894 of climate change on debris-flow activity in the Swiss Alps. *Climate Change*, 122: 141-  
895 155.
- 896 Sæmundsson, K. 1979: Outline of the geology of Iceland. *Jökull*, 29. 7-28.
- 897 Sæmundsson, K. 1974: Evolution of the axial rifting zone in Northern Iceland and the  
898 Tjörnes Fracture Zone, *Geol. Soc. Am. Bull.* 85, 495-504.
- 899 Sæmundsson, K., Kristjánsson, L., McDougal, I. & Warkins, N.D. 1980: K-Ar dating,  
900 geological and paleomagnetic study of a 5-km lava succession in Northern Iceland.  
901 *Jour. Geoph. Research*, 85, B7, 3628-3646.
- 902 Sæmundsson, Þ., Pétursson, H.G. & Decaulne, A. 2003: Triggering factors for rapid  
903 mass-movements in Iceland. In D. Rickenman & C.I. Chen (eds): *Debris-Flow Hazards*  
904 *Mitigation: Mechanics, Prediction, and Assessment*, vol. 1, 167-178.
- 905 Sæmundsson, Þ. & Decaulne, A. 2007: Meteorological triggering factors and threshold  
906 conditions for shallow landslides and debris-flow activity in Iceland. In: V.R. Schaefer,  
907 R.L. Schuster & A.K. Turner (Eds.): *First North American Landslide Conference*, Vail  
908 Colorado, AEG Publication No. 23, 1475-1485.
- 909 Sæmundsson, Þ., Arnalds, O., Kneisel, C. Jónsson, H.P. & Decaulne, A. 2012: The  
910 Orravatnsrústir palsa site in Central Iceland – Palsas in an Aeolian sedimentation  
911 environment. *Geomorphology*, 167-168, 13-20.
- 912 Sæmundsson, Þ., Helgason, J.K. & Pétursson, H.P. 2013: The debris slide in the  
913 Móafellshyrna Mountain on the 20th of September 2012. Was it triggered by intense

- 914 precipitation and earthquake activity or simply by melting of the permafrost? 8th IAG  
915 International Conference on Geomorphology - August 27th to 31st.
- 916 Sæmundsson, Þ., Helgason, J.K. & Pétursson, H.P. 2014a: The melting of mountain  
917 permafrost and the Móafellshyrna debris slide in Northern Iceland. 31st Nordic  
918 Geological Winter Meeting. 8-10 January 2014, Lund University.
- 919 Sæmundsson, Þ., Helgason, J.K. & Pétursson, H.P. 2014b: Decline of mountain  
920 permafrost and the occurrence of recent large debris slides in Iceland. European  
921 Geosciences Union, General Assembly 2014, Vienna, Austria, 27 April – 02 May 2014.
- 922 Tatard, L., Grasso, J.R., Helmstetter, A. & Garambois, A. 2010: Characterization and  
923 comparison of landslide triggering in different tectonic and climatic settings, J.  
924 Geophys. Res., 115, F04040, doi:10.1029/2009JF001624.
- 925 Thorarinsson, S. 1937: Das Dalvik-Beben in Nord-island, 2. juni 1934. Geografiska  
926 Annaler, 19, 232-277.
- 927 Thorarinsson, S. 1951: Notes on patterned ground in Iceland. Geografiska Annaler 33:  
928 144–156.
- 929 Thorarinsson, S., Einarsson, T. & Kjartansson, G. 1959. On the geology and  
930 geomorphology of Iceland. Geografiska Annaler, Vol. 41. No 2/3. pp. 135-169.
- 931 Thorarinsson S. 1964: Additional notes on patterned ground in Iceland with a particular  
932 reference to icewedge polygons. Biuletyn Peryglacjalny 14: 327–336.
- 933 Thordarson, T. & Hoskuldsson, A. 2002: Iceland. Classic Geology in Europe 3. Terra  
934 Publishing, Harpenden, UK, 200 pp.
- 935 Thorhallsdottir, Th.E. 1994: Effects of changes in groundwater level on palsas in Central  
936 Iceland. Geografiska Annaler A76, 161–167.
- 937 Thorhallsdottir, Th.E. 1996: Seasonal and annual dynamics of frozen ground in the  
938 central highland of Iceland. Arctic and Alpine Research 28, 237–243.
- 939 Thorhallsdottir, Th.E. 1997: Tundra ecosystems of Iceland. In: Wiegolaski, F.E. (Ed.),  
940 Polar and Alpine Tundra. Elsevier, Amsterdam, pp. 85–96.
- 941 Tryggvason, K., Huseby, E. & Stefánsson, R. 1983: Seismic image of the hypothesized  
942 Icelandic hot spot. Tectonophysics 100, 97–118.
- 943 Tveito, O.E., Førland, E., Heino, R., Hanssen-Bauer, I., Alexandersson, H., Dahlström,  
944 B.A., Drebs, Kern-Hansen, C., Jónsson, T., Vaarby Laursen, E. & Westman, Y. 2000:  
945 Nordic temperature maps. Norwegian Meteorological Institute, Oslo, Report, 4 09/00,  
946 28 pp.
- 947 Walter, M., & Joswig, M. 2008: Seismic monitoring of fracture processes from a creeping  
948 landslide in the Vorarlberg Alps, Geophys. Res. Abstr., 10, 09212.
- 949 Watkins, N.D. & Walker, G.P.L. 1977: Magnetostratigraphy of Eastern-Iceland. American  
950 Journal of Science, 277: 513–584.
- 951 Wirz, V., Beutel, J., Buchli, B., Gruber, S. & Limpach, P. 2013: Temporal characteristics  
952 of different cryosphere-related slope movements in high mountains. In Landslide  
953 Science and Practice (pp. 383-390). Springer Berlin Heidelberg.
- 954 Wolfe, C.J., Bjarnason, I.Th., VanDecar, J. C. & Solomon, S. C. 1997: Seismic structure  
955 of the Iceland mantle plume. Nature 385, 245–247.

956 Yin, Y., Wang, F. and Sun, P. 2009: Landslide hazards triggered by the 2008 Wenchuan  
957 earthquake, Sichuan, China. *Landslides* Vol 6 (2), pp. 139-152.