



Synthesis of new calixarene phosphine oxides and their extraction properties in ionic liquids

Stanislav Miroshnichenko, Dariia Ternova, Isabelle Billard, Ali Ouadi, Olga Klimchuk, Vitaly Kalchenko

► To cite this version:

Stanislav Miroshnichenko, Dariia Ternova, Isabelle Billard, Ali Ouadi, Olga Klimchuk, et al.. Synthesis of new calixarene phosphine oxides and their extraction properties in ionic liquids. Phosphorus, Sulfur, and Silicon and the Related Elements, 2011, 18th International Conference on Phosphorus Chemistry (ICPC2010), 186 (4), pp.903-905. hal-02271897

HAL Id: hal-02271897

<https://hal.science/hal-02271897>

Submitted on 27 Aug 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Synthesis of new calixarene phosphine oxides and their extraction properties in ionic liquids

Isabelle Billard

► To cite this version:

Isabelle Billard. Synthesis of new calixarene phosphine oxides and their extraction properties in ionic liquids. *Phosphorus, Sulfur, and Silicon and the Related Elements*, Taylor

Francis, 2011. hal-02271897

HAL Id: hal-02271897

<https://hal.archives-ouvertes.fr/hal-02271897>

Submitted on 27 Aug 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SYNTHESIS OF NEW CALIXARENE-PHOSPHINE OXIDES AND THEIR EXTRACTION PROPERTIES IN IONIC LIQUIDS

Stanislav Miroshnichenko,¹ Dariia Ternova,¹ Isabelle Billard,²
Ali Ouadi,² Olga Klimchuk,^{1,2} and Vitaly Kalchenko¹

¹Institute of Organic Chemistry, NASU, Kyiv, Ukraine

²Institut Pluridisciplinaire Hubert Curien, DRS/CHNU, Strasbourg, France

Abstract New calix[4]arenes functionalized by $\text{Alk}_2\text{P}(\text{O})\text{CH}_2$ binding groups and *N*-methylimidazolium groups at the opposite rims of the macrocyclic skeleton have been synthesized and characterized. Their extraction properties towards of Eu^{3+} and Am^{3+} in ionic liquid were investigated.

Keywords Calixarenes; ionic liquids; phosphine oxides

INTRODUCTION

The use of nuclear power has led to the accumulation of great amounts of both high- and medium-level activity radioactive wastes, which require efficient methods of separation and safe disposal. The main method for the decontamination of radioactive wastes is liquid–liquid extraction of hazardous radionuclides followed by burial or transmutation. A modern approach in the design of efficient and selective extractants of radioanuclides is based on the functionalization of calixarene platform by several P=O containing fragments, capable of cooperatively binding metal cations.¹ Room temperature ionic liquids (RTIL) are promising solutions for the extraction of the radionuclides from spent nuclear fuel.²

RESULTS AND DISCUSSION

We present in this article the synthesis of a series of new *cone*-shaped calix[4]arenes possessing $\text{CH}_2\text{P}(\text{O})\text{Alk}_2$ metallocomplexing groups at the macrocyclic wide and narrow rim and *N*-methylimidazolium groups at the narrow rim. Extraction properties of the calixarene synthesized and dissolved in ionic liquid were investigated.

Calix[4]arenes **3–10** functionalized by four or eight $\text{Alk}_2\text{P}(\text{O})\text{CH}_2$ binding groups have been synthesized by the Arbuzov reaction of tetrachloromethylcalix[4]arenes **1, 2** with dialkylphosphinites.

Received 29 July 2010; accepted 20 September 2010.

Address correspondence to Isabelle Billard, Institut Pluridisciplinaire Hubert Curien, DRS/CHNU, 23 rue du loess, 67037, Strasbourg Cedex 2, France. E-mail: Isabelle.billard@ires.in2p3.fr and Vitaly I. Kalchenko, Institute of Organic Chemistry Chemistry, NAS of Ukraine, 5 Murmanska str., 02660, Kyiv-94, Ukraine. E-mail: vik@ioch.kiev.ua





Figure 1 Synthesis of calixarene phosphine oxides.

Figure 2 Extraction of Eu^{3+} , UO_2^{2+} , Am^{3+} by phosphine oxides from HNO_3 (0.3M) into N -butyl- N -methylimidazolium \times Tf_2N . D—distribution ratio of the metal cations between the ionic liquid and HNO_3 phases; TAPO—triptylphosphine oxide.

NEW CALIXARENE-PHOSPHINE OXIDES

3

30 Tetra(*N*-methylimidazolium) derivatives **11**, **12** were prepared by the reaction of
31 bromobutoxycalixarenes **8**, **10** with *N*-methylimidazole. Metathetical reactions of bromides
32 **11**, **12** with LiN(SO₂CF₃)₂ in DMF gave new quaternary salts **13**, **14** (Figure 1).

33 Extraction properties of some calixarene phosphine oxides towards Eu³⁺, UO₂²⁺,
34 and Am³⁺ were investigated in ionic liquid *N*-butyl-*N*-methylimidazolium × Tf₂N (Figure
35 2). The calixarene tetraphosphine oxides **5–7** are more efficient extractants than industrial
36 trialkylphosphine oxides. Calixarene **5** demonstrates high selectivity in UO₂²⁺ extraction.

37 REFERENCES

- 38 1. Smirnov, I.; Karavan, M.; Babain, V.; Kvasnitskiy, I.; Stoyanov, E.; Miroshnichenko, S. *Radiochim. Acta* **2007**, *95*, 97–102.
39 2. Ouadi, A.; Gadenne, B.; Hesemann, P.; Moreau, J. J. E.; Billard, I.; Gaillard, C.; Mekki, S.;
40 Moutiers, G. *Chem. Eur. J.* **2006**, *12*, 3074–3081.
41