

Morphology of fore-reef slopes and terraces, Takapoto Atoll (Tuamotu Archipelago, French Polynesia, central Pacific): The tectonic, sea-level and coral-growth control

Lucien F. Montaggioni, Antoine Collin, Dorothée James, Bernard Salvat, Bertrand Martin-Garin, Gilles Siu, Marguerite Taiarui, Yannick Chancerelle

▶ To cite this version:

Lucien F. Montaggioni, Antoine Collin, Dorothée James, Bernard Salvat, Bertrand Martin-Garin, et al.. Morphology of fore-reef slopes and terraces, Takapoto Atoll (Tuamotu Archipelago, French Polynesia, central Pacific): The tectonic, sea-level and coral-growth control. Marine Geology, 2019, pp.106027. 10.1016/j.margeo.2019.106027. hal-02276200

HAL Id: hal-02276200

https://hal.science/hal-02276200

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



- 1 Morphology of fore-reef slopes and terraces, Takapoto
- 2 Atoll (Tuamotu Archipelago, French Polynesia, central
- 3 Pacific): the tectonic, sea-level and coral-growth control
- 4 Lucien .F. Montaggioni ^{a,*}, Antoine Collin ^b, Dorothée James ^b,
- 5 Bernard Salvat ^c, Bertrand Martin-Garin ^a, Gilles Siu ^d,
- 6 Marguerite Taiarui ^d, Yannick Chancerelle ^d
- 7 a Aix Marseille Univ, CNRS, IRD, Coll France, CEREGE,
- 8 13331 Marseille, France
- 9 b PSL-École pratique des hautes études (EPHE), CNRS 6554
- 10 LTG, Labex Corail, Dinard, France
- 11 c PSL-École pratique des hautes études (EPHE), CNRS, UPVD,
- 12 USR3278, Labex Corail, Université de Perpignan, 66000
- 13 Perpignan, France
- 14 d PSL Research University: EPHE-UPVD-CNRS, USR 3278
- 15 CRIOBE BP 1013, 98729 Papetoai, Moorea, French Polynesia
- * *Corresponding author*
- 17 Email address: montaggioni@cerege.fr
- 18 Postal address:
- 19 Aix–Marseille université
- 20 CEREGE UM 34
- 21 Saint-Charles
- 22 CS80249 (case 67)
- 23 3, Place Victor Hugo

F-13331 Marseille cedex 3

25 Abstract

26 A dataset of 41 bathymetric profiles detected by satelline 27 imagery and echo-sounding to a maximum depth of 38 m along 28 the fore-reef slopes of Takapoto Atoll (Tuamotu Plateau, 29 French Polynesia, central Pacific) were qualitatively and 30 quantitatively analyzed in order to identify the main 31 geomorphological scuba-diving features. In addition, investigations from a selected site resulted in defining coral 32 33 composition and cover rate. Six distinct fore-reef sectors were morphologically differentiated on the basis of seaward 34 35 extension and slope gradient. Only the north end, and a 36 restricted area of the mid-south-east side extend seawards over 37 an average distance of 150 m, with a relatively low gradient 38 (23°-38°). Most of the outer-slope areas are narrow (less than 39 100 m on average) and steep (average slope equals to 45° to up 40 to 70°). Geomorphic features include spur-and-groove systems, 41 sub-horizontal to gently dipping terraces (23–45 m wide) 42 ending seawards in vertical steps (< 10 m high) and drop-offs 43 (> 10 m high), outer-reef crest-like reliefs (0.5 m-1.5 m high), 44 pinnacles (2 m-17 m high), coral heads (< 2 m high), trenchs 45 and concave scars. The recognized seven terrace levels occur at the respective depths of about -5 m, -10 m, -15 m, -20 m, 46 -24 m, -30.5 m and -38 m. Based on comparison with 47

48 subsurface chronostratigraphy of drilled reef sequences from 49 some other Pacific atolls, the terraces are interpreted as 50 remnants of reef tracts formed during Pleistocene interglacial, 51 high sea-stands, respectively during MIS substage 5e (about 52 -125 ka), MIS substages 7a (about -195 ka), 7c (about -210 53 ka), 7e (about -240 ka) and 9e (-330 ka). After depth 54 correction of ground lowering during periods of emergence and 55 subsidence effects, terrace elevations were expected to indicate 56 eustatic sea-level positions at each identified high sea-stand, at 57 approximately 7 m (MIS 5e) and 2.2 m (MIS 7a) above present 58 sea level and 9.5 m (MIS 7c), 11 m (MIS 7e) and 8.6 m (MIS 59 9e) below present sea level. During the post-glacial sea-level 60 rise, from about 9.5 ky, the studied portions of fore-reef slopes 61 were re-flooded. It was assumed that coralgal deposits in the 62 form of thin veneers, coral heads, outer-reef crest-like reliefs 63 and pinnacles have simply overlapped pre-existing structures. 64 Locally, the fore-reef zones are carved by trenchs, i.e. incipient 65 detachment faults, and scars, i.e. imprints of former landslides. 66 Similarly, some drop-offs may be regarded as fault walls. The 67 prevailing direction of faulting-collapsing surfaces is 68 consistent with the SW-NE orientation of the main axis of the 69 atoll and submarine ridges rising up from the north-western 70 floor of the Tuamotu Plateau. Analysis of the composition and 71 distribution of coral communities to -30 m depth revealed 72 high, downslope-increasing coral coverage (24 % to 67 % on

- 73 average) and the sharp replacement of robust-encrusting
- 74 *Pocillopora* by sub-massive to plate-like *Porites* from depths
- 75 greater than -10 m. Significant inter-regional differences were
- 76 detected with atolls in west Pacific, probably due to local
- 77 ecological constraints.

Highlights

78

- 79 A detailed survey of morphological features of upper fore-
- reef slopes from a Pacific atoll is provided for the first time,
- using satellite-based records.
- 82 Five submerged terrace levels among the seven detected are
- regarded as related to distinct Pleistocene high sea stands
- are identified.
- 85 At a large scale, the morphogenetic evolution of the atoll
- fore-reef slopes is mainly controlled by SW-NE trending
- 87 regional-scale, tectonic structures.
- 88 At a small scale, living coral communities with low
- diversity, but high cover contribute significantly to atoll
- 90 fore-reef shaping.
- 91 **Keywords:** submerged terraces, fore-reef slopes, Pleistocene
- 92 high stands, atoll, central Pacific.

93 1 Introduction

94	Coral reefs, typical ecosystems of many tropical coastlines, are
95	known to be reliable indicators of climate, environmental (see
96	Montaggioni and Braithwaite, 2009; Phinney et al., 2013 for
97	reviews) and relative sea-level changes (see reviews by
98	Montaggioni and Braithwaite, 2009; Murray-Wallace and
99	Woodroffe, 2014; Woodroffe and Webster, 2014; Camoin and
100	Webster, 2015). Intensive submarine exploration has given
101	evidence that reefs have developed at times of lower sea levels,
102	and at present located along continental and island shelves at
103	different depths. A number of submerged reefs were described
104	in the Pacific Ocean (Australia: Abbey et al., 2011; Webster et
105	al., 2009; Linklater et al., 2015 – French Polynesia:
106	Montaggioni et al., 1987; Cabioch et al., 2008; Camoin et al.,
107	2012; Seard et al., 2013 – Hawaii: Faichney et al., 2009, 2010
108	and references herein - Papua New Guinea: Webster et al.,
109	2004b; Ryukyus: Arai et al., 2016), in the Indian Ocean (Dullo
110	et al., 1998; Rovere et al., 2018; and references herein) and the
111	western Atlantic (Lightly et al., 1978; James and Ginsburg,
112	1979; Blanchon et al., 2002; Khanna et al., 2017; and
113	references herein). Submerged reefs have proven to provide
114	valuable information on relative sea-level and tectonic histories
115	(Webster et al., 2004a,b, 2006; Cabioch et al., 2008; Faichney
116	et al., 2010; Abbey et al., 2011; Rovere et al., 2018) and sub-
117	aerial erosional patterns as well (Purdy and Winterer, 2001).

118 The origin of submerged reefs and associated topographic 119 anomalies – terraces, breaks-in-slope, pinnacles and steps, has 120 been intensively discussed (Hopley, 2006; Beaman et al., 2008; 121 Abbey et al., 2011; Rovere et al., 2018). Reef tracts were 122 speculated to form during either the last deglacial sea-level rise 123 (-20 ka to -8 ka) or early, during late to mid-Pleistocene 124 variations of sea level. They may be multi-generational, 125 resulting from stacked successive growth and erosional 126 episodes over several cycles of submergence and sub-aerial 127 exposure, finally overlain by last deglacial deposits. Especially, 128 terraces are interpretated to be constructional in origin or to 129 have been shaped by erosive wave action during sea-level sea-130 stands lower than present. Similarly, steps may result from 131 seaward accretional processes during stillstands, step and drop-132 off recession during low stands or faulting. Contrary to shelf 133 slopes of continents and high volcanic islands, those of atolls 134 have been little explored. Earlier bathymetric surveys using 135 single beam echo-sounding have been conducted from the end 136 of the forties on a number of Pacific atolls (Ladd et al., 1970; 137 Winterer, 1976; Smith et al., 1994; Wong et al., 1999). 138 Recently, investigations using shallow-water multibeam 139 mapping, multispectral satellite-, optical- and acoustic-based 140 imagery off several atolls from Pacific US overseas territories, 141 provided new detailed insights into shelf morphology (for 142 instance, see Poppenga et al., 2018). In French Polynesia,

143 shallow-water (0 to -30 m) bathymetric investigations were 144 initiated at Moruroa (Chevalier et al. 1968).. At Takapoto, the 145 first underwater surveys were carried out by Kühlmann and 146 Chevalier (1986). A series of terraces and breaks-in-slope occur 147 at different depths to about -40 m downslope. From satellite-148 derived near-shore bathymetry, Collin et al. (2018) provided 149 the first high-resolution outer-slope morphometry of the eastern 150 side of Takapoto. 151 In addition to geomorphology, the characterisation of coral 152 communities living along the upper fore-reef slopes is an 153 important step in terms of paleo-water depth reconstructions. 154 The earlier works on the compositional and distributional 155 patterns of coral communities at Takapoto were carried out by 156 Chevalier and Denizot (1979), Kühlmann (1982), Bouchon 157 (1983) and Kühlmann and Chevalier (1986). 158 In this study, are presented both morphological and biological 159 datasets from the foreslopes of Takapoto Atoll in the Tuamotu 160 Archipelago (French Polynesia, south-central Pacific). Analysis 161 of satellite-based datasets gains insight into the detailed spatial 162 distribution of submerged geomorphic features. Scuba-diving 163 investigations reveal richness and cover rate of coral 164 populations. 165 The aims of this study are to: (1) identify the main geomorphic 166 features present on the shelf slopes and quantitatively describe 167 their morphology; (2) define the depth distributional trends of 168 these features, especially those of terraces; (3) explore 169 scenarios of their possible nature, mode and timing of 170 formation, based on chronostratigraphic comparisons with 171 other atoll sites; (4) determine the main depth-related 172 compositional and distributional attributes of coral 173 communities.

2 **Environmental setting**

175 2.1 Geological background

174

176

177

178

179

180

181

182

183

185

186

187

188

189

190

191

French Polynesia lies in the south-central Pacific and extends over about 5 500 000 km², from 134°28'W to 154°40'W and from 7°50'S to 27°S (Fig. 1a). Its sea floor includes a broad zone, the so-called 'Pacific Superswell' which is 250 to about 1000 m shallower compared to the depth predicted for its age by classical seafloor subsidence models (McNutt and Fischer, 1987; McNutt, 1998; Adam et al., 2014). It is typified by a great number of volcanoes, aseismic volcanic plateaus and 184 seamounts, aligned along south-east to north-west linear (N115°E) trends, consistent with the predicted direction of absolute motion of the Pacific plate (Duncan and Clague, 1985). By contrast, the underlying oceanic crust exhibits structural features, including south-west to north-east-oriented, transform faults and magnetic lineations (Patriat et al., 2002) and volcanic ridges (Jordahl et al, 2004), and thus, oblique to the hot-spot-generated island trails (Ito et al., 1995).

- 192 The Tuamotu Plateau, part of the oceanic superswell,
- encompasses an area of approximately 1500 km long and 800
- km large (Fig. 1b). It does not exceed 2000 m in bathymetry
- and is composed of basaltic flows, postulated to have erupted
- 196 near the East Pacific spreading centre. The plateau is known to
- 197 be aged of 70 (Okal and Cazenave, 1985) to less than 50
- million years (Schlanger et al., 1984).
- 199 Accretion of the north-west atoll-bearing volcanoes is
- attributed to the passage of two hotspots 50–30 million years
- ago as they migrated beneath the Pacific plate. However, the
- 202 morphology and orientation of a number of linear volcanic
- features onto the plateau were controlled by tectonic structures
- of the underlying lithosphere. (Ito et al., 1995). Carbonate
- deposition over the oldest volcanoes is likely to have begun as
- early as 49 million years (Jackson and Schlanger, 1976).
- 207 2.2 Location of the study area.
- 208 Located between 14°33'S-14°43'S and 145°08'W-145°16'W,
- 209 Takapoto Atoll rises from the northwestern portion of the
- Tuamotu Plateau (Fig. 1b). Its total surface area is 74 km²,
- about 17 km long and 5.5 km wide. The atoll is elongate,
- 212 oriented NE-SW, with an approximately 350 m wide rim
- 213 enclosing a -45 m-deep lagoon. The outer part of the atoll rim
- 214 is a living reef system, subdivided into three zones, from the
- sea inwards (Chevalier et al., 1979; Salvat, 1981; Salvat and

216 Richard, 1985; Laboute, 1985; Kühlmann and Chevalier, 1986; 217 Montaggioni, 1981, personal observation). The outer reef zone 218 comprises a spur-and-groove system, extending downwards to 219 about 6-10 m, followed by sub-horizontal to gently sloping 220 terraces to depths of 10-15 m, previously interpreted as the top 221 of the last interglacial high-stand reef (Montaggioni et al., 222 2018). Below this, low gradient to sub-vertical slopes are 223 developing to depths greater than 60 m. The emergent, 224 innermost parts of the spurs are capped by an algal crest, about 225 20 m wide, reaching 0.20 m to 0.50 m above low tide level. 226 The climate in the northern Tuamotu is tropical, warm and 227 humid. The rainy, austral summer lasts from November to 228 April and the relatively cooler, and dryer, austral winter from 229 May to October. Air temperatures vary between 23 °C and 230 30 °C. Annual rainfall averages 1500 mm. Tides are a twice 231 daily regime and microtidal, 0.5 m in average amplitude and 232 0.70 m in maximum at spring tide. As in all Tuamotu islands, 233 Takapoto is influenced by the south-east trade-winds, blowing 234 from the ENE sectors for 70 % of the year and for 20 % from 235 the SE sector. In the austral winter, 2.5-m high swells are 236 generated by strong trade-winds. During the austral summer, 237 trade-winds are significantly less active and generate moderate-238 energy wave conditions, locally disturbed by storm events 239 initiated either by ENSO-related cyclones or by storms originating from northern latitudes (Andréfoüet et al., 2012). 240

- 241 The eastward migration of warmer sea surface temperatures
- 242 during ENSO episodes promotes the displacement of tropical
- 243 storms further east into the Tuamotu region (Larrue and
- 244 Chiron, 2010).
- Little is known about frequency of extreme wave hazards, i.e.
- 246 tropical cyclone- and tsunami-generated. Over periods of
- several centuries, the number of cyclonic events has probably
- 248 not exceeded 2-3 per century. The effects of cyclones and
- storms in shaping atoll morphology at the long-term scale is
- 250 poorly documented (Montaggioni et al., 2018) onto atoll
- surfaces (Harmelin-Vivien and Laboute, 1986). Similarly, there
- is little information about the frequency of tsunami occurrences
- 253 in the Tuamotu. Based on numerical modelling and direct
- 254 observations, the Tuamotu islands seem not to have
- significantly suffered from tsunami impacts compared to some
- high-volcanic islands in French Polynesia (Sladen et al., 2007).

257 3 Methods and data

- 258 3.1 Acquisition and analysis of geomorphic data
- 259 Pleiades-1 satellite imagery was collected over Takapoto Atoll
- on September, 22 2016 at 20 h 01 min 03 sec (UTC). Orbiting at
- 261 694 km altitude, Pleiades-1 sensor provides four spectral
- 262 wavebands (blue: 430 nm-550 nm, green: 490 nm-610 nm,
- 263 red: 600 nm-720 nm, and infrared: 750 nm-950 nm) at 2 m
- pixel size. The processed imagery totaled 72 294 672 pixels in

265 the form of 9759 lines and 7408 columns. Given the sun-266 sensor-scene trigonometry, a geometric correction was 267 necessary to project multispectral imagery in UTM 6 South 268 within RGPF local datum. For the sake of standardization and 269 further comparison, a radiometric correction was carried out. It 270 firstly converted top-of-atmosphere (TOA) digital number into 271 TOA radiance using calibration factors, secondly TOA 272 radiance into down-of-atmosphere (DOA) radiance by 273 compensating for atmosphere absorption, and thirdly DOA 274 radiance into DOA reflectance by sun irradiance normalization. 275 In parallel, a sonar campaign, aiming at calibrating-validating a 276 depth model, was conducted, from September, 25 to 26, 2016, 277 to acquire water depth soundings over both lagoon and ocean 278 parts (Fig. 2a). Fieldwork week was close enough to imagery 279 date to be confident with an absence of bathymetric 280 disturbances between two periods. An array of 276 082 281 geolocated measurements was compiled across 104.4 km over 282 16 h 44 min recordings. The dataset ranged from 0.54 m to 283 180.85 m with an average of 9.90 m \pm 10.02 m. The sonar 284 device, installed on a fishboat, was a 200 kHz echosounder 285 Lowrance HD7 leveraging a 10 Hz GPS unit. Water level 286 driven by tide was corrected using the nearest French Marine 287 gauge in Rangiroa atoll. 288 The bathymetry has been extracted based on the optical law, 289 showing that electro-magnetic radiation is absorbed by water in respect to the wavelength. Pleiades-1 infrared waveband enabled to discriminate land from sea areas, owing to the strong water absorption of the infrared. Three remaining visible Pleiades-1 wavebands (blue, green and red) were used to elucidate the bathymetry based on the ratio transform (Stumpf et al., 2003):

$$Z = a_1 \left(\frac{\ln(R_w(\lambda_i) - R_\infty(\lambda_i))}{\ln(R_w(\lambda_i) - R_\infty(\lambda_i))} \right) - a_0 \tag{1}$$

297 where a_0 is the intercept for a null water depth (0 m), a_1 is the 298 modeled slope converting the ratio to absolute bathymetry, $R_{\rm w}$ 299 is the DOA reflectance related to the waveband λ_i , and R_{∞} is the 300 DOA reflectance above no-bottom water. Both intercept and 301 slope stemmed from soundings, whose values were re-adjusted 302 by an offset linked with the tide level observed at the satellite 303 collection time. 304 very strong agreement between waterborne 305 observations and spaceborne modeled depths was reached $(R^2 = 0.95, RMSE = 0.17 m$, see Collin et al. 2018 for more 306 307 details). Since the model accuracy declined with depth, a 308 truncature at 38 m was applied. The formula, underlying the 309 model, was thereafter implemented to every pixel, in order to 310 build a digital depth model spanning the entire scene captured 311 by the satellite. 312 Finally, a set of 41 bathymetric profiles was selected (Fig. 2b) 313 and visually inspected to define the type and spatial distribution 314 of morphological features imaged on the atoll shelf slopes. 315 Especially, criteria used to define terraces include the 316 identification of flat-topped to low-gradient surfaces and major 317 breaks-in-slope across the island shelves. The geomorphic 318 features are classified according to their sizes, from larger-scale 319 (hectometres to kilometres) to finer-scale (metres to 320 decametres). The geomorphological classification used in this 321 study is adapted from previously published descriptions and in 322 common usage (see Beaman et al., 2008 and Addey et al., 2011 323 for reviews). Table 1 and figures 2b and 3 present the different 324 geomorphic features. 325 Shelf-slope inclination and size, elevation and width of the 326 different geomorphic features were manually measured from 327 the recorded transects. To measure terrace width, slope 328 extension and inclination, and graphically compare the depth 329 ranges of the successive terrace levels, positive reliefs, such as 330 pinnacles and coral heads, were masked.

331 3.2 Acquisition and analysis of biologic data

332

333

334

335

336

337

The study site is located at the south end of the atoll, close to the village of Fakatopatere and profile T23. Digital images of the quadrats were acquired using a SLR Nikon D300s body camera with aquatica housing and equipped with a Nikkor 15 mm (AF 15 mm f/ 2.8) lens and two SB 105 Nikonos strobes. Pictures were taken at -5 m, -10 m, -15 m, -20 m, -25 m, and

—30 m depth running downslope. At each depth site, 4 pictures were taken perpendicular to the bottom surface with a frequency of 1 picture every 30 seconds by a scuba diver swimming at low speed (1 m per 5 sec). Reef bottom to camera lens distance was approximately 5 m. Digital pictures were processed on computer screen using the point intercept method (50 points—picture). Frequency of coral/non-coral substrate was assessed on each picture by identifying the benthic category present under each point. Except in some cases, coral taxa were identified at the genus level. Amount of living coral cover was obtained by calculating the proportion of points occupied by living coral colonies against the total number of points in a given processed surface area.

4 Results

- *4.1 Morphometry of fore-reef features.*
- Visual examination of the 41 bathymetric profiles (T1 to T41)
- revealed significant variability in distribution, shape and size of
- 355 fore-reef features.
- 356 4.2 Gross morphology
- 357 Six morphologically distinct sectors can be delineated around
- 358 the atoll, based on seaward extension and mean slope gradients
- of the bathymetric profiles (Figs. 4 and 5).

360 The north-end sector (T1 to T5) appears to be the widest with 361 deepest points at 30 m to 36 m, at a distance of 143 m to 190 m 362 from the modern reef-crest line (Fig.4a and 5a). It has relatively 363 gently dipping slopes (23° to 38°) and numerous sub-horizontal 364 terraces, 7 m to 45 m wide. Usually, terraces terminate in steps, 365 1.80 m to 10 m high. There is a prominent line of pinnacles, 366 with elevations of 2.70 m to 17 m. 367 The eastern sector (T11 to T16) exhibits steeply dipping slopes 368 (45°-58°) punctuated by narrow (less than 15 m) terraces 369 ending in steps, 2 m to 15 m high (Figs. 4b and 5b). The 370 deepest points are at 30 m to 37 m, 80 m to 110 m far from the 371 modern reef-crest line. Pinnacles are usually rare or scattered, 372 and not exceed 3.50 m in average elevation. Coral heads are 373 occasional. 374 The south-end sector (T19 to T25) is characterized by steeply 375 dipping slopes (45°-65°) frequently ending in vertical drop-376 offs (Figs. 4c and 5c). The lowest depths recorded range 377 between 26 m to 36 m, at the distance of 70 m to 135 m from 378 the modern reef-crest line. Terrace width varies between 3 m 379 and 15 m. Step height ranges from 2 m to 17 m. Pinnacles are 380 absent or uncommon. 381 The north-western sector (T31 to T41) exhibits a gradient of 382 45° to up to 70°. The deepest points recorded range between 383 6 m and 24 m, at 80 m to 105 m far from the reef front line.

384 There are successive, narrow terraces (2 m–15 m wide) and low 385 steps (1.80 m–12 m). Pinnacles and coral heads are occasional. 386 The steepest foreslopes are located in the north-eastern and 387 south-western sectors of the atoll (T6 to T10 and T26 to T30, 388 respectively) (Fig. 4e and 5e). The mean slope angle varies 389 between 64° and up to 75° (Fig. 6). Along the north-eastern 390 margin, the base of the profiles does not exceed depths of 26 m 391 to 36 m, at the distance of 40 m to 55 m from the modern reef-392 crest line. Along the south-western side, the deepest detected 393 isobath ranges between 9.5 m and 14 m only, at a distance of 394 40 m-50 m from the modern outer reef-crest line. In both 395 sectors, the slope is interrupted by narrow terraces, usually less 396 than 7 m wide ending in low steps and drop-offs, ranging from 397 2 m to 15 m in elevation. The height of pinnacles does not 398 exceed 8.5 m. Coral heads are common. 399 There is locally an atypical zone at the mid-south-eastern side 400 (T17, T18) along which a gently dipping (25°-35°) slope ends 401 in drop-offs not higher than 7 m (Fig. 4a and 5e). The zone 402 between the reef crest and the 38 m isobath extends for a length 403 of 120 m to 210 m. Pinnacles are common, with heights of 2 m

405 4.3 Terrace levels.

to 4.80 m.

404

The morphology of the Takapoto outer-reef slopes is typified by several sets (or levels) of terraces. These lie at relatively

- recurrent depths from the sea surface to about 38 m and have a relatively spatial continuity. Seven morphologically distinct levels of terrace (TL1 to TL7) were recognized (Fig. 7).
- 411 4.3.1 Terraces TL1
- 412 As in most of reef zones facing the open ocean, the outer reef 413 crest extends into a seaward dipping terrace, which is the top 414 surface of the spur-and-groove system (Fig. 8). Locally, this 415 terrace is missing or poorly developed, ending in abruptly 416 dipping (50°-76°) or vertical drop-offs, especially along the 417 north-eastern, eastern (windward) and south-western (leeward) 418 sites. This means that the development of the spur-and-groove 419 system is not directly dependent upon the location and 420 orientation of the foreslopes, since locally lacking from both 421 some windward and leeward sectors. The main driving factor 422 seems to the slope angle; spurs and grooves are usually 423 restricted to more gently dipping slopes, irrespective of 424 exposure to winds and waves. At its seaward end, the depth of 425 TL1 edges range between 2 m and 7 m, with an average depth 426 of $5.24 \text{ m} \pm 2.0 \text{ m}$.
- 427 4.3.2 Terraces TL 2
- This terrace, identified from 34 profiles, is found at the average depth of 10.1 m ± 2.4 m (Figs. 6, 8). It is missing from narrow, steeply dipping slopes and drop-offs (profiles T6 to T10, T22,

- T23, T27 to T30). The surfaces seem generally to have a
- bumpy morphology generated by thick and widespread coral
- layers or metre-high coral heads, locally oriented perpendicular
- 434 to the terrace margins. In most cases, the seaward margin of
- TL2 ends in a steep break-in-slope. Another point to note is the
- presence of outer reef-crest-like morphologies, less than 2 m
- high, at the seaward margin of the terraces.
- 438 4.3.3 Terraces TL 3
- On 28 profiles, a third set of terraces is identified at an average
- depth of 14.7 m \pm 3.8 m. These terraces are missing from a
- number of transects because of local strong steepness (T4, T8,
- 442 T9, T10, T20, T26 to T33, T37, T39 to T41).
- 443 4.3.4 Terraces TL4
- Another level of terraces is encountered at an average depth of
- 20.3 m \pm 2.3 m on 20 out of the 41 profiles. Yet, the strong
- steepness of the fore-reef slopes makes the mid-fore-reef zone
- deeper than 20 m not accessible, as seen on T21 to T30, T33,
- 448 T34, T35, T36, T39 to T41. Despite high coral cover rate, the
- occurrence of TL4 can be inferred from T17 to T20 based on
- sharp changes in slope.

- 451 4.3.5 Terraces TL5
- The relevant terraces are found from 21 sites at an average
- depth of $-24 \text{ m} \pm 2 \text{ m}$. These are missing from the transects T6,
- 454 T13 à T17, T26 to T41.
- 455 4.3.6 Terraces TL6
- 456 These terraces are inferred from seven profiles showing gently
- shelving surfaces (T3, T11, T14 to T17, T20, and T23) and
- 458 located at an average depth of 30.5 m \pm 3 m.
- 459 4.3.7 Terraces TL7
- The deepest terrace recorded is at 38 m, only from T17 and
- 461 T18. Examining patterns among profiles show similarities and
- 462 contrasts in the number and width of terraces. In order to
- determine the number of terraces within a given width range, a
- 464 frequency histogram was generated (Fig. 9). Among the 112
- terraces identified from the 41 profiles, most have widths less
- 466 than 20 m.
- 467 Furthermore, the width of terraces appears to be partly
- 468 correlated to the length of profiles (Fig. 10) which is slope-
- gradient dependent. Where profile lengths do not exceed 80 m,
- 470 most terrace widths are less than 8 m. On profiles as long as
- 471 100 m-110 m, a number of terraces can be as wide as 15 m to
- 472 20 m. The profiles longer than 130 m have one-quarter of their
- terraces, 25 m to 45 m wide.

474 4.4 Pinnacles

- Steep-sided, high reliefs, found on terraces, were interpretated as isolated pinnacles. These are widepread along gently inclined slopes (profiles T1 to T6), but less common on steeply dipping slopes. Their elevation and diameter range from about 2.30 m to 15 m and 2 m to 12 m respectively. However, it is neccessary to remain cautious about inferring the possible lateral extent of a pinnacle line from transects.
- 482 4.5 Drop-offs, trenchs, and scars
- Drop-offs of 11 m to 13 m high are encountered at several shelf 483 484 locations (T13, T19, T21, T26, T27, T32, T37) at different 485 depth ranges, from -9 m to -35 m deep. Trenchs are observed 486 on profiles T24 and T31 (Fig. 5d), in the form of 6 m to 17 m-487 deep indentations, at depths of -6 to -12 m and -12.5 to -29.5 488 m respectively. Concave scars, interpreted as collapsed margin 489 portions, are found at the south-eastern side and along the 490 western side of the atoll (Fig. 2a).
- 491 4.6 Quantitative assessment of coral communities
- Living coral cover increases downslope from 24 %–25 % at depths of –5 m and –10 m to 42 %, 55 %, 67 %, and 50 % at depths of –15 m, –20 m, –25 m, and –30 m respectively (Table 2, Fig. 11). Robust-branching *Pocillopora* (mainly, *P. verrucosa*) is the dominant genus at the upper fore-reef

497 settings, representing 10 % to 26 % of coral coverage, from 498 -5 m to -10 m deep. At depths of -25 m to -30 m, 499 pocilloporids do not exceed 2 %-4 %. Sub-massive to plate-500 like Porites (P. rus) becomes the prominent coral form from 501 15 m deep, locally covering 50 % to up to 85 % of the 502 substrate. Other scleractinian corals belong to Acropora, 503 Porites cf. lobata, Leptastrea, and Montipora, each of them not 504 exceeding 2 % to 4 % of the total coral cover. It is noteworthy 505 that along steeply dipping slopes and drop-offs, modern coral 506 buildups have significantly re-shaped antecent substrate 507 morphology (Fig. 12)

508 **5 Discussion**

509

5.1 Mode and timing of terrace formation

510 As pointed out by Rovere et al. (2018), in absence of 511 radiochronological constraints, the age and formational patterns 512 of marine terraces remain speculative. However, different 513 possible scenarios for their mode and timing of formation can 514 be discussed, especially since two processes, respectively 515 constructional or erosional, may have been involved in terrace 516 shaping, either prior to the last glacial maximum (-23 ka to 517 -20 ka-interval) over the past 300 ky or afterwards, during the 518 last deglaciation. 519 From a contructional point of view, most submerged terraces 520 identified in this study, especially those from gently sloping

521 shelves (profiles T1 to T5, T17, T25, T35) may have developed 522 according to one of the two following models. First, these may 523 result from backstepping of reef bodies developed during the 524 last deglacial rising sea level or previous deglaciation times. In 525 these cases, reef-related terraces would have been generated by 526 punctuated sea-level jump events that have forced reef bodies 527 to drown and backstep upslope through time, rather than 528 develop upwards from a gradual and continuous sea-level rise. 529 Secondly, the terraces may have been shaped during one of 530 high sea-stands that have occurred over the late to mid-531 Pleistocene. In all scenarios, terraces represent the still visible, 532 outermost surfaces of reef edges of former reef tracts. 533 From an erosional point of view, submerged terraces may result 534 from marine planation during episodes of slowing in sea-level 535 rise or stillstands during the last deglaciation. In this scenario, 536 terraces would be wave-cut platforms, as suggested in the 537 Maldives for the last deglaciation (Rufin-Soler et al., 2013; 538 Green et al., 2014). 539 At Takapoto, as everywhere in French Polynesia, during the 540 post-glacial transgressive phase, sea was at a level 541 corresponding to the deepest isobath recorded in the profiles 542 (-38 m) by -9.5 ka. Sea level continued to rise at a rapid and relatively constant rate of 10 mm · y⁻¹ before reaching its 543 544 present position by about -6 ka (Bard et al., 1996, 2010). 545 Slowly rising sea level is known to provide the much-needed

accommodation space for enhanced vertical reef accretion while lateral accretion is dependent of the relative time spent at sea level. According to the mean vertical reef accretion rates $(4-5 \text{ mm} \cdot \text{y}^{-1})$ estimated for the early Holocene (see Dullo, 2005; Montaggioni, 2005; Hubbard and Dullo, 2016; Gischler and Hudson, 2019, for reviews), the rate of sea-level rise was most likely too high to allow coral growth to maintain pace with sea level or to prograde oceanwards. As observed on the mid-shelf slopes at Tahiti (Seard et al., 2013), during the postglacial transgression, corals have rather built isolated, straightwalled pinnacles or thin veeners covering the Pleistocene antecedent topography than typical, flat-topped fringing reefs. A similar scheme was described from Mayotte Island (Comoros, Western Indian Ocean) by Dullo et al. (1998). By analogy, at Takapoto, the major morphological characteristics of the fore-reef slopes are assumed to dominantly reflect those of the antecedent substrate upon which coral buildups have grown more recently. Accordingly, the core of most terraced structures is probably late to mid Pleistocene in age. The height of steps is probably indicative of the thickness of the related reef unit. Additionally, a sea-level rise at the rate of 10 mm \cdot y⁻¹ appears inappropriate to allow shelf slopes to be abrased or excavated. Similarly, during the penultimate (from MIS 6b to MIS 5e) and oldest (MIS 10 to MIS 9, MIS 8 to MIS 7) deglaciation episodes, sea level rose at rates similar to or higher

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571 than those recorded from the last deglaciation (Siddall et al., 572 2007), precluding any significant constructional, abrasional or 573 excavational event to have occurred. Accordingly, the identified terraces and associated reef units were assumed to 574 575 have formed during high sea stands. 576 Some terraces possess outer crest-like reliefs of 0.5-1.5 m in 577 elevation and 3-6 m in width at the seaward margin (for 578 intance, see Fig. 5a, b). Given their specific location on top of 579 the outermost part of submerged reef edges, these features are 580 interpreted to be isolated coral heads or laterally extending, 581 continuous ridge-like, coralgal buildups developed 582 windward, high-energy settings during the Holocene course of 583 sea-level. Contrary to most of pinnacles described along the 584 shelf slopes at Takapoto, their upward development would have 585 been drastically interrupted in response to the rapid sea-level 586 rise. In some way, these are « give-up » reefal features in the 587 sense of Neumann and Macintyre (1985). Morphologically 588 speaking, these buildups are similar to windward, high-energy, 589 coralline algal ridges (Adey, 1978, 1986; Montaggioni and 590 Braithwaite, 2009), typical of Pacific atoll rims (Chevalier et 591 al., 1969), particularly at Takapoto (Chevalier et al., 1979; 592 Chevalier and Salvat, 1976). Submerged terraces with raised 593 coralgal ridges at their outer extent are known from a number 594 of Pacific reef sites: Australian Great Barrier Reef (Beaman et 595 al., 2008; Abbey et al., 2011; Webster et al., 2018), Hawaii

596 (Webster, 2004a) and Papua New Guinea (Webster et al., 597 2004b, 2009), Ryukyu Islands (Arai et al., 2016) and the 598 Marquesas (Cabioch et al., 2008). These terraces are thought 599 primarily to be remnants of coralgal piles developed at the 600 seaward margins of incipient fringing reefs during the post-601 glacial times, but not capable of keeping pace with rising sea 602 level. 603 At Takapoto, terraces TL2 exhibit locally low, coral-built 604 ridges oriented perpendicular to the reef-edge line and 605 alternating with elongated, sand-filled grooves, similar to 606 spurs-and-groove reliefs (Fig. 11b and Montaggioni, personal 607 observations, 1981). Accordingly, TL2 are regarded as outer-608 reef edge features. By analogy, the terraces identified 609 downslope (TL3 to TL7) may be also interpreted to be 610 remnants of paleo-reef-edge surfaces. 611 Along both gently inclined terraces and shelf slopes, pinnacles 612 are common (for instance, see Fig. 5a-f). These features were 613 also described on a number of submerged tropical shelves, 614 mostly on the Australian Great Barrier Reef where they are 615 usually associated with modern juvenile reefs (Hopley, 1982). 616 According to Abbey et al. (2011), during the last deglaciation, 617 terraces would not be ideal settings for extensive vertical coral 618 accretion, due to their narrowness. Similarly, at Takapoto, 619 during the final course of the rising sea level, from about 620 9.5 ka, and despite the subsequent increase in accommodation

space, coral growth was unable to produce, flat-topped, aggrading reefs.

5.2 Faulting and submarine landslides

623

624 The present data reveal that, at both windward and leeward 625 settings, except in its north extremity and mid-south-east, part 626 the Takapoto rim is bounded by steep fore-reef slopes, with gradients up to 45° or vertical submarine cliffs. There are no 627 628 significant differences in the overall fore-reef morphology 629 between windward and leeward sides of the atoll. The steepest 630 fore-reef slopes were found as well along the western, leeward 631 coast as the eastern, windward one (Figs.4 and 5). This 632 suggests that the swell regime from the east is not the main 633 driver of foreslope evolution. 634 The atoll seaward margins are nearly linear as opposed to most 635 of the curved high volcanic island margins found in the nearby 636 Society and Marquesas archipelagos, thereby suggesting 637 structural failure. Faulting control in shelf shaping is further 638 supported by both the presence of vertical drop-offs, higher 639 than 10 m, and trenchs, 6 m to 17 m deep, cutting through the 640 upper parts of the shelf slopes. The trenchs are considered to be 641 incipient, normal faults. Similar features were reported from 642 Tikehau (Faure and Laboute, 1984), and Moruroa (Chevelier et 643 al., 1969). The overall tectonic evolution of the Tuamotu 644 plateau, appears to be partly driven by extensional processes in

645 relation to major, regional fracture zones. The fault 646 distributional patterns of nearby, high-carbonate Makatea 647 Island (Montaggioni, 1985) are quite similar, oriented in a 648 south-west to north-east (35°N-50°E) direction. It is 649 noteworthy that the lagoons of Takapoto and other atolls of the 650 outer Northern most Tuamotu island-belt (Takaroa, Ahe, 651 Manihi, Raroia, Takume), together with that of southermost 652 Amanu atoll, are aligned according the same direction. These 653 atolls are superimposed upon ridges, that are expected to have 654 been formed from erupting fissures influenced by the 655 underlying lithosphere fractures (Jordahl et al., 2004). The 656 dominant fault system affecting the western and eastern shelf 657 slopes at Takapoto is clearly oriented parallell to the atoll and 658 ridge alignment. By contrast, the island shelves at the north 659 end, which developed outwards at right-angles to the SW-NE 660 fault system, have probably escaped large-scale collapsing, thus 661 explaining their unusual morphology, i.e. large seaward 662 development, high density of pinnacles and lack of high 663 escarpments. 664 Scars in the normally convex seaward margins of atolls and 665 reef barriers as well were usually interpretated as resulting from 666 submarine landslides generated by slope failure (Fairbridge, 667 1950; Keating, 1987, 1998; Keating and McGuire, 2000; Riyaz 668 and Park, 2008; Terry and Goff, 2012, 2017; Webster et al., 669 2016; Poupardin et al., 2017; Puga-Bernabéu et al., 2017). As

emphazised by Terry and Goff (2017), submarine landslides
can significantly modify gross atoll morphology. In addition,
giant landslides can generate tsunamis (Webster et al., 2016)
and subsequent, partial submersion of atoll rims (Poupardin et
al., 2017). Although rare, especially in the Tuamotu (Sladen et
al. 2017), landslide-generated tsunamis can contribute to atoll
shaping and island formation.

677 5.3 Significance of terraces in terms of sea level

678 In absence of a chronological frame, as a first approximation, 679 the submerged terraces identified at Takapoto may refer to 680 higher or lower Pleistocene sea-stands relative to present sea 681 level. Hovewer, over the last 500,000 years, sea level during 682 interstadials and glacial, low stands were systematically at 683 depths greater than -30 m (see Murray-Wallace and 684 Woodroffe, 2014 for reviews), especially at atoll-subsiding 685 sites (Camoin et al., 2001). Accordingly, at Takapoto, all the 686 terrace sets were assumed to refer to interglacial, high-sea 687 stands. 688 Morphological and chronostratigraphical correlations between 689 Takapoto and some other atolls used as reference sites in the 690 Pacific can be attempted to help attributing a given terrace level 691 to a specific highstand. All reference cores used herein were

extracted from outer reef-flat settings (Fig. 13).

692

693 On Moruroa Atoll, there is an upper, gently sloping (10°) 694 terrace, about 60 m wide, extending from about -4 m to -10 m 695 deep. It is interrupted by a break-in-slope followed by a steeper 696 (25°) mid-shelf topography (Chevalier et al., 1969). Similar 697 morphologies were described from Enewetak and Bikini Atolls, 698 Marshall Islands (Tracey and Ladd, 1974), Tarawa, Gilbert 699 Islands (Marshall and Jacobsen, 1985) and Yongle Atoll, Xisha 700 Islands, South China Sea (Guozhong et al., 1990). The break-701 in-slope at approximately -10 m deep was regarded as 702 superficially translating the presence of an unconformity, i.e. 703 subaerial exposure surface. This surface is encountered beneath 704 the modern outer reef-flat and reef-edge zones at depths of 705 −10 m to −14 m at Enewetak (Tracey and Ladd, 1974; Lincoln 706 and Schlanger, 1991; Ristvet et al., 1977; Szabo et al., 1985), 707 -8 m to −10 m at Bikini (Emery et al., 1954; Tracey and Ladd, 708 1974), -11 m to -17 m at Tarawa (Marshall and Jacobsen, 709 1985), and -16 m to -17.3 m at Yonge Atoll (Jian et al., 1997). 710 Uranium-series dating of coral material collected above and below this unconformity revealed that this surface separates an 711 712 overlying Holocene reef unit not older than 9 ka, from a late 713 interglacial (MIS 5e) unit, dated at $-120 \text{ ka} \pm 10 \text{ ka}$ to 714 $-134 \text{ ka} \pm 5 \text{ ka}$, as observed in Moruroa, Enewetak, Bikini, and 715 Tarawa (Fig. 13). The thickness of the Holocene reef sequences 716 ranges approximately between 12 m and 17 m. At Takapoto, 717 the uppermost reef unit, spur-shaped, approximates a maximun

718 thickness of 10 m. By analogy with the reference atolls, it is 719 considered to be Holocene in age, probably not older than 9 ka. 720 On the basis of morphological and structural similarities 721 between the upper parts of outer-edge zones among atolls, 722 terraces TL2 were thoughh to be geomorphic analogs of the 723 unconformity surfaces atop of MIS 5e reef units. During the relevant high sea-level episode (-128 ka to -116 ka; Muhs, 724 725 2002), sea level was between 5.5 m to 9 m above psl (Dutton 726 and Lambeck, 2012). Based on statistical modelling, there 727 would have a 95 % confidence level that global sea level, 728 irrespective of local controls, peaked at 6.6 m above psl and a 729 67 % probability that sea level was higher than 8 m above psl 730 during MIS 5e, but it was unlikely to have been higher than 731 9.4 m (Kopp et al., 2009). 732 A second subaerial exposure surface was found in the same 733 reference cores at Moruroa, Enewetak, Bikini and Yongle, at 734 depths of -18 m to -22 m, -22 m to -23 m, -24 m to -27 m, 735 and 30 m respectively (Fig. 13). The reef sequence capped by 736 this surface yielded U-Th ages averaging -215 ka at Moruroa 737 (Camoin et al., 2001) thus indicating that the relevant reefs 738 have developed during the penultimate interglacial (MIS 7) 739 high stand. No date is available from other atoll sites. On these 740 atolls, the depth of the second unconformity relative to psl is 741 locally greater than that of terraces TL3 (14.7 m \pm 3.8 m), but 742 closer to the level of terraces TL4 (20.3 m \pm 2.3 m) and TL 5

743 $(24 \text{ m} \pm 2 \text{ m})$. It is obvious that attempting to link the terraced 744 surfaces TL3, TL4, and TL5 to MIS 7 substages result in 745 significant uncertainties, in part, induced by differences in local 746 erosional and tectonic patterns (see sections 4 and 5 stage). 747 Among the reference atolls, the total thickness of reef units 748 deposited during the entire MIS 7 stage is similar, varying from 749 11 m to 15 m. At Takapoto, the thickness of the intervals 750 delineated by TL3-TL4, TL4-TL5, and TL5-TL6 range 751 between 5 m and 7 m only. Such thickness values appear to be 752 too low to account for the entire MIS 7 depositional history of 753 Takapoto when compared to other atoll sites and with regard to 754 the thermal subsidence constraints of the atoll (see section 5 755 below). On the contrary, integrating the whole TL3–TL6 depth 756 interval (average = 16 m) provides a greater reliance on 757 depositional patterns for the entire MIS 7 episode in the studied 758 area. Furthermore, while there is often an inability to 759 distinguish one substage to another within the penultimate and 760 older interglacials from cored reef sequences (Camoin et al., 761 2001; Montaggioni et al., 2015), identification of distinct sea-762 level pauses from a series of closely stepped terraces seems 763 achievable if assuming that terrace formation is the result of 764 successive backstepping events during successive high-stand 765 substages. Each terrace may correspond to the outer-edge 766 surface of one single backstepped reef tract as observed in a number of reef provinces worldwide (Barbados: Fairbanks, 767

768 1979; the Marquesas: Cabioch et al., 2008 – Hawaii: Sanborn 769 et al., 2010 - Texas: Khann et al., 2017 - Great Barrier Reef, Australia: Webster et al., 2018). Accordingly, the best 770 771 interpretative scenarios for TL3, TL4, and TL5 are the 772 following. These terraces may correlate respectively with MIS 773 7a (-201.5 ka to -189.7 ka), MIS 7c highstand (-217.2 ka to 774 -206 ka) and MIS 7e (-248.9 ka to -231 ka) (Figs.13 and 14). 775 Few coastal landforms referring to MIS 9 (-335.6 ka to 776 -316.8 ka), especially in reef provinces, are documented, most 777 likely due to lower preservation potential and reduced 778 accessibility (Murray-Wallace and Woodroffe, 2014). At 779 Moruroa, a third solution unconformity was recognized in 780 vertical cores between 29 m to 32 m below psl. It caps a 7-781 11 m thick, reef unit (Buigues, 1997). This unit is regarded as 782 the lateral equivalent of reef-margin deposits recovered from oblique boreholes and dated at $-332 \text{ ka} \pm 12 \text{ ka}$ (MIS 9e 783 784 substage) by Camoin et al., (2001). On Enewetak and Bikini, a 785 third unconformity was penetrated at 36.2-42.5 m (Ristvet et 786 al., 1977; Szabo et al., 1985), and at 49 m below psl (Tracey 787 and Ladd, 1974) respectively. In the latter site, a coral sample 788 from the underlying interval gave an age older than 400 ka, 789 apparently inconsistent with local stratigraphy. To maintain 790 consistency with the overall development history of Takapoto 791 submerged terraces, TL6 are approximated to refer to MIS 9e.

- The belonging of terraces TL7 to a given MIS stage is facing considerable uncertainty and therefore is not discussed.
- 794 5.4 Effects of subaerial erosion
- 795 Another prerequisite to define a paleo-relative sea-level 796 position from given terrace levels is quantifying the impact of 797 subaerial erosion on the morphological evolution of terraces 798 during times of emergence. From estimates of the amount of 799 reef rocks destroyed over the last 120 ky, the rates of carbonate 800 ground lowering from some Pacific atolls and reliefs were thought to range between 0.011 mm \cdot y⁻¹ and 0.202 mm \cdot y⁻¹, 801 with mean rates of $0.11 \text{ mm} \cdot \text{y}^{-1}$ (Paulay and McEdward, 802 803 1990). These values are similar to those calculated $(0.12 \text{ mm} \cdot \text{y}^{-1} \text{ on average})$ by Purdy and Winterer (2001) from 804 805 Enewetak Atoll for the same time span in response to local 806 rainfall regime. Since present-day annual rainfall at Takapoto is around 1500 mm · y⁻¹, quite similar to that recorded at 807 808 Enewetak (Purdy and Winterer, 2001), a mean value of $0.12 \text{ mm} \cdot \text{y}^{-1}$ is taken as the maximum solution rate during 809 810 Pleistocene low sea stands in the north-western Tuamotu.
- 811 5.5 Subsidence
- In the Tuamotu, the thermal sinking rate of the underlying oceanic crust is lower than the usual decaying exponential rate (about $0.32 \text{ mm} \cdot \text{y}^{-1}$ – $0.35 \text{ mm} \cdot \text{y}^{-1}$: Menard, 1969; Korenaga

815 and Korenaga, 2008) as a function of increasing age as for most 816 of oceanic volcanic domains (Patriat et al., 2002). The Tuamotu 817 plateau. In relation to its age (47 to 55 million years old), the 818 Tuamotu plateau is considered to be near isostatic equilibrium 819 since it was emplaced (Patriat et al., 2002). Moreover, for the 820 north-western Tuamotu islands, earth dynamic topography 821 models predicted about 2-3 m fall since the last 125 ky, which 822 is included in the subsidence signal (Austermann et al., 2017). 823 Quantitative constraints on the possible rate of subsidence at 824 Takapoto can be provided on the basis of the following 825 assumptions: (1) subsidence is expected to approximate 826 linearity for the past 300 ky; (2) TL2- related reef unit formed 827 during MIS 5e high stand; (3) at that time, sea level peaked at 828 +7 m above psl (Rovere et al., 2016); (4) the rate of sub-aerial erosion averaged 0.12 mm \cdot y⁻¹; (5) the time span during which 829 830 MIS 5e reef was exposed lasted 106.5 ky, from 116 ka to 831 9.5 ka, i.e. from the termination of the MIS 5e substage to the 832 time of re-flooding of the shelf slope from the deepest detected 833 isobath. As a result, the ground lowering of the TL2-related 834 reef pile during emergence can be estimated at around 12 m. If 835 correct, given TL2 is now at about 10 m deep, the subsidence 836 amplitude would be about 5 m. The calculated subsidence rate 837 is $0.047 \text{ mm} \cdot \text{y}^{-1}$ for the past 125 ky. This rate is in complete accord with that estimated (0.045 mm \cdot y⁻¹) from Pukapuka 838 839 and Rakahanga Atolls (Gray et al., 1992), the origin of which is

linked to the formation of the Manihiki Plateau as old as the
Tuamotu basement (Mahoney, 1987). Similar thermal
subsidence rates, around 0.035 mm · y⁻¹ to 0.050 mm · y⁻¹
were inferred from other mid-plate low-lying carbonate islands
originated during the early Cenozoic (Schlanger et al., 1987;
Quinn, 1991; Gischler et al., 2008).

5.6 Identification of paleo-relative high sea-levels

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

Paleo-sea-level reconstructions over the last 300 ky were attempted to be achieved on the basis of the vertical distribution of terrace levels at Takapoto (Table 3). The rates of sub-aerial erosion $(0.12 \text{ mm} \cdot \text{y}^{-1})$ and thermal subsidence $(0.047 \text{ mm} \cdot \text{y}^{-1})$ previously determined were used in the calculations, assuming that similar environmental and tectonic conditions have prevailed during all considered high sea stands. The position of a paleosea level at a given high stand (SL) was calculated as follows: SL = S + E - TL where S is the product of the amount of time during which a given terraced-reef tract has subsided and of the subsidence rate, subsidence operating as soon as reef growth has initiated; E is the produit of the amount of time during which a given reef pile has been emerged and undergone ground lowering and of the erosion rate, considering that duration of emergence is the time interval between the termination of a given high stand and the start of the following re-flooding and subsequent overlapping by a new

reef generation. TL is the present depth at which a given terrace is found (Table 3).

866 Terraces TL3, as remains of MIS 7a reef surfaces, would have 867 suffered a ground lowering of 7.40 m and a drowning of 9.50 m 868 since they have formed, resulting in a deepening of -16.90 m 869 relative to their initial position. Based on its present location at 870 an average depth of $14.70 \text{ m} \pm 3.8 \text{ m}$, the paleo-sea level is 871 estimated to have been at 2.2 m \pm 3.8 m above psl. Regarding 872 terraces TL4, as referred to MIS 7c, ground-lowering and 873 subsidence amplitudes were assessed to be 1.20 m and 9.60 m 874 respectively. The resultant drop did not exceed 10.80 m while 875 these features are found at present at $-20.30 \text{ m} \pm 2.3 \text{ m}$. This 876 means that the related past sea level was at 9.50 m± 2.3 m 877 below psl. Two alternative scenarios can account for the 878 evolutionary history of MIS 7e reefs represented by terraces 879 TL5. In the case where this reef generation was overlapped by 880 younger MIS 7c reefs, the time of emergence has not exceeded 881 11 ky and the thickness of the reef pile released by solution 882 ranged around 1.30 m. If sea level was lower, MIS 7e reefs 883 may have remained exposed over 31 ky. Accordingly, ground 884 lowering of MIS 7e deposits would reach 3.70 m. In both 885 situations, subsidence was 10.30 m. The resulting terrace 886 deepening ranges between 13.0 m and 15.40 m. Given terraces 887 TL5 are now located at a depth of $-24 \text{ m} \pm 2 \text{ m}$, the MIS 7e sea 888 level may have been at approximately 11 m or $8.60 \text{ m} \pm 2 \text{ m}$

889 below psl. The morphological evolution of reefs TL6 is 890 summarized as follows. Sub-aerial erosion would have 891 removed 9.20 m while sinking motion was 15.80 m in 892 amplitude. Finally, terraces TL6 were deepened by 25 m. With 893 respect to their present depth location, the corresponding reef 894 tops were believed to have originally cropped out at a paleo-sea 895 level between 5.80 m \pm 3 m below psl. 896 The sea-level positions derived from fore-reef 897 morphometry at Takapoto appear to be in good agreement with 898 some predicted on the basis of different proxies in different 899 seas (Fig. 14). Regarding the elevations at which sea level 900 peaked during the MIS 7 substages, reconstructions gave 901 evidence for only two prominent high sea levels, MIS 7a and 902 7e, close to or higher than present, between -12 m and +4 m903 relative to psl. These findings are based on U/Th dating of 904 speleothems in Bahamian caves (Li et al., 1989; Lundberg and 905 Ford, 1994), open-system uranium-series ages (Thompson et 906 al., 2003; Thompson and Goldstein, 2005) and ESR ages 907 (Schellmann and Radtke, 2004) of corals collected from 908 uplifted reef terraces at Barbados, stratigraphic and radiometric 909 constraints in the Bahamas and Bermuda (Hearty and Kindler, 910 1995), paired Mg/Ca and oxygene isotope measurements of 911 planktonic foraminifera cooled in a core from the Cocos Ridge 912 (Lea et al., 2002), sea water oxygen isotope in the Red Sea 913 (Siddall et al., 2003) and coupling of the Red Sea sea-level

914 signal with Asian monsoon records (Grant et al., 2014). By 915 contrast, past sea-level reconstructions derived from oxygene 916 isotope analysis of speleothems (Bard et al., 2002; Dutton et 917 al., 2009) would indicate that sea level during the MIS 7 stage 918 maintained continuously below psl, at a depth of around -18 m. 919 Similar results, suggesting that paleo-sea level during the 920 penultimate interglacial stage has fluctuated between 10 m and 921 17 m below psl, were gained from the analysis of inclined reef 922 cores at Moruroa (Camoin et al., 2001). During the MIS 9 923 stage, sea level is inferred to have peaked at levels close to 924 present (Siddall et al., 2006; Rohling et al., 2009; Murray-925 Wallace and Woodroffe, 2014).

926 5.7 Comparison with the composition of coral communities I
 927 in some other Tuamotu and Pacific atolls

928 Compositional and distributional patterns of corals along outer-929 reef slopes from atolls, are known to be controlled, not only by 930 bathymetry and water agitation, but also by reef orientation, 931 and slope gradient (Wells, 1956; Kühlmann, 1982; Bouchon, 932 1983; Faure and Laboute, 1984; Laboute, 1985; Kühlmann and 933 Chevalier, 1986; Maragos, 1994). At Takapoto, as observed in 934 both windward and leeward sides, Pocillopora remains 935 dominating within the upper -10 m to -15 m along gently 936 sloping spurs (TL1), and terraces TL2, while Porites achieves 937 prominence from -15 m to -30 m, along steeper to sub-vertical 938 slopes (Bouchon, 1983; Kühlmann and Chevalier, 1986). 939 Similar trends were reported from the nearby Tikehau Atoll 940 (Faure and Laboute, 1984). 941 Coral cover rates from other Pacific atolls are poorly 942 documented (Wells, 1954; Maragos, 1994). The only detailed, 943 quantitative coral censuses of seaward outer margins were 944 carried out on Mili and Rongelap Atolls (Marshall Islands, 945 central Pacific) at different depths between -5 m and -20 m 946 (Pinca et al., 2006). Porites is by far the most abundant genus 947 (14-62 %) in both leeward and windward sites. Surprisingly, 948 robust branching Pocillopora, 12 % to up to 20 % in cover 949 rates, are restricted to lagoons, apparently missing from open-950 sea environments. By contrast, on some other Marshall Islands 951 (Bikini, northern atolls), as on Takapoto, *Pocillopora* species 952 are amongst the most common outer-reef dwellers, especially 953 in the upper fore-reef parts (Wells, 1954; Maragos, 1994). 954 These observations indicate there can have marked, within-955 region to inter-regional differences in coral composition, 956 maybe due to. competitive interactions between coral groups 957 (Lenihan et al., 2011). Anyway, in the study areas, coral 958 growths contribute intensively to shaping of upper fore-reef 959 slopes to depths greater than 40 m. 960 The present study confirmed the reliability of robust-branching 961 Pocillopora verrucosa as a relative sea-level marker as previously demonstrated in some Indo-Pacific reef sites 962

963 (Montaggioni and Faure, 1997; Montaggioni et al., 1997; 964 Cabioch et al., 1999; Camoin et al., 2001). Composite P. 965 verrucosa-dominated frameworks appeared to be restricted to 966 the 0–10 m depth range.

Conclusions

967

968

969

970

971

972

974

975

976

977

978

979

980

981

982

983

984

985

986

Detailed geomorphological analysis of satellite-based bathymetric profiles obtained from fore-reef zones all around Takapoto Atoll, using spaceborne imagery, together with direct underwater observations of fine-scale geomorphic features and coral community composition, allows the following 973 conclusions to be drawn.

> 1. Six distinct fore-reef sectors were delineated based on seaward extension to the detected slope depths and mean slope gradients. The lowest gradient ($< 30^{\circ}-35^{\circ}$) and most extended (close to 200 m) sectors are situated at the northern atoll end and along a short portion of the mid-south-eastern side. It exhibits wide terraces and low steps. The north-eastern sector displays highest gradient (68° on average) and narrowest (90 m) slopes and narrow terraces. The remaining eastern and northwestern sectors, together with south-western and southern ends have steep slopes (52° to 57° on average), extending seawards over 90 m to 100 m on average, with narrow terraces and high steps.

2. Among the described geomorphic features, of specific interest are terraces, outer-reef crest-like reliefs, pinnacles, steps and drop-offs, scars and trenches to reconstruct the development history of fore-reef zones. Seven levels of submerged terraces were identified at depths averaging -5 m, -10 m, -15 m, -20 m, -24 m, -30.5 m, and -38 m respectively. These range from about 45 m to 2 m in width. The shallowest terrace level refers to the present-day surfaces of spurs. In lack of any radiometric dating, the deeper terrace levels were postulated to be remnants of a series of Pleistocene reef tracts, severely altered by sub-aerial erosion during periods of emergence, and finally overlapped by coral veneers during the post-glacial marine transgression, from about 9.5 ky to present. This scenario is in great part based on comparisons with subsurface reef chronostratigraphy from several Pacific atolls. The rates of sub-aerial erosion and linear subsidence were at $0.12 \text{ mm} \cdot \text{y}^{-1}$ and $0.047 \text{ mm} \cdot \text{y}^{-1}$ estimated respectively from comparisons with calculated surface lowering of Pacific atolls and usually accepted sea-level position at 125 ka. After depth correction of ground lowering and of subsidence effects, the interglacial high sea-stands during which each reef tract has formed and the position of the corresponding sea level were

987

988

989

990

991

992

993

994

995

996

997

998

999

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

tentatively defined. The 10-m, 14.5m, 20-m, 24-m and 30.5-m terraces were expected to relate to the MIS 5e, 7a, 7c, 7e and 9e respectively, despite substantial uncertainties on elevation estimates.

Outer reef crest-like features, less than 2 m high, occur frequently at the seaward edge of terraces. Similarly, steep-walled pinnacles are commun at different places and depths on terraces and slopes. Both were considered to be remains of coralgal buildups, unable to compensate with the post-glacial sea-level rise.

Steps and drop-offs, a few metres to more than 15 m high, are presumably remains of the foreslopes of pre-existing reef tracts or preserved scars of partially collapsed fore-reef slopes. Trenchs cutting the fore-reef zone were found locally and represent detachment fissures. The faces of the western and eastern fore-reef slopes are aligned in south-west to north-east direction, consistent with the main axis orientation of the atoll. This direction is that of atoll-bearing, submarine volcanic ridges discovered in the north-western part of the Tuamotu plateau, generated from and shaped by underlying lithospheric fractures.

3. Compositional and distributional patterns of coral populations at the study site are in agreement with previous investigations in the Tuamotu atolls. Robust

branching *Pocillopora* forms appear to be the main builders on the uppermost, gently dipping parts of forereef zones, to the depth of about 10 m. This explains why this species is regarded as diagnostic in terms of paleo-bathymetry reconstruction. Pocilloporids are replaced by sub-massive to plate-shaped *Porites*, profusely growing from –15 m to –35 m deep at least. Concerning biogeographic distribution of corals across the central Pacific, there are noticeable differences between the Marshall and Tuamotu islands, probably controlled mainly by local ecological constraints. However, along upper fore-reef zones, coral growths appear to play a key role in controlling superficial morphogenesis.

Acknowledgments

The present work was supported by the French national research agency (CNRS) under the STORISK research project (N° ANR-15-CE03-0003) coordinated by Virginie Duvat. Field work has benefited from logistical support and help from the research station of the Office of Marine Resources in Takapoto, Gaby Maiti Haumani. Bathymetric, satellite and echo-sounder-based records were provided by AC and treated by DJ. Data on the composition and distribution of coral communities were provided by GS and MT, and exploited by YC. LM, the corresponding author, analysed

- and interpreted the data, and wrote the manuscript with
- imput from the co-authors. Figures were conceived by LM
- 1064 and BM-G.

1065 References

- Abbey, E., Webster, J.M., Beaman, R.J., 2011. Geomorphology
- of submerged reefs on the shelf edge of the Great Barrier
- Reef: the influence of oscillating Pleistocene sea-levels.
- 1069 Mar. Geol. 288, 61–78.
- 1070 Adam, C., Yoshida, M., Suetsugu, D., Fukao, Y., Cadio, C.,
- 1071 2014. Geodynamic modeling of the South Pacific
- superswell. Phys. Earth Planet. Interiors 229, 24–39.
- 1073 Adey, W. H., 1978. Algal ridges of the Caribbean Sea and West
- 1074 Indies. Phycologia 17, 361–367.
- 1075 Adey, W.H., 1986. Coralline algae as indicators of sea level. In:
- 1076 Van de Plassche, O. (ed). Sea-level Research: A manual
- for the Collection and Evaluation of Data, Springer,
- 1078 Dordretch, p.229–280.
- 1079 Andréfouet, S., Ardhuin, F., Queffeulou, P., Le Gendre, R.,
- 1080 2012. Island shadow effect and the wave climate of the
- 1081 Western Tuamotu Archipelago (French Polynesia) inferred
- from altimetry and numerical model data. Mar. Pollut.
- 1083 Bull. 65, 415–424.
- 1084 Arai, K., Matsuda, H., Sasaki, K., Machiyama, H., Yamaguchi,
- 1085 T., Inoue, T., Sato, T., Takayanagi, H., Iryu, Y., 2016. A
- newly discovered submerged reef on the Miyako-Sone

- 1087 platform, Ryukyu Island Arc, northwestern Pacific. Mar.
- 1088 Geol. 373, 49–54.
- 1089 Austermann, J., Mitrovica, J.X., Huybers, P., Rovere, A., 2017.
- 1090 Detection of a dynamic topography signal in last
- interglacial sea-level records. Sci. Adv. 3, e 1700457.
- 1092 Bard, E., Hamelin, B., Arnold, M., Montaggioni, L.F., Cabioch,
- 1093 G., Faure, G., Rougerie, F., 1996. Deglacial sea-level
- record from Tahiti corals and the timing of global
- meltwater discharge. Nature 382, 241–244.
- 1096 Bard, E., Antonioli, F., Silenzi, S., 2002. Sea-level during the
- penultimate interglacial period based on a submerged
- stalagmite from Argentarola Cave (Italy). Earth Planet.
- 1099 Sci. Lett. 196, 135–146.
- 1100 Bard, E., Hamelin, B., Delanghe-Sabatier, D., 2010. Deglacial
- meltwater pulse 1B and Younger Dryas sea-levels
- revisited with boreholes at Tahiti. Science 327, 1235-
- 1103 1237.
- 1104 Beaman, R.J., Webster, J.M., Wust, R.A.J., 2008. New evidence
- for drowned shelf edge reefs in the Great Barrier Reef,
- 1106 Australia. Mar. Geol. 247, 17–34.
- 1107 Blanchon, P., Jones, B., Ford, D.C., 2002. Discovery of
- submerged relic reef and shoreline off Grand Cayman:
- further evidence for an early Holocene jump in sea level.
- 1110 Sedim. Geol. 147, 253–270.
- 1111 Bouchon, C., 1983. Les peuplements de scléractiniaires de
- 1112 l'atoll de Takapoto (Polynésie Française). J. Soc.
- 1113 Océanistes 39, 35–42.

- 1114 Buigues, D., 1997. Geology and hydrology of Mururoa and
- 1115 Fangataufa, French Polynesia. In: Vacher, H.L., Quinn,
- 1116 T.M. (eds). Geology and Hydrology of Carbonate Islands,
- Elsevier, Amsterdam, Develop. Sediment. 54, 433–452
- 1118 Cabioch, G., Montaggioni, L.F., Faure, G., Ribaud-Laurenti, A.
- 1119 1999. Reef coralgal assemblages as recorders of
- paleobathymetry and sea level changes in the Indo-Pacific
- 1121 province. Quat. Sci. Rev. 18, 1681–1695.
- 1122 Cabioch, G., Montaggioni, L.F., Norbert, F., Seard, C., Sallé,
- E., Payri, C., Pelletier, B., Paterne, M., 2008. Successive
- reef depositional events along the Marquesas foreslopes
- 1125 (French Polynesia) since 26 ka. Mar. Geol. 254, 18–34.
- 1126 Camoin, G.F., Ebren, Ph., Eisenhauer, A., Bard, E., Faure, G.,
- 1127 2001. A 300,000-yr coral reef record of sea-level changes,
- Mururoa Atoll (Tuamotu Archipelago, French Polynesia).
- Palaeogeogr. Palaeoclimatol. Palaeoecol. 175, 325–341.
- 1130 Camoin, G.F., Seard, C., Deschamps P., Webster, J.M., Abbey,
- 1131 E., Braga, J.C., Iryu, Y., Durand, N., Bard, E., Hamelin,
- B., Yokoyama, Y., Thomas, A.L. Henderson, G.M.,
- Dussoulliez, Ph., 2012. Reef response to sea-level and
- environmental changes during the last deglaciation:
- 1135 Integrated Ocean Drilling Program Expedition 310, Tahiti
- 1136 Sea Level. Geology 40, 643–646.
- 1137 Camoin, G. F., Webster, J.M., 2015. Coral reef response to
- 1138 Quaternary sea-level and environmental changes: state-of-
- 1139 the art. Sedimentology 61, 401–428.
- 1140 Chevalier, J.-P., Salvat, B., 1976. Eude géomorphologique de
- 1141 l'atoll fermé de Taiaro. Cah. Pac. 19, 169–201.

- 1142 Chevalier, J.-P., Denizot, M., 1979. Les organismes
- 1143 constructeurs de l'atoll de Takapoto. J. Soc. Océan. 35,
- 1144 30–34.
- 1145 Chevalier, J.-P., Denizot, M., Mougin, J.L., Plessis, Y., Salvat,
- B., 1968. Etude géomorphologique et bionomique de
- 1147 l'atoll de Mururoa (Tuamotu). Cah. Pacifique 12, 1–144.
- 1148 Chevalier, J.-P., Denizot, M., Ricard, M., Salvat, B., Sournia,
- 1149 A., Vasseur, P., 1979. Géomorphologie de l'atoll de
- Takapoto. J. Soc. Océanistes 35, 9–18.
- 1151 Collin, A., Duvat, V., Pillet, V., Salvat, B., James, D., 2018.
- 1152 Understanding interactions between shoreline changes and
- reef slope morphometry on Takapoto Atoll (French
- 1154 Polynesia). J. Coast.Res. SI 496–500.
- 1155 Dullo, W.Ch., 2005. Coral growth and reef growth: a brief
- 1156 review. Facies 51, 33–48.
- 1157 Dullo, W.Ch., Camoin, G.F., Blomeier, D., Colonna, M.,
- Eisenhauer, A., Faure, G., Casanova, J., Thomassin, B.A.,
- 1159 1998. Morphology and sediments of the fore-slopes of
- 1160 Mayotte, Comoro Islands: direct observations from a
- submersible. In: Camoin, G.F., Davies, P.J. (eds). Reefs
- and Carbonate Platforms in the Pacific and Indian Oceans,
- Intern. Assoc. Sedimentologists, Spec. Publ. 27, 217–238.
- 1164 Duncan, R. A., Clague, D. A., 1985. Pacific plate motion
- recorded by linear volcanic chains. In: Nairn, A.E.A. et al.
- 1166 (eds). The Ocean Basins and Margins, 7A, The Pacific
- Ocean, Plenum, New York, pp. 89–121.
- 1168 Dutton, A., Bard, E., Antonioli, F., Esat, T.M., Lambeck, K.,
- McCulloch, M.T., 2009. Phasing and amplitude of sea-

- 1170 level and cimate change during the penultimate
- interglacial. Nature Geosciences DOI: 10.1038/ngeo470
- 1172 Dutton, A., Lambeck, K., 2012. Ice volume and sea level during
- the last interglacial. Science 337, 216–219.
- 1174 Emery, K.O., Tracey, J.I., Jr., Ladd, H.S., 1954. Geology of
- Bikini and nearby atolls: U.S. Geol. Surv. Prof. Paper 260
- 1176 A, 1–265 pp.
- 1177 Faichney, I.D.E., Webster, J.M., Clague, D., Kelley, Ch.,
- Appelgate, B., Moore, J.G., 2009. The morphology and
- distribution of submerged reefs. In the Maui-Nui
- 1180 Complex, Hawaii: new insights into their evolution since
- the early Pleistocene. Mar. Geol. 265, 130–145.
- 1182 Faichney, I.D.E., Webster, J.M., Clague, D.A., Paduan, J.B.,
- Fullagar, P.D., 2010. Unravaling the tilting history of the
- submerged reefs surrounding Oahu and the Maui_Nui
- 1185 Complex, Hawaii. Geochem. Geophys. Geosyst. 11,
- 1186 007002, DOI: 10.1029/2010GC003044
- 1187 Fairbanks, R.G., 1979. A 17,000-year glacio-eustatic sea level
- record: influence of glacial melting rates on the Younger
- Dryas event and deep-ocean circulation. Nature 342, 637-
- 1190 642.
- 1191 Fairbridge, R.W., 1950. Landslide patterns on oceanic volcanoes
- 1192 and atolls. The Geogr. J. 115, 84–88.
- 1193 Faure, G., Laboute, P., 1984. Formations récifales 1: Définitions
- des unités récifales et distribution des principaux
- peuplements de scléractiniaires. In: L'atoll de Tikehau
- 1196 (Archipel des Tuamotu, Polynésie française), Premiers

- 1197 Résultats. Notes et Doc. Océanogr., Office Rech. Sci.
- 1198 Techn. Outre-Mer 22, 108–136.
- 1199 Gischler, E., Hudson, J.H., Pisera, A., 2008. Late Quaternary
- reef growth and sea level in the Maldives (Indian Ocean).
- 1201 Mar. Geol. 250, 104–113.
- 1202 Gischler, E., Hudson, J.H., 2019. Holocene tropical reef
- 1203 accretion and lagoon sedimentation: a quantitative
- approach to the influence of sea-level rise, climate and
- subsidence (Belize, Maldives, French Polynesia). The
- Depositional Record 5: https://doi.org/10.1002/dep2.62.
- 1207 Grant, K.M., Rohling, E.J., Ramsey, C.B., Cheng, H., Edwards,
- 1208 R.L., Florindo, F., Heslope, D., Marra, F., Roberts, A.P.,
- Tamisiea, M.E., Williams, F., 2014. Sea-level variability
- over five glacial cycles. Nature Comm. 5:5076. DOI:
- 1211 10.1038/ncomms6076.
- 1212 Gray, S.C, Hein, J.R., Hausmann, R., Radtke, U., 1992.
- Geochronology and subsurface stratigraphy of Pukapuka
- and Rakahanga atolls, Cook Islands: late Quaternary reef
- growth and sea level history. Palaeogeogr. Palaeoclim.
- 1216 Palaeoecol 91, 377–393.
- 1217 Green, A.N., Cooper, J.A.G., Salzmann, L., 2014. Geomorphic
- 1218 and stratigraphic signals of postglacial meltwater pulses
- on continental shelves. Geology 42, 151–154.
- 1220 Guozhong, W., Bingquan, L., Songqing, Q., 1990. Sedimentary
- characteristics of coral reefs in the northern part of the
- 1222 South China Sea. Atoll Res. Bull. 346, 1–21.

- 1223 Harmelin-Vivien, M.L., Laboute, P., 1986. Catastrophic impact
- of hurricanes on atoll outer reef slopes in the Tuamotu
- 1225 (French Polynesia). Coral Reefs 5, 55-62.
- Hearty, P.J., Kindler, P., 1995. Sea-level high stand chronology
- from stable carbonate platforms (Bermuda and Bahamas).
- 1228 J.Coast. Res.11, 675–689.
- 1229 Hopley, D., 1982. The Geomorphology of the Great Barrier
- Reef: Quaternary Development of Coral Reefs. Wiley-
- 1231 Interscience, New-York, 453 pp.
- 1232 Hopley, D., Smithers, S.G., Parnell, K.E., 2007. The
- Geomorphology of the Great Barrier Reef: Development,
- Diversity and Change. Cambridge Univ. Press, Cambridge,
- 1235 UK, 532 pp.
- 1236 Hubbard, D. K., Dullo, W.Ch., 2016. The Changing Face of
- Reef Building. In: Hubbard, D. K., Rogers, C. S., Lipps, J.
- 1238 H., Stanley, G. D (eds). Coral Reefs at the Crossroads.
- 1239 Coral Reefs of the World, 6. Springer, Dordrecht, pp.
- 1240 127–153.
- 1241 Ito, G., McNutt, M., Gibson, R.L., 1995. Crustal structure of the
- Tuamotu Plateau, 15°S, and implications for its origin. J.
- 1243 Geophys. Res. 100, 8097-8114.
- 1244 Jackson, E.D., Schlanger, S.O., 1976. Regional syntheses, Line
- 1245 Islands chain, Tuamotu Island chain and Manihiki Plateau,
- 1246 central Pacific Ocean. Init Repts, DSDP 33, 915–927.
- 1247 James, N.P., Ginsburg, R.N., 1979. The seaward margin of
- 1248 Belize Barrier and Atoll Reefs. Intern. Assoc.
- 1249 Sedimentologists, Spec. Publ. 3, Blackwell, Oxford, 191
- 1250 pp.

- 1251 Jian, L., Zhizheng, Y., Chunrui, H., Xinbo, L., Gaosheng, Q.,
- 1252 1997. Meteoric diagenesis in Pleistocene reef limestones
- of Xisha Islands, China. J. Asian Earth Sci. 15, 465–476.
- 1254 Jordahl, K., Caress, D., McNutt, M., Bonneville, A., 2004. Sea-
- floor topography and morphology of the Superswell
- Region. In: Hekinian, R., Stoffers, P., Cheminée, J.-L.
- 1257 (eds). Oceanic Hotspots: Intraplate submarine Magmatism
- and Tectonism, Springer, Berlin, pp. 9-28.
- 1259 Keating, B.H., 1987. Structural failure and drowning of
- Johnston Atoll, central Pacific Basin. In: Keating, B.H.,
- 1261 Fryer, P., Batiza, R., Boehlert, G.W. (eds). Seamounts,
- 1262 Islands and Atolls. American Geophys. Union,
- Washington, D.C., Geophys. Monograph Ser. 43, pp. 40-
- 1264 60.
- 1265 Keating, B.H., 1998. Side-scan sonar images of submarine
- landslides on the flanks of atolls and guyots. Mar.
- 1267 Geodesy 21, 129–145.
- 1268 Keating, B.H., McGuire, J.M., 2000. Island failures and
- 1269 associated tsunami hazards. Pure Appl. Geophys. 157,
- 1270 899–955.
- 1271 Khanna, P., Droxler, A., Nittrouer, J.A., Tunnell, J., Shirley,
- Th. C., 2017. Coralgal reef morphology records
- punctuated sea-level rise during the last deglaciation.
- 1274 Nature Comm., 8: 1046. DOI/ 10.1038/s41467-017-00966-
- 1275 x.
- 1276 Kopp, R.E., Simmons, F.J., Mitrovica, J.X., Maloof, A.C.,
- 1277 Oppenheimer, M., 2009. Probablistic assessment of sea

- 1278 level during the last interglacial stage. Nature 462, 863-
- 1279 867.
- 1280 Korenaga, T., Korenaga, J., 2008. Subsidence of normal oceanic
- 1281 lithosphere, apparent thermal expansivity, and sea floor
- flattening. Earth Planet. Sci. Lett. 268, 41–51.
- 1283 Kühlmann, H.H., 1982. Zusammensetzung und ökologie von
- 1284 Tiefwasser-Korallenassoziationen. Wissenschaftliche
- 1285 Zeitschrift der Humbolt-Universität zu Berlin. Math-Nat.
- 1286 R, 31, 233–244.
- 1287 Kühlmann, H.H., Chevalier, J.P., 1986. The corals
- 1288 (Scleractinians and Hydrocorals) of the Takapoto Atoll,
- Tuamotu Islands: ecological aspects. Mar. Ecol. 7, 75-
- 1290 104.
- 1291 Laboute, P., 1985. Evaluation des dégats causés par les passages
- des cyclones de 1982-1983 en Polynésie Française sur les
- pentes externes des atolls de Tikehau et de Takapoto
- 1294 (Archipel des Tuamotu). Proc. Fifth Internat. Coral Reef
- 1295 Congr. 3, 323–329.
- 1296 Ladd, H.S., Tracey, Jr., I.J., Gross, M.G., 1970. Deep drilling
- on Midway Atoll. US Geol. Surv. Prof. Papers 680-A, A1-
- 1298 A22.
- 1299 Larrue, S., Chiron, Th., 2010. Les îles de Polynésie Française
- 1300 face à l'aléa cyclonique. Vertigo 10.
- 1301 https://doi.org/10.4000/vertigo.10558.
- 1302 Lea, D.W., Martin, P.A., Pak, D.K., Spero, H.J., 2002.
- Reconstructing a 350 history of sea level using planktonic
- Mg/Ca and oxygen isotope record from a Cocos Ridge
- 1305 core. Quat. Sci. Rev. 21, 283–293.

- 1306 Lenihan, H.S., Holbrook, S.J., Schmitt, R.J., Brooks, A.J., 2011.
- 1307 Influence of corallivory, competition, and habitat
- structure on coral community shifts. Ecology 92, 1959-
- 1309 1971. doi.org/10.1890/11-0108.1
- 1310 Li., W.-X., Lundberg, J., Dickin, A.P., Ford, D.C., Schwarcz,
- H.P., McNutt, R., Williams, D., 1989. High precision
- mass spectrometric uranium-series dating of cave deposits
- and implications for paleoclimate studies. Nature 339,
- 1314 564–536.
- 1315 Lighty, R.G., Macintyre, I. G., Stuckenrath, R., 1978.
- Submerged early Holocene barrier reef, south-east Florida
- 1317 shelf. Nature 276, 59-60.
- 1318 Lincoln, J.M., Schlanger, S.O., 1991. Atoll stratigraphy as
- record of sea level changes: problems and prospects. J.
- 1320 Geophys. Res. 96, 6727–6752.
- 1321 Linklater, M., Brooke, B. P., Hamylton, S. M., Nichol, S. L.,
- Woodroffe, C. D., 2015. Submerged fossil reefs
- discovered beyond the limit of modern reef growth in the
- Pacific Ocean. Geomorphology 246, 579–588.
- 1325 Lundberg, J., Ford, D.C., 1994. Late Pleistocene sea level
- change in the Bahamas from mass spectrometric U-series
- dating of submerged speleothem. Quat. Sci. Rev. 13, 1-
- 1328 14.
- 1329 Mahoney, J.J., 1987. An Isotopic Survey of Pacific Oceanic
- 1330 Plateaus: Implications for their Nature and Origin. In:
- Keating, B.H., Fryer, P., Batiza, R., Boehlert, G.W. (eds)
- 1332 Seamounts, Islands and Atolls. American Geophys. Union,

- Washington, D.C., Geophys. Monogr. Ser. 43, pp. 207-
- 1334 220.
- 1335 Maragos, J.E., 1994. Description of reefs and corals for the
- 1336 1988 protected area survey of the northern Marshall
- 1337 Islands. Atoll Res. Bull. 419, 88 pp.
- 1338 Marshall, J.F., Jacobsen, G., 1985. Holocene growth of a mid-
- Pacific atoll: Tarawa, Kiribati. Coral Reefs 4, 11–14.
- 1340 McNutt, M., Fischer, K.M., 1987. The South Pacific Superswell.
- In: Keating, B.H., Fryer, P., Batiza, R., Boehlert, G.W.
- 1342 (eds). Seamounts, Islands and Atolls. America Geophys.
- Union, Washington, D.C., Geophys. Monograph Ser. 43,
- pp. 25–34.
- 1345 McNutt, M., 1998. Superswells, Rev. Geophys. 36, 211–244.
- 1346 Menard, H.W., 1969. Elevation and subsidence of oceanic crust.
- 1347 Earth Planet. Sci. Lett. 6, 275–284.
- 1348 Montaggioni, L.F., 1985. Makatea Island. Proc. Fifth Int. Coral
- 1349 Reef Congress, Tahiti, 1, 103–158.
- 1350 Montaggioni, L.F., 2005. History of Indo-Pacific coral reef
- systems since the last deglaciation: development patterns
- and controlling factors. Earth Sci. Rev. 71, 1–75.
- 1353 Montaggioni, L.F., Gabrié, C., Naim, O., Payri, C., Richard, G., Salvat,
- B., 1987. The seaward margin of Makatea, an uplifted
- carbonate island (Tuamotu, central Pacific). Atoll. Res. Bull.
- 1356 299, 1-18.
- 1357 Montaggioni, L.F., Faure, G., 1997. Response of coral reef
- 1358 commununities to sea level rise: A Holocene model from
- 1359 Mauritius (Western Indian Ocean). Sedimentology 44, 1053-
- 1360 1070.

- 1361 Montaggioni, L.F., Cabioch, G., Camoin, G.F., Bard, E., Ribaud-
- 1362 Laurenti, A., Faure, G., Déjardin, P., Récy, J., 1997. Continuous
- record of reef growth over the past 14 ky on the mid-Pacific island
- 1364 of Tahiti. Geology 25, 555-558.
- 1365 Montaggioni L.F., Braithwaite, C.J.R., 2009. Quaternary Coral Reef
- 1366 Systems: Development processes and controlling factors.
- Developments in Marine Geology 5. Elsevier, Amsterdam, 532
- 1368 pp.
- 1369 Montaggioni, L.F., Borgomano, J., Fournier, F., Granjeon, D.
- 1370 2015. Quaternary atoll development: New insights from
- the two-dimensionam stratigraphic forward modeling of
- Mururoa Island (Central Pacific). Sedimentology 62, 466-
- 1373 500.
- 1374 Montaggioni, L.F., Salvat, B., Aubanel, A., Eisenhauer, A.,
- Martin-Garin, B., 2018. The mode and timing of windward
- reef-island accretion in relation with Holocene sea-level
- change: a case study from Takapoto Atoll, French
- Polynesia. Geomorphology 318, 320–335.
- 1379 Muhs, D.R., 2002. Evidence for the timing and duration of the
- last interglacial period from the high-precision uranium-
- series ages of corals on tectonically stable coastlines.
- 1382 Quat. Res. 58, 36–40.
- 1383 Müller, R.D., Sdrolias, M., Gaina, C., Roest, W.R., 2008. Age,
- spreading rates, and spreading assymmetry of the world's
- ocean crust. Geochem. Geophys. Geosyst. 9, Q04006,
- 1386 doi:10.1029/2007GC001743
- 1387 Murray-Wallace, C.V., Woodroffe, C.D., 2014. Quaternary sea-
- 1388 level changes. Cambrige Univ. Press, Cambridge, UK, 484
- 1389 pp.

- 1390 Neumann, A.C., Macintyre, I., 1985. Reef response of sea level
- rise: keep-up, catch-up or give-up. Proc. Fifth Int. Coral
- 1392 Reef Congr. 3, 105–110.
- Okal, E.A., Cazenave, J., 1985. A model for the plate tectonic
- evolution of the east-central Pacific based on Seasat
- investigations. Earth Planet Sc Lett 72, 99–116.
- 1396 Patriat, M., Kingelhoefer, F., Aslanian, D., Contrucci, I.,
- Gutscher, M.-A., Talandier, J., Avedik, F., Francheteau,
- J., Weigel, W., 2002. Deep crustal structure of the
- Tuamotu plateau and Tahiti (French Polynesia) based on
- seismic refraction data. Geophys. Res. Lett. 29, 388–391.
- 1401 Paulay, G., McEdward, L.R., 1990. A simulation model of reef
- island morphology: the effects of sea level fluctuations,
- growth, subsidence and erosion. Coral Reefs 9, 51–62.
- 1404 Phinney, J.T., Hoegh-Guldberg, O., Kleypas, J., Skirving, W.
- 1405 (eds), 2013. Coral Reefs and Climate Change: Science and
- 1406 Management. American Gephys. Union, Coast. Estuarines
- 1407 Stud., Wiley, New-York, 61, 244 pp.
- 1408 Pinca, S., Beger, M., Muller, B., Reeves, E., White, M., 2006.
- 1409 Community analysis of reef habitats in the Marshall
- 1410 Islands. Proc. Tenth Int. Coral Reef Symp. Okinawa,
- 1411 Japan, 300–305.
- 1412 Poppenga, S.K., Palaseanu-Lovejov, M., Gesch, D.B.
- Danielson, J.J, Tyler, D.J., 2018. Evaluating the potential
- for near-shore bathymetry on the Majuro Atoll, Republic
- of the Marshall Islands, using Landsat 8 and WorldView-3
- 1416 Imagery. US Geol. Surv. Scientific Investigations Report
- 1417 2018–5024, 14 pp.

- 1418 Poupardin, A., Henrich, Ph., Frère, A., Imbert, D., Hébert, H.,
- 1419 Flouzat, M., 2017. The 1979 submarine landsline-
- 1420 generated tsunami in Mururoa, French Polynesia. Pure
- 1421 Appl. Geophys. 174, 3293–3311.
- 1422 Puga-Bernabéu, A., Beaman, R.J., Webster, J.M., Thomas, A.L.,
- Jacobsen, G., 2017. Gloria Knolls slide: a prominent
- submarine landslide complex on the Great Barrier Reef
- margin of north-eastern Australia. Mar. Geol. 385, 68-83.
- 1426 Purdy, E.G., Winterer, E.L., 2001. Origin of atoll lagoons.
- 1427 American Geol. Soc. Bull. 113, 837–854.
- 1428 Quinn, T.M., 1991. The history of post-Miocene sea level
- change: inferences from stratigraphic modeling of
- 1430 Enewatok Atoll. J. Geophys. Res. 96, 6713–6725.
- 1431 Ristvet, B.L., Couch, Jr., R.F., Tremba, E.L., 1980. Late
- 1432 Cenozoic solution unconformities at Enewetak Atoll.
- 1433 Geol. Soc. America Bull. Abstracts Programs 12, 510.
- 1434 Riyaz, M., Park, K.H., 2008. Reef slope failure in the north-
- 1435 eastern corner of Malé, Maldives. Proc. Eleventh Coral
- Reef Symp. Fort Lauderdale, Florida, session 1, 31–33.
- 1437 Rohling, E.J., Grant, K., Bolshaw, M., Roberts, A.P., Siddall,
- 1438 M., Hembelen, Ch., Kucera, M., 2009. Antarctic
- temperature and global sea level closely coupled over the
- past five glacial cycles. Nat. Geosci. 2, 500–504.
- 1441 Rovere, A., Khanna, P., Bianchi, C.N, Droxler, A.W., 2018.
- 1442 Submerged reef terraces in the Maldivian Archipelao
- 1443 (Indian Ocean). Geomorphology 317, 218–232.

- 1444 Rufin-Soler, C., Mörner, N.-A., Laborel, J., Collina-Girard, J.,
- 1445 2013. Submarine morphology in the Maldives and
- Holocene sea-level rise. Mar. Biodivers. 45, 599–600.
- 1447 Salvat, B., 1981. Geomorphology and marine ecology of the
- 1448 Takapoto Atoll. Proc. Fourth Intern. Coral Reef Symp.,
- 1449 Univ. Philippines, Manila, 1, 503-509.
- 1450 Salvat, B., Richard, G., 1985. Takapoto Atoll, Tuamotu
- 1451 Archipelago. Proc. Intern. Coral Reef Congr. 1, pp. 323-
- 1452 378.
- 1453 Sanborn, K.L., Webster, J.M., Yokoyama, Y., Dutton, A., Braga,
- J.C., Clague, D.A., Paduan, J.B., Wagner, D., Rooney,
- J.J., Hansen, J.R., 2010. Evidence of Hawaiian coral reef
- drowning in response to meltwater pulse 1A. Quat. Sci.
- 1457 Rev. 175, 60–72.
- 1458 Schellmann, G., Radtke, U., 2004. A revised morpho- and
- chronostratigraphy of the late and middle Pleistocene
- 1460 coral reef terraces on southern Barbados (West Indies).
- 1461 Earth Sci. Rev. 64, 157–187.
- 1462 Schlanger, S. O., Garcia, M. O., Keating, B. H., Naughton, J. J.,
- Sager, W. W., Haggerty, J. A., Philpotts, J. A., Duncan,
- R. A., 1984. Geology and Geochronology of the Line
- 1465 Islands, J. Geophys. Res. 89, 11261–11272.
- 1466 Schlanger, S.O., Campbell, J.F. Jackson, M.W., 1987. Post-
- Eocene subsidence of the Marshall Islands recorded by
- drowned atolls on Harrie and Sylvania guyots. In:
- 1469 Keating, B.H., Fryer, P., Batiza, R., Boehlert, G.W. (eds)
- 1470 Seamounts, Islands and Atolls Geophys. Monogr.
- 1471 American Geophys. Union 43, 165–174.

- 1472 Seard, C., Borgomano, J., Granjeon, D., Camoin, G.F., 2013.
- 1473 Impact of environmental parameters on coral reef
- development and drowning: forward modelling of the last
- deglacial reefs from Tahiti (French Polynesia; IODP
- 1476 Expedition # 310. Sedimentology 60, 1357–1388.
- 1477 Siddall, M., Rohling, E.J., Almogi-Labin, A., Hemleben, Ch.,
- 1478 Meischner, D., Schmelzer, I., Smeed, D.A., 2003. Sea-
- 1479 level fluctuations during the last glacial cycle. Nature
- 1480 423, 853–858.
- 1481 Siddall, M., Chappell, J., Potter, E.K., 2006. Eustatic sea level
- during past interglacials. In: Sirocko, F., Litt, T.,
- 1483 Claussen, M., Sanchez-Goni, M.F. (eds) The climate of
- past interglacials, Elsevier, Amsterdam, p. 75-92.
- 1485 Siddall, M., Chappell, J., Potter, E.K., 2007. Eustatic sea-level
- during past interglacials. Dev. Quat. Sci. 7, 75-92.
- 1487 Sladen, A., Hébert, H., Schindelé, F., Reymond, D., 2007.
- 1488 Evaluation of far-field tsunami hazard in French Polynesia
- 1489 based on historical and numerical simulations. Nat.
- 1490 Hazards Earth Syst. Sci. 7, 195–206.
- 1491 Smith, R., Young, S., Biribo, N., 1994. Bathymetric, seismic
- and alternative sand and gravel resource surveys, Tarawa
- 1493 Atoll, Kiribati. SOPAC Preliminary Report 72, 14 pp.
- 1494 Stumpf, R., Holderied K., Sinclair, M., 2003. Determination of
- water depth with high-resolution satellite imagery over
- variable bottom types. Limnol. Oceanogr. 48, 547–556.
- 1497 Szabo, B.J., Tracey, Jr., J.I., Goter, E.R., 1985. Age of
- subsurface stratigraphic intervals in the Quaternary of
- Enewetak Atoll, Marshall Islands. Quat. Res. 23, 54–61.

- 1500 Terry, J.P., Goff, J., 2012. One hundred and thirty years since
- Darwin: reshaping the theory of atoll formation. The
- 1502 Holocene 23, 615–619.
- 1503 Terry, J.P., Goff, J., 2017. Comment on 'Late Cenozoic sea
- level and the rise of modern rimmed atolls' by Toomey et
- al. (2016), Palaeogeogr. Palaeoclim. Palaeoecol. 451: 73-
- 1506 83; Palaeogeogr. Palaeoclim. Palaeoecol. 469, 156–158.
- 1507 Thompson, W.G., Goldstein, S.L., 2005. Open-system coral ages
- reveal persistent suborbital sea-level cycles. Science 308,
- 1509 401–404.
- 1510 Thompson, W.G, Spiegelman, M.W., Goldstein, S.L., Speed,
- 1511 R.P., 2003. An open-system model for U-series age
- determinations of fossil corals. Earth Planet. Sci. Lett.
- 1513 210, 365–381.
- 1514 Tracey, Jr., J.I., Ladd, H.S., 1974. Quaternary history of
- 1515 Eniwetok and Bikini atolls, Marshall Islands. Proc.
- 1516 Second Intern. Coral Reef Symp., Great Barrier Reef
- 1517 Committee, Brisbane, 2, 537–550.
- 1518 Webster, J.M., et al., 2004a. Drowning of the -150 m reef off
- Hawaii: a casualty of global meltwater pulse 1A? Geology
- **1520** 32, 249–252.
- Webster, J.M., Wallace, L., Silver, E., Applegate, B., Potts, D.,
- Riker-Coleman, K., Gallup, C., 2004b. Drowned carbonate
- platforms in the Huon Gulf, Papua New Guinea. Geochem.
- 1524 Geophys. Geosyst. 5, Q11008,
- 1525 doi:10.1029/2004GC000726.
- Webster, J.M., Clague, D.A., Braga, J.C., Spalding, H., Renema,
- W., Kelley, Ch., Applegate, B., Smith, J.R., Paull, C.K.,

- Moore, J.G., Potts, D., 2006. Drowned coralline algal
- dominated deposits off Lanai, Hawaii; carbonate accretion
- and vertical tectonics over the last 30 ka. Mar. Geol. 225,
- 1531 223.
- Webster, J.M., Braga, J.C., Clague, D.A., Gallup, C., Hein, J.R.,
- Potts, D.C., Renema, W., Riding, R., Riker-Coleman, K.,
- Silver, E., Wallace, L.M., 2009. Coral reef evolution on
- rapidly subsiding margins. Global Planet. Change 66,
- 1536 129–148.
- 1537 Webster, J.M., George, N.P.G., Beaman, R.J., Hill, J., Puga-
- 1538 Bernabéu, A., Hinestrosa, G., Abbey, E.A., Daniell, J.,
- 1539 2016. Submarine landslides on the Great Barrier Reef
- shelf edge and upper slope: a mechanism for generating
- tsunamis on the north-east Australian coast? Mar. Geol.
- **1542** 371, 120–129.
- Webster, J.M., Braga, J.C., Humblet, M., Potts, D.C. Iryu, Y.,
- Yokoyama, Y., Fujita, K., Bourillot, R., Esat, T.M.,
- 1545 Fallon S., Thompson, W.G., Thomas, A.L., Kan, H.,
- 1546 McGregor, H.V., Hinestrosa, G., Obrochta, S.P.,
- Lougheed, B.C., 2018. Response of the Great Barrier Reef
- to sea-level and environmental changes over the past
- 1549 30,000 years. Nat. Geosci. 11, 426–432.
- Wells, J.W., 1954. Recent corals of the Marshall Islands. U.S.
- 1551 Geol. Surv. Prof. Paper. 260–1: 385–486.
- Winterer, E.L., 1976. Bathymetry and and regional tectonic
- setting of the Line Islands Chain. Initial Reports, DSDP
- **1554** 33, 25, 731–747.

Woodroffe, C.D., Webster, J.M., 2014. Coral reefs and sea-level change. Mar. Geol. 352, 248-267.
Wong, F.L., Hein, J.R., Mosier, D.L., 1999. Gridded bathymetric data for the Republic of the Marshall Islands.
USGS Miscellaneous Field Stud. Map, US Geol. Surv. MF 2324.

Table captions

- 1563 **Table 1.** Geomorphic features identified along the fore-reef
- slopes of Takapoto Atoll.
- 1565 **Table 2.** Coral cover rate at different depths along fore-reef
- 1566 slopes (-5 m, -10 m, -15 m, -20 m, -25 m, and -30 m), south
- end, Fakatopatere area, Takapoto Atoll.
- **Table 3.** Depth distribution of terrace levels (TL3 to TL6)
- 1569 identified from fore-reef slopes at Takapoto Atoll, and
- interpreted in terms of paleo-sea level positions.
- 1571 The following parameters are given: time duration (ky) of MIS
- substage; duration (ky) during which terraces were exposed to
- sub-aerial erosion; time at which terraces were re-fooded by
- 1574 sea-level rise (ka); amplitude of subsidence and ground
- lowering (metres); resulting deepening (metres) of terraces;
- interpretative eustatic paleo-high sea stands.

1577 8 Figure captions


- 1578 **Fig. 1.** (a) Location map of Takapoto Atoll in the Tuamotu
- 1579 Archipelago (French Polynesia, Central Pacific). (b)
- 1580 Bathymetric and structural map of the northern area of the
- 1581 Tuamotu Plateau. Are specifically indicated the location of
- atolls (Manihi, Takaroa, Ahe, Raroia, Takume) whose major
- axis of elongation is oriented south-west-north-east, consistent
- with that of Takapoto and a number of nearby submarine


1585 volcanic ridges (SVR). Background map from Google Earth 1586 Pro 1587 **Fig. 2.** Google Earth Pro aerial maps of Takapoto Atoll. (a) 1588 Location of water-depth sounding tracks along the eastern outer 1589 margin of the atoll and landslide-related scars. (b) Location of 1590 satellite-derived bathymetric profiles (T1 to T41) around the 1591 atoll fore-reef slope, running from the modern outer reef crest 1592 to the deepest detected toe. Colors refer to profile length. 1593 Fig. 3. Virtual bathymetric profile showing the different 1594 geomorphic features identified along the fore-reef slopes at 1595 Takapoto. 1596 Fig. 4. Plots of bathymetric profiles from the six fore-reef 1597 sectors delineated around the atoll, on the basis of profile 1598 length and gradient. See figure 2b for location of profiles. 1599 Fig. 5. Selected bathymetric profiles from each delineated fore-1600 reef sector showing local geomorphic attributes. See figure 2b 1601 for location of profiles. 1602 Fig. 6. 3-D reconstruction of a portion of the WSW side at 1603 Takapoto Atoll, based on satellite imagery surveys. The image 1604 portion is lying between the profiles T26 and T27 – see figure 1605 2b for location of profiles. Successively, from inland 1606 oceanwards are shown: the atoll island, the reef flat zone 1607 bounded outwards by an algal ridge, the spur-and-groove


- 1608 system (terrace TL1), two identified terrace lines, TL2 and
- 1609 TL3, respectively at depths of about -10 m and -14 m, an
- 1610 expected terrace line TL4, at about -20 m ending into a drop-
- off about 15 m high.
- 1612 Fig. 7. Depth location of terraces identified from each
- bathymetric profile along the fore-reef slopes at Takapoto. A
- total of 112 individual terraces, including 96 well-defined and
- 1615 16 expected, were counted. These are distributed among 7
- 1616 terrace levels, at recurrent depths of about -5 m, -10 m, -15 m,
- -20 m, -24 m, -30.5 m, and -38 m.
- 1618 **Fig. 8.** Google Earth Pro aerial view of a portion of the eastern
- atoll margin, showing from the rim seawards, the outer reef-flat
- zone, algal reef crest, spur-and-groove system (terrace TL1),
- gently dipping terrace TL2 abruptly ending into dark waters.
- 1622 **Fig. 9.** Relative frequency and percentage of terraces plotted by
- width range, and calculated from 112 terraces identified from
- levels TL 2 to TL7.
- The histogram is left-skewed, expressing high frequency of
- terraces less than 15 m wide.
- 1627 Fig. 10. Plot of terrace width as a function of the detected
- length of bathymetric profiles.
- Are plotted 112 terraces identified from levels TL2 to TL7.


- 1630 Fig. 11. Close-up pictures of coral communities at different
- depths from −5 m to −30 m, fore-reef slope, Fakatopatere area,
- south end of Takapoto.
- 1633 (a) –5 m depth: spur surface partly covered by robust branching
- 1634 Pocillopora (P. verrucosa).
- 1635 (b) -10 m depth: terrace colonized by coral head mostly
- 1636 composed of *Pocillopora verrucosa*, together with some
- branching, cluster-like *Pocillopora elegans*.
- 1638 (c) –15 m depth: up to 1 m, thick coral cover, mainly composed
- of plate-shaped to encrusting colonies of *Porites rus*.
- 1640 (d) –20 m depth: coral patch dominated by plate-like *Porites*
- 1641 rus.
- 1642 (e) and (f): -25 m and -30 m deep: drop-offs densely encrusted
- by plate-like *Porites rus*.
- 1644 **Fig. 12.** Wide-angle view of drop-offs between –28 m and
- 1645 –32 m, showing dense cover of *Porites rus*.
- 1646 Fig. 13. Depth ranges of terrace levels (TL1 to TL6) identified
- 1647 from fore-reef slopes at Takapoto, compared with stratigraphic
- position of surface unconformities in interpretative core
- sections from Moruroa, Enewetak, Bikini, Tarawa, and Yongle
- atolls. Absolute dates are indicated locally on the right side of
- Moruroa, Enewetak, and Tarawa cores.


- 1652 **Fig. 14.** Eustatic sea level curve for the last 350 ky (from Grant
- 1653 et al., 2014) with position of terrace levels (TL1 to TL6)
- identified from fore-reef slopes at Takapoto.
- Symbols for terrace levels: red coloured = present-day depth
- range; green coloured = former depth range after correction of
- subsidence and ground lowering.
- 1658 Correspondance between terrace levels and marine isotope
- 1659 stages: TL1 = MIS 1; TL2 = MIS 5e; TL3 = MIS 7a;
- 1660 TL4 = MIS 7c; TL5 = MIS 7e; TL6 = MIS 9e.





































Types	Definitions
Fore-reef zone	the area extending from the seaward margin of the atoll rim to the deepest detected isobath
Shelf slope	the area extending beyond the lowest limits of the spur-and-groove system (-10 m deep on average) to the deepest detected isobaths
Reef edge	the outermost, top part of a reef tract
Spur-and-groove system	the uppermost part of a fore-reef zone formed by a series of parallel, linear ridges (spurs) alternating with narrow channels (grooves), oriented perpendicular to the reef-crest line and extending down to average depths of -10 m
Terrace	a relatively flat-topped or gently dipping surface, which is bounded seawards by a steeper descending slope or step and inwards by a steeper ascending slope
Outer-reef crest like-relief	about 1 m-high, strictly located to the seaward outer margin of a reef tract or terrace
Drop-off	a relatively high (up to 10 m), vertical surface facing the open sea
Step	a low (less than 10 m), steeply inclined to vertical surface, and usually located at the seaward end of a terrace
Pinnacle	an isolated, composite, tower-shaped relief up to 2 m high, both terraces and slopes
Coral head	a relief less than 2 m high, occasionally composed of one single large colonies, but usually composite in nature, located at different places on terraces and slopes
Trench	narrow, vertical cut deeply dissecting the fore-reef zone
Scar	bight-shaped indentations along the outer reef margin



Terrace levels	Present depth (m)	Expected MIS stages	Time duration (ky)	Reflooding time (ka)	duration of emergence (ky)	Amplitude of ground lowering (m)	Amplitude of subsidence (m)	Deepening of terraces (m)	Past sea-level elevation (m)
TL3	14.7 ± 3.8	7a	−201.5 to −189.7	128	61.7	7.4	9.5	16.9	$2.0 \pm 3.8 \text{ apsl}$
TL4	20.3 ± 2.3	7c	-217.2 to -206.0	196	10.0	1.2	9.6	10.8	$9.5 \pm 2.3 \text{ bpsl}$
TL5 24.0 ± 2.0	$.0 \pm 2.0$ 7e	-248.9 to -231.0	(1) 220	11.0	1.3	11.7	13.0	$11.0 \pm 2.0 \text{ bpsl}$	
			(2) 200	31.0	3.7	11.7	15.4	$8.6 \pm 2.0 \text{ bpsl}$	
TL6	30.5 ± 3.0	9	-335.6 to -316.8	240	76.8	9.2	15.8	25.0	5.5 ± 3.0

Rate of sub-aerial erosion = 0.12 mm/y

(1) MIS 7e reef overlapped by MIS 7c reef

Rate of subsidence = 0.047 mm/y

apsl : above present sea-level bpsl : below present sea-level (2) No overlapping