



**HAL**  
open science

# The immunobiological interplay between *Pseudosuccinea columella* resistant/susceptible snails with *Fasciola hepatica*: Hemocytes in the spotlight

Annia Alba, David Duval, Jorge Sánchez, Ana B Perez, Silvain Pinaud,  
Richard Galinier, Antonio Vázquez, Benjamin Gourbal

## ► To cite this version:

Annia Alba, David Duval, Jorge Sánchez, Ana B Perez, Silvain Pinaud, et al.. The immunobiological interplay between *Pseudosuccinea columella* resistant/susceptible snails with *Fasciola hepatica*: Hemocytes in the spotlight. *Developmental and Comparative Immunology*, 2020, 102, pp.103485. 10.1016/j.dci.2019.103485 . hal-02279344

**HAL Id: hal-02279344**

**<https://hal.science/hal-02279344>**

Submitted on 20 Jan 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



14 **Abstract**

15 The *Fasciola hepatica*/*Pseudosuccinea columella* interaction in Cuba involves a unique pattern of phenotypes;  
16 while most snails are susceptible, some field populations are naturally resistant to infection and parasites are  
17 encapsulated by snail hemocytes. Thus, we investigated the hemocytes of resistant (R) and susceptible (S) *P.*  
18 *columella*, in particular morphology, abundance, proliferation and *in vitro* encapsulation activity following  
19 exposure to *F. hepatica*. Compared to susceptible *P. columella*, hemocytes from exposed resistant snails  
20 showed increased levels of spreading and aggregation (large adherent cells), proliferation of circulating blast-  
21 like cells and encapsulation activity of the hemocytes, along with a higher expression of the cytokine granulin.  
22 By contrast, there was evidence of a putative *F. hepatica*-driven inhibition of host immunity, only in  
23 susceptible snails. Additionally, (pre-)incubation of naïve hemocytes from *P. columella* (R and S) with different  
24 monosaccharides was associated with lower encapsulation activity of *F. hepatica* larvae. This suggests the  
25 involvement in this host-parasite interaction of lectins and lectins receptors (particularly related to mannose  
26 and fucose sensing) in association with hemocyte activation and/or binding to *F. hepatica*.

27 **Keywords:** host-parasite interaction; proliferation; parasite encapsulation; glycan; granulin; immune  
28 modulation

29

## 30 **1. Introduction**

31 *Fasciola hepatica* is a digenetic trematode parasite with worldwide distribution and long-established in many  
32 countries that causes an important liver disease known as fascioliasis, a severe burden particularly for bovine  
33 and ovine livestock, and an emerging disease affecting 17 million people (Mas-Coma et al., 2009). Small  
34 freshwater snails are a key factor in the parasite life cycle, several species of the gastropod family Lymnaeidae,  
35 including *Pseudosuccinea columella*, transmit this trematode (Andrews, 1999). Therefore, the study of the  
36 particular interactions established between the parasite and each host species may give insights into the ways  
37 of transmission, epidemiology and vector-control opportunities, and reveal aspects of snail internal defenses  
38 toward broader understanding of animal immunity as afforded by comparative immunology.

39 The *F. hepatica*/*P. columella* system in Cuba presents an attractive and accessible model to study  
40 immunobiology within a trematode/snail interaction because two distinct phenotypes exist in naturally  
41 occurring Cuban snails. While most *P. columella* populations are susceptible to the parasite infection and thus  
42 support transmission, some field populations are resistant to *F. hepatica* (Gutiérrez et al., 2003b). To our  
43 knowledge, this is the only lymnaeid snails/*F. hepatica* system in which persistent, naturally occurring  
44 resistance to the parasite exists within the same snail species (Gutiérrez et al., 2003b, Gutiérrez et al., 2003a),  
45 and that is effective irrespective of the geographical origin or the infective load of the parasite isolate (Calienes  
46 et al., 2004, Vázquez et al., 2014, Alba et al., 2018). Hemocytes (circulating cells specialized in immune  
47 response) of resistant snails encapsulate *F. hepatica* larvae shortly after penetration and well-formed capsules  
48 can be observed, by histological sections, as early as 24 hours (h) following exposure (Gutiérrez et al., 2003b).  
49 The observed cellular immune response is supposed to mediate parasite killing and resolution of infection, and  
50 has been associated typically with the resistant phenotype of *P. columella* (Gutiérrez et al., 2003b).

### 51 **1.1 Hemocytes: the “vedettes” of the cellular immune response in mollusks**

52 Host responses against invading pathogens are mainly linked with cellular response activation. In  
53 invertebrates, hemocytes figure among the main arsenal of the innate immune system, playing crucial roles in  
54 both effecting cellular cytotoxicity and releasing humoral defense factors (Hillyer, 2016, Pila et al., 2016b).  
55 Notably, innate immune cells are highly conserved among vertebrates and various invertebrate phyla  
56 (Buchmann, 2014, Hartenstein, 2006). Even though the designations vary among the different taxa (*e.g.*  
57 amebocytes, hemocytes, coelomocytes, granulocytes, monocytes, macrophages), these are macrophage-like  
58 cells with similar phagocyte functions (Buchmann, 2014), that are thought to have evolutionary diverged from  
59 a common type of ancestral progenitor cell-type, the hemocytoblast (Hartenstein, 2006).

60 In mollusks, as in other invertebrates, circulating hemocytes are the leading effectors of the cellular immune  
61 response, and beyond immune-surveillance, phagocytosis (Cueto et al., 2015, Ataev et al., 2016) and  
62 encapsulation (Gutiérrez et al., 2003b), their functions include cytotoxic reactions (Humphries and Yoshino,  
63 2008) and the production of a plethora of molecules involved in signal transduction, regulation of immune  
64 defenses or immune effector activities (Baeza-Garcia et al., 2010, Galinier et al., 2013, Pila et al., 2016a). To  
65 date, no consensus exists regarding classification of hemocytes lineages, morphotypes and subtypes in  
66 mollusks, due to the use of diverse criteria for differentiation and the lack of standardized biological markers,  
67 see Pila et al. (2016b). However, a general description of the hemocytes based on their spreading capabilities  
68 defines two main types: (a) cells that typically spread, forming filopodia or lamellopodia in the process,  
69 believed to be the mature effectors in phagocytosis and encapsulation processes and that constitute the  
70 majority (if not all) of hemocyte populations, and (b) cells that do not or minimally spread when placed on  
71 artificial surfaces, often called round cells or blast-like cells (Loker, 2010). The production, differentiation and  
72 activation of these immune cells are essential and regulated by various molecular signaling pathways, as yet to  
73 be characterized in detail, to respond effectively to different stimuli (Pila et al., 2016b, Zhang et al., 2016).

74 Encapsulation of an invading parasite by snail hemocytes, as observed particularly, in resistant *P. columella*-*F.*  
75 *hepatica* interaction, is an important defense response that involves the isolation of the parasite from host  
76 signals and nutrients, and generates a secure and controlled environment for the hemocytes to release  
77 cytotoxic and humoral effectors that will help in destroying the parasite (Loker et al., 1982, Coustau et al.,  
78 2015). Several studies, involving snail species from a diverse phylogenetic background, report that increased  
79 numbers of circulating hemocytes associates with lower parasite susceptibility (e.g. Gorbushin and Iakovleva,  
80 2008, Larson et al., 2014, Oliveira et al., 2010) or where parasite killing is effected by hemocyte-mediated  
81 cytotoxicity (e.g. Sullivan and Yeung, 2011, Gutiérrez et al., 2003b, Coustau et al., 2015). Therefore, the study  
82 of hemocytes is essential to fully understand the immunological interplay between snails and trematode  
83 parasites.

84 Here, we explore immune features of the hemocytes of resistant (R) and susceptible (S) *P. columella* following  
85 parasitic challenge. We compare morphology, total numbers, proliferative activity, and *in vitro* encapsulation  
86 capacity of hemocytes in each phenotype on naïve snails and after *F. hepatica* exposure. In addition, as  
87 granulins among other cytokines offer crucial assistance for hemocyte activation, proliferation and  
88 differentiation within the context of snail-trematode interaction (Pila et al. 2016a), we analyzed its  
89 relative expression in *F. hepatica*-exposed S and R-*P. columella* snails. The present study provides novel  
90 insights into the immunobiology of *P. columella* – *F. hepatica* interaction.

## 91 **2. Material and methods**

### 92 ***2.1 Laboratory-reared susceptible and resistant P. columella snails***

93 All snails used were obtained from the Laboratory of Malacology of the Institute of Tropical Medicine “Pedro  
94 Kouri”, Cuba. Susceptible (S) snail cultures were established with parental *P. columella* sampled from two field  
95 populations, S-Negrines and S-Aurora, located in Havana and Mayabeque province, respectively. The resistant  
96 (R) snail cultures (R-La Coca and R-La Palma) were established with snails originating from La Coca locality  
97 (Havana province) and, from La Palma, (Pinar del Rio province). These R- *P. columella snails* display the  
98 phenotypic marker of typical mantle pigmentation patterns, associated with resistant populations (Gutiérrez  
99 et al., 2003b, Alba et al., 2019). All snails were reared according to Sánchez et al. (1995), in Petri dishes,  
100 containing growing algae in 26°C de-chlorinated water supplemented with crushed shells as calcium carbonate  
101 source. The snails were fed on the algae *ad libitum* relocating snails to new Petri dishes with cultured algae  
102 once the food was depleted.

### 103 ***2.2 Experimental exposures of snails to F. hepatica***

104 *Fasciola hepatica*, adult flukes were obtained from a local cattle abattoir at Jaruco (Mayabeque province) and  
105 kept alive for 4h in a solution of 0.85% NaCl, 5% glucose (Sigma, USA), for egg laying. Eggs were preserved at  
106 4°C until use.

107 For complete maturation, eggs of *F. hepatica* were incubated for 15 days in distilled water, in total darkness, at  
108 28°C. At day 15, mature eggs induced to hatch by exposure to light to obtain miracidia. Experimental exposure  
109 to the parasite were carried out using five freshly hatched miracidia per snail, according to Vázquez et al.  
110 (2014).

111 Because the cellular response in the resistant host occurs shortly after parasite penetration, exposed snails  
112 were sampled at different time points (depending on the experiments, as described below) within the first 96  
113 h after parasitic exposure. Additionally, for each of the four *P. columella* strains, a group of 30 exposed snails  
114 was used to monitor the outcome of infection and checked for parasite larvae by dissection at 25 days post  
115 exposure. As expected, no infection developed in resistant snails from R-La Coca and R-La Palma, while  
116 showed prevalence of *F. hepatica* infection in S-Aurora and S-Negrines populations was 76.3% and 54.67%,  
117 respectively.

### 118 ***2.3 Hemolymph sampling***

119 Five-to-six weeks old lab-reared *P. columella* snails (shell sizes ranged from 5.1 to 6.9 cm), sexually mature  
120 adult as indicated by egg laying were used to obtain hemolymph. Snail shells were cleaned with de-chlorinated  
121 water and cotton swabs to remove any surface debris and mucus, and wiped dry. Snails were bled by cardiac  
122 puncture (Bayne et al., 1980) using sterile pipette tips to collect hemolymph samples, including hemolymph  
123 that pooled on the shell, from individual snails.

## 124 **2.4 Morphology-based analysis of resistant/susceptible *P. columella* hemocytes**

### 125 *2.4.1 Flow cytometry*

126 Hemocytes from naïve snails were analyzed in a MACS Flow Cytometer equipment (Miltenyl Biotech), using  
127 the MACS Quantity™ v. 2.5. (2003) software. Data acquisition was performed at a flow rate of 1 µL/s, with  
128 analysis of 10,000 events. Hemolymph samples were pooled using 5 µL of freshly collected hemolymph from  
129 each of four snails, see section 2.3) per strain combined with 180 µL of saline snail buffer (SSB: 5 mM HEPES,  
130 3.7 mM NaOH, 36 mM NaCl, 2 mM KCl, 2 mM MgCl<sub>2</sub>, 4 mM CaCl<sub>2</sub>; pH 7.9). Samples were placed in conical  
131 skirted polypropylene tubes and kept on ice until measurement to minimize cellular adhesion. Flow cytometry  
132 was performed using a blank sample containing only SSB and five replicate samples per strain. Cells were gated  
133 by size (forward scattering; FSC) and complexity (side scattering; SSC) and visualized on SSC (linear)/FSC  
134 (linear) density dot plots.

### 135 *2.4.2 Confocal Fluorescence microscopy*

136 Hemolymph samples collected from naive snails (0h) and at 24h (time point at which well-formed capsules of  
137 hemocytes surrounding *F. hepatica* larvae are observed) post-exposure were applied onto polystyrene slides  
138 (Evergreen) and incubated at room temperature for 45 min in a humid chamber to allow for cell  
139 sedimentation, adherence and spreading to form hemocyte layers. Hemocytes were fixed with  
140 paraformaldehyde 4% for 10 min and samples were rinsed with phosphate saline buffer (PBS: 8 g/L NaCl, 0,2  
141 g/L KCl, 1,15 g/L Na<sub>2</sub>HPO<sub>4</sub>, 0,2 g/L KH<sub>2</sub>PO<sub>4</sub>, pH 7,2) for 30 s. Each sample was incubated (30 min, room  
142 temperature) with 500 µL of Alexa Fluor® 488 Phalloidin (Invitrogen; 40X stock solution in PBS following  
143 manufacture guidelines) diluted 1/40 in PBS, for actin labelling. The excess of phalloidin was removed by  
144 rinsing with PBS (30 s) and 500 µL of 4',6-diamidino-2-phenylindole (DAPI: Invitrogen; 1 mg/mL stock solution  
145 in H<sub>2</sub>O) diluted 1/1000 in PBS was added to the slides for DNA (nucleus) staining and incubated for 2 min.  
146 Samples were washed in PBS (30 s), dried to air and mounted in fluorescence mounting medium (Dako). The  
147 stained hemocytes were inspected using a Zeiss LSM 700 confocal microscope. The associated software Zen

148 (2008) enabled the adjustment of acquisition parameters. The images were processed with ImageJ 1.41o.  
149 Based on staining profiles, the overall percentage of different morphotypes (blast cells versus large  
150 hemocytes) was estimated for naïve (0h) and at 24h post-exposure for snails of each strain.

### 151 **2.5 Hemocyte counts post-exposure of *P. columella* to *F. hepatica***

152 The hemolymph from individual snails (see section 2.3) was mixed 1:5 with SSB, loaded on a Neubauer  
153 Improved hemocytometer, leaving the hemocytes to settle for 5 min. Total number of cells was counted in five  
154 1 mm squares at 200X magnification and cell numbers were calculated as cells/ $\mu$ L of hemolymph. Cell counts  
155 were performed at 6 time points: 0 h (naïve snails), 6 h, 12 h, 24 h, 48 h and 96 h after exposure of the snails to  
156 *F. hepatica* miracidia. Samples of hemolymph of nine snails per strain were individually counted at each time  
157 point. Counts were obtained from two independent experiments.

### 158 **2.6 Assessment of in vitro hemocyte proliferation**

159 The proliferation activity of circulating hemocytes from R- and S- *P. columella* strains was determined *in vitro*  
160 by assaying for DNA synthesis with the Click-iT<sup>®</sup> EdU Imaging kit (Invitrogen) according to manufacturer's  
161 instructions. Briefly, 5  $\mu$ L of freshly collected hemolymph from naïve (0h) and 24h-exposed snails (n = 6  
162 snails/time point/strain) were applied as individual samples onto polystyrene slides (Evergreen) and diluted  
163 1:2 with SSB. The EdU solution was added to each sample (0.1 mM final concentration) and incubated for 3h at  
164 room temperature in a humid chamber. Afterwards, cells were fixed with 4% formaldehyde (100  $\mu$ L) for 15  
165 min, washed twice with 3% bovine serum albumin in PBS (PBS-BSA) and permeabilized with 100  $\mu$ L of 0.5%  
166 Triton X-100 for 20 min. After washing twice with PBS-BSA for 30 s, samples were incubated for 30 min in the  
167 dark in freshly prepared Click-iT solution. DAPI was used for counterstaining of nuclei (see section 2.4 above),  
168 and samples were mounted in fluorescence mounting medium (Dako). Hemocytes were inspected  
169 microscopically using epifluorescence and phase-contrast to characterize morphology of proliferating (EdU-  
170 marked) cells and to record the overall number of cells (DAPI-stained nucleus) and the number of proliferating  
171 cells (EdU-marked nucleus) for each R and S snail strains (n = 4) under naïve conditions (0h) and at 24h post-  
172 exposure.

### 173 **2.7 Granulin expression in *P. columella* resistant/susceptible after *F. hepatica* exposure**

174 The full transcriptome of *P. columella*, assembled de novo from 75 nt paired-end Illumina RNAseq, sequenced  
175 from triplicate samples of two (R and S) strains (Alba et al. 2019; RNAseq data is available in the Sequence Read  
176 Archive of NCBI under the following accession numbers: Submission number [SUB5947916](#); BioProject:



177 PRJNA555222: Pseudosuccinea columella RNA sequencing; BioSample: SAMN12305757: Pseudosuccinea  
178 columella RNAseq (TaxID: 31228)), was used for RT-qPCR primers design. Briefly, the sequence of the  
179 proliferation-related cytokine granulin from *Biomphalaria glabrata* snails (GenBank accession HQ661843.1) was  
180 used in blast searches to identify the granulin ortholog gene in *P. columella* transcriptome (TR27784|c1\_g1/4;  
181 transcriptome available at <http://ihpe.univ-perp.fr/acces-aux-donnees/>). The ortholog transcript sequence  
182 was verified through analysis of the primary protein sequence and functional domains with Blastx, ExPasy and  
183 CDD servers and was deposited in the GenBank (accession number MN244715). Primer3Plus software was used  
184 to design several sets of primers for RT-qPCR primers from the *P. columella* granulin sequence, further  
185 evaluating the primer design with OligoProperties Calculator (Kibbe, 2007).  
186 Initial testing led to selection of a primer set that generated a single qPCR product with an efficiency higher  
187 than 1.9 for the quantification of granulin expression (F: 5' TGTCTGCCCTTGCTTACATTCA, R: 5'  
188 GCAAGGGTATTTACCTTTGATTTC). The nuclear 18S ribosomal gene of *P. columella* (GenBank accession  
189 EU241866.1) was used as standard for relative quantification (F: 5' AGGGACGAGTGGCGTTAGC, R:  
190 5'CTTCGTGCTAGGGATTGGGG).

191 *RNA extraction*: Adult S and R *P. columella* snails were used at five time points: naive snails (0h) and 6 h, 12 h,  
192 24 h and 96 h post-exposure to *F. hepatica*. RNA samples were collected from 30 snails from each of the four  
193 (R and S) strains for every time point. After separation from the shell, snail bodies were immediately placed in  
194 individual vials containing RNA*later*<sup>®</sup> (Ambion) solution at a ratio of 1 mg of snail : 10 µL of RNA*later*. Samples  
195 were kept overnight at 4°C and then stored at -20°C until use. RNA*later* was discarded and whole snail tissues  
196 were pooled into three biological replicates, each of 10 individuals per time point, immediately frozen in liquid  
197 nitrogen and homogenized using a laboratory mixer mill MM400 (Retsch). RNA was extracted in TRIzol<sup>®</sup>  
198 Reagent (Ambion) according to manufacturer guidelines. Following treatment with Turbo DNA-free<sup>TM</sup> kit  
199 (Ambion) to remove residual genomic DNA from the samples, the quantity of RNA was estimated  
200 spectrophotometrically.

201 *RT-qPCR*: RNA (0.5 µg) of each biological replicate was reverse transcribed to cDNA using random primers and  
202 the Maxima<sup>™</sup> H Minus First Strand cDNA Synthesis kit (Thermo Fisher), simultaneously including RT negative  
203 controls (RT-) for sample that were stored at -20°C until use as (negative) controls. Then, qPCRs were  
204 performed in a LightCycler<sup>®</sup> 480 Instrument II system (Roche) using 10 µL reactions consisting of 1X of SYBR  
205 Green Master Mix Takyon<sup>®</sup> qPCR kit (Eurogentec), 0.5 µM of each primer, 2 µL of 10-fold diluted cDNA as  
206 template and molecular grade water. Three technical replicates for each biological replicate were done.  
207 Negative controls contained water or RT- products instead of cDNA template. Standard curves of cDNA from

208 naïve snails (two-fold diluted from 1/10 to 1/640) were generated to calculate the primer efficiencies. The  
209 amplification program was set as: denaturation at 95°C for 10 s followed by 35 cycles of hybridization at 60°C  
210 for 20 s and elongation at 72°C for 30 s. The relative quantity (RQ) of granulin was estimated versus the  
211 housekeeping gene (18S) using the LightCycler® 480 Instrument II software v. 1.5.0 (Roche), considering and  
212 efficiency of 2. The resulting RQ of each time point post-exposure were normalized against the RQ calculated  
213 at naïve conditions (0h) for each strain.

## 214 **2.8 Functional characterization: *in vitro* encapsulation assays of *F. hepatica* sporocyst**

215 Hemolymph from naïve (0h) and 24h-exposed R and S snails was obtained by cardiac puncture (see section  
216 2.3) to assess the propensity of snail hemocytes to encapsulate *F. hepatica* sporocysts *in vitro*. The activity of  
217 the cells toward parasitic larvae was tested under two conditions: (1) naïve (0h) hemocytes were (pre-  
218 )incubated with monosaccharides (*i.e.* the experiment combined pre-incubation of the hemocytes and  
219 encapsulation of the parasite in presence of the monosaccharides), to assess the effect of specific glycans on  
220 the encapsulation activity, and (2) hemocytes were collected from snails at 24h post-*F. hepatica* exposure to  
221 evaluate for variations in the performance of the cells after parasitic challenge. Untreated, naïve control  
222 hemocytes were used as baseline for comparing the results from the two experimental conditions. Six snails  
223 for the R and S strains were used and results were independently recorded for the different assays and *P.*  
224 *columella* strains.

225 Briefly, hemolymph samples (20 µL) from naïve and parasite-exposed (24h) snails were individually added to  
226 wells of 96-well culture plates previously filled with sterile SSB (110 µL/well). The plates were kept in ice to  
227 prevent aggregation and attachment of the hemocytes until all samples were added to the wells. Counting  
228 hemocytes and assessment of cell viability (Trypan blue exclusion) were performed in a Neubauer chamber,  
229 with aliquots taken from 2 randomly selected wells for each experimental group. Aliquot of naïve hemocytes  
230 were pre-incubated separately by addition of one of each of the following monosaccharides: D-manose, D-  
231 glucose, L-fucose and D-galactose (from Sigma), at a final concentration of 150 mM (20 µL of 1M stock) under  
232 agitation (100 rpm) for 30 min, at 27°C. To standardize the volumes for all aliquots, 20 µL SSB was added to  
233 wells containing hemocytes that were not treated with monosaccharides. Afterwards, one newly *in vitro*-  
234 transformed *F. hepatica* sporocyst obtained according to Gourbal et al. (2008) was placed in each well (in 20  
235 µL) with the help of a stereoscope, and plates were then incubated at 27°C, for 2h, in agitation (100 rpm).  
236 Then, encapsulation was evaluated for each condition. Based on pilot studies, categories had been established  
237 to score the state and thicknesses of the hemocytes capsules around the parasite larvae into five grades (from

238 1 to 5: from almost none to complete encapsulation), to express and compare observed differences in  
239 encapsulation on a semi-quantitative scale, as done elsewhere (Kryukova et al., 2014, Loker et al., 1989).

## 240 **2.9 Data analysis**

241 Means and standard deviations were plotted for each value of cell count per strain, discarding outlier values  
242 (points beyond standard deviation). Data was checked for normality (Shapiro-Wilk test) and variance  
243 homogeneity (Levene test) and Factorial ANOVAs were carried out to assess the significance between the  
244 amount of hemocytes (dependent variable) with the snail phenotype and time post-infection as factors  
245 (section 2.5). A post hoc Tukey test followed the ANOVAs was used to obtain a multiple comparison matrix for  
246 each mean. The percentage of specific morphotypes and of proliferative hemocytes per strain per time point  
247 were compared by Fisher's exact Tests (sections 2.4.2 and 2.6). A Kruskal-Wallis Test and Duncan's Tests were  
248 used to estimate statistical differences between RQs of granulins at each time point in each strain (section 2.7).  
249 The encapsulation grade showed by the hemocytes from each strain was associated to the assayed conditions  
250 (incubation with monosaccharides or parasitic challenge) by means of a multivariate correspondence analysis  
251 using the previously established qualitative scale (see section 2.8). All analyses were performed in Statistica  
252 v.12 (StatSoft. Inc., Tulsa, OK, USA 2014). Differences were considered statistically significant for values of  $P <$   
253 0.05.

## 254 **3. Results**

### 255 **3.1 Morphological observations of *P. columella* hemocytes**

256 Two well-defined morphotypes of circulating cells, differing in size/granulocytic content and size/spreading  
257 capacity, were observed in the hemolymph of both S and R strains of *P. columella* (Figure 1). Flow cytometry  
258 analysis performed on cells from naïve snails showed two main, dense clusters of *P. columella* hemocytes  
259 segregated by size and granulometry: large cells highly granular and smaller cells with lower cytoplasmic  
260 complexity (Figure 1 A, B). Consistent with the flow cytometry results, inspection of from phalloidin-DAPI  
261 stained adherent hemocyte preparations also showed two morphotypes (i) large adherent cells (7 to 15  $\mu\text{m}$ )  
262 displaying pseudopodia and (ii) roundly-shaped small cells (3 to 5  $\mu\text{m}$ ), with high nucleus/cytoplasm ratio and  
263 low cytoplasmic complexity, here called blast-like cells, in both R and S phenotypes of *P. columella* (Figure 1 C,  
264 D).

265 Extensive evaluation indicated that large adherent hemocytes are the main morphotype, and these hemocytes  
266 assume a typically spreading morphology with formation of pseudopods, similarly for samples obtained from

267 naïve snails of both R and S phenotypes (Figure 2 A-D). However, qualitative differences were observed from  
268 large hemocytes from S and R snails at 24h-post exposure to *F. hepatica* (Figure 2 E-H). Specifically, large  
269 adherent hemocytes of R snails tended to aggregate, and displayed increased length of broad-based  
270 cytoplasmic projections, extending these into a sheet-like spreading pattern (Figure 2 E, F). By contrast, large  
271 adherent cells from S snail strains appeared more scattered, without a clear tendency to aggregate and  
272 showed a discrete spreading morphology with shorter pseudopodia, as in the case of S-Aurora snails (Figure 2  
273 G, H). Intra and inter-strain variability regarding the morphology of large hemocytes morphology was more  
274 pronounced in S snails (Figure 2). The percentage of large adherent hemocytes was always higher in R snails  
275 (ranging from 89 to 97%) compared to snails from S strains snails (62 to 78%), irrespective of experimental  
276 conditions (at 0h RR vs. SS Fisher's Exact Tests  $P < 0.05$ ; at 24h R-La Palma vs. SS Fisher's Exact Tests  $P < 0.05$ ).  
277 This difference in relative percentage of this large hemocyte morphotype between resistant and susceptible  
278 snails was also indicated by dot plots of flow cytometry profiles (see Figure 1 A, B).

279 Overall, the percentages of the blast-like cell morphotype ranged from 3 to 36% of the total cell counts in  
280 naïve *P. columella*; (Figure 2A-D), with significantly higher percentages of blast-like cells in susceptible snails (S-  
281 Aurora - 26.3%, S-Negrines – 36.5%) compared to resistant strains (R-La Coca – 3.7%, R-La Palma – 10.1%;  
282 Fisher's Exact Tests  $P < 0.05$ ). At 24h post-exposure, an increase of blast-like cells was observed in resistant  
283 strains (11 and 11.2% in R-La Coca and R-La Palma, respectively) that was significant in R-La Coca (Fisher's  
284 Exact Tests  $P = 0.014$ ).

### 285 **3.2 Hemocyte counts, proliferation rate and granulin expression following exposure to *F. hepatica***

286 The cellular response of susceptible and resistant *P. columella* to *F. hepatica* displayed basically the same trend  
287 at early times post-exposure characterized by a gradual increase of hemocyte counts starting within the first  
288 hours to reach a maximum at 24h post parasite-exposure, and beginning to decrease thereafter (Figure 3A).  
289 Nevertheless, the actual number of hemocytes was significantly higher in R snails compared to S isolates, with  
290 cell counts almost two-fold higher at 24h post-exposure ( $P < 0.001$ ; Figure 3A). Relative to naïve control R  
291 snails, the number of circulating hemocytes in exposed-resistant snails was still significantly higher at 48 and  
292 96h post-exposure (Figure 3A).

293 Positive EdU staining of hemocytes, indicative of DNA synthesis, was recorded for resistant snails, 24 h post  
294 exposure (R-La Coca - 6.25%; R-La Palma - 7.14%,  $P = 0.8496$ ), suggesting the active proliferation within the  
295 circulating hemocyte population (Figure 3B). Conversely, no EdU staining was observed in hemocytes from S-  
296 Aurora snails, either naïve or at 24h post-exposure. A single individual snail from the S-Negrines strain 24h

297 post exposure) displayed EdU-signal indicative of proliferation in a modest 1.96% of hemocytes. This  
298 proportion, however, was significantly lower ( $P < 0.05$ ) than the numbers of proliferative hemocytes of R-*P.*  
299 *columella* strains. EdU marked-cells were always of the small and round-shape hemocyte morphotype,  
300 referred here as blast-like cells (Figure 3B, lower panel).

301 Lastly, a significant increase in expression of the cytokine granulin was recorded only from resistant *P.*  
302 *columella*, early after parasite exposure (6h;  $P = 0.041490$ ). By contrast, in susceptible snails, granulin  
303 expression decreased below constitutive levels at 12h and 24h post parasite exposure ( $P < 0.05$ ; Figure 3C).  
304 Comparatively, the relative expression of granulin between R and S phenotypes accounted for an almost 3-fold  
305 difference, at 24h post-exposure. Additionally, an increase of expression was recorded for both phenotypes  
306 four days after parasite exposure, however, significant differences were also observed between resistant and  
307 susceptible snails at this time point (Figure 3C).

### 308 **3.3 *In vitro* encapsulation assays**

309 The observations from pilot *in vitro* assays (Figure 4A), were used to establish a quantitative scale to compare  
310 the encapsulation grade of *F. hepatica* sporocysts by *P. columella* hemocytes under different conditions, (pre-)  
311 incubation of naïve hemocytes with monosaccharides or *F. hepatica* challenge in a multivariate  
312 correspondence analysis (Figure 4B). Encapsulation activities of naïve hemocytes from resistant strain snails  
313 were higher (strong association with Grade 4) than those from snail of the susceptible strains (association with  
314 Grade 3). In addition, encapsulation grade 5, corresponding to the observation of a thickest capsule of cells  
315 around the larvae (Figure 4A), was associated only with hemocytes from R-strain snails at 24h post exposure to  
316 *F. hepatica* (Figure 4B). Notably, while naïve hemocytes from susceptible *P. columella* strongly associated with  
317 an encapsulation grade 3, no clear pattern was observed from hemocytes at 24h post exposure with  
318 encapsulation activities ranging between grades 2 to 4 (Figure 4B).

319 On the other hand, (pre-)incubation of naïve hemocytes with monosaccharides was associated with lower  
320 encapsulation degrees (Figure 4B). Particularly, (pre-)incubation with D-mannose was associated with  
321 encapsulation grade 1 regardless of the phenotype (R or S) and of *P. columella* population. In both susceptible  
322 populations, (pre-)incubation with L-fucose was also associated with the lowest encapsulation grades (Figure  
323 4B).

## 324 **4. Discussion**

### 325 **4.1 Hemocytes from naïve *P. columella* snails: morphological and functional observations**

326 In the present study, the analysis of differences in spreading properties, size and cytoplasmic complexity (see  
327 Loker (2010) of *P. columella* hemocytes revealed two main morphotypes: i) non-adherent small and roundly-  
328 shaped hemocytes with low cytoplasmic complexity, characteristic features described for blast-like cells and ii)  
329 large adherent cells with higher cytoplasmic complexity. The latter morphotype may potentially be further  
330 segregated into granulocytes and hyalinocytes after thorough analyses of cell granularity, similar to hemocytes  
331 in *Biomphalaria* spp. and *Lymnaea palustris* (Russo and Lagadic, 2000, Cavalcanti et al., 2012), but such  
332 classification requires additional staining and functional approaches.

333 In any case, distinct cell morphologies are proposed to reflect certain functional differences among snail  
334 hemocytes, representing either different subtypes of cells or different developmental stages of a single lineage  
335 of hemocytes (Boisseaux et al., 2016, Matricon-Gondran and Letocart, 1999, Mohandas et al. 1992). In this  
336 sense, high numbers of large, adherent (spreading) hemocytes in the hemolymph of *B. glabrata* snails were  
337 suggested to be a predictive parameter of lower susceptibility to *Schistosoma mansoni* (Larson et al., 2014). In  
338 the present study, large adherent cells were highly predominant in snail from both R- versus S- *P. columella*  
339 strains. Additionally, naïve hemocytes from R snails were observed to have higher encapsulation activity of *F.*  
340 *hepatica* sporocysts compared to both S strains. Both results suggest that R- *P. columella* snails have a higher  
341 constitutive baseline of circulating hemocytes more competent of effective immunity against this parasite than  
342 those from S-*P. columella* snails (Figure 5). A likely molecular basis for this enhanced immune competence is  
343 provided by a previous comparative transcriptomic study that disclosed, compared to S-strain snails, a  
344 significantly greater resource allocation in naïve *P. columella* snails from resistant strains to express an overall  
345 greater abundance of transcripts encoding for several pathogen-interacting molecules, pro-inflammatory  
346 signaling pathways and cytokines (for details, see Alba et al., 2019).

347 Interestingly, in both *P. columella* phenotypes, the involvement of molecules with different carbohydrate  
348 specificities (lectins) appears to be fundamental for hemocyte functionality; monosaccharides generally led to  
349 lower encapsulation grades (see Figure 4B). Similarly, adhesion of cells from the *B. glabrata* embryonic (Bge)  
350 cell line to *S. mansoni* sporocyst was inhibited by carbohydrate compounds, including mannose-6-phosphate,  
351 heparin, dextran sulfate, and mainly, fucoidan (Castillo and Yoshino, 2002). The significantly diminished  
352 encapsulation activity of *P. columella* hemocytes due to incubation with mannose or fucose (Figure 4B) may be  
353 explained with two non-exclusive hypotheses: i) monosaccharides saturate glycan-binding molecules that  
354 function as hemocyte receptors for sensing exogenous entities, thereby involved in non-self recognition  
355 (binding and subsequent immune activation) of the complex carbohydrate surface motifs of a parasite like *F.*

356 *hepatica*, or ii) monosaccharide binding alters hemocyte activity through over-activation of particular glycan-  
357 binding molecules that are involved in the normal regulation of hemocyte activity.

358 Lectins, and their activation by specific carbohydrate ligands, have fundamental roles in host-parasite  
359 interaction, either associated with parasite binding/recognition (Adema et al., 1997, Yoshino et al., 2001, Wu  
360 et al., 2017) or as target for parasite-driven immune modulation/suppression (Guasconi et al., 2011, Rodríguez  
361 et al., 2015, Rodríguez et al., 2017). In *B. glabrata*, stimulation with mannose, fucose and galactose promotes  
362 the generation of reactive oxygen species by hemocytes (Hahn et al., 2000), fucosylated glycans are crucial for  
363 interaction of *S. mansoni* sporocyst with *Bge* cells (Castillo et al., 2007), and fucoidan promotes division and  
364 proliferation of hemocyte precursor cells (Sullivan et al., 2014). By contrast, exposure of *B. glabrata* to  
365 carbohydrate PAMPS (pathogen-associated molecular patterns), like immunogenic fucoidan, led  
366 downregulation of expression in the amebocyte producing organ of many genes, including several that are  
367 directly involved in immune response (Zhang et al., 2016). In addition, fucosylated-motifs are proposed to  
368 participate along with other carbohydrates in molecular (glycan) mimicry employed by *S. mansoni* parasites to  
369 evade the immunity of *B. glabrata* (Lehr et al., 2008, Yoshino et al., 2013). It is of note that evasion by *F.*  
370 *hepatica* of immune responses from the mammalian definitive hosts is thought to involve carbohydrates like  
371 mannose, fucose and galactose (Rodríguez et al., 2017, Rodríguez et al., 2015, Guasconi et al., 2011).

372 Immune-recognition of different groups of digenetic trematodes appears to require detection of unique  
373 carbohydrates. In contrast to the highly fucosylated tegumental antigens of *S. mansoni* sporocysts, (Johnston  
374 and Yoshino, 1996, Peterson et al., 2009), intramolluscan *F. hepatica* sporocysts display no fucose motifs, but  
375 mainly N-acetyl-D-glucosamine and  $\alpha$ -mannose, with minor proportions of galactose and N-acetyl-D-  
376 galactosamine at the surface (Georgieva et al. 2016). Although the parasite larvae may present other  
377 fucosylated glycans without a terminal fucose--galactose-motif, mannose glycans, because of significant  
378 presence at the parasite surface may be particularly important for non-self recognition/binding of *F. hepatica*  
379 sporocyst by snail hemocytes. The strong inhibitory impact of mannose on encapsulation of *F. hepatica*  
380 sporocysts by snail hemocytes from both R- and S- strains also points at a central role of mannose and  
381 mannose-receptors in the *P. columella-F. hepatica* interaction.

382 It is of note that hemocyte (pre-)incubation with monosaccharides (galactose, glucose and L-fucose in  
383 particular) impacted individual snails of both R- and S-strains differently, producing different intrapopulation  
384 patterns with regards to encapsulation grade; Figure 4b. This observation points at the occurrence of  
385 qualitative/quantitative differences concerning lectin receptors for specific carbohydrates on the surface of  
386 naïve hemocytes among *P. columella* strains and phenotypes. Similar differences in hemocyte surface

387 determinants have been reported for susceptible and resistant *B. glabrata* snails, suggesting that these may  
388 contribute to determine the ultimate phenotypic response after parasite infection (Agner and Granath, 1995,  
389 Yoshino et al., 2013). This notion is supported by the demonstration of significant differences in abundance of  
390 several immune-factors involved in antigen recognition and pro-inflammatory signaling pathways (C-type  
391 lectins, CD109, and mannose-binding protein C, protein kinase C) between R and S phenotypes of *P. columella*  
392 prior exposure to *F. hepatica*; Alba et al., 2019). Adding this result to the previous line of thought, an hemocyte  
393 surface enriched, for example, in lectins potentially involved in parasite recognition (*e.g.* through mannose  
394 interaction) and in the further activation of a protective response, as it appears to be the case of resistant  
395 snails, could increase the chances for cell-parasite interaction to occur. The latter was demonstrated here as  
396 naïve hemocytes from both resistant populations showed higher encapsulation grades than those from  
397 susceptible snails (Figure 4B). In contrast, enriched hemocytes on receptors with regulatory activities are more  
398 likely to be driven towards a less responsive phenotype upon stimulation by certain parasite antigens involved  
399 in immune modulatory mechanism of host defenses.

## 400 **4.2 Differential immune-related patterns behind *P. columella* phenotypes during *F. hepatica*** 401 **challenge**

### 402 *4.2.1 Resistant snails: higher cellular immune competence following exposure*

403 Considering the major role of cellular immunity, at least in part, in resolving *F. hepatica* infections in resistant  
404 *P. columella* (Gutiérrez et al., 2003b), the significantly higher hemocyte numbers, particularly of large  
405 spreading hemocytes, (compared to susceptible snails) could facilitate the orchestration and maintenance of  
406 an effective immunoprotective response against the parasite. Indeed, low numbers of hemocytes may  
407 associate with increased susceptibility to infection in the *B. glabrata* – *S. mansoni* model (Oliveira et al., 2010,  
408 Larson et al., 2014), and immune stimulation leads to increased hemocyte counts in parasite-resistant *B.*  
409 *glabrata* (Noda, 1992, Sullivan et al., 2004). Significantly, hemocyte counts reached a significantly maximum  
410 increase in resistant *P. columella* at 24h-post parasite challenge which is also the same time point at which  
411 hemocyte encapsulation of *F. hepatica* sporocyst is complete in resistant snails (see Gutiérrez et al., 2003b).

412 Although hematopoiesis in snails is generally believed to occur in a primary hematopoietic site, like the APO of  
413 planorbid snails (Pila et al., 2016b), concordant with previous reports of proliferation of circulating hemocytes  
414 from lymnaeid snails (Dikkeboom et al., 1988), proliferative blast-like hemocytes were recorded from the  
415 hemolymph of *P. columella* snails from R-strains in response to infection. These mitotic events in the immune  
416 cells of resistant snails represent a further differential response versus the susceptible phenotype. This



417 increase in hemocyte proliferation may directly contribute to the overall significant increase of cell counts in  
418 resistant *P. columella* during infection. While the functional role of blast-like cells remains unclear, perhaps  
419 these results indicate a potential stem-cell capacity in *P. columella*, considering that blast-like cells in molluscs  
420 from other taxa (caenogastropoda and bivalves) were proposed to be an immature stem cell-like  
421 developmental stage (Gorbushin and Iakovleva, 2008, Wang et al., 2012). Alternatively, blast-like cells have  
422 been also associated with crucial roles in pathogen interaction during infection through the production of  
423 molecular signals (Cima et al., 2000, Travers et al., 2008, Portet et al., 2018).

424 However, an increase in the number of circulating hemocytes and/or proliferative rate does not always  
425 provide increased resistance, (for example consider parasite-snail associations of *Himasthla elongate* -  
426 *Littorina littorea* (Gorbushin and Iakovleva, 2008) and *F. hepatica* – susceptible *P. columella* (Ribeiro et al.,  
427 2017). For effective encapsulation to occur, hemocytes must recognize and adhere to the surface of a non-self  
428 target and to each other, and form layers of overlapping cells to surround and isolate the parasite (Loker et al.,  
429 1982). At 24h-post exposure, the large adherent hemocytes from R-strain *P. columella* snails showed a strong  
430 *in vitro* tendency to aggregate into sheet-like cell layers of spreading, cohesive hemocytes that are  
431 interconnected by an abundance of numerous and long, broad pseudopods (see Figure 2), likely behaviors that  
432 contributes to effective cellular immune response against *F. hepatica*. Remarkably, the highest encapsulation  
433 grade was strongly associated with hemocytes from R snails at 24h post-exposure suggesting that immune  
434 activation in response to *F. hepatica* further enhances the considerable baseline immune effector capabilities  
435 of hemocytes from R strains of *P. columella* (see Figure 4B).

436 Lastly, resistant *P. columella* significantly increased the expression of granulin, a proliferative cytokine,  
437 following exposure to *F. hepatica*. This evolutionary-conserved cytokine regulates hematopoiesis and  
438 inflammation (Jian et al., 2013), *B. glabrata* granulin not only promotes hemocytes proliferation but also drives  
439 differentiation of these cells towards a more adherent and highly phagocytic phenotype, more prompt to  
440 cellular response and associated with a lowered susceptibility to *S. mansoni* infection (Pila et al., 2016a). Thus,  
441 as result of overall higher expression during 24h after parasite challenge observed here in R-snails (Figure 3C),  
442 granulin may function in proliferation and regulation of the phenotypical and functional changes of hemocytes  
443 (Figures 2-4) also observed in R-*P. columella* - *F. hepatica* interaction. Of course, functional validation through  
444 gene silencing/editing is needed to demonstrate the role of granulin in the activation of *P. columella*  
445 hemocytes following *F. hepatica* exposure. In addition, multiple determinants and molecular signals from both  
446 host and parasite probably contribute to the different profiles recorded from parasite-exposed *P. columella*  
447 and these remain to be elucidated also.

#### 448 4.2.2 Susceptible snails: evidence of parasite-driven immune modulation

449 Study of hemocytes from susceptible *P. columella* provided insights that contrast strongly with the indications  
450 of potent cellular immunity from resistant snails. Following infection with *F. hepatica*, hemocyte layers from  
451 exposed-susceptible snails showed that large hemocytes remained mostly scattered, displaying slender or  
452 short pseudopodia compared to naïve cells. In addition, no significant proliferative activity was recorded in  
453 blast-like cells (or any other hemocytes). Compared to naïve snails, there was a significant decrease in  
454 expression levels of the cytokine granulin at 12 and 24h post exposure, and this might relate to the differential  
455 morphological and functional patterns in exposed-hemocytes from susceptible *P. columella*. Possibly, the  
456 depressed immune features evident from S-strains after infection is relate to modulation of snail host  
457 immunity by the parasite, through as yet to be characterized signals (molecules) that only (or mostly) interact  
458 with molecular targets of the susceptible phenotype snails. Trematode countermeasures against host defenses  
459 responses include direct immune modulation (interference), and this has been reported for several parasite-  
460 host models, including *S. mansoni*/*E. paraensei* – *B. glabrata* (e.g. DeGaffé and Loker, 1998, Loker et al., 1992,  
461 Zahoor et al., 2008), *H. elongata* – *L. littorea* (Iakovleva et al., 2006), *Trichobilharzia regenti* – *Radix lagotis*  
462 (Skála et al., 2014) and *Plagiorchis mutationis* – *Lymnaea stagnalis* (Kryukova et al., 2014). In case of *F.*  
463 *hepatica*, parasite antigens are known to mediated potent immune suppression/modulation in the definitive  
464 host (e.g. Prowse et al., 2002, Hamilton et al., 2009, Dowling et al., 2010), and the results of this study indicate  
465 that *F. hepatica* also interferes with immune function in the susceptible snail intermediate host.

466 It is also significant that no clear pattern was evident regarding encapsulation activity of hemocytes from  
467 parasite-exposed snails of the susceptible *P. columella* strains, particularly S-Aurora, with encapsulation grades  
468 ranging between 2 to 4. This differential impact of parasite infection upon hemocyte immune function may be  
469 explained by different degrees of compatibility, sensitivity to parasite-mediated interference among snails of  
470 the same susceptible population. In this sense, while both the S-Aurora and S-Negrines populations of *P.*  
471 *columella* are susceptible, infection does not develop following exposure to *F. hepatica*, in 20% and 45% of  
472 snails, respectively. Genotype-genotype interactions may drive different infection outcomes for single,  
473 individual level host-parasite interactions. Such a compatibility polymorphism has been applied to the *B.*  
474 *glabrata* – *S. mansoni* model (see Theron and Coustau, 2005) and is proposed to also occur in susceptible *P.*  
475 *columella* – *F. hepatica* system (Vázquez et al., 2014, Alba et al., 2018).

#### 476 4.3 A model for immunobiology of resistant *P. columella* – *F. hepatica* interactions

477 About the genetic bases of the interplay: Remarkably, the resistant *P. columella* field populations always defeat  
478 *F. hepatica* infection, regardless of the parasitic load (Alba et al. 2018) or the geographic origin of the parasitic  
479 strain used in the challenge, either Cuban (e.g. Calienes et al., 2004, Vázquez et al., 2014) or more  
480 geographically-distant (see Alba et al. 2018). This suggest that resistance in *P. columella* is not tightly restricted  
481 to a host genotype – parasite genotype interaction, especially because *F. hepatica* is recognized as a  
482 genetically diverse parasite (Cwiklinsky et al., 2015, Vázquez et al., 2016). The hypothesis postulated here is  
483 that the outcome (infection versus non-infection) after *F. hepatica* penetration is determined by  
484 absence/scarcity or presence/abundance of a major contributing factor, possibly associated in *P. columella*  
485 with a single locus or few low-diversified loci. This major host factor would be involved proximately or  
486 ultimately in the immunobiological interplay, interacting/interfering with factor/function(s) determined by a  
487 single or few highly conserved loci of the parasite, that are fundamentally crucial for establishment within  
488 snails, that accordingly are displayed by most, if not all, *F. hepatica* strains. While all snails of resistant  
489 populations likely are selected for genotypes that possess the above-mentioned major contributing factor, it  
490 appears that lack of this trait in other genetic lineages determines susceptibility in *P. columella*. Recent studies  
491 have demonstrated the influence of allelic variations in a single 1 Mb sized-genomic region containing 15  
492 protein coding loci of *B. glabrata* (i.e. Guadalupe resistance complex) susceptibility to schistosome infection  
493 (Allan et al., 2017) and its effect on other biological roles like microbiome diversity (Allan et al., 2018b), and  
494 humoral (Allan et al., 2019) and cellular (Allan and Blouin, 2018a) immune properties of the hemolymph.  
495 However, the outcome and intensity of infection in *S-P. columella* snails is ultimately dictated by yet other  
496 molecular determinants, possibly coded by different loci with several allelic variants as to explain the proposed  
497 polymorphism of compatibility. This is a plausible theory because several steps ranging from snail host finding,  
498 parasite establishment and development, and molecular determinants (including immune factors) determine  
499 compatibility in any host-parasite interaction (Chen, 1968, Combes, 2001, Tennessen et al., 2015, Portet et al.,  
500 2017).

501 About the evolutionary and mechanistic bases of resistance: The limited information concerning the molecular  
502 and physiological dialogue between *F. hepatica* and lymnaeid host snails challenges detailed speculations  
503 about particular molecules and mechanisms coevolving and codetermining the outcome of infection.  
504 However, *F. hepatica* is presumed to mount efforts to effect parasite-driven immune modulation/suppression  
505 in the intermediate hosts, that will particularly affect susceptible but not resistant *P. columella* strains (see  
506 section 4.2.2 for discussion). One of the major differences between both phenotypes at the molecular and  
507 cellular level is the existence of enhanced immune responsiveness prior infection in resistant snails which  
508 could have been influenced from host-parasite coevolutionary dynamics. Recently, it has been proposed that

509 coevolution of parasites with specific immune components (receptors for detection/interaction or effectors for  
510 elimination) may alter energy allocation in the host to optimize defense against infection. Especially, if co-  
511 evolutionary feedback between parasite and host populations renders induced immune responses less  
512 efficient due to parasite counter-evolution, then the host will be driven to heavily invest in constitutive  
513 immunity (Kamiya et al., 2016). In this sense, *F. hepatica* driven-immune modulation as a mean for parasite  
514 establishment would certainly pose a difficult challenge to overcome for a snail host if it relies mainly on  
515 induced immunity. Thus, an emphasis toward enhanced constitutive immune responsiveness could enable the  
516 host to counter immediately, preventing the parasite to settle and activate an arsenal of immune modulatory  
517 mechanisms. Recalling what was previously discussed, *F. hepatica* immunomodulatory/immunosuppressive  
518 strategies may be the above-mentioned major highly conserved function fundamentally crucial for  
519 establishment within snails, whereas the major factor determining resistance in *P. columella* circumvents  
520 parasite's major infective function possibly by proximately or ultimately participating in the constitutively  
521 increased resource allocation to immune defense (e.g. higher expression levels of pathogen interacting  
522 molecules, pro-inflammatory signaling pathways, cytokines, antimicrobial and anti-oxidant molecules; Alba et  
523 al., 2019). Apart of the nature of the antigen, the lineage/state of the host cells and the environment in which  
524 the antigen is 'seen' by the immune system are crucial to decide the ultimate response thus, a differential  
525 molecular and cellular baseline between *P. columella* phenotypes may contribute from the start to the  
526 "insensibility" of resistant snails towards parasite immune modulation mechanisms. Similarly, constitutive  
527 differences between *B. glabrata* strains, susceptible and experimentally-selected for resistance to infection  
528 with *Echinostoma caproni*, were proposed as the base for refractoriness by preventing immunosuppression  
529 efforts of the parasite (Humbert and Coustau, 2001). For instance, prior infection, enriched hemocytes in  
530 surface determinants directly committed to non-self recognition and/or pro-activation signaling upon  
531 stimulation in resistant snails may render them more prone to trigger protective anti-parasitic responses than  
532 to be driven into a refractory or suppressed immunophenotype, as appears to be the case in *S-P. columella*  
533 strains. As it was previously suggested, differences in hemocyte surface determinants are likely to occur  
534 between resistant and susceptible snails and the higher propensity by naïve hemocytes to encapsulate *F.*  
535 *hepatica* sporocysts (see Figure 4B) constitutes an experimental evidence of an enhanced immune  
536 responsiveness prior infection in the resistant *P. columella* phenotype.

537 Finally, the indications for a highly efficient amplification of immune activities following parasite exposure in R-  
538 compared to *S-P. columella* needs to be interpreted as the conjunction of both factors: the enhanced baseline  
539 immune capabilities that contribute to recognize non-self and to trigger a potent protective response, and the

540 hijacked of *F. hepatica* immunosuppressive strategies that influence parasite's infective capacity to  
541 migrate/colonize or to develop inside the host.

542 Figure 5 presents a model that integrates the current knowledge regarding molecular and functional  
543 determinants that may contribute the immunobiological bases for resistance in this interesting host-parasite  
544 interplay between *F. hepatica* in *P. columella* snails. Future studies aiming at comparing the transcriptomic  
545 dynamics during *F. hepatica* – *P. columella* resistant/susceptible interactions will help focus further  
546 investigations and bring forward new elements to confirm and/or expand this model.

## 547 **Conclusions**

548 Comparative approaches were used to explore the immunobiology behind interactions of predictably resistant  
549 and susceptible phenotypes of *P. columella* with *F. hepatica*. After (pre-)incubation with different  
550 monosaccharides, decreased encapsulation activities of *F. hepatica* sporocysts by naïve hemocytes, were  
551 observed, irrespective of the *P. columella* phenotype. This suggests the involvement of carbohydrate-binding  
552 receptor molecules (lectins) in regulation of functional behavior of these cells towards the parasite. Future  
553 investigations will be aimed at in-depth exploration of the lectin – ligand interactome that governs the  
554 hemocyte response to the parasite. Whereas naïve hemocytes from resistant *P. columella* strains showed  
555 higher encapsulation grades compared to susceptible snails, parasitic challenge also brought about significant  
556 phenotypical and functional differences at the hemocyte level between R and S snail phenotypes. These  
557 changes may further enhance the differential resistant and susceptible phenotypes of *P. columella* in relation  
558 to *F. hepatica* infection. Specifically, parasite infection resulted in immune modulation, rendering anti-parasite  
559 immunity in susceptible snails even less effective while *F. hepatica* exposure boosted hemocyte activities in  
560 resistant *P. columella*, possibly augmenting the rapid immune-elimination of parasites that associates with the  
561 resistant phenotype of these snail strains. The augmented response was characterized (but may not be limited  
562 to) a significant increase of overall hemocyte counts, enhanced proliferation of blast-like cells, morphological  
563 changes of large adherent cells, a greater *in vitro* encapsulation activity from hemocytes (the latter also  
564 observed *in vivo*; Gutiérrez et al., 2003b), and a significantly increased transcription of granulin, a cytokine  
565 involved in proliferative, activation and differentiation of hemocytes.

566 In summary, snails from resistant strains of *P. columella* have an inherent ability to initiate and amplify a  
567 powerful and protective response following *F. hepatica* exposure. This ability is likely due to an enhanced  
568 allocation of resources to defense in resistant *P. columella* at baseline level and after parasitic exposure.

569 **Acknowledgments**

570 This study is set within the framework of the "Laboratoires d'Excellences (LABEX)" TULIP (ANR-10-LABX-41).  
571 Authors acknowledged Dr. Anaïs Portet for the insights provided on the proliferation assay. We would also like  
572 to thank two anonymous reviewers for their time and insightful observations. A special acknowledge to Dr.  
573 Coen Adema for constructive inputs and invaluable help during editorial processes.

574 **Funding sources**

575 Partial financial support for this investigation was provided by the subventions granted to AA by the French  
576 Embassy in Cuba. BG was supported by ANR JCJC INVIMORY (number ANR 13-JSV7-0009) from the French  
577 National Research Agency (ANR).

578 **Author's contributions:**

579 AA, DD, JS, ABP, AAV, BG designed and performed the experiments and analysis, and participated in the  
580 reviewing process. AA, AAV, JS participated in snail breeding, parasite exposures, hemocyte quantification and  
581 *in vitro* assays (experiment and analysis). AA, ABP, DD: conducted the cytometer experiments, cell staining and  
582 morphological descriptions. DD, RG, SP participated in qRT-PCR experiments. AA, DD, BG drafted the  
583 manuscript. All authors read and approved the final version of the manuscript. All data generated or analyzed  
584 during this study are included in this article.

585 **Declarations of interest:**

586 Author's declare that there no conflict of interest exists

587 **References**

588 Adema, C.M., Hertel, L.A., Miller, R.D., Loker, E.S. 1997. A family of fibrinogen-related proteins that  
589 precipitates parasite-derived molecules is produced by an invertebrate after infection. Proc Natl Acad Sci USA.  
590 94, 8691-8696.

591 Agner, A.E., Granath, W.O.J. 1995. Hemocytes of schistosome-resistant and -susceptible *Biomphalaria glabrata*  
592 recognize different antigens on the surface of *Schistosoma mansoni* sporocysts. J Parasitol. 81, 179-186.

593 Alba, A., Tetrau, G., Chaparro, C., Sánchez, J., Vázquez, A.A., Gourbal, B. 2019. Natural resistance to *Fasciola*  
594 *hepatica* (Trematoda) in *Pseudosuccinea columella* snails: a review from literature and insights from  
595 comparative “omic” analyses. Dev Comp Immunol. doi: 10.1016/j.dci.2019.103463.

596 Alba, A., Vázquez, A.A., Sánchez, J., Duval, D., Hernández, H., Sabourin, E., Vittecoq, M., Hurtrez-Boussés, S.,  
597 Gourbal, B. 2018. *Fasciola hepatica* - *Pseudosuccinea columella* interaction: effect of increasing parasite doses,  
598 successive exposures and geographic origin on the infection outcome of naturally-resistant and susceptible  
599 snails from Cuba. Parasite Vect. 11, 559.

600 Allan, E.R.O., Blouin, M.S. 2018a. Allelic variation partially regulates galactose-dependent hydrogen peroxide  
601 release from circulating hemocytes of the snail *Biomphalaria glabrata*. Fish Shellfish Immunol. 72, 111–116.

602 Allan, E.R.O., Tennessen, J.A., Bollmann, S.R., Hanington, P.C., Bayne, C.J., Blouin, M.S. 2017b. Schistosome  
603 infectivity in the snail, *Biomphalaria glabrata*, is partially dependent on the expression of Grctm6, a  
604 Guadeloupe Resistance Complex protein, PLoS Neglected Trop Dis. 11, e0005362.

605 Allan, E.R.O., Tennessen, J.A., Sharpton, T.J., Blouin, M.S. 2018b. Allelic variation in a single genomic region  
606 alters the microbiome of the snail *Biomphalaria glabrata*. J Hered. 109, 604–609.

607 Allan, E.R.O., Yang, L., Tennessen, J.A., Blouin, M.S. 2019. Allelic variation in a single genomic region alters the  
608 hemolymph proteome in the snail *Biomphalaria glabrata*. Fish Shellfish Immunol. 88: 301-307.

609 Andrews, S.J. 1999. The life cycle of *Fasciola hepatica*. In: DALTON, J.P. (ed.) Fasciolosis. UK: CAB International.

610 Ataev, G.L., Prokhorova E.E., Kudryavtsev, I.V., Polevshchikov, A.V. 2016. The influence of trematode infection  
611 on the hemocyte composition in *Planorbarius corneus* (Gastropoda, Pulmonata). Invertebr Survival J. 13, 164–  
612 171.

613 Baeza-Garcia, A., Pierce, R.J., Gourbal, B., Werkmeister, E., Colinet, D., Reichhart, J.M., Dissous, C., Coustau, C.  
614 2010. Involvement of the cytokine MIF in the snail host immune response to the parasite *Schistosoma*  
615 *mansoni*. PLoS Pathog. 6, e1001115.

616 Bayne, C.J., Buckley, P.M., Dewan, P.C. 1980. Macrophage-like hemocytes of resistant *Biomphalaria glabrata*  
617 are cytotoxic for sporocysts of *Schistosoma mansoni* in vitro. J Parasitol. 66, 413–419.

618 Boisseaux, P., Delignette-Muller, M. L., Abbaci, K., Thomas, H., Garric, J. 2016. Analysis of hemocytes in  
619 *Lymnaea stagnalis*: Characterization and effects of repeated hemolymph collections. Fish Shellfish Immunol.  
620 57, 116-126.

621 Buchmann, K. 2014. Evolution of innate immunity: clues from invertebrates via fish to mammals. Front  
622 Immunol. 5, 459.

623 Calienes, A.F., Fraga, J., Pointier, J.P., Yong, M., Sánchez, J., Coustau, C., Gutiérrez, A., Théron, A. 2004.  
624 Detection and genetic distance of resistant populations of *Pseudosuccinea columella* (Mollusca: Lymnaeidae)  
625 to *Fasciola hepatica* (Trematoda: Digenea) using RAPD markers. Acta Trop. 92, 83-87.

626 Castillo, M.G., Wu, X.J., Dinguirard, N., Nyame, A.K., Cummings, R.D., Yoshino, T.P. 2007. Surface membrane  
627 proteins of *Biomphalaria glabrata* embryonic cells bind fucosyl determinants on the tegumental surface of  
628 *Schistosoma mansoni* primary sporocysts. J Parasitol. 93, 832-840.

629 Castillo, M.G., YOSHINO, T.P. 2002. Carbohydrate inhibition of *Biomphalaria glabrata* embryonic (Bge) cell  
630 adhesion to primary sporocysts of *Schistosoma mansoni*. Parasitology. 125, 513-525.

631 Cavalcanti, M.G.S., Filho, F.C., Mendonca, A.M.B., Duarte, G.R., Barbosa, C.C.G.S., De Castro, C.M.M.B., Alves,  
632 L.C., Brayner, F.A. 2012. Morphological characterization of hemocytes from *Biomphalaria glabrata* and  
633 *Biomphalaria straminea*. Micron. 43, 285-291

634 Cima, F., Matozzo, V., Marin, M.G., Ballarin, L. 2000. Haemocytes of the clam *Tapes philippinarum*:  
635 morphofunctional characterization. Fish Shellfish Immunol. 10, 677-693.

636 Combes, C. 2001. Parasitism. The Ecology and Evolution of Intimate Interactions., Chicago, USA, University of  
637 Chicago Press.

638 Coustau, C., Gourbal, B., Duval, D., Yoshino, T.P., Adema, C.M., Mitta, G. 2015. Advances in gastropod  
639 immunity from the study of the interaction between the snail *Biomphalaria glabrata* and its parasites: A  
640 review of research progress over the last decade. Fish Shellfish Immunol. 46, 5-16.

641 Cueto, J.A., Rodriguez, C., Vega, I.A., Castro-Vazquez, A. 2015. Immune defenses of the invasive Apple Snail  
642 *Pomacea canaliculata* (Caenogastropoda, Ampullariidae): phagocytic hemocytes in the circulation and the  
643 kidney. PLoS One. 10, e0123964.



644 Cwiklinsky, K., Dalton, J.P., Dufresne, P.J., La Course, J., Williams, D.J.L., Hodgkinson, J., Paterson, S. 2015. The  
645 *Fasciola hepatica* genome: gene duplication and polymorphism reveal adaptation to the host environment and  
646 the capacity for rapid evolution. *Genome Biol.* 16, 71; doi: 10.1186/s13059-015-0632-2.

647 Chen, T.C. 1968. The compatibility and incompatibility concept as related to trematodes and molluscs. *Pacific*  
648 *Sci.*, XXII, 141-156.

649 DeGaffé, G., Loker, E.S. 1998. Susceptibility of *Biomphalaria glabrata* to infection with *Echinostoma paraensei*:  
650 correlation with the effect of parasite secretory-excretory products on host hemocyte spreading. *J Invertebr*  
651 *Pathol.* 71, 64-72.

652 Dikkeboom, R., Tijnagel, J.M., Van Der Knaap, W.P. 1988. Monoclonal antibody recognized hemocyte  
653 subpopulations in juvenile and adult *Lymnaea stagnalis*: functional characteristics and lectin binding. *Dev*  
654 *Comp Immunol.* 12, 17-32.

655 Dowling, D.J., Hamilton, C.M., Donnelly, S., La Course, J., Brophy, P.M., Dalton, J., O'Neill, S.M. 2010. Major  
656 secretory antigens of the helminth *Fasciola hepatica* activate a suppressive dendritic cell phenotype that  
657 attenuates Th17 cells but fails to activate Th2 immune responses. *Infect Immun.* 78, 793-801.

658 Galinier, R., Portela, J., Moné, Y., Allienne, J.F., Henri, H., Delbecq, S., Mitta, G., Gourbal, B., Duval, D. 2013.  
659 Biomphalysin, a new  $\beta$  pore-forming toxin involved in *Biomphalaria glabrata* immune defense against  
660 *Schistosoma mansoni*. *PLoS Pathog.* 9, e1003216.

661 Georgieva, K., Georgieva, L., Mizinska-Boevska, Y., Stoitsova, S.R. 2016. Study of surface carbohydrates in  
662 *Galba truncatula* tissues before and after infection with *Fasciola hepatica*. *Mem Inst Oswaldo Cruz.* 111, 475-  
663 483.

664 Gorbushin, A.M., Iakovleva, N.V. 2008. The enigma of the haemogram left-shift in periwinkles infected with  
665 trematodes. *Fish Shellfish Immunol.* 24, 745-751.

666 Gourbal, B.E., Guillou, F., Mitta, G., Sibille, P., Thèron, A., Pointier, J.P., Coustau, C. 2008. Excretory-secretory  
667 products of larval *Fasciola hepatica* investigated using a two-dimensional proteomic approach. *Mol Biochem*  
668 *Parasit.* 161, 63-66.

669 Guasconi, L., Serradell, M.C., Garro, A.P., Iacobelli, L., Masih, D.T. 2011. C-type lectins on macrophages  
670 participate in the immunomodulatory response to *Fasciola hepatica* products. *Immunology.* 133, 386-396.

671 Gutiérrez, A., Pointier, J.P., Fraga, J., Jobet, E., Modat, S., Pérez, R.T., Yong, M., Sanchez, J., Loker, E.S., Théron,  
672 A. 2003a. *Fasciola hepatica*: identification of molecular markers for resistant and susceptible *Pseudosuccinea*  
673 *columella* snail hosts. *Exp Parasitol.* 105, 211-218.

674 Gutiérrez, A., Pointier, J.P., Yong, M., Sánchez, J., Théron, A. 2003b. Evidence of phenotypic differences  
675 between resistant and susceptible isolates of *Pseudosuccinea columella* (Gastropoda: Lymnaeidae) to *Fasciola*  
676 *hepatica* (Trematoda: Digenea) in Cuba. *Parasitol Res.* 90, 129-34.

677 Hahn, U.K., Bender, R.C. & Bayne, C.J. 2000. Production of reactive oxygen species by hemocytes of  
678 *Biomphalaria glabrata*: carbohydrate-specific stimulation. *Dev Comp Immunol.* 24, 531-541.

679 Hamilton, C.M., Dowling, D.J., Loscher, C.E., Morphew, R.M., Brophy, P.M., O'Neill, S.M. 2009. The *Fasciola*  
680 *hepatica* tegumental antigen suppresses dendritic cell maturation and function. *Infect Immun.* 77, 2488-2498.

681 Hartenstein, V. 2006. Blood cells and blood cell development in the animal kingdom. *Annu Rev Cell Dev Biol.*  
682 22, 677-712.

683 Hillyer, J.F. 2016. Insect immunology and hematopoiesis. *Dev Comp Immunol.* 58, 102-118.

684 Humbert, E., Coustau, C. 2001. Refractoriness of host haemocytes to parasite immunosuppressive factors as a  
685 putative resistance mechanism in the *Biomphalaria glabrata*-*Echinostoma caproni* system. *Parasitology.* 122,  
686 651-660.

687 Humphries, J.E., Yoshino, T.P. 2008. Regulation of hydrogen peroxide release in circulating haemocytes of the  
688 planorbid snail *Biomphalaria glabrata*. *Dev Comp Immunol.* 32, 554-562.

689 Iakovleva, N.V., Shaposhnikova, T.G., Gorbushin, A.M. 2006. Rediae of echinostomatid and heterophyid  
690 trematodes suppress phagocytosis of haemocytes in *Littorina littorea* (Gastropoda: Prosobranchia). *Exp*  
691 *Parasitol.* 113, 24-29.

692 Jian, J., Konopka, J., Liu, C. 2013. Insights into the role of progranulin in immunity, infection, and inflammation.  
693 *J Leukoc Biol.* 93, 199-208.

694 Johnston, L. A., Yoshino, T.P. 1996. Analysis of lectin- and snail plasma-binding glycopeptides associated with  
695 the tegumental surface of the primary sporocysts of *Schistosoma mansoni*. *Parasitology.* 112, 469-479.

696 Kamiya, T., Oña, L., Wertheim, B., Van Doorn, G.S. 2016. Coevolutionary feedback elevates constitutive  
697 immune defence: a protein network model. *BMC Evol Biol.* 16, 92.

698 Kibbe, W.A. 2007. OligoCalc: an online oligonucleotide properties calculator. *Nucleic Acids Res.* 35, 43–46.

699 Kryukova, N.A., Yurlova, N.I., Rastyagenko, N.M., Antonova, E.V., Glupov, V.V. 2014. The influence of  
700 *Plagiorchis mutationis* larval infection on the cellular immune response of the snail host *Lymnaea stagnalis*. *J*  
701 *Parasitol.* 100, 284–287.

702 Larson, M.K., Bender, R.C., Bayne, C.J. 2014. Resistance of *Biomphalaria glabrata* 13-16-R1 snails to  
703 *Schistosoma mansoni* PR1 is a function of haemocyte abundance and constitutive levels of specific transcripts  
704 in haemocytes. *Int J Parasitol.* 44, 343-353.

705 Lehr, T., Beuerlein, K., Doenhoff, M.J., Grevelding, C.G., Geyer, R. 2008. Localization of carbohydrate  
706 determinants common to *Biomphalaria glabrata* as well as to sporocysts and miracidia of *Schistosoma*  
707 *mansoni*. *Parasitology.* 135, 931-42.

708 Loker, E.S. 2010. Gastropod immunobiology. *Adv Exp Med Biol.* 708, 17-43.

709 Loker, E.S., Bayne, C.J., Buckley, P.M., Kruse, K.T. 1982. Ultrastructure of encapsulation of *Schistosoma*  
710 *mansoni* mother sporocysts by hemocytes of juveniles of the 10-R2 strain of *Biomphalaria glabrata*. *J Parasitol.*  
711 68, 84-94.

712 Loker, E.S., Boston, M.E., Bayne, C.J. 1989. Differential adherence of M line *Biomphalaria glabrata* hemocytes  
713 to *Schistosoma mansoni* and *Echinostoma paraensei* larvae, and experimental manipulation of hemocyte  
714 binding. *J Invertebr Pathol.* 54, 260-268.

715 Loker, E.S., Cimino, D.F., Hertel, L.A. 1992. Excretory-secretory products of *Echinostoma paraensei* sporocysts  
716 mediate interference with *Biomphalaria glabrata* hemocyte functions. *J Parasitol.* 78, 104-115.

717 Mas-Coma, S., Bargues, M.D., Valero, M.A. 2009. Chapter 2. *Fasciola*, lymnaeids and human fascioliasis, with a  
718 global overview on disease transmission, epidemiology, evolutionary genetics, molecular epidemiology and  
719 control. *Adv Parasitol.* 69, 41-146.

720 Matricon-Gondran, M., Letocart, M. 1999. Internal defenses of the snail *Biomphalaria glabrata*. *J Invertebr*  
721 *Pathol.* 74, 235-247.

722 Mohandas, A., Adema, C.M., Van Der Knaap, W.P.W. & Sminia, T. 1992. The effect of haemolymph extraction  
723 on distribution of lysosomal enzymes in *Lymnaea stagnalis* haemocytes: a cytochemical study. *Comp Haematol*  
724 *Int.* 2, 61–67.

725 Noda, S. 1992. Effects of excretory-secretory products of *Echinostoma paraensei* larvae on the hematopoietic  
726 organ of M-line *Biomphalaria glabrata* snails. *J Parasitol.* 78, 512–517.

727 Oliveira, A.L.D., Levada, P.M., Zanotti-Magalhaes, T., Magalhaes, L.A., Ribeiro-Paes, J.T. 2010. Differences in  
728 the number of hemocytes in the snail host *Biomphalaria tenagophila*, resistant and susceptible to *Schistosoma*  
729 *mansoni* infection. *Genet Mol Res.* 9, 2436-2445.

730 Peterson, N.A., Hokke, C.H., Deelder, A.M., Yoshino, T.P. 2009. Glycotope analysis in miracidia and primary  
731 sporocysts of *Schistosoma mansoni*: differential expression during the miracidium-to-sporocyst  
732 transformation. *Int J Parasitol.* 39, 1331-1344.

733 Pila, E.A., Gordy, M.A., Phillips, V.K., Kabore, A.L., Rudko, S.P., Hanington, P.C. 2016a. Endogenous growth  
734 factor stimulation of hemocyte proliferation induces resistance to *Schistosoma mansoni* challenge in the snail  
735 host. *PNAS.* 113, 5305-5310.

736 Pila, E.A., Sullivan, J.T., Wu, X.Z., Fang, J., Rudko, S.P., Gordy, M.A., Hanington, P.C. 2016b. Haematopoiesis in  
737 molluscs: A review of haemocyte development and function in gastropods, cephalopods and bivalves. *Dev*  
738 *Comp Immunol.* 58, 119-128.

739 Portet, A., Pinaud, S., Tetreau, G., Galinier, R., Cosseau, C., Duval, D., Grunau, C., Mitta, G., Gourbal, B. 2017.  
740 Integrated multi-omic analyses in *Biomphalaria-Schistosoma* dialogue reveal the immunobiological significance  
741 of FREP-SmPoMuc interaction. *Dev Comp Immunol.* 75, 16-27.

742 Portet, A., Galinier, R., Pinaud, S., Portela, J., Nowacki, F., Gourbal, B., Duval, D. 2018. BgTEP: An antiprotease  
743 involved in innate immune sensing in *Biomphalaria glabrata*. *Front Immunol.* 9, 1206.

744 Prowse, R.K., Chaplin, P., Robinson, H.C., Spithill, T.W. 2002. *Fasciola hepatica* cathepsin L suppresses sheep  
745 lymphocyte proliferation in vitro and modulates surface CD4 expression on human and ovine T cells. *Parasite*  
746 *Immunol.* 24, 57-66.

747 Ribeiro, V.M.A., Coaglio, A.L., Pereira Oliveira, F.L., Pereira, C.A.J., Lima, W.S. 2017. Dynamics of haemocytes  
748 from *Pseudosuccinea columella* circulating infected by *Fasciola hepatica*. *Braz J Vet Parasitol.* 26, 411-418.

749 Rodríguez, E., Kalay, H., Noya, V., Brossard, N., Giacomini, C., Van Kooyk, Y., García-Vallejo, J.J., Freire, T. 2017.  
750 *Fasciola hepatica* glycoconjugates immunoregulate dendritic cells through the Dendritic Cell-Specific  
751 Intercellular adhesion molecule-3-Grabbing Non-integrin inducing T cell anergy. *Sci Reports.* 7, 46748.

752 Rodríguez, E., Noya, V., Cervi, L., Chiribao, M.L., Brossard, N., Chiale, C., Carmona, C., Giacomini, C., Freire, T.  
753 2015. Glycans from *Fasciola hepatica* modulate the host immune response and TLR-induced maturation of  
754 dendritic cells. PLoS Negl Trop Dis. 9, e0004234.

755 Russo, J., Lagadic, L. 2000. Effects of parasitism and pesticide exposure on characteristics and functions of  
756 hemocyte populations in the freshwater snail *Lymnaea palustris* (Gastropoda, Pulmonata). Cell Biol Toxicol. 16,  
757 15-30.

758 Sánchez, R., Perera, G., Sánchez, J. 1995. Cultivo de *Fossaria cubensis* (Pfeiffer) (Pulmonata: Lymnaeidae)  
759 hospedero intermediario de *Fasciola hepatica* (Linnaeus) en Cuba. Rev Cubana Med Trop. 47, 71-3.

760 Skála, V., Cerníková, A., Jindrová, Z., Kasny, M., Vostry, M., Walker, A.J., Horák, P. 2014. Influence of  
761 *Trichobilharzia regenti* (Digenea: Schistosomatidae) on the defence activity of *Radix lagotis* (Lymnaeidae)  
762 haemocytes. PLoS One. 9, e111696.

763 Sullivan, J.T., Belloir, J.A., Beltran, R.V., Grivakis, A., Ransone, K.A. 2014. Fucoidan stimulates cell division in the  
764 amebocyte-producing organ of the schistosome-transmitting snail *Biomphalaria glabrata*. J Invertebr Pathol.  
765 123, 6-13.

766 Sullivan, J.T., Pikios, S.S., Alonzo, A.Q. 2004. Mitotic responses to extracts of miracidia and cercariae of  
767 *Schistosoma mansoni* in the amebocyte-producing organ of the snail intermediate host *Biomphalaria glabrata*.  
768 J Parasitol. 90, 92–96.

769 Sullivan, J.T., Yeung, J.T. 2011. Incompatibility between miracidia of *Schistosoma mansoni* and *Helisoma duryi*  
770 occurs at two stages: penetration and intramolluscan establishment. J Parasitol. 97, 743-745.

771 Tennessen, J.A., Bonner, K.M., Bollmann, S.R., Johnstun, J.A., Yeh, J.Y., Marine, M., Tavalire, H.F., Bayne, C.J.,  
772 Blouin, M.S. 2015. Genome-wide scan and test of candidate genes in the snail *Biomphalaria glabrata* reveal  
773 new locus influencing resistance to *Schistosoma mansoni*. PLoS Negl Trop Dis. 9, e0004077.

774 Théron, A., Coustau, C. 2005. Are *Biomphalaria* snails resistant to *Schistosoma mansoni*? J Helminthol. 79, 189-  
775 191.

776 Travers, M.A., Mirella Da Silva, P., Le Go, N., Marie, D., Donval, A., Huchette, S., Koken, M., Paillard, C. 2008.  
777 Morphologic, cytometric and functional characterisation of abalone (*Haliotis tuberculata*) haemocytes. Fish  
778 Shellfish Immunol. 24, 400-411.

779 Vázquez, A.A., Lounnas, M., Sánchez, J., Alba, A., Milesi, A., Hurtrez-Boussés, S. 2016. Genetic and infective  
780 diversity of the common liver fluke *Fasciola hepatica* (Trematoda: Digenea) from Cuba. *J Helminthol.* 14, 1-7.

781 Vázquez, A.A., Sánchez, J., Pointier, J.P., Théron, A., Hurtrez-Boussés, S. 2014. *Fasciola hepatica* in Cuba:  
782 compatibility of different isolates with two intermediate intermediate hosts, *Galba cubensis* and  
783 *Pseudosuccinea columella*. *J Helminthol.* 88, 434-440.

784 Wang, Y., Hu, M., Chiang, M.W., Shin, P.K., Cheung, S.G. 2012. Characterization of subpopulations and  
785 immune-related parameters of hemocytes in the green-lipped mussel *Perna viridis*. *Fish Shellfish Immunol.* 32,  
786 381–390.

787 Wu, X.J., Dinguirard, N., Sabat, G., Lui, H.D., Gonzalez, L., Gehring, M., Bickham-Wright, U., Yoshino, T.P. 2017.  
788 Proteomic analysis of *Biomphalaria glabrata* plasma proteins with binding affinity to those expressed by early  
789 developing larval *Schistosoma mansoni*. *PLoS Pathog.* 13, e1006081.

790 Yoshino, T.P., Boyle, J.P., Humphries, J.E. 2001. Receptor-ligand interactions and cellular signaling at the host-  
791 parasite interface. *Parasitology.* 123 Suppl, S143-157.

792 Yoshino, T.P., Wu, X.J., Gonzalez, L.A., Hokke, C.H. 2013. Circulating *Biomphalaria glabrata* hemocyte  
793 subpopulations possess shared schistosome glycans and receptors capable of binding larval glycoconjugates.  
794 *Exp Parasitol.* 133, 28-36.

795 Zahoor, Z., Davies, A. J., Kirk, R.S., Rollinson, D., Walker, A.J. 2008. Disruption of ERK signaling in *Biomphalaria*  
796 *glabrata* defence cells by *Schistosoma mansoni*: implications for parasite survival in the snail host. *Dev Comp*  
797 *Immunol.* 32, 1561-1571.

798 Zhang, S.M., Loker, E.S., Sullivan, J.T. 2016. Pathogen-associated molecular patterns activate expression of  
799 genes involved in cell proliferation, immunity and detoxification in the amebocyte-producing organ of the snail  
800 *Biomphalaria glabrata*. *Dev Comp Immunol.* 56, 25-36.

801

802

803

## 804 **Legends to figures**

805 **Figure 1.** Dot plots (A-B; pooled hemolymph from 4 snails/strain) and DAPI/Phalloidin stained (C-D) naïve  
806 hemocytes from *Pseudosuccinea columella* susceptible and resistant to *Fasciola hepatica*. Green circles in A-B  
807 indicates the two main morphotypes of cells, segregated by size and cytoplasmic complexity. Arrows in C-D  
808 point at the small-round shape cell with high nucleus/cytoplasm ratio morphotype; phalloidin (actin labelling,  
809 green) and DAPI (nucleus labelling, blue).

810 **Figure 2.** Confocal microscopy images of *Pseudosuccinea columella* hemocytes from naïve (upper panel, 0h)  
811 and *F. hepatica*-challenged (lower panel, 24h post-exposure), resistant and susceptible snails from four  
812 different populations. Arrows in A-D point at the small-roundly shape cell with high nucleus/cytoplasm ratio  
813 morphotype. Cell staining was performed with Phalloidin (actin labelling, green) and DAPI (nucleus labelling,  
814 blue).

815 **Figure 3.** (A) Number of circulating hemocytes in the hemolymph of resistant and susceptible *Pseudosuccinea*  
816 *columella* at different time points post-exposure to *Fasciola hepatica* miracidia. B) Hemocytes of 24h-  
817 challenged resistant snails with DAPI-labelled nucleus (in blue) in some of which active proliferation (EdU  
818 labelling, green), corresponding only to small- roundly shape cells (arrow, lower panel), was observed. C)  
819 Relative quantity of granulin expression in resistant and susceptible *P. columella* at different time points post  
820 exposure to *F. hepatica*. In bars, normalized RQs at each time point against the RQ in naïve conditions (0h) for  
821 each strain; different letters represent statistical significance among experimental points  $P < 0.05$ .

822 **Figure 4.** A) Quantitative scale used to represent the observed differences (encapsulation grade: 1 to 5) on the  
823 *in vitro* encapsulation of *Fasciola hepatica* sporocyst by *Pseudosuccinea columella* hemocytes. B) Association  
824 (through multivariate correspondence analysis), of the encapsulation grade of hemocytes from resistant (R)  
825 and susceptible (S) *P. columella* with the assayed conditions (incubation of naïve hemocytes with  
826 monosaccharides or *F. hepatica* challenge). *P. columella* populations: LP: La Palma, LC: La Coca, AU: Aurora,  
827 NE: Negrines; Monosaccharides: man: mannose, fuc: fucose, gal: galactose, glu: Glucose; Time-points of  
828 experimental infection: 0h: naïve hemocytes, 24h: 24h-challenged hemocytes.

829 **Figure 5.** Integrative model of the immune response in resistant (R) *Pseudosuccinea columella* snails toward  
830 the digenean *Fasciola hepatica*. The significant immune competence in naïve snails prior to infection possibly  
831 helps to negate and avoid *F. hepatica*-driven immune modulation, leading to enhanced immune

832 responsiveness and to activation of specific pathways resulting in parasite elimination through hemocytic  
833 encapsulation. During interaction, the early molecular crosstalk between parasite-specific antigens and host  
834 receptors increases the hematopoiesis of circulating blast-like cells, the immune activation of large-adherent  
835 hemocytes to assume a more active morphotype and stimulates the production of the cytokine granulins. The  
836 constitutive high expression levels of protein kinase C (PKC) and of several Toll-like receptors (TLR) in naïve  
837 resistant snails may facilitate orchestration of the induced immune response that is triggered upon parasite  
838 penetration. Likewise, the abundant expression of macrophage migration inhibitory factor (MIF) prior to  
839 infection may also contribute to the immune stimulation of hemocytes. All the latter may contribute to build  
840 up the cellular immune reaction; at 24h post exposure the increase in the encapsulation activity of the  
841 hemocytes results in the visualization of well-formed capsules surrounding the parasite that will eventually  
842 leads to the parasite death. In this sense, the abundance of superoxide dismutase and nitric oxide synthase  
843 along with anti-parasite specific humoral or intracellular immune effectors (e.g. ferritin, lipopolysaccharide-  
844 binding protein/bactericidal permeability-increasing protein), as reported in resistant *P. columella* prior  
845 infection, might positively influence parasite clearance following exposure. Man: mannose, ROS: reactive  
846 oxygen species, RNS: reactive nitrogen species.

847


848

849



850 Figure 1

851



852

853


854

855

856

857 Figure 2

858




859

860

861

862 Figure 3

863





864

865

866

867 Figure 4




868

869

870

871 Figure 5

872



873

874