



HAL
open science

Abstracting Linear Equation Systems

Emilie Allart, Joachim Niehren, Cristian Versari

► **To cite this version:**

Emilie Allart, Joachim Niehren, Cristian Versari. Abstracting Linear Equation Systems. 2020. hal-02279942v2

HAL Id: hal-02279942

<https://hal.science/hal-02279942v2>

Preprint submitted on 15 May 2020 (v2), last revised 24 Jun 2022 (v9)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Abstracting Linear Equation Systems

2 **Emilie Allart**

3 BioComputing CRISAL - Centre de Recherche en Informatique, Signal et Automatique de Lille
4 (CRISAL) - UMR 9189,
5 Université de Lille
6 emilie.allart@univ-lille.fr

7 **Joachim Niehren**

8 LINKS - Linking Dynamic Data CRISAL - Centre de Recherche en Informatique, Signal et
9 Automatique de Lille (CRISAL) - UMR 9189, Inria Lille - Nord Europe,
10 BioComputing CRISAL - Centre de Recherche en Informatique, Signal et Automatique de Lille
11 (CRISAL) - UMR 9189

12 **Cristian Versari**

13 BioComputing CRISAL - Centre de Recherche en Informatique, Signal et Automatique de Lille
14 (CRISAL) - UMR 9189,
15 Université de Lille

16 — Abstract —

17 We show how to compute finite abstractions of linear equations systems over the reals. We develop
18 a general method based on elementary modes, that can be applied to the various abstractions into
19 finite structures, including the sign abstraction as used in abstract interpretation based program
20 analysis, and the difference abstraction as used in change prediction algorithms for flux networks in
21 systems biology.

22 **2012 ACM Subject Classification**

23 **Keywords and phrases** Linear equation systems, abstract interpretation, systems biology.

24 **Digital Object Identifier** 10.4230/LIPIcs...

25 **1** Introduction

26 Systems of linear equations serve in abstract interpretation to abstract from the concrete
27 semantics of programs with arithmetic operations [4, 10]. In systems biology, systems of
28 linear equations are used to describe the fluxes of a reaction network in a steady state [13, 14].
29 Both applications raise a very similar question on how to compute an abstraction for linear
30 equations systems. For program analysis based on abstract interpretation [15], one might ask
31 for instance, whether the value of a variable x in a program is strictly positive if the value
32 of another variable y was. This question is about the sign abstraction of the \mathbb{R} -solution set
33 of a linear equation system (that itself abstracts from the programs semantics). In systems
34 biology [11, 17, 5, 1] one might want to know for a given flux network, whether a flux x must
35 increase (resp. decrease) if some other flux y does. This question concerns the difference
36 abstraction of a pair of \mathbb{R} -solutions of the equation system, that is of an \mathbb{R}^2 -solution of the
37 system of linear equations where addition and multiplication are defined component-wise.

We therefore study the question of how to compute finite abstractions of the solution set
of a system of linear equation over the reals. Given an homomorphism $h : \mathbb{R}^k \rightarrow \Delta$ between
 Σ -structures where $\Sigma = \{+, *, 0, 1\}$ and Δ is finite, and a linear equation system ϕ with
integer coefficients, the question is how to compute the h -abstraction of the solution set of ϕ .
If V is the set of variables of ϕ , then the set to be computed is:

$$h \circ \text{sol}^{\mathbb{R}^k}(\phi) = \{h \circ \alpha \mid \alpha : V \rightarrow \mathbb{R}^k, \text{s.t. } \mathbb{R}^k, \alpha \models \phi\}$$



© Emilie Allart and Joachim Niehren and Cristian Versari;
licensed under Creative Commons License CC-BY

Leibniz International Proceedings in Informatics

LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

38 This problem generalizes on both questions above. The instance for the first question concerns
 39 the sign abstraction $h_{\mathbb{S}} : \mathbb{R} \rightarrow \{-1, 0, 1\}$ which satisfies $h_{\mathbb{S}}(r) = 1$ if $r > 0$, $h_{\mathbb{S}}(r) = -1$ if
 40 $r < 0$ and otherwise $h_{\mathbb{S}}(0) = 0$. The second instance concerns the difference abstraction
 41 $h_{\Delta_3} : \mathbb{R}^2 \rightarrow \{\uparrow, \downarrow, \sim\}$, where \uparrow stands for increase, \downarrow for decrease, and \sim for no change. It
 42 satisfies $h_{\Delta_3}(r, r') = \uparrow$ if $r < r'$, $h_{\Delta_3}(r, r') = \downarrow$ if $r > r'$, and otherwise $h_{\Delta_3}(r, r') = \sim$.

43 We next illustrate the difficulty of the problem by showing how to compute $h_{\mathbb{S}} \circ \text{sol}^{\mathbb{R}}(\phi)$
 44 from ϕ with the existing methods. Note that the set $\text{sol}^{\mathbb{R}}(\phi)$ cannot be enumerated since it
 45 is infinite. Instead we can enumerate the set of sign assignments $\alpha : V \rightarrow \{-1, 0, 1\}$ which
 46 is finite. Furthermore, since $h_{\mathbb{S}}(x) = -1$ is equivalent to $x < 0$ and $h_{\mathbb{S}}(x) = 1$ to $0 < x$,
 47 the system $\phi \wedge \bigwedge_{x \in V} h_{\mathbb{S}}(x) = \alpha(x)$ is equivalent to a system of linear equations and strict
 48 inequations. The satisfiability of such a system can be decided in at most exponential time [6].
 49 However, this method is not feasible in practice given that the number of sign assignments
 50 grows exponentially with the number of variables. So the question is whether there exists a
 51 more efficient algorithm for computing $h_{\mathbb{S}} \circ \text{sol}^{\mathbb{R}}(\phi)$.

52 From John's overapproximation theorem [11, 17], we know for any homomorphism
 53 $h : S \rightarrow \Delta$ between Σ -structures and any negation-free first-order Σ -formula ϕ that $h \circ$
 54 $\text{sol}^S(\phi) \subseteq \text{sol}^{\Delta}(\phi)$. We call ϕ h -exact if and only if $h \circ \text{sol}^S(\phi) = \text{sol}^{\Delta}(\phi)$. If Δ is finite, then
 55 we can compute for any h -exact formula ϕ the abstraction $h \circ \text{sol}^S(\phi)$ by computing $\text{sol}^{\Delta}(\phi)$
 56 with finite domain constraint programming [18].

57 In a first step, we show that any integer linear matrix equation $Ax=0$ can be transformed
 58 into some \mathbb{R}_+ -equivalent Σ -formula that is quasi-positive and quasi-triangular and thus
 59 $h_{\mathbb{B}}$ -exact. This transformation is based on the computation of elementary modes [16, 8, 9, 19]
 60 – the extreme rays of the cone $\text{sol}^{\mathbb{R}^+}(Ax=0)$ – that can be done in practice by various libraries
 61 from computational geometry [3]. The conversion may take exponential time in the worst
 62 case, but is often well-behaved. The correctness of the conversion relies on the fact that the
 63 Σ -algebras \mathbb{R}_+ and \mathbb{B} permit unique division by nonzero natural numbers.

64 In the second step, we introduce $h_{\mathbb{B}}$ -mixed systems, which generalize on systems of linear
 65 equations, positive polynomial equations $p=0$ and inequations $p \neq 0$ where polynomial p has
 66 no constant term. We then show that $h_{\mathbb{B}}$ -mixed systems can be converted to an $h_{\mathbb{B}}$ -exact
 67 formula too. In order to do so, extend on the results from the first step by introducing
 68 the notion of $h_{\mathbb{B}}$ -invariant Σ -formulas, which subsume the polynomial equations $p=0$ and
 69 inequations $p \neq 0$ for all positive polynomials p without constant terms.

70 In a third step, we rewrite linear equations systems ϕ into $h_{\mathbb{B}}$ -mixed formulas ϕ' , based on
 71 the two previous steps, so that sign abstraction of $h_{\mathbb{S}} \circ \text{sol}^{\mathbb{R}}(\phi)$ can be computed from boolean
 72 abstraction $h_{\mathbb{B}} \circ \text{sol}^{\mathbb{R}}(\phi') = \text{sol}^{\mathbb{B}}(\phi')$. It is then sufficient to compute the boolean solution set
 73 $\text{sol}^{\mathbb{B}}(\phi')$ by finite domain constraint programming. The rewriting approach based the results
 74 for $h_{\mathbb{B}}$ -mixed systems presented here was applied recently by the authors [1] to the difference
 75 abstraction $h_{\Delta_3} : \mathbb{R}_+^2 \rightarrow \{\uparrow, \downarrow, \sim\}$ and a refinement thereof into a finite Σ -structure with 6
 76 elements. This procedure was implemented and applied successfully for change prediction in
 77 systems biology. It illustrates that our results presented here do provide a general framework
 78 enabling the computation of various finite abstractions of linear equation systems.

79 We illustrate our results by applying the sign abstraction for program analysis based on
 80 abstract interpretation. We consider the Python implementation in Fig. 1 of the function
 81 `integral` : $\mathbb{R}^2 \rightarrow \mathbb{R}$ with parameter `f` : $\mathbb{R} \rightarrow \mathbb{R}$. A call `integral(a, step)` computes the
 82 approximation of the integral $\int_0^a \mathbf{f}(x)dx$ with stepwidth `step`. Abstract interpretation applied
 83 to this program with a polyhedral abstract domain may produce the following first-order

```

def integral(a: float, step: float):
    if a < 0: raise ValueError('This should never happen')
    if step > a:
        return 0
    else:
        return step * f(a) + integral(a - step, step)

```

■ **Figure 1** Python function approximating the integral $\int_0^a \mathbf{f}(x)dx$ for a given function $\mathbf{f} : \mathbb{R} \rightarrow \mathbb{R}$.

84 formula ϕ_{integral} :

$$\begin{aligned}
 & (\text{throw_exception} = 1 \iff a < 0) \wedge (\text{do_recursion} = 1 \iff \text{step} \leq a) \wedge \\
 & a_{\text{rec}} = a - \text{step} \wedge \text{step}_{\text{rec}} = \text{step}
 \end{aligned}$$

86 This formula uses the following variables: a flag *throw_exception* that is true on exception
 87 throwing; a flag *do_recursion* that is true when a recursive call is made; two variables a_{rec} ,
 88 step_{rec} representing the parameters passed recursively to `integral`.

89 In order to know whether an exception may be thrown, we are interested in the sign
 90 abstraction for this formula $h_{\mathbb{S}} \circ \text{sol}^{\mathbb{R}}(\phi_{\text{integral}})$. According to John's Theorem [11], this
 91 abstraction can be overapproximated by $\text{sol}^{\mathbb{S}}(\phi_{\text{integral}})$ which in turn can be computed
 92 by finite domain constraint programming. However, this approximation does not rule out
 93 that a_{rec} may be strictly negative when *do_recursion* is true, although this condition is not
 94 possible when running the program. Conversely, this is correctly reflected by the abstraction
 95 of its abstract interpretation $h_{\mathbb{S}} \circ \text{sol}^{\mathbb{R}}(\phi_{\text{integral}})$, that can be computed by converting it to a
 96 $h_{\mathbb{B}}$ -mixed system which is then solved with the methods presented above.

97 2 Preliminaries

98 *Sets.* We start with usual notation from set theory. Let \mathbb{N} be the set natural numbers and
 99 \mathbb{R}_+ the set of positive real numbers, both including 0. For any set A and $n \in \mathbb{N}$, the set
 100 of n -tuples of elements in A is denoted by A^n . The i -th projection function on n -tuples of
 101 elements in A , where $1 \leq i \leq n$ is the function $\pi_i : A^n \rightarrow A$ such that $\pi_i(a_1, \dots, a_n) = a_i$ for
 102 all $a_1, \dots, a_n \in A$. If A is finite the number of elements of A is denote by $|A|$.

103 *Projections and Pairs.* The projection $\pi_a(f)$ of a function $f : A \rightarrow B$ is its restriction $\alpha_{|A \setminus \{a\}}$.
 104 The projection of a set F of functions $f : A \rightarrow B$ is $\pi_a(F) = \{\pi_a(f) \mid f \in F\}$. Furthermore,
 105 we define the pair function $f^2 : A^2 \rightarrow B^2$ such that $f^2(a_1, a_2) = (f(a_1), f(a_2))$.

106 *Σ -Algebras and Σ -Structures.* We next recall the usual notions of Σ -algebras and of homomor-
 107 phism between Σ -algebras. Let $\Sigma = \cup_{n \geq 0} F^{(n)} \uplus C$ be a ranked signature. The elements of
 108 $f \in F^{(n)}$ are called the n -ary function symbols of Σ and the elements in $c \in C$ its constants.

109 ► **Definition 1.** A Σ -algebra $S = (\text{dom}(S), \cdot^S)$ consists of a set $\text{dom}(S)$ and an interpretation
 110 \cdot^S such that $c^S \in \text{dom}(S)$ for all $c \in C$, and $f^S : \text{dom}(S)^n \rightarrow \text{dom}(S)$ for all $f \in F^{(n)}$.

111 In order to generalize Σ -algebras to Σ -structures, we consider n -ary function symbols as
 112 $n+1$ -ary relation symbols.

113 ► **Definition 2.** A Σ -structure $\Delta = (\text{dom}(\Delta), \cdot^{\Delta})$ consists of a set $\text{dom}(\Delta)$ and an interpre-
 114 tation \cdot^{Δ} such that $c^{\Delta} \in \text{dom}(\Delta)$ for all $c \in C$ and $f^{\Delta} \subseteq \text{dom}(\Delta)^{n+1}$ for all $f \in F^{(n)}$.

115 Clearly, any Σ -algebra is also a Σ -structure. Note also that symbols in $F^{(0)}$ are interpreted
 116 as monadic relations, i.e., as subsets of the domain, in contrast to constants in C that are
 117 interpreted as elements of the domain.

d	d'	$d +^{\mathbb{S}} d'$	$d *^{\mathbb{S}} d'$	d	d'	$d +^{\mathbb{S}} d'$	$d *^{\mathbb{S}} d'$	d	d'	$d +^{\mathbb{S}} d'$	$d *^{\mathbb{S}} d'$
-1	1	{-1, 0, 1}	-1	0	1	1	0	1	1	1	1
-1	0	-1	0	0	0	0	0	1	0	1	0
-1	-1	-1	1	0	-1	-1	0	1	-1	{-1, 0, 1}	-1

■ **Figure 2** Interpretation of Σ -structure of signs \mathbb{S} .

118 ▶ **Definition 3.** A homomorphism between two Σ -structures S and Δ is a function $h :$
 119 $dom(S) \rightarrow dom(\Delta)$ such that for $c \in C$, $n \in \mathbb{N}$, $f \in F^{(n)}$, and $s_1, \dots, s_{n+1} \in dom(S)$:

- 120 1. $h(c^S) = c^\Delta$, and
 121 2. if $(s_1, \dots, s_{n+1}) \in f^S$ then $(h(s_1), \dots, h(s_{n+1})) \in f^\Delta$.

122 We can convert any $n + 1$ -ary relation to a n -ary set valued functions. In this way any
 123 n -function is converted to a n -ary set valued n -functions. In other words, functions of type
 124 $D^n \rightarrow D$ are converted to functions of type $D^n \rightarrow 2^D$ where $D = dom(\Delta)$. In set-valued
 125 notation, the second condition on homomorphism can then be rewritten equivalently as
 126 $h(f^S(s_1, \dots, s_n)) \subseteq f^\Delta(h(s_1), \dots, h(s_n))$. A homomorphism for Σ -algebras thus satisfies
 127 $h(c^S) = c^\Delta$ and $h(f^S(s_1, \dots, s_n)) = f^\Delta(h(s_1), \dots, h(s_n))$.

128 ▶ **Definition 4.** A Σ -abstraction is a homomorphism $h: S \rightarrow \Delta$ between Σ -structures such
 129 that $dom(\Delta) \subseteq dom(S)$.

130 3 The Boolean and the Sign Abstraction

131 Throughout the paper we will use the signature $\Sigma = F^{(2)} \uplus C$ with two binary function
 132 symbols in $F^{(2)} = \{+, *\}$, and two constants $C = \{0, 1\}$. In the Σ -algebras that we will
 133 consider the functions $+$ and $*$ will be associative and commutative, with neutral element 0
 134 and 1 respectively.

135 ▶ **Example 5.** The set of positive reals \mathbb{R}_+ can be turned into a Σ -algebra, in which the
 136 functions symbols are interpreted as addition of positive reals $+^{\mathbb{R}_+}$, multiplication of positive
 137 reals $*^{\mathbb{R}_+}$. The constants are interpreted by themselves $0^{\mathbb{R}_+} = 0$ and $1^{\mathbb{R}_+} = 1$.

138 ▶ **Example 6.** The set of Booleans $\mathbb{B} = \{0, 1\} \subseteq \mathbb{R}_+$ equally defines a Σ -algebra. There,
 139 the function symbols are interpreted as a disjunction $+^{\mathbb{B}} = \vee^{\mathbb{B}}$ and conjunction $*^{\mathbb{B}} = \wedge^{\mathbb{B}}$ on
 140 Booleans. The constants are interpreted by themselves $0^{\mathbb{B}} = 0$ and $1^{\mathbb{B}} = 1$.

141 In order to abstract positive real numbers into booleans, we can define a function $h_{\mathbb{B}} : \mathbb{R}_+ \rightarrow \mathbb{B}$
 142 such that $h_{\mathbb{B}}(0) = 0$ and $h_{\mathbb{B}}(r) = 1$ if $r > 0$.

143 ▶ **Lemma 7.** The function $h_{\mathbb{B}} : \mathbb{R}_+ \rightarrow \mathbb{B}$ is a Σ -abstraction between Σ -algebras.

144 ▶ **Example 8.** The set of signs $\{-1, 0, 1\} \subseteq \mathbb{R}$ can be turned into a Σ -structure $\mathbb{S} =$
 145 $(\{-1, 0, 1\}, \cdot^{\mathbb{S}})$ with the interpretation $+^{\mathbb{S}}$ and $*^{\mathbb{S}}$ given in Fig. 2. The constants are interpreted
 146 by themselves $0^{\mathbb{S}} = 0$ and $1^{\mathbb{S}} = 1$. Note that all $+^{\mathbb{S}}$ contains $(-1, 1, -1)$, $(-1, 1, 1)$ and
 147 $(-1, 1, 0)$ meaning that the sum of a strictly negative and a strictly positive real has a sign in
 148 $-1 +^{\mathbb{S}} 1$, so it may either be strictly positive, strictly negative, or zero. For this reason, \mathbb{S} is
 149 a Σ -structure but not a Σ -algebra.

150 We define the sign abstraction $h_{\mathbb{S}} : \mathbb{R} \rightarrow \mathbb{S}$ such that $h_{\mathbb{S}}(0) = 0$, $h_{\mathbb{S}}(r) = -1$ for all strictly
 151 negative reals $r < 0$ and $h_{\mathbb{S}}(r) = 1$ for all strictly positive reals $r > 0$.

152 ▶ **Lemma 9.** $h_{\mathbb{S}} : \mathbb{R} \rightarrow \mathbb{S}$ is a Σ -abstraction into a Σ -structure (that is not a Σ -algebra).

$$\begin{aligned}
e \in \mathcal{E}_\Sigma & ::= x \mid c \mid e \odot e' && \text{where } c \in C = \{0, 1\} \text{ and } \odot \in F^{(2)} = \{+, *\} \\
\phi \in \mathcal{F}_\Sigma & ::= e \overset{\circ}{=} e' \mid \exists x. \phi \mid \phi \wedge \phi \mid \neg \phi && \text{where } x \in \mathcal{V}
\end{aligned}$$

■ **Figure 3** First-order Σ -expressions and Σ -formulas, where $\Sigma = C \cup F^{(2)}$.

4 Abstractions and First-Order Logic

We recall the first-order logic for Σ -structures and recall John's theorem [11] on how to overapproximate the Σ -abstraction first-order Σ -formulas.

We fix a set of variables \mathcal{V} (for instance $\mathcal{V} = \mathbb{N}$). The variables in \mathcal{V} will be ranged over by x and y . The abstract syntax in Fig. 3 define the set of first-order expressions $e \in \mathcal{E}_\Sigma$ which are constructed from the function symbols in the signature Σ and the variables in \mathcal{V} . A Σ -equation is a pair $e \overset{\circ}{=} e'$ where $e, e' \in \mathcal{E}_\Sigma$. A first-order formula $\phi \in \mathcal{F}_\Sigma$ is constructed from Σ -equations with the usual first-order connectives. As shortcuts, we define the formula $true =_{\text{def}} 1 \overset{\circ}{=} 1$ and for any sequence of formulas ϕ_1, \dots, ϕ_n we define $\bigwedge_{i=1}^n \phi_i$ as $\phi_1 \wedge \dots \wedge \phi_n$ which is equal to $true$ if $n = 0$. Furthermore, we define formulas $e \neq 0$ by $\neg e \overset{\circ}{=} 0$.

The semantics of first-order logic is standard. Let S be a Σ -structure and $\alpha : V \rightarrow \text{dom}(S)$ be a variable assignment. For any expressions $e \in \mathcal{E}_\Sigma$ and variable assignment $\mathcal{V}(e) \subseteq V$, the semantics defines a subset of $\llbracket e \rrbracket^{\alpha, S} \subseteq \text{dom}(S)$, and for any formula $\phi \in \mathcal{F}_\Sigma$ with $\mathcal{V}(\phi) \subseteq V$ a truth value $\llbracket \phi \rrbracket^{\alpha, S} \in \mathbb{B}$. Expressions $e, e' \in \mathcal{E}_\Sigma$ may be set valued in the case that S is not a Σ -algebra. Therefore, the equality symbol $\overset{\circ}{=}$ will be interpreted as nondisjointness, i.e., $e \overset{\circ}{=} e'$ is true if and only if $\llbracket e \rrbracket^{\alpha, S} \cap \llbracket e' \rrbracket^{\alpha, S} \neq \emptyset$. If S is a Σ -algebra, then both sets will be singletons. Therefore, the equality symbol $\overset{\circ}{=}$ is indeed interpreted as equality for Σ -algebra, but not for Σ -structures. See Fig. 4 of the appendix for the details.

The set of solutions of a formula $\phi \in \mathcal{F}_\Sigma$ over a Σ -algebra S with respect to a set of variables V that contains $\mathcal{V}(\phi)$ is defined by $\text{sol}_V^S(\phi) = \{\alpha : V \rightarrow \text{dom}(S) \mid \llbracket \phi \rrbracket^{\alpha, S} = 1\}$. If $V = \mathcal{V}(\phi)$ we omit the index V , i.e., $\text{sol}^S(\phi) = \text{sol}_V^S(\phi)$.

► **Lemma 10 Quantification is projection.** $\text{sol}^S(\exists x. \phi) = \pi_x(\text{sol}^S(\phi))$.

Let $h : S \rightarrow \Delta$ be a Σ -abstraction and $V \subseteq \mathcal{V}$. For any subset of assignments R of type $V \rightarrow \text{dom}(S)$ we define $h \circ R = \{h \circ \alpha : V \rightarrow \text{dom}(\Delta) \mid \alpha \in R\}$.

► **Theorem 11 John's Overapproximation Theorem [1, 17, 11].** For any Σ -abstraction $h : S \rightarrow \Delta$ between Σ -structures and negation-free formula $\phi \in \mathcal{F}_\Sigma$: $h \circ \text{sol}^S(\phi) \subseteq \text{sol}^\Delta(\phi)$.

We only give a sketch of the proof. Let $\alpha : V \rightarrow \text{dom}(S)$. For any expression $e \in \mathcal{E}_\Sigma$ such that $\mathcal{V}(e) \subseteq V$ we can show that $h(\llbracket e \rrbracket^{\alpha, S}) = \llbracket e \rrbracket^{h \circ \alpha, \Delta}$ by induction on the structure of e . It then follows for any positive formula $\phi \in \mathcal{F}_\Sigma$ with $\mathcal{V}(\phi) \subseteq V$ that $\llbracket \phi \rrbracket^{\alpha, S} \subseteq \llbracket \phi \rrbracket^{h \circ \alpha, \Delta}$. This is equivalent to that: $\{h \circ \alpha \mid \alpha \in \text{sol}_V^S(\phi)\} \subseteq \text{sol}_V^\Delta(\phi)$ and thus $h \circ \text{sol}_V^S(\phi) \subseteq \text{sol}_V^\Delta(\phi)$.

5 Linear Equation Systems and Elementary Modes

We are interested in systems of Σ -equation where $\Sigma = \{+, *, 1, 0\}$. The base case will be homogeneous linear equations systems with natural coefficients, which capture linear matrix integer equations $\mathbf{Ax} = 0$. We will show that elementary modes [16, 8, 9, 19] can be used to make linear integer matrix equations quasi-positive and strongly-triangular.

We also need systems of polynomial equations, with natural coefficients and no constant term, that are nonlinear. For any natural n and expression $e, e_1, \dots, e_n \in \mathcal{E}_\Sigma$, we define the

XX:6 Abstracting Linear Equation Systems

190 expression $\prod_{i=1}^n e_i = e_1 * \dots * e_n$, which is equal to 1 if $n = 0$ and to $\sum_{i=1}^n e_i = e_1 + \dots + e_n$
 191 which is equal to 0 if $n = 0$. Furthermore, let $e^n = \prod_{i=1}^n e$ and $ne =_{\text{def}} \sum_{i=1}^n e$.

192 ► **Definition 12.** A Σ -equation is called positive if it has the form $e \stackrel{\circ}{=} 0$ and quasi-positive if
 193 it has the form $e \stackrel{\circ}{=} ny$, where $n \in \mathbb{N}$, $y \in \mathcal{V}$, and $e \in \mathcal{E}_\Sigma$.

194 This definition makes sense, since all constants in Σ -expressions are positive and all operators
 195 of Σ -expressions preserve positivity. Note also that any positive equation is quasi-positive
 196 since the constant 0 is equal to the polynomial $0y$. A system of Σ -equations is a conjunctive
 197 formula of the form $\bigwedge_{i=1}^n e_i \stackrel{\circ}{=} e'_i$ in \mathcal{F}_Σ . We call a system of Σ -equations positive respectively
 198 quasi-positive if all its equations are.

199 A polynomial (with natural coefficients) is an expression of the form $\sum_{j=1}^l n_j \prod_{k=1}^{i_j} x_{j,k}^{m_{j,k}}$
 200 where l and i_j are naturals, $x_{1,1}, \dots, x_{l,i_l}$ variables, all $n_j \neq 0$ naturals called the coefficients,
 201 and all $m_{j,k} \neq 0$ naturals called the exponents. The products $\prod_{k=1}^{i_j} x_{j,k}^{m_{j,k}}$ are called the
 202 monomials of the polynomial.

203 ► **Definition 13.** A polynomial $\sum_{j=1}^l n_j \prod_{k=1}^{i_j} x_{j,k}^{m_{j,k}}$ with natural coefficients $n_j \neq 0$ has no
 204 constant term if none of its monomials is equal to 1, i.e., $i_j \neq 0$ for all $1 \leq j \leq l$. It is linear
 205 if all its monomials are variables, i.e. $i_j = 1$ and $m_{j,1} = \dots = m_{j,i_j} = 1$ for all $1 \leq j \leq l$.

206 Note that any linear polynomial has the form $\sum_{j=1}^l n_j x_{j,1}$ where l and all $n_j \neq 0$ are naturals
 207 and all $x_{j,1}$ are variables. In particular, linear polynomials do not have a constant term.
 208 Furthermore, note that the constant 0 is equal to the linear polynomial with $l = 0$. A
 209 polynomial equation is a Σ -equation $p \stackrel{\circ}{=} p'$ between polynomials. A (homogeneous) linear
 210 equation is a polynomial equation with linear polynomials, so without constant terms. A
 211 linear equation system is a system of linear equations.

212 An (homogeneous) linear integer matrix equation has the form $A\mathbf{y} \stackrel{\circ}{=} 0$ where A is an
 213 $n \times m$ matrix of integers for some naturals m, n such that $\mathbf{y} \in \mathcal{V}^m$. Any linear integer
 214 matrix equation can be turned into a linear equation system with natural coefficients, by
 215 bringing the negative coefficients on the right-hand side. For instance, the linear integer
 216 matrix equation on the right corresponds to the linear equation system with natural coefficients $3x \stackrel{\circ}{=} 0 \wedge 2x \stackrel{\circ}{=} 5y$.

$$\begin{pmatrix} 3 & 0 \\ 2 & -5 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \stackrel{\circ}{=} 0$$

217 This system is quasi-positive, but not positive since $5y$ appears on a right-hand side.
 218 More generally, the linear equation system for an linear integer matrix equation $A\mathbf{y} \stackrel{\circ}{=} 0$ is
 219 positive if and only if all integers in A are positive, and quasi-positive, if each line of A
 220 contains at most one negative integer. Furthermore, the above linear equation system is
 221 triangular in the following sense, but not strongly triangular:

222 ► **Definition 14.** We call a quasi-positive system of Σ -equations triangular if it has the form
 223 $\bigwedge_{l=1}^n e_l \stackrel{\circ}{=} n_l y_l$ such that the variables y_l are l -fresh for all $1 \leq l \leq n$, i.e., $y_l \notin \mathcal{V}(\bigwedge_{i=1}^{l-1} e_i \stackrel{\circ}{=} e'_i)$
 224 and if $n_l \neq 0$ then $y_l \notin \mathcal{V}(e_l)$. We call the quasi-positive polynomial system strongly-triangular
 225 if it is triangular and satisfies $n_l \neq 0$ for all $1 \leq l \leq n$.

226 Consider a linear integer matrix equation $A\mathbf{y} \stackrel{\circ}{=} 0$. If A is positive and triangular, then the
 227 corresponding linear equation system is positive and triangular too. For being quasi-positive
 228 and strongly-triangular, the integers below the diagonal of A must negative, those on the
 229 diagonal must be strictly negative, and those on the right of the diagonal must be positive.

230 ► **Theorem 15 Elementary Modes.** For any system of linear equations ϕ with natural
 231 coefficients, one can compute in at most exponential time an \mathbb{R}_+ -equivalent formula $\exists \mathbf{x}.\phi'$
 232 such that ϕ' is a quasi-positive strongly-triangular system of linear equations with natural
 233 coefficients and \mathbf{x} the sequence of variables on the left hand sides of the equations.

234 The theorem applies in particular to the linear equation systems of integer matrix
 235 equations $A\mathbf{y} \stackrel{\circ}{=} 0$. It shows that there exists a matrix E of naturals, a vector of naturals \mathbf{n} ,
 236 and a vector of fresh variables \mathbf{x} , such that $A\mathbf{y} \stackrel{\circ}{=} 0$ is \mathbb{R}_+ -equivalent to $\exists \mathbf{x}. E\mathbf{x} \stackrel{\circ}{=} \mathbf{n}\mathbf{y}$.

237 **Proof** Consider a system ϕ of linear equations with natural coefficients. Geometrically,
 238 the solution space of ϕ over the reals is a linear subspace of $\mathbb{R}^{\mathcal{V}(\phi)}$. When restricted to
 239 positive reals, as we do, this linear subspace is to be intersected with the positive cone
 240 $\mathbb{R}_+^{\mathcal{V}(\phi)}$. Therefore, $\text{sol}^{\mathbb{R}_+}(\phi)$ is a rational cone obtained by intersecting finitely many half-
 241 spaces: its H-representation is defined by the inequations $A\mathbf{y} \leq 0 \wedge A\mathbf{y} \geq 0 \wedge \mathbf{y} \geq 0$. The
 242 elementary modes of ϕ are the extreme rays of this cone and allow its V-representation.
 243 Up to normalization there are finitely many such extreme rays. Moreover, since the cone
 244 is rational and ϕ is homogeneous, the elementary modes can be normalised so that the
 245 V-representation contains only integer coefficients. The normalized elementary modes will be
 246 vectors of naturals in $\mathbb{N}^{\mathcal{V}(\phi)}$. Let e_1, \dots, e_n be the set of all normalized elementary modes
 247 in some arbitrary total order. Let E be the matrix with columns e_1, \dots, e_n . Let \mathbf{y} be the
 248 sequence of all variables of $\mathcal{V}(\phi)$ in some arbitrary total order. By construction, the variable
 249 in \mathbf{y} are pairwise distinct. According to the normalized V-representation of the system, every
 250 point of the cone is a positive linear combination of the elementary modes $\exists \mathbf{x}. E\mathbf{x} \stackrel{\circ}{=} \mathbf{n}\mathbf{y}$, where
 251 \mathbf{n} contains the normalization factors. The linear system $E\mathbf{x} \stackrel{\circ}{=} \mathbf{n}\mathbf{y}$ is quasi-positive, since
 252 E and \mathbf{n} are positive. The variables in \mathbf{x} can be chosen freshly, and thus pairwise distinct
 253 with \mathbf{y} . The linear system $E\mathbf{x} \stackrel{\circ}{=} \mathbf{n}\mathbf{y}$ is strongly-triangular, since each variable of \mathbf{y} occurs in
 254 exactly one equation and never on the left. Therefore, we can define ϕ' as $E\mathbf{x} \stackrel{\circ}{=} \mathbf{n}\mathbf{y}$. The
 255 computation of the elementary modes and thus of E can be done in at most exponential
 256 time in the size of ϕ by Motzkin's double description method [16, 8, 9].

257 6 Abstraction Exactness

258 John's overapproximation Theorem 11 shows that the set of solutions over the abstract
 259 domain $\text{sol}^{\Delta}(\phi)$ is an approximation by the abstraction of the concrete solution set $h(\text{sol}^S(\phi))$
 260 for any abstraction $h : S \rightarrow \Delta$ from concrete to abstract structure and any positive first-order
 261 formula ϕ . We say ϕ is h -exact if even equality holds.

262 **► Definition 16 Exactness.** Let $h : S \rightarrow \Delta$ be a Σ -abstraction, ϕ a Σ -formula and $V \supseteq \mathcal{V}(\phi)$.
 263 We call ϕ h -exact with respect to V if $h(\text{sol}_V^S(\phi)) = \text{sol}_V^{\Delta}(\phi)$. We call ϕ h -exact if ϕ is
 264 h -exact with respect to $\mathcal{V}(\phi)$.

265 Our next objective is to study the preservation of h -exactness by logical operators. The
 266 main difficulty of this paper is the fact that h -exactness is not preserved by conjunction.
 267 Nevertheless, as we will show next, it is preserved by disjunction and existential quantification.
 268 For the case of disjunction, we need a basic property of union which fails for intersection.

269 **► Lemma 17.** Let V be a set of variables, R_1 and R_2 be subsets of assignments of type
 270 $V \rightarrow \text{dom}(S)$ and $h : S \rightarrow \Delta$ be a Σ -abstraction. $h \circ (R_1 \cup R_2) = h \circ R_1 \cup h \circ R_2$.

271 **► Proposition 18.** The disjunction of h -exact formulas is h -exact.

272 **► Lemma 19 Projection commutes with abstraction.** For any Σ -abstraction $h : S \rightarrow \Delta$,
 273 subset R of assignments of type $V \rightarrow S$, and variable $x \in \mathcal{V}$: $h \circ \pi_x(R) = \pi_x(h \circ R)$.

274 **► Proposition 20 Quantification preserves exactness.** For any surjective Σ -abstraction
 275 $h : S \rightarrow \Delta$ and formula $\exists x.\phi \in \mathcal{F}_{\Sigma}$, if ϕ is h -exact then $\exists x.\phi$ is h -exact.

XX:8 Abstracting Linear Equation Systems

276 We next study the h -exactness for strongly-triangular systems of Σ -equations, under the
277 condition that h is an abstraction between Σ -algebras with unique division.

278 ► **Lemma 21 Singleton property.** *If S is a Σ -algebra, $e \in \mathcal{E}_\Sigma$, and $\alpha : \mathcal{V}(e) \rightarrow S$ a variable
279 assignment, then the set $\llbracket e \rrbracket^{\alpha, S}$ is a singleton.*

280 A Σ -algebra is a Σ -structure with the singleton property. Let ele be the function that
281 maps any singleton to the element that it contains.

282 ► **Definition 22.** *We say that a Σ -structure S has unique division, if it satisfies the first-order
283 formula $\forall x. \exists \overset{\circ}{=}^1 y. ny \overset{\circ}{=} x$ for all nonzero natural number $n \in \mathbb{N}$.*

284 Clearly, the Σ -algebras \mathbb{R}_+ and \mathbb{B} have unique division. For any element s of the domain
285 of a structure S with unique division and any nonzero natural number $n \in \mathbb{N}$, we denote by
286 $\frac{s}{n}$ the unique element of $\{\alpha(y) \mid \alpha \in sol^S(ny \overset{\circ}{=} z), \alpha(z) = s\}$.

► **Lemma 23.** *Let $\phi \in \mathcal{F}_\Sigma$ be a Σ -formula and S a Σ -algebra with unique division. For
nonzero natural number n , variable $y \notin \mathcal{V}(\phi)$, and expression $e \in \mathcal{E}_\Sigma$ with $\mathcal{V}(e) \subseteq \mathcal{V}(\phi)$:*

$$sol^S(\phi \wedge ny \overset{\circ}{=} e) = \{\alpha[y / \frac{ele(\llbracket e \rrbracket^{\alpha, S})}{n}] \mid \alpha \in sol^S(\phi)\}$$

287 ► **Proposition 24.** *Let $\phi \in \mathcal{F}_\Sigma$ a Σ -formula, $n \neq 0$ a natural number, $e \in \mathcal{E}_\Sigma$ a Σ -expression
288 with $\mathcal{V}(e) \subseteq \mathcal{V}(\phi)$ and $y \notin \mathcal{V}(\phi)$ and the Σ -abstraction $h : S \rightarrow \Delta$ with S and Δ two Σ -algebras
289 with unique division. Then if ϕ is h -exact implies that $\phi \wedge e \overset{\circ}{=} ny$ is h -exact.*

Sketch of Proof. We can show that $h(ele(\llbracket e \rrbracket^{\alpha, S})) = ele(\llbracket e \rrbracket^{h \circ \alpha, \Delta})$ and $h(\frac{s}{n}) = \frac{h(s)}{n}$. Hence:

$$\begin{aligned} h \circ sol^S(\phi \wedge e \overset{\circ}{=} ny) &= h \circ \{\alpha[y / \frac{ele(\llbracket e \rrbracket^{\alpha, S})}{n}] \mid \alpha \in sol^S(\phi)\} && \text{by Lemma 23} \\ &= \{(h \circ \alpha)[y / h(\frac{ele(\llbracket e \rrbracket^{\alpha, S})}{n})] \mid \alpha \in sol^S(\phi)\} && \text{elementary} \\ &= \{\sigma[y / h(\frac{ele(\llbracket e \rrbracket^{\sigma, S})}{n})] \mid \sigma \in sol^\Delta(\phi)\} && h\text{-exactness of } \phi \\ &= \{\sigma[y / \frac{h(ele(\llbracket e \rrbracket^{\sigma, S}))}{n}] \mid \sigma \in sol^\Delta(\phi)\} \\ &= \{\sigma[y / \frac{ele(\llbracket e \rrbracket^{h \circ \sigma, \Delta})}{n}] \mid \sigma \in sol^\Delta(\phi)\} \\ &= sol^\Delta(\phi \wedge e \overset{\circ}{=} ny) && \text{by Lemma 23} \end{aligned}$$

290 ► **Proposition 25.** *Let $h : S \rightarrow \Delta$ be a Σ -abstraction between Σ -algebras with unique division.
291 Then any strongly-triangular system of Σ -equations with natural coefficients is h -exact.*

292 **Sketch of Proof** By induction on the number of equations and Proposition 24.

293 We notice that Proposition 25 remains true for triangular systems that are not strongly-
294 triangular. This will follow with further results from the next section (Theorem 37 and
295 Proposition 31) requiring a different argument.

296 ► **Theorem 26 Exactness.** *Quasi-positive strongly-triangular polynomial systems are $h_{\mathbb{B}}$ -exact.*

297 **Proof** The Σ -algebras \mathbb{R}_+ and \mathbb{B} have unique division, so we can apply Proposition 25.

298 The Elementary Modes Theorem 15 shows that any integer matrix equation $A\mathbf{x} \overset{\circ}{=} 0$ is
299 \mathbb{R}_+ -equivalent to some quasi-positive strongly-triangular linear equation system. We can
300 thus apply Theorem 26 to obtain the following corollary.

301 ► **Corollary 27.** *Any matrix integer equation can be converted in at most exponential time
302 to some \mathbb{R}_+ -equivalent $h_{\mathbb{B}}$ -exact Σ -formula.*

303 This corollary permits us to compute the $h_{\mathbb{B}}$ -abstraction of an integer matrix equation by
304 computing the \mathbb{B} -solutions of the \mathbb{R}_+ -equivalent $h_{\mathbb{B}}$ -exact formula. For computing abstractions
305 between structures without unique division we need to strengthen this result.

7 Abstraction Invariance

The essential problem approached by the paper is that conjunctions of two h -exact formulas may not be h -exact. The situation changes when assuming the following notion of h -invariance for at least one of the two formulas.

► **Definition 28 Invariance.** *Let $h : S \rightarrow \Delta$ be a Σ -abstraction and $V \subseteq \mathcal{V}$ a subset of variables. We call a subset R of variable assignments of type $V \rightarrow \text{dom}(S)$ h -invariant iff:*

$$\forall \alpha, \alpha' : V \rightarrow \text{dom}(S). (\alpha \in R \wedge h \circ \alpha = h \circ \alpha' \implies \alpha' \in R).$$

We call a Σ -formula ϕ h -invariant if its solution set $\text{sol}^S(\phi)$ is.

The relevance of the notion of invariance for exactness of conjunctions – that we will formalize in Proposition 31 – is due to the the following lemma:

► **Lemma 29.** *If either R_1 or R_2 are h -invariant then: $h \circ (R_1 \cap R_2) = h \circ R_1 \cap h \circ R_2$.*

Sketch of Proof. The one inclusion is straightforward without invariance. For the other inclusion, we can assume with loss of generality that R_1 is h -invariant. So let $\beta \in h \circ R_1 \cap h \circ R_2$. Then there exist $\alpha_1 \in R_1$ and $\alpha_2 \in R_2$ such that $\beta = h \circ \alpha_1 = h \circ \alpha_2$. By h -invariance of R_1 it follows that $\alpha_1 \in R_2$. So $\alpha_1 \in R_1 \cap R_2$, and hence, $\beta \in h \circ (R_1 \cap R_2)$.

We continue with an algebraic characterization of h -invariance. Given a Σ -abstraction $h : S \rightarrow \Delta$, and a set R of variable assignments to $\text{dom}(\Delta)$, we define the left-decomposition of R with respect to h as the following set of variable assignments to $\text{dom}(S)$:

$$h \circ R \quad =_{\text{def}} \{ \alpha \mid h \circ \alpha \in R \}$$

Clearly, $R \subseteq h \circ (h \circ R)$. The inverse inclusion characterizes the h -invariance of R .

► **Lemma 30 Algebraic characterization.** *A subset of R variables assignment of type $V \rightarrow \text{dom}(S)$ is h -invariant for an Σ -abstraction $h : S \rightarrow \Delta$ iff $h \circ (h \circ R) \subseteq R$.*

► **Proposition 31 Exactness is preserved by conjunction when assuming invariance.** *Let h be a surjective Σ -abstraction. If ϕ_1 and ϕ_2 are h -exact Σ -formulas and ϕ_1 or ϕ_2 are h -invariant then the conjunction $\phi_1 \wedge \phi_2$ is h -exact.*

Proof Let ϕ_1 and ϕ_2 be h -exact Σ -formulas. We assume without loss of generality that ϕ_1 is h -invariant. Let $V = \mathcal{V}(\phi_1 \wedge \phi_2)$. Since $\mathcal{V}(\phi_2) \subseteq V$ the set $\text{sol}_V^S(\phi_2)$ is h -invariant too by Lemma 54. We can now show that $\phi_1 \wedge \phi_2$ is h -exact as follows:

$$\begin{aligned} h \circ \text{sol}^S(\phi_1 \wedge \phi_2) &= h \circ (\text{sol}_V^S(\phi_1) \cap \text{sol}_V^S(\phi_2)) \\ &= h \circ \text{sol}_V^S(\phi_1) \cap h \circ \text{sol}_V^S(\phi_2) \quad \text{by Lemma 29} \\ &= \text{sol}_V^\Delta(\phi_1) \cap \text{sol}_V^\Delta(\phi_2) \quad \text{by } h\text{-exactness of } \phi_1 \text{ and } \phi_2 \text{ wrt } V \\ &= \text{sol}^\Delta(\phi_1 \wedge \phi_2) \end{aligned}$$

Our next objective is to show that h -invariant formulas are closed under conjunction, disjunction, and existential quantification. The two former closure properties rely on the following two algebraic properties of abstraction decomposition.

► **Lemma 32.** *For any Σ -abstraction $h : S \rightarrow \Delta$, any subsets of assignments of type $V \rightarrow \text{dom}(S)$ R_1 and R_2 and V a subset of variables:*

$$\blacksquare \quad h \circ (R_1 \cap R_2) = h \circ R_1 \cap h \circ R_2.$$

XX:10 Abstracting Linear Equation Systems

331 ■ $h \ominus (R_1 \cup R_2) = h \ominus R_1 \cup h \ominus R_2.$

332 ► **Lemma 33 Intersection and union preserve invariance.** *Let $h : S \rightarrow \Delta$ be a Σ -abstraction.*
 333 *Then the intersection and union of any two h -invariant subsets R_1 and R_2 of variables*
 334 *assignments of type $V \rightarrow \text{dom}(S)$ is h -invariant.*

335 ► **Lemma 34 Projection commutes with left-decomposition.** $h \ominus \pi_x(R) = \pi_x(h \ominus R).$

336 ► **Proposition 35 Invariance is preserved by conjunction, disjunction, and quantification.**
 337 *If h is a surjective abstraction then the class of h -invariant FO-formulas is closed under*
 338 *conjunction, disjunction, and existential quantification.*

339 We do not know whether negation preserves h -invariance in general, but for finite Δ it
 340 can be shown that if ϕ is h -exact and h -invariant, then $\neg\phi$ is h -exact and h -invariant too.

341 ► **Proposition 36.** *Let h be a surjective Σ -abstractions. Then the class of h -exact and h -*
 342 *invariant Σ -formulas is closed under conjunction, disjunction and existential quantification.*

343 ► **Theorem 37 Invariance.** *Any positive polynomial equation $p \stackrel{\circ}{=} 0$ such that p has no constant*
 344 *term is $h_{\mathbb{B}}$ -exact and $h_{\mathbb{B}}$ -invariant.*

Sketch of Proof. Any positive polynomial equation $p \stackrel{\circ}{=} 0$ such that p has no constant
 term and only positive coefficients has the form $\sum_{j=1}^l n_j \prod_{k=1}^{i_j} x_{j,k}^{m_{j,k}} \stackrel{\circ}{=} 0$ where $l \geq 0$, and
 $n_j, i_j, m_{j,k} > 0$. We can now show that for both algebras $S \in \{\mathbb{B}, \mathbb{R}_+\}$ that:

$$\text{sol}^S(p \stackrel{\circ}{=} 0) = \text{sol}^S\left(\bigwedge_{j=1}^l \bigvee_{k=1}^{i_j} x_{j,k} \stackrel{\circ}{=} 0\right)$$

345 Since the formulas $x_{j,k} \stackrel{\circ}{=} 0$ are $h_{\mathbb{B}}$ -exact and $h_{\mathbb{B}}$ -invariant, the theorem follows from the closure
 346 properties of Proposition 36.

8 Boolean Abstractions of $h_{\mathbb{B}}$ -Mixed Systems

348 In this section we prove our main result stating how to compute the $h_{\mathbb{B}}$ -abstraction of the
 349 \mathbb{R}_+ -solution set of a $h_{\mathbb{B}}$ -mixed systems exactly.

350 ► **Definition 38.** *A $h_{\mathbb{B}}$ -mixed system is a formula in \mathcal{F}_{Σ} of the form $\exists \mathbf{z}. \phi \wedge \phi'$ where ϕ is a*
 351 *linear equation system and ϕ' a $h_{\mathbb{B}}$ -invariant and $h_{\mathbb{B}}$ -exact first-order formula.*

352 Note that linear equation systems $A\mathbf{y} \stackrel{\circ}{=} 0$, with A an integer matrix and \mathbf{y} a sequence of
 353 pairwise distinct variables, need not to be $h_{\mathbb{B}}$ -exact, if A is not positive. However, any linear
 354 equation systems of this shape is \mathbb{R}_+ -equivalent to some quasi-positive strongly-triangular
 355 polynomial system, as shown by the Elementary Modes Theorem 15. And quasi-positive
 356 strongly-triangular polynomial system were shown to be $h_{\mathbb{B}}$ -exact by Exactness Theorem 26.

357 ► **Theorem 39 Main.** *Any $h_{\mathbb{B}}$ -mixed system can be converted in exponential time to an*
 358 *\mathbb{R}_+ -equivalent Σ -formula that is $h_{\mathbb{B}}$ -exact.*

359 **Proof** Consider a $h_{\mathbb{B}}$ -mixed system $\exists \mathbf{x}. (\phi \wedge \phi')$ where ϕ is a linear equation system and
 360 ϕ' a first-order formula that is $h_{\mathbb{B}}$ -exact and $h_{\mathbb{B}}$ -invariant. Based on the Elementary Modes
 361 Theorem 15, the linear equation system ϕ can be transformed in exponential time to the
 362 form $\exists \mathbf{z}. \phi''$ where ϕ'' is a quasi-positive strongly-triangular system of linear equations. Such
 363 polynomial equation systems are $h_{\mathbb{B}}$ -exact by Theorem 26, and so is ϕ'' . The Invariance
 364 Proposition 31 shows that the conjunction $\phi'' \wedge \phi'$ is $h_{\mathbb{B}}$ -exact too, since ϕ' was assumed to
 365 be $h_{\mathbb{B}}$ -exact and $h_{\mathbb{B}}$ -invariant. Finally, $h_{\mathbb{B}}$ -exactness is preserved by existential quantification
 366 by Proposition 20, so the formula $\exists \mathbf{x}. \exists \mathbf{z}. \phi'' \wedge \phi'$ is $h_{\mathbb{B}}$ -exact too.

367 ► **Corollary 40.** *The $h_{\mathbb{B}}$ -abstraction of the \mathbb{R}_+ -solution set of an $h_{\mathbb{B}}$ -mixed system ϕ , that is*
 368 *$h_{\mathbb{B}} \circ \text{sol}^{\mathbb{R}_+}(\phi)$, can be computed in at most exponential time in the size of the system ϕ .*

369 The algorithm from the proof Corollary 40 can be improved so that it becomes sufficiently
 370 efficient for practical use. For this the two steps with exponential worst case complexity must
 371 be made polynomial for the particular instances. First note that the computation of the
 372 elementary modes (Theorem 15) is known to be computationally feasible. Various algorithms
 373 for this purpose were implemented [7, 12, 2, 3] and applied successfully to problems in
 374 systems biology [9]. The second exponential step concerns the enumeration of all boolean
 375 variable assignments. This enumeration may be avoided by using constraint programming
 376 techniques for computing the solution set $\text{sol}^{\mathbb{B}}(\phi'')$. For those $h_{\mathbb{B}}$ -mixed systems for which
 377 both steps can be done in polynomial time, we can compute the boolean abstraction of
 378 the \mathbb{R}_+ -solution set in polynomial time too. The practical feasibility of this approach was
 379 demonstrated recently at an application to knockout prediction in systems biology [1], where
 380 previously only over-approximations could be computed.

381 9 Computing Sign Abstractions

382 We next show how to compute the sign abstraction $h_{\mathbb{S}} \circ \text{sol}^{\mathbb{R}}(\phi)$ of the \mathbb{R} -solutions set of a
 383 linear equation system ϕ . For this, we convert ϕ into a first-order formula ϕ' based on our
 384 main Theorem 39 such that $h_{\mathbb{S}} \circ \text{sol}^{\mathbb{R}}(\phi)$ can be computed from $\text{sol}^{\mathbb{B}}(\phi')$ in polynomial time.
 385 In order to do so, we relate in a first step the sign abstraction to the boolean abstraction,
 386 then show in a second step that this relationship can be defined in first-order logic, so that
 387 our Main Theorem for the boolean abstraction can be applied.

In the first step, we relate the sign abstraction to the boolean abstraction. For doing so,
 we decompose any real number r into two positive numbers, its negative part $\ominus(r)$ and its
 positive part $\oplus(r)$, such that if $r \geq 0$ then $\ominus(r) = 0$ and $\oplus(r) = r$ and otherwise $\ominus(r) = -r$
 and $\oplus(r) = 0$. The decomposition function $\text{dec} : \mathbb{R} \rightarrow (\{0\} \times \mathbb{R}_+) \cup (\mathbb{R}_+ \times \{0\})$ is defined as
 follows for $r \in \mathbb{R}$:

$$\text{dec}(r) = (\ominus(r), \oplus(r))$$

388 This function is a bijection, so it has an inverse function $\text{dec}^{-1} : (\{0\} \times \mathbb{R}_+) \cup (\mathbb{R}_+ \times \{0\}) \rightarrow \mathbb{R}$,
 389 which satisfies $\text{dec}^{-1}((r_1, r_2)) = r_2 - r_1$ for all pairs (r_1, r_2) in its domain.

390 ► **Lemma 41 Decomposition.** $h_{\mathbb{S}} = \text{dec}^{-1} \circ h_{\mathbb{B}}^2 \circ \text{dec}$

391 **Proof** If r is negative then $\text{dec}^{-1}(h_{\mathbb{B}}^2(\text{dec}(r))) = \text{dec}^{-1}(h_{\mathbb{B}}^2((-r, 0))) = \text{dec}^{-1}((h_{\mathbb{B}}(-r), 0)) =$
 392 $-h_{\mathbb{B}}(-r) = h_{\mathbb{S}}(r)$. Otherwise if r is positive then $\text{dec}^{-1}(h_{\mathbb{B}}^2(\text{dec}(r))) = \text{dec}^{-1}(h_{\mathbb{B}}^2((0, r))) =$
 393 $\text{dec}^{-1}((0, h_{\mathbb{B}}(r))) = h_{\mathbb{B}}(r) = h_{\mathbb{S}}(r)$.

We will show in a first step that first-order formulas over the reals can be rewritten, such
 that interpretation over the positive reals is enough. We start with a definition of positivity
 of reals in first-order logic. For any variable $x \in \mathcal{V}$ we define the formulas $\text{pos}(x) \in \mathcal{F}_{\Sigma}$ by:

$$\text{pos}(x) =_{\text{def}} \exists z. x \stackrel{\circ}{=} z * z$$

394 Clearly, if $\alpha \in \text{sol}^{\mathbb{R}}(\text{pos}(x))$ then $\alpha(x) \in \mathbb{R}_+$. We can use this formula to relate \mathbb{R}_+ -solutions
 395 to \mathbb{R} -solutions of particular formulas.

396 ► **Definition 42.** *A formula $\phi \in \mathcal{F}_{\Sigma}$ is called domain-positive if ϕ has the form $\phi' \wedge$
 397 $\bigwedge_{y \in \mathcal{V}(\phi')} \text{pos}(y)$ as well as all formulas ϕ'' for which $\exists x. \phi''$ is a subformula of ϕ .*

398 ► **Lemma 43.** *All domain-positive formulas $\phi \in \mathcal{F}_{\Sigma}$ satisfy $\text{sol}^{\mathbb{R}_+}(\phi) = \text{sol}^{\mathbb{R}}(\phi)$.*

XX:12 Abstracting Linear Equation Systems

399 **Proof** By induction on the structure of formulas. The induction step is straightforward
400 from due to the fact that $pos(y)$ imposes the positivity of the value y .

401 We next show how to make first-order formulas domain-positive based on the decompo-
402 sition of real numbers into two positive numbers presented earlier. We fix two generators
403 of fresh variable $\nu_{\ominus}, \nu_{\oplus} : \mathcal{V} \rightarrow \mathcal{V}$. For any $x \in \mathcal{V}$, the intention is that $\nu_{\oplus}(x)$ stands
404 for the positive part of x and $\nu_{\ominus}(x)$ for its negative part. We will preserve the invariant
405 that $x = \nu_{\oplus}(x) - \nu_{\ominus}(x)$ and $\nu_{\oplus}(x) * \nu_{\ominus}(x) = 0$. Furthermore, let $\nu : \mathcal{V} \rightarrow \mathcal{V}^2$ such that
406 $\nu(x) = (\nu_{\oplus}(x), \nu_{\ominus}(x))$ for all $x \in \mathcal{V}$.

► **Proposition 44 Positivity.** *For any formula $\phi \in \mathcal{F}_{\Sigma}$ we can compute in linear time a formula $dec_{\nu}(\phi) \in \mathcal{F}_{\Sigma}$ such that:*

$$dec \circ sol^{\mathbb{R}}(\phi) = \{\alpha^2 \circ \nu|_{\mathcal{V}(\phi)} \mid \alpha \in sol^{\mathbb{R}+}(dec_{\nu}(\phi) \wedge \bigwedge_{x \in \mathcal{V}(\phi)} \nu_{\oplus}(x) * \nu_{\ominus}(x) \stackrel{\circ}{=} 0)\}$$

407 *Furthermore, if ϕ was a linear equation system, then $dec_{\nu}(\phi)$ is a $h_{\mathbb{B}}$ -mixed system.*

Proof We can assume without loss of generality that all equations of ϕ are flat, i.e., of the form $x \stackrel{\circ}{=} x_1 + x_2$, $x \stackrel{\circ}{=} x_1 * x_2$, $x \stackrel{\circ}{=} 0$, or $x \stackrel{\circ}{=} 1$. We define the formulas $dec_{\nu}(\phi), dec'_{\nu}(\phi) \in \mathcal{F}_{\Sigma}$ for all formulas ϕ in \mathcal{F}_{Σ} with flat equations recursively as follows:

$$\begin{aligned} dec_{\nu}(\exists x.\phi) &= \exists \nu_{\ominus}(x). \exists \nu_{\oplus}(x). dec_{\nu}(\phi) & dec_{\nu}(\phi \wedge \phi') &= dec_{\nu}(\phi) \wedge dec_{\nu}(\phi') \\ &\wedge \nu_{\oplus}(x) * \nu_{\ominus}(x) \stackrel{\circ}{=} 0 & dec_{\nu}(\neg\phi) &= \neg dec_{\nu}(\phi) \\ &\wedge pos(\nu_{\oplus}(x)) \wedge pos(\nu_{\ominus}(x)) \\ dec_{\nu}(x \stackrel{\circ}{=} x_1 + x_2) &= & dec_{\nu}(x \stackrel{\circ}{=} x_1 * x_2) &= \\ \nu_{\oplus}(x) + \nu_{\ominus}(x_1) + \nu_{\ominus}(x_2) \stackrel{\circ}{=} & & \nu_{\oplus}(x) + \nu_{\oplus}(x_1) * \nu_{\ominus}(x_2) + \nu_{\ominus}(x_1) * \nu_{\oplus}(x_2) \stackrel{\circ}{=} & \\ \nu_{\ominus}(x) + \nu_{\oplus}(x_1) + \nu_{\oplus}(x_2) & & \nu_{\ominus}(x) + \nu_{\oplus}(x_1) * \nu_{\oplus}(x_2) + \nu_{\ominus}(x_1) * \nu_{\ominus}(x_2) & \\ dec_{\nu}(x \stackrel{\circ}{=} 0) &= \nu_{\oplus}(x) \stackrel{\circ}{=} \nu_{\ominus}(x) & dec_{\nu}(x \stackrel{\circ}{=} 1) &= \nu_{\oplus}(x) \stackrel{\circ}{=} \nu_{\ominus}(x) + 1 \end{aligned}$$

The rewriting for equations with multiplication relies on the distributivity law. We also use inverses for addition in the structure of the reals. Let

$$dec'_{\nu}(\phi) = dec_{\nu}(\phi) \wedge \bigwedge_{x \in \mathcal{V}(\phi)} \nu_{\oplus}(x) * \nu_{\ominus}(x) \stackrel{\circ}{=} 0 \wedge pos(\nu_{\ominus}(x)) \wedge pos(\nu_{\oplus}(x)).$$

408 We can show by induction on the structure of formulas that all formulas $dec'_{\nu}(\phi)$ are domain-
409 positive and satisfy $dec \circ sol^{\mathbb{R}}(\phi) = \{\alpha^2 \circ \nu|_{\mathcal{V}(\phi)} \mid \alpha \in sol^{\mathbb{R}}(dec'_{\nu}(\phi))\}$. The proposition follows
410 with \mathbb{R}_+ instead of \mathbb{R} from Lemma 43 and the domain-positivity of $dec'_{\nu}(\phi)$.

► **Theorem 45 Computing Sign Abstractions.** *For any linear equation system $\phi \in \mathcal{F}_{\Sigma}$ we can compute in at most exponential time a formula $\phi' \in \mathcal{F}_{\Sigma}$ such that:*

$$h_{\mathbb{S}} \circ sol^{\mathbb{R}}(\phi) = \{[y/\beta(\nu_{\oplus}(y)) - \beta(\nu_{\ominus}(y))] \mid y \in \mathcal{V}(\phi') \mid \beta \in sol^{\mathbb{B}}(\phi')\}$$

Proof Let $\phi \in \mathcal{F}_{\Sigma}$ be a system of linear equations. The formula $dec_{\nu}(\phi) \wedge \bigwedge_{x \in \mathcal{V}(\phi)} \nu_{\oplus}(x) * \nu_{\ominus}(x) \stackrel{\circ}{=} 0$ is a $h_{\mathbb{B}}$ -mixed system by Positivity Proposition 44, so that we can apply the Main Theorem 39. Therefore, we can compute in at most exponential time an \mathbb{R}_+ -equivalent formula ϕ' that is $h_{\mathbb{B}}$ -exact.

$$\begin{aligned} &h_{\mathbb{S}} \circ sol^{\mathbb{R}}(\phi) \\ &= dec^{-1} \circ h_{\mathbb{B}}^2 \circ dec \circ sol^{\mathbb{R}}(\phi) && \text{by Decomposition Lemma 41} \\ &= dec^{-1} \circ h_{\mathbb{B}}^2 \circ \{\alpha^2 \circ \nu|_{\mathcal{V}(\phi)} \mid \alpha \in sol^{\mathbb{R}+}(dec_{\nu}(\phi) \wedge \bigwedge_{x \in \mathcal{V}(\phi)} \nu_{\oplus}(x) * \nu_{\ominus}(x) \stackrel{\circ}{=} 0)\} && \text{by Positivity Proposition 44} \\ &= dec^{-1} \circ h_{\mathbb{B}}^2 \circ \{\alpha^2 \circ \nu|_{\mathcal{V}(\phi')} \mid \alpha \in sol^{\mathbb{R}+}(\phi')\} && \text{where } \phi' \text{ from Main Theorem 39} \\ &= \{dec^{-1} \circ \beta^2 \circ \nu|_{\mathcal{V}(\phi')} \mid \beta \in sol^{\mathbb{B}}(\phi')\} && \text{by } h_{\mathbb{B}}\text{-exactness of } \phi' \\ &= \{[y/\beta(\nu_{\oplus}(y)) - \beta(\nu_{\ominus}(y))] \mid y \in \mathcal{V}(\phi') \mid \beta \in sol^{\mathbb{B}}(\phi')\} && \text{by definition of } dec^{-1} \end{aligned}$$

411

412 ——— **References** ———

- 413 **1** Emilie Allart, Joachim Niehren, and Cristian Versari. Computing difference abstractions of
414 metabolic networks under kinetic constraints. In Luca Bortolussi and Guido Sanguinetti,
415 editors, *Computational Methods in Systems Biology - 17th International Conference, CMSB*
416 *2019, Trieste, Italy, September 18-20, 2019, Proceedings*, volume 11773 of *Lecture Notes*
417 *in Computer Science*, pages 266–285. Springer, 2019. URL: [https://doi.org/10.1007/](https://doi.org/10.1007/978-3-030-31304-3_14)
418 [978-3-030-31304-3_14](https://doi.org/10.1007/978-3-030-31304-3_14), doi:10.1007/978-3-030-31304-3_14.
- 419 **2** David Avis and Charles Jordan. mpls: A scalable parallel vertex/facet enumeration code.
420 *Mathematical Programming Computation*, 10(2):267–302, 2018.
- 421 **3** Roberto Bagnara, Patricia M. Hill, and Enea Zaffanella. The parma polyhedra library: Toward
422 a complete set of numerical abstractions for the analysis and verification of hardware and
423 software systems. *Sci. Comput. Program.*, 72(1-2):3–21, 2008. URL: [https://doi.org/10.](https://doi.org/10.1016/j.scico.2007.08.001)
424 [1016/j.scico.2007.08.001](https://doi.org/10.1016/j.scico.2007.08.001), doi:10.1016/j.scico.2007.08.001.
- 425 **4** Patrick Cousot and Nicolas Halbwachs. Automatic discovery of linear restraints among
426 variables of a program. In Alfred V. Aho, Stephen N. Zilles, and Thomas G. Szymanski,
427 editors, *Conference Record of the Fifth Annual ACM Symposium on Principles of Programming*
428 *Languages, Tucson, Arizona, USA, January 1978*, pages 84–96. ACM Press, 1978. URL:
429 <https://doi.org/10.1145/512760.512770>, doi:10.1145/512760.512770.
- 430 **5** François Coutte, Joachim Niehren, Debarun Dhali, Mathias John, Cristian Versari, and
431 Philippe Jacques. Modeling Leucine’s Metabolic Pathway and Knockout Prediction Improving
432 the Production of Surfactin, a Biosurfactant from *Bacillus Subtilis*. *Biotechnology Journal*,
433 10(8):1216–34, August 2015. URL: <https://hal.inria.fr/hal-01153704>, doi:10.1002/
434 [biot.201400541](https://hal.inria.fr/hal-01153704).
- 435 **6** Lloyd L. Dines. On positive solutions of a system of linear equations. *Annals of Mathematics*,
436 28(1/4):386–392, 1926. URL: <http://www.jstor.org/stable/1968384>.
- 437 **7** K Fukuda. cdd. c: C-implementation of the double description method for computing all
438 vertices and extremal rays of a convex polyhedron given by a system of linear inequalities.
439 *Department of Mathematics, Swiss Federal Institute of Technology, Lausanne, Switzerland*,
440 1993.
- 441 **8** Komei Fukuda and Alain Prodon. Double description method revisited. In Michel Deza,
442 Reinhardt Euler, and Ioannis Manoussakis, editors, *Combinatorics and Computer Science*,
443 pages 91–111, Berlin, Heidelberg, 1996. Springer Berlin Heidelberg.
- 444 **9** Julien Gagneur and Steffen Klamt. Computation of elementary modes: a unifying framework
445 and the new binary approach. *BMC bioinformatics*, 5(1):1, 2004.
- 446 **10** Philippe Granger. Static analysis of linear congruence equalities among variables of a program.
447 In Samson Abramsky and T. S. E. Maibaum, editors, *TAPSOFT’91: Proceedings of the*
448 *International Joint Conference on Theory and Practice of Software Development, Brighton,*
449 *UK, April 8-12, 1991, Volume 1: Colloquium on Trees in Algebra and Programming (CAAP’91)*,
450 volume 493 of *Lecture Notes in Computer Science*, pages 169–192. Springer, 1991. URL:
451 https://doi.org/10.1007/3-540-53982-4_10, doi:10.1007/3-540-53982-4_10.
- 452 **11** Mathias John, Mirabelle Nebut, and Joachim Niehren. Knockout Prediction for Reaction
453 Networks with Partial Kinetic Information. In *14th International Conference on Verification,*
454 *Model Checking, and Abstract Interpretation*, pages 355–374, Rom, Italy, January 2013. URL:
455 <http://hal.inria.fr/hal-00692499>.
- 456 **12** Steffen Klamt, Jörg Stelling, Martin Ginkel, and Ernst Dieter Gilles. FluxAnalyzer:
457 exploring structure, pathways, and flux distributions in metabolic networks on inter-
458 active flux maps. *Bioinformatics*, 19(2):261–269, 01 2003. URL: [https://doi.org/](https://doi.org/10.1093/bioinformatics/19.2.261)
459 [10.1093/bioinformatics/19.2.261](https://doi.org/10.1093/bioinformatics/19.2.261), arXiv:[https://academic.oup.com/bioinformatics/](https://academic.oup.com/bioinformatics/article-pdf/19/2/261/1059937/190261.pdf)
460 [article-pdf/19/2/261/1059937/190261.pdf](https://academic.oup.com/bioinformatics/article-pdf/19/2/261/1059937/190261.pdf), doi:10.1093/bioinformatics/19.2.261.

- 461 **13** K. Lotz, A. Hartmann, E. Grafahrend-Belau, and B.H. Schreiber, F.and Junker. Elementary
462 flux modes, flux balance analysis, and their application to plant metabolism. *Plant Metabolism.*
463 *Methods in Molecular Biology (Methods and Protocols)*, 2014.
- 464 **14** Costas D. Maranas and Ali R. Zomorodi. *Flux Balance Analysis and LP Problems*, chap-
465 ter 3, pages 53–80. Wiley-Blackwell, 2016. URL: [https://onlinelibrary.wiley.com/doi/](https://onlinelibrary.wiley.com/doi/abs/10.1002/9781119188902.ch3)
466 [abs/10.1002/9781119188902.ch3](https://onlinelibrary.wiley.com/doi/abs/10.1002/9781119188902.ch3), arXiv:[https://onlinelibrary.wiley.com/doi/pdf/10.](https://onlinelibrary.wiley.com/doi/pdf/10.1002/9781119188902.ch3)
467 [1002/9781119188902.ch3](https://onlinelibrary.wiley.com/doi/pdf/10.1002/9781119188902.ch3), doi:10.1002/9781119188902.ch3.
- 468 **15** Antoine Miné. A few graph-based relational numerical abstract domains. In Manuel V.
469 Hermenegildo and Germán Puebla, editors, *Static Analysis, 9th International Symposium,*
470 *SAS 2002, Madrid, Spain, September 17-20, 2002, Proceedings*, volume 2477 of *Lecture*
471 *Notes in Computer Science*, pages 117–132. Springer, 2002. URL: [https://doi.org/10.1007/](https://doi.org/10.1007/3-540-45789-5_11)
472 [3-540-45789-5_11](https://doi.org/10.1007/3-540-45789-5_11), doi:10.1007/3-540-45789-5_11.
- 473 **16** T.S. Motzkin, H. Raiffa, GL. Thompson, and R.M. Thrall. The double description method. In
474 H.W. Kuhn and A.W.Tucker, editors, *Contributions to theory of games*, volume 2. Princeton
475 University Press, 1953.
- 476 **17** Joachim Niehren, Cristian Versari, Mathias John, François Coutte, and Philippe Jacques.
477 Predicting Changes of Reaction Networks with Partial Kinetic Information. *BioSystems*,
478 149:113–124, July 2016. URL: <https://hal.inria.fr/hal-01239198>.
- 479 **18** Andrea Rendl, Tias Guns, Peter J. Stuckey, and Guido Tack. Minisearch: A solver-independent
480 meta-search language for minizinc. In Gilles Pesant, editor, *Principles and Practice of*
481 *Constraint Programming - 21st International Conference, CP 2015, Cork, Ireland, August*
482 *31 - September 4, 2015, Proceedings*, volume 9255 of *Lecture Notes in Computer Science*,
483 pages 376–392. Springer, 2015. URL: https://doi.org/10.1007/978-3-319-23219-5_27,
484 doi:10.1007/978-3-319-23219-5_27.
- 485 **19** Dr.Jürgen Zanghellini, David E. Ruckerbauer, Michael Hanscho, and Christian Jungreuthmayer.
486 Elementary flux modes in a nutshell: Properties, calculation and applications. *Biotechnology*
487 *Journal*, pages 1009–1016, 2013.

Interpretation of expressions as sets of elements $\llbracket e \rrbracket^{\alpha, S} \subseteq \text{dom}(S)$, where S is a Σ -structure and $\alpha : V \rightarrow \text{dom}(S)$ where V contains all free variables.

$$\llbracket c \rrbracket^{\alpha, S} = c^S \quad \llbracket x \rrbracket^{\alpha, S} = \{\alpha(x)\} \quad \llbracket e \odot e' \rrbracket^{\alpha, S} = \cup \{s \odot^S s' \mid s \in \llbracket e \rrbracket^{\alpha, S}, s' \in \llbracket e' \rrbracket^{\alpha, S}\}$$

Interpretation of formulas as truth values $\llbracket \phi \rrbracket^{\alpha, S} \in \mathbb{B}$:

$$\begin{aligned} \llbracket e \overset{\circ}{=} e' \rrbracket^{\alpha, S} &= \begin{cases} 1 & \text{if } \llbracket e \rrbracket^{\alpha, S} \cap \llbracket e' \rrbracket^{\alpha, S} \neq \emptyset \\ 0 & \text{else} \end{cases} & \llbracket \phi \wedge \phi' \rrbracket^{\alpha, S} &= \llbracket \phi \rrbracket^{\alpha, S} \wedge^{\mathbb{B}} \llbracket \phi' \rrbracket^{\alpha, S} \\ \llbracket \neg \phi \rrbracket^{\alpha, S} &= \neg^{\mathbb{B}}(\llbracket \phi \rrbracket^{\alpha, S}) & \llbracket \exists x. \phi \rrbracket^{\alpha, S} &= \begin{cases} 1 & \text{if exists } s \in \text{dom}(S). \\ & \llbracket \phi \rrbracket^{\alpha[x/s], S} = 1 \\ 0 & \text{else} \end{cases} \end{aligned}$$

■ **Figure 4** Semantics of Σ -expressions and formulas over a Σ -structure S with respect to a variable assignment $\alpha : V \rightarrow \text{dom}(S)$.

A Proofs for Section 3 (The Boolean and the Sign Abstraction)

► **Lemma 7.** *The function $h_{\mathbb{B}} : \mathbb{R}_+ \rightarrow \mathbb{B}$ is a Σ -abstraction between Σ -algebras.*

Proof. For all $r, r' \in \mathbb{R}_+$ we have:

$$\begin{aligned} h_{\mathbb{B}}(r +^{\mathbb{R}_+} r') = 1 &\Leftrightarrow r +^{\mathbb{R}_+} r' \neq 0 \Leftrightarrow r \neq 0 \vee r' \neq 0 \Leftrightarrow h_{\mathbb{B}}(r) = 1 \vee h_{\mathbb{B}}(r') = 1 \\ h_{\mathbb{B}}(r *^{\mathbb{R}_+} r') = 1 &\Leftrightarrow r *^{\mathbb{R}_+} r' \neq 0 \Leftrightarrow r \neq 0 \wedge r' \neq 0 \Leftrightarrow h_{\mathbb{B}}(r) = 1 \wedge h_{\mathbb{B}}(r') = 1 \end{aligned}$$

Hence $h_{\mathbb{B}}(r +^{\mathbb{R}_+} r') = h_{\mathbb{B}}(r) +^{\mathbb{B}} h_{\mathbb{B}}(r')$ and $h_{\mathbb{B}}(r *^{\mathbb{R}_+} r') = h_{\mathbb{B}}(r) *^{\mathbb{B}} h_{\mathbb{B}}(r')$. Finally, for both constants $c \in C$ we have that $h_{\mathbb{B}}(c^{\mathbb{R}_+}) = h_{\mathbb{B}}(c) = c = c^{\mathbb{B}}$. ◀

► **Lemma 9.** *$h_{\mathbb{S}} : \mathbb{R} \rightarrow \mathbb{S}$ is a Σ -abstraction into a Σ -structure (that is not a Σ -algebra).*

Proof. For any $r, r' \in \mathbb{R}$ the second condition for homomorphism follows for all $\odot \in F^{(2)}$: $h_{\mathbb{S}}(r \odot r') \in h_{\mathbb{S}}(r) \odot^{\mathbb{S}} h_{\mathbb{S}}(r')$. And for all constants $c \in C$ we have by definition that $h_{\mathbb{S}}(c^{\mathbb{R}}) = c^{\mathbb{S}}$. ◀

B Proofs for Section 4 (Abstractions and First-Order Logic)

The following two lemmas are classical. Let R be a subset of assignments of type $V' \rightarrow \text{dom}(S)$ and $V \cap V' = \emptyset$ two subsets of \mathcal{V} . We define: $\text{ext}_V^S(R) = \{\alpha \cup \alpha' \mid \alpha' : V' \cup V \rightarrow \text{dom}(S), \alpha \in R\}$.

► **Lemma 46 Cylindrification.** *If $V \cap \mathcal{V}(\phi) = \emptyset$ then: $\text{sol}_{V \cup \mathcal{V}(\phi)}^S(\phi) = \text{ext}_V^S(\text{sol}^S(\phi))$.*

Proof. We can show for all expressions $e \in \mathcal{E}_{\Sigma}$ with $\mathcal{V}(e)$ disjoint to V and any variables assignment $\alpha : \mathcal{V}(e) \cup V \rightarrow \text{dom}(S)$ that $\llbracket e \rrbracket^{\alpha, S} = \llbracket e \rrbracket^{\alpha|_{\mathcal{V}(e)}, S}$. This is by induction on the structure of expressions. It follows for all formulas $\phi \in \mathcal{F}_{\Sigma}$ such that $\mathcal{V}(\phi)$ disjoint from V and $\alpha : \mathcal{V}(\phi) \cup V \rightarrow \text{dom}(S)$ that $\llbracket \phi \rrbracket^{\alpha, S} = \llbracket \phi \rrbracket^{\alpha|_{\mathcal{V}(\phi)}, S}$. This is by induction on the structure of formulas. It implies the lemma. ◀

► **Lemma 10 Quantification is projection.** *$\text{sol}^S(\exists x. \phi) = \pi_x(\text{sol}^S(\phi))$.*

Proof. This is straightforward from the semantics of existential quantifiers: $\text{sol}^S(\exists x. \phi) = \{\alpha|_{\mathcal{V}(\phi) \setminus \{x\}} \mid \alpha \in \text{sol}^S(\phi)\} = \pi_x(\text{sol}^S(\phi))$. ◀

509 **C Proofs for Section 6 (Abstraction Exactness)**

510 In order to do so we first show that h -exactness is preserved when adding variables. For
 511 this we have to assume that the abstraction h is surjective, which will be the case of all
 512 abstraction of interest.

513 ► **Lemma 47 Variable extension preserves exactness.** *Let $h : S \rightarrow \Delta$ be a Σ -abstraction
 514 that is surjective, $\phi \in \mathcal{F}_\Sigma$ a formula, and $V \supseteq \mathcal{V}(\phi)$. Then the h -exactness of ϕ implies the
 515 h -exactness of ϕ with respect to V .*

516 **Proof.** Essentially this follows from that solutions of formulas can be extended arbitrarily to
 517 variables that do not appear freely in the formula, as stated by the following claim.

518 ► **Claim 48.** *Any variable assignment $\sigma : V \rightarrow \Delta$ satisfies $\sigma \in h \circ \text{sol}_V^S(\phi)$ iff $\sigma|_{\mathcal{V}(\phi)} \in$
 519 $h \circ \text{sol}^S(\phi)$.*

520 For the one direction let $\sigma \in h \circ \text{sol}_V^S(\phi)$. Then there exists $\alpha \in \text{sol}_V^S(\phi)$ such that
 521 $\sigma = h \circ \alpha$. Since $V \supseteq \mathcal{V}(\phi)$ it follows that $\alpha|_{\mathcal{V}(\phi)} \in \text{sol}^S(\phi)$. Furthermore $\sigma|_{\mathcal{V}(\phi)} = h \circ \alpha|_{\mathcal{V}(\phi)}$
 522 and thus $\sigma|_{\mathcal{V}(\phi)} \in h \circ \text{sol}^S(\phi)$.

523 For the other direction let $\sigma|_{\mathcal{V}(\phi)} \in h \circ \text{sol}^S(\phi)$. Then there exists $\alpha \in \text{sol}^S(\phi)$ such that
 524 $\sigma|_{\mathcal{V}(\phi)} = h \circ \alpha$. For any $y \in V \setminus \mathcal{V}(\phi)$ let $s_y \in \text{dom}(S)$ be such that $h(s_y) = \sigma(y)$. Such
 525 values exists since h is surjective. Now define $\alpha' = \alpha[y/s_y \mid y \in V \setminus \mathcal{V}(\phi)]$. Since $V \supseteq \mathcal{V}(\phi)$ it
 526 follows that $\alpha' \in \text{sol}_V^S(\phi)$. Furthermore, $\sigma = h \circ \alpha'$, so $\sigma \in h \circ \text{sol}_V^S(\phi)$. ◀

527 ► **Lemma 17.** *Let V be a set of variables, R_1 and R_2 be subsets of assignments of type
 528 $V \rightarrow \text{dom}(S)$ and $h : S \rightarrow \Delta$ be a Σ -abstraction. $h \circ (R_1 \cup R_2) = h \circ R_1 \cup h \circ R_2$.*

Proof. This lemma follows from the following equivalences:

$$\begin{aligned} \beta \in h \circ (R_1 \cup R_2) &\Leftrightarrow \exists \alpha. \alpha \in R_1 \cup R_2 \wedge \beta = h \circ \alpha \\ &\Leftrightarrow \exists \alpha. (\alpha \in R_1 \vee \alpha \in R_2) \wedge \beta = h \circ \alpha \\ &\Leftrightarrow \exists \alpha. (\alpha \in R_1 \wedge \beta = h \circ \alpha) \vee (\alpha \in R_2 \wedge \beta = h \circ \alpha) \\ &\Leftrightarrow \beta \in h \circ R_1 \vee \beta \in h \circ R_2 \\ &\Leftrightarrow \beta \in h \circ R_1 \cup h \circ R_2 \end{aligned}$$

529 ◀

530 ► **Proposition 18.** *The disjunction of h -exact formulas is h -exact.*

Proof. Let ϕ_1 and ϕ_2 be negation free formulas that are h -exact. Let $V = \mathcal{V}(\phi_1) \cup \mathcal{V}(\phi_2)$.
 Lemma 47 shows that ϕ_1 and ϕ_2 are also h -exact with respect to the extended variable set
 V , i.e., for both $i \in \{1, 2\}$:

$$h \circ \text{sol}_V^S(\phi_i) = \text{sol}_V^\Delta(\phi_i)$$

The h -exactness of the disjunction $\phi_1 \vee \phi_2$ can now be shown as follows:

$$\begin{aligned} h \circ \text{sol}^S(\phi_1 \vee \phi_2) &= h \circ (\text{sol}_V^S(\phi_1) \cup \text{sol}_V^S(\phi_2)) \\ &= h \circ \text{sol}_V^S(\phi_1) \cup h \circ \text{sol}_V^S(\phi_2) && \text{by Lemma 17} \\ &= \text{sol}_V^\Delta(\phi_1) \cup \text{sol}_V^\Delta(\phi_2) && \text{by } h\text{-exactness of } \phi_1 \text{ and } \phi_2 \text{ wrt. } V \\ &= \text{sol}^\Delta(\phi_1 \vee \phi_2) \end{aligned}$$

531 ◀

532 ► **Lemma 19 Projection commutes with abstraction.** For any Σ -abstraction $h : S \rightarrow \Delta$,
 533 subset R of assignments of type $V \rightarrow S$, and variable $x \in \mathcal{V}$: $h \circ \pi_x(R) = \pi_x(h \circ R)$.

534 **Proof.** For all $\alpha : V \rightarrow \text{dom}(S)$ we have $h \circ \pi_x(\alpha) = h \circ \alpha|_{V \setminus \{x\}} = (h \circ \alpha)|_{V \setminus \{x\}} = \pi_x(h \circ \alpha)$. ◀

535 ► **Proposition 20 Quantification preserves exactness.** For any surjective Σ -abstraction
 536 $h : S \rightarrow \Delta$ and formula $\exists x.\phi \in \mathcal{F}_\Sigma$, if ϕ is h -exact then $\exists x.\phi$ is h -exact.

Proof. Let ϕ be h -exact. By definition ϕ is h -exact with respect to V . Since h is assumed to be surjective, Lemma 47 implies that ϕ is h -exact with respect to $V \cup \{x\}$ (independently of whether x occurs freely in ϕ or not). Hence:

$$\begin{aligned} h(\text{sol}^S(\exists x.\phi)) &= h(\pi_x(\text{sol}^S(\phi))) && \text{by Lemma 10} \\ &= \pi_x(h(\text{sol}^S(\phi))) && \text{by Lemma 19} \\ &= \pi_x(\text{sol}^\Delta(\phi)) && \text{since } \phi \text{ is } h\text{-exact} \\ &= \text{sol}^\Delta(\exists x.\phi) && \text{by Lemma 10} \end{aligned}$$

537

538 ► **Lemma 21 Singleton property.** If S is a Σ -algebra, $e \in \mathcal{E}_\Sigma$, and $\alpha : \mathcal{V}(e) \rightarrow S$ a variable
 539 assignment, then the set $\llbracket e \rrbracket^{\alpha, S}$ is a singleton.

540 **Proof.** By induction on the structure of expressions $e \in \mathcal{E}$:

541 **Case** of constants $c \in \{0, 1\}$. The set $\llbracket c \rrbracket^{\alpha, S} = \{c^S\}$ is a singleton.

542 **Case** of variables $x \in \mathcal{V}$. The set $\llbracket x \rrbracket^{\alpha, S} = \{\alpha(x)\}$ is a singleton.

Case $e_1 \odot e_2$ where $e_1, e_2 \in \mathcal{E}_\Sigma$ and $\odot \in \{+, *\}$.

$$\llbracket e_1 \odot e_2 \rrbracket^{\alpha, S} = \{s \odot^S s' \mid s \in \llbracket e_1 \rrbracket^{\alpha, S}, s' \in \llbracket e_2 \rrbracket^{\alpha, S}\}$$

543 This set is a singleton since $\llbracket e_1 \rrbracket^{\alpha, S}$ and $\llbracket e_2 \rrbracket^{\alpha, S}$ are singletons by induction hypothesis,
 544 meaning that $s \odot^S s'$ is also a singleton since S is a Σ -algebra. ◀

► **Lemma 23.** Let $\phi \in \mathcal{F}_\Sigma$ be a Σ -formula and S a Σ -algebra with unique division. For
 nonzero natural number n , variable $y \notin \mathcal{V}(\phi)$, and expression $e \in \mathcal{E}_\Sigma$ with $\mathcal{V}(e) \subseteq \mathcal{V}(\phi)$:

$$\text{sol}^S(\phi \wedge ny \overset{\circ}{=} e) = \{\alpha[y / \frac{\text{ele}(\llbracket e \rrbracket^{\alpha, S})}{n}] \mid \alpha \in \text{sol}^S(\phi)\}$$

545 **Proof.** We fix some $\alpha : \mathcal{V}(\phi) \rightarrow \text{dom}(S)$ arbitrarily. Since S is a Σ -algebra, $\llbracket e \rrbracket^{\alpha, S}$ is a
 546 singleton and $\mathcal{V}(e) \subseteq \mathcal{V}(\phi)$, $\text{ele}(\llbracket e \rrbracket^{\alpha, S})$ is defined uniquely. Furthermore S has unique
 547 division, so that $\frac{\text{ele}(\llbracket e \rrbracket^{\alpha, S})}{n}$ is well defined element of $\text{dom}(S)$. Therefore and since $y \notin \mathcal{V}(\phi)$,
 548 $\alpha[y / \frac{\text{ele}(\llbracket e \rrbracket^{\alpha, S})}{n}]$ is the unique solution of the equation $ny \overset{\circ}{=} e$ that extends on α .

549 First we prove the inclusion “ \supseteq ”. Let $\alpha \in \text{sol}^S(\phi)$, $y \notin \mathcal{V}(\phi)$, and $\alpha[y / \frac{\text{ele}(\llbracket e \rrbracket^{\alpha, S})}{n}]$ is a
 550 solution of $ny \overset{\circ}{=} e$, it follows that $\alpha[y / \frac{\text{ele}(\llbracket e \rrbracket^{\alpha, S})}{n}]$ is a solution of $\phi \wedge ny \overset{\circ}{=} e$.

551 Second, we prove the inverse inclusion “ \subseteq ”. Let $\alpha \in \text{sol}^S(\phi \wedge ny \overset{\circ}{=} e)$. Since $\alpha[y / \frac{\text{ele}(\llbracket e \rrbracket^{\alpha, S})}{n}]$
 552 is the unique solution of the equation $ny \overset{\circ}{=} e$ that extends on $\alpha' = \alpha|_{\mathcal{V}(\phi)}$ it follows that
 553 $\alpha(y) = \frac{\text{ele}(\llbracket e \rrbracket^{\alpha, S})}{n}$ so that $\alpha = \alpha'[y / \frac{\text{ele}(\llbracket e \rrbracket^{\alpha, S})}{n}]$ while $\alpha' \in \text{sol}^S(\phi)$. ◀

554 ► **Proposition 24.** Let $\phi \in \mathcal{F}_\Sigma$ a Σ -formula, $n \neq 0$ a natural number, $e \in \mathcal{E}_\Sigma$ a Σ -expression
 555 with $\mathcal{V}(e) \subseteq \mathcal{V}(\phi)$ and $y \notin \mathcal{V}(\phi)$ and the Σ -abstraction $h : S \rightarrow \Delta$ with S and Δ two Σ -algebras
 556 with unique division. Then if ϕ is h -exact implies that $\phi \wedge e \overset{\circ}{=} ny$ is h -exact.

557 **Proof.** Let $e \in \mathcal{E}_\Sigma$ a Σ -expression.

558 ▶ **Claim 49.** For any $\alpha : V \rightarrow \mathbb{R}_+$ with $V \supseteq \mathcal{V}(e) : h(\text{ele}(\llbracket e \rrbracket^{\alpha, S})) = \text{ele}(\llbracket e \rrbracket^{h \circ \alpha, \Delta})$.

559 This can be seen as follows. For any $\alpha : \mathcal{V}(\phi) \rightarrow S$ Theorem 11 on homomorphism yields
 560 $h(\llbracket e \rrbracket^{\alpha, S}) \subseteq \llbracket e \rrbracket^{h \circ \alpha, \Delta}$. Since S and Δ are both Σ -algebras, the sets $\llbracket e \rrbracket^{\alpha, S}$ and $\llbracket e \rrbracket^{h \circ \alpha, \Delta}$ are
 561 both singletons by Lemma 21, so that $h(\text{ele}(\llbracket e \rrbracket^{\alpha, S})) = \text{ele}(\llbracket e \rrbracket^{h \circ \alpha, \Delta})$.

562 ▶ **Claim 50.** For any $s \in \text{dom}(S)$ and $n \neq 0$ a natural number: $h(\frac{s}{n}) = \frac{h(s)}{n}$.

563 Since S is assumed to have unique division $s' = \frac{s}{n}$ is well-defined as the unique element
 564 of $\text{dom}(S)$ such that $\underbrace{s' +^S \dots +^S s'}_n = s$. Hence, $h(\underbrace{s' +^S \dots +^S s'}_n) = h(s)$ and since h is a
 565 homomorphism, it follows that $\underbrace{h(s') +^\Delta \dots +^\Delta h(s')}_n = h(s)$. Since Δ is assumed to have
 566 unique division, this implies that $h(s') = \frac{h(s)}{n}$.

The Proposition can now be shown based on these two claims. Let ϕ be h -exact, $y \notin \mathcal{V}(\phi)$,
 and $\mathcal{V}(e) \subseteq \mathcal{V}(\phi)$. We have to show that $\phi \wedge ny \stackrel{\circ}{=} e$ is h -exact too:

$$\begin{aligned}
 h \circ \text{sol}^S(\phi \wedge e \stackrel{\circ}{=} ny) &= h \circ \{ \alpha [y / \frac{\text{ele}(\llbracket e \rrbracket^{\alpha, S})}{n}] \mid \alpha \in \text{sol}^S(\phi) \} && \text{by Lemma 23} \\
 &= \{ (h \circ \alpha) [y / h(\frac{\text{ele}(\llbracket e \rrbracket^{\alpha, S})}{n})] \mid \alpha \in \text{sol}^S(\phi) \} && \text{elementary} \\
 &= \{ \sigma [y / h(\frac{\text{ele}(\llbracket e \rrbracket^{\alpha, S})}{n})] \mid \sigma \in \text{sol}^\Delta(\phi) \} && h\text{-exactness of } \phi \\
 &= \{ \sigma [y / \frac{h(\text{ele}(\llbracket e \rrbracket^{\alpha, S}))}{n}] \mid \sigma \in \text{sol}^\Delta(\phi) \} && \text{by Claim 50} \\
 &= \{ \sigma [y / \frac{\text{ele}(\llbracket e \rrbracket^{h \circ \alpha, \Delta})}{n}] \mid \sigma \in \text{sol}^\Delta(\phi) \} && \text{by Claim 49} \\
 &= \text{sol}^\Delta(\phi \wedge e \stackrel{\circ}{=} ny) && \text{by Lemma 23}
 \end{aligned}$$

567

568 ▶ **Proposition 25.** Let $h : S \rightarrow \Delta$ be a Σ -abstraction between Σ -algebras with unique division.
 569 Then any strongly-triangular system of Σ -equations with natural coefficients is h -exact.

570 **Proof.** Any strongly-triangular system of equations has the form $\bigwedge_{i=1}^n e_i \stackrel{\circ}{=} n_i y_i$ where n and
 571 $n_i \neq 0$ are naturals and y_i is i -fresh for all $1 \leq i \leq n$. The proof is by induction on n . In
 572 the case $n = 0$, the conjunction is equal to true which is h -exact since $h(\text{sol}^S(\text{true})) = h(\{\})$
 573 $= \text{sol}^\Delta(\text{true})$. In the case $n > 0$, we have by induction hypothesis that $\bigwedge_{j=1}^{i-1} e_j \stackrel{\circ}{=} n_j y_j$ is
 574 h -exact. Since $n_i \neq 0$ it follows from Proposition 24 that that $e_i \stackrel{\circ}{=} n_i y_i \wedge \bigwedge_{j=1}^{i-1} e_j \stackrel{\circ}{=} n_j y_j$ is
 575 h -exact. ◀

576 **D Proofs for Section 7 (Abstraction Invariance)**

577 ▶ **Lemma 29.** If either R_1 or R_2 are h -invariant then: $h \circ (R_1 \cap R_2) = h \circ R_1 \cap h \circ R_2$.

Proof. The one inclusion is straightforward without invariance:

$$\begin{aligned}
 h \circ (R_1 \cap R_2) &= \{ h \circ \alpha \mid \alpha \in R_1, \alpha \in R_2 \} \\
 &\subseteq \{ h \circ \alpha \mid \alpha \in R_1 \} \cap \{ h \circ \alpha \mid \alpha \in R_2 \} \\
 &= h \circ R_1 \cap h \circ R_2
 \end{aligned}$$

578 For the other inclusion, we can assume with loss of generality that R_1 is h -invariant. So let
 579 $\beta \in h \circ R_1 \cap h \circ R_2$. Then there exist $\alpha_1 \in R_1$ and $\alpha_2 \in R_2$ such that $\beta = h \circ \alpha_1 = h \circ \alpha_2$. By
 580 h -invariance of R_1 it follows that $\alpha_1 \in R_2$. So $\alpha_1 \in R_1 \cap R_2$, and hence, $\beta \in h \circ (R_1 \cap R_2)$. ◀

581 ► **Lemma 30 Algebraic characterization.** *A subset of R variables assignment of type $V \rightarrow$*
 582 *$dom(S)$ is h -invariant for an Σ -abstraction $h : S \rightarrow \Delta$ iff $h \circlearrowleft (h \circ R) \subseteq R$.*

583 **Proof.** “ \Rightarrow ”. Let R be h -invariant and $\alpha \in h \circlearrowleft (h \circ R)$. Then there exists $\alpha' \in R$ such that
 584 $h \circ \alpha = h \circ \alpha'$. The h -invariance of R thus implies that $\alpha \in R$.

585 “ \Leftarrow ”. Suppose that $h \circlearrowleft (h \circ R) \subseteq R$. Let $\alpha, \alpha' : V \rightarrow dom(S)$ such that $h \circ \alpha = h \circ \alpha'$ and
 586 $\alpha \in R$. We have to show that $\alpha' \in R$. From $h \circ \alpha = h \circ \alpha'$ and $\alpha \in R$ it follows that
 587 $\alpha' \in h \circlearrowleft (h \circ R)$ and thus $\alpha' \in R$ as required. ◀

588 ► **Lemma 51 Variable extension preserves invariance.** *Let h be a surjective abstraction*
 589 *and R a subset of functions of type $V' \rightarrow dom(S)$ and V a subset of variables disjoint from*
 590 *V' . If R is h -invariant then $ext_V^S(R)$ is h -invariant too.*

591 **Proof.** This will follow straightforwardly from the characterization of h -invariance in Lemma
 592 30 and the following two claims:

593 ► **Claim 52.** *If h is surjective then $h \circ ext_V^S(R) = ext_V^\Delta(h \circ R)$.*

594 This follows from $h \circ ext_V^S(R) = \{h \circ \alpha \mid \alpha \in ext_V^S(R)\} = ext_V^\Delta(\{h \circ \alpha' \mid \alpha' \in R\})$ where we
 595 use the surjectivity of h in the last step.

596 ► **Claim 53.** *$h \circ ext_V^\Delta(R') = ext_V^S(h \circlearrowleft R')$ for any subset R' of functions of type $V' \rightarrow dom(\Delta)$.*

$$\begin{aligned} h \circ ext_V^\Delta(R') &= \{\alpha : V \cup V' \rightarrow dom(S) \mid h \circ \alpha \in ext_V^\Delta(R')\} \\ &= \{\alpha : V \cup V' \rightarrow dom(S) \mid h \circ \alpha|_{V'} \in R'\} \\ &= ext_V^S(\{\alpha' : V' \rightarrow dom(S) \mid h \circ \alpha' \in R'\}) \\ &= ext_V^S(h \circlearrowleft R') \end{aligned}$$

597

598 ► **Lemma 54.** *Let $h : S \rightarrow \Delta$ be a surjective Σ -abstraction, ϕ be a Σ -formula, and $V \supseteq \mathcal{V}(\phi)$.*
 599 *Then the h -invariance of ϕ implies the h -invariance of $sol_V^S(\phi)$.*

600 **Proof.** This follows from the cylindrification Lemma 46 and that extension preserves h -
 601 invariance as shown in Lemma 51. ◀

602 ► **Lemma 32.** *For any Σ -abstraction $h : S \rightarrow \Delta$, any subsets of assignments of type*
 603 *$V \rightarrow dom(S)$ R_1 and R_2 and V a subset of variables:*

604 ■ $h \circlearrowleft (R_1 \cap R_2) = h \circlearrowleft R_1 \cap h \circlearrowleft R_2$.

605 ■ $h \circlearrowleft (R_1 \cup R_2) = h \circlearrowleft R_1 \cup h \circlearrowleft R_2$.

Proof. The case for unions follows straightforwardly from the definitions:

$$\begin{aligned} h \circlearrowleft (R_1 \cup R_2) &= \{\alpha \mid h \circ \alpha \in R_1 \cup R_2\} \\ &= \{\alpha \mid h \circ \alpha \in R_1 \vee h \circ \alpha \in R_2\} \\ &= \{\alpha \mid h \circ \alpha \in R_1\} \cup \{\alpha \mid h \circ \alpha \in R_2\} \\ &= h \circlearrowleft R_1 \cup h \circlearrowleft R_2 \end{aligned}$$

The case of intersection is symmetric:

$$\begin{aligned} h \circlearrowleft (R_1 \cap R_2) &= \{\alpha \mid h \circ \alpha \in R_1 \cap R_2\} \\ &= \{\alpha \mid h \circ \alpha \in R_1 \wedge h \circ \alpha \in R_2\} \\ &= \{\alpha \mid h \circ \alpha \in R_1\} \cap \{\alpha \mid h \circ \alpha \in R_2\} \\ &= h \circlearrowleft R_1 \cap h \circlearrowleft R_2 \end{aligned}$$

XX:20 Abstracting Linear Equation Systems

606

607 ► **Lemma 33 Intersection and union preserve invariance.** *Let $h : S \rightarrow \Delta$ be a Σ -abstraction.*
 608 *Then the intersection and union of any two h -invariant subsets R_1 and R_2 of variables*
 609 *assignments of type $V \rightarrow \text{dom}(S)$ is h -invariant.*

610 **Proof.** This follows from the algebraic characterization Lemma 30 for invariance, in combi-
 611 nation with the algebraic properties of composition and decomposition given in Lemmas 17,
 612 29, and 32. ◀

613 ► **Lemma 34 Projection commutes with left-decomposition.** $h \circ \pi_x(R) = \pi_x(h \circ R)$.

614 **Proof.** For all $\alpha : V \rightarrow \text{dom}(\Delta)$ we have $h \circ \pi_x(\alpha) = h \circ \alpha|_{V \setminus \{x\}} = (h \circ \alpha)|_{V \setminus \{x\}} = \pi_x(h \circ \alpha)$. ◀

615 ► **Proposition 35 Invariance is preserved by conjunction, disjunction, and quantification.**
 616 *If h is a surjective abstraction then the class of h -invariant FO-formulas is closed under*
 617 *conjunction, disjunction, and existential quantification.*

618 **Proof.** Let $h : S \rightarrow \Delta$ be a Σ -abstraction.

Case of conjunction: Let ϕ_1 and ϕ_2 be h -invariant and $V = \mathcal{V}(\phi_1 \wedge \phi_2)$. By Lemma 54 the sets $\text{sol}_V^S(\phi_1)$ and $\text{sol}_V^S(\phi_2)$ are both h -invariant, and so by Lemma 33 is their intersection. Hence:

$$\begin{aligned} h \circ (h \circ \text{sol}^S(\phi_1 \wedge \phi_2)) &= h \circ (h \circ (\text{sol}_V^S(\phi_1) \cap \text{sol}_V^S(\phi_2))) \\ &\subseteq \text{sol}_V^S(\phi_1) \cap \text{sol}_V^S(\phi_2) && \text{by } h\text{-invariance and Lemma 30} \\ &= \text{sol}^S(\phi_1 \wedge \phi_2) \end{aligned}$$

619 By Lemma 30 in the other direction, this implies that $\phi_1 \wedge \phi_2$ is h -invariant.

620 **Case of disjunction:** Analogous to the case of conjunction.

Case of existential quantification:

$$\begin{aligned} h \circ (h \circ \text{sol}^S(\exists x.\phi_1)) &= h \circ (h \circ \pi_x(\text{sol}^S(\phi_1))) && \text{by Lemma 10} \\ &= h \circ (\pi_x(h \circ \text{sol}^S(\phi_1))) && \text{by Lemma 19} \\ &= \pi_x(h \circ (h \circ \text{sol}^S(\phi_1))) && \text{by Lemma 34} \\ &\subseteq \pi_x(\text{sol}^S(\phi_1)) && \text{by } h\text{-invariance of } \phi_1 \text{ and Lemma 30} \\ &= \text{sol}^S(\exists x.\phi_1) && \text{by Lemma 10} \end{aligned}$$

621 By Lemma 30, this implies that $\exists x.\phi_1$ is h -invariant. ◀

622 ► **Proposition 36.** *Let h be a surjective Σ -abstractions. Then the class of h -exact and h -*
 623 *invariant Σ -formulas is closed under conjunction, disjunction and existential quantification.*

624 **Proof.** Closure under conjunction follows from Propositions 31 and 35, closure under disjunc-
 625 tion from Propositions 18 and 35, and closure under existential quantification by Propositions
 626 20 and 35. ◀

627 ► **Theorem 37 Invariance.** *Any positive polynomial equation $p \stackrel{\circ}{=} 0$ such that p has no constant*
 628 *term is $h_{\mathbb{B}}$ -exact and $h_{\mathbb{B}}$ -invariant.*

629 **Proof.** Consider a positive polynomial equation $p \stackrel{\circ}{=} 0$ such that p has no constant term and
 630 only positive coefficients. Thus p has the form $\sum_{j=1}^l n_j \prod_{k=1}^{i_j} x_{j,k}^{m_{j,k}} \stackrel{\circ}{=} 0$ where $l \geq 0$, and
 631 $n_j, i_j, m_{j,k} > 0$.

632 ▶ **Claim 55.** For both algebras $S \in \{\mathbb{B}, \mathbb{R}_+\}$: $\text{sol}^S(p \stackrel{\circ}{=} 0) = \text{sol}^S(\bigwedge_{j=1}^l \bigvee_{k=1}^{i_j} x_{j,k} \stackrel{\circ}{=} 0)$.

633 The polynomial has values zero if and only if all its monomials do, that is: $\prod_{k=1}^{i_j} x_{j,k}^{m_{j,k}} = 0$
 634 for all $1 \leq j \leq l$. Since constant terms are ruled out, we have $i_j \neq 0$. Furthermore, we
 635 assumed for all polynomials that $m_{j,k} \neq 0$. So for all $1 \leq j \leq l$ there must exist $1 \leq k \leq i_j$
 636 such that $x_{j,k} = 0$.

637 ▶ **Claim 56.** The equation $x \stackrel{\circ}{=} 0$ is $h_{\mathbb{B}}$ -exact and $h_{\mathbb{B}}$ -invariant.

This is straightforward from the definitions. With these two claims we are now in the position to prove the lemma. Since the class of $h_{\mathbb{B}}$ -exact and $h_{\mathbb{B}}$ -invariant formulas is closed under conjunction and disjunction by Proposition 36, it follows from by Claim 56 that $\bigwedge_{j=1}^l \bigvee_{k=1}^{i_j} x_{j,k} \stackrel{\circ}{=} 0$ is both $h_{\mathbb{B}}$ -exact and $h_{\mathbb{B}}$ -invariant. Since this formula is equivalent over \mathbb{R}_+ to the polynomial equation by Claim 55, the $h_{\mathbb{B}}$ -invariance carries over to $p \stackrel{\circ}{=} 0$. The $h_{\mathbb{B}}$ -exactness also carries over based on the equivalence for both structures \mathbb{R}_+ and \mathbb{B} :

$$\begin{aligned} h_{\mathbb{B}} \circ \text{sol}^{\mathbb{R}_+}(p \stackrel{\circ}{=} 0) &= h_{\mathbb{B}} \circ \text{sol}_V^{\mathbb{R}_+}(\bigwedge_{j=1}^l \bigvee_{k=1}^{i_j} x_{j,k} \stackrel{\circ}{=} 0) && \text{by Claim 55 for } \mathbb{R}_+ \\ &= \text{sol}^{\mathbb{B}}(\bigwedge_{j=1}^l \bigvee_{k=1}^{i_j} x_{j,k} \stackrel{\circ}{=} 0) && \text{by } h_{\mathbb{B}} \text{ exactness} \\ &= \text{sol}^{\mathbb{B}}(p \stackrel{\circ}{=} 0) && \text{by Claim 55 for } \mathbb{B}. \end{aligned}$$

638

639 **E** Proofs for Section 8 (Boolean Abstractions of $h_{\mathbb{B}}$ -Mixed Systems)

640 ▶ **Corollary 40.** The $h_{\mathbb{B}}$ -abstraction of the \mathbb{R}_+ -solution set of an $h_{\mathbb{B}}$ -mixed system ϕ , that is
 641 $h_{\mathbb{B}} \circ \text{sol}^{\mathbb{R}_+}(\phi)$, can be computed in at most exponential time in the size of the system ϕ .

642 **Proof.** Given an $h_{\mathbb{B}}$ -mixed system ϕ , we can apply Theorem 39 to compute in at most
 643 exponential time an \mathbb{R}_+ -equivalent formula ϕ'' that is $h_{\mathbb{B}}$ -exact. It is then sufficient to
 644 compute $\text{sol}^{\mathbb{B}}(\phi'')$ in exponential time in the size of ϕ . This can be done in the naive
 645 manner, that is by evaluating the formula ϕ'' – which may be of exponential size – over all
 646 possible boolean variable assignments – of which there may be exponentially many. For each
 647 assignment the evaluation can be done in PSPACE and thus in exponential time. The overall
 648 time required is thus a product of two exponentials, which remains exponential. ◀