



HAL
open science

Generalized logarithmic Hardy-Littlewood-Sobolev inequality

Jean Dolbeault, Xingyu Li

► **To cite this version:**

Jean Dolbeault, Xingyu Li. Generalized logarithmic Hardy-Littlewood-Sobolev inequality. International Mathematics Research Notices, inPress, 10.1093/imrn/rnz324 . hal-02281279v3

HAL Id: hal-02281279

<https://hal.science/hal-02281279v3>

Submitted on 24 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Generalized logarithmic Hardy-Littlewood-Sobolev inequality

Jean Dolbeault¹ and Xingyu Li¹

¹ CEREMADE (CNRS UMR n° 7534), PSL university, Université Paris-Dauphine,
Place de Lattre de Tassigny, 75775 Paris 16, France

Correspondence to be sent to: dolbeaul@ceremade.dauphine.fr

This paper is devoted to logarithmic Hardy-Littlewood-Sobolev inequalities in the two-dimensional Euclidean space, in presence of an external potential with logarithmic growth. The coupling with the potential introduces a new parameter, with two regimes. The attractive regime reflects the standard logarithmic Hardy-Littlewood-Sobolev inequality. The second regime corresponds to a reverse inequality, with the opposite sign in the convolution term, which allows us to bound the free energy of a drift-diffusion-Poisson system from below. Our method is based on an extension of an entropy method proposed by E. Carlen, J. Carrillo and M. Loss, and on a nonlinear diffusion equation.

1 Main result and motivation

On \mathbb{R}^2 , let us define the *density of probability* $\mu = e^{-V}$ and the *external potential* V by

$$\mu(x) := \frac{1}{\pi(1+|x|^2)^2} \quad \text{and} \quad V(x) := -\log \mu(x) = 2 \log(1+|x|^2) + \log \pi \quad \forall x \in \mathbb{R}^2.$$

We shall denote by $L_+^1(\mathbb{R}^2)$ the set of a.e. nonnegative functions in $L^1(\mathbb{R}^2)$. Our main result is the following *generalized logarithmic Hardy-Littlewood-Sobolev inequality*.

Theorem 1.1. For any $\alpha \geq 0$, we have that

$$\int_{\mathbb{R}^2} f \log \left(\frac{f}{M} \right) dx + \alpha \int_{\mathbb{R}^2} V f dx + M(1-\alpha)(1+\log \pi) \geq \frac{2}{M}(\alpha-1) \iint_{\mathbb{R}^2 \times \mathbb{R}^2} f(x)f(y) \log|x-y| dx dy \quad (1)$$

for any function $f \in L_+^1(\mathbb{R}^2)$ with $M = \int_{\mathbb{R}^2} f dx > 0$. Moreover, the equality case is achieved by $f_\star = M\mu$ and f_\star is the unique optimal function for any $\alpha > 0$. \square

With $\alpha = 0$, the inequality is the classical *logarithmic Hardy-Littlewood-Sobolev inequality*

$$\int_{\mathbb{R}^2} f \log \left(\frac{f}{M} \right) dx + \frac{2}{M} \iint_{\mathbb{R}^2 \times \mathbb{R}^2} f(x)f(y) \log|x-y| dx dy + M(1+\log \pi) \geq 0. \quad (2)$$

In that case f_\star is an optimal function as well as all functions generated by a translation and a scaling of f_\star . As long as the parameter α is in the range $0 \leq \alpha < 1$, the coefficient of the right-hand side of (1)

December 24, 2019 – File: LogHLS.tex

Keywords: logarithmic Hardy-Littlewood-Sobolev inequality; drift-diffusion-Poisson equation; entropy methods; nonlinear parabolic equations

MSC 2010: 26D10; 46E35; 35K55

is negative and the inequality is essentially of the same nature as the one with $\alpha = 0$. It can indeed be written as

$$\int_{\mathbb{R}^2} f \log\left(\frac{f}{M}\right) dx + \alpha \int_{\mathbb{R}^2} V f dx + M(1-\alpha)(1+\log\pi) + \frac{2}{M}(1-\alpha) \iint_{\mathbb{R}^2 \times \mathbb{R}^2} f(x)f(y) \log|x-y| dx dy \geq 0.$$

For reasons that will be made clear below, we shall call this range the *attractive range*.

If $\alpha = 1$, the inequality is almost trivial since

$$\int_{\mathbb{R}^2} f \log\left(\frac{f}{M}\right) dx + \int_{\mathbb{R}^2} V f dx = \int_{\mathbb{R}^2} f \log\left(\frac{f}{f_\star}\right) dx \geq 0 \quad (3)$$

is a straightforward consequence of Jensen's inequality. Now it is clear that by adding (2) multiplied by $(1-\alpha)$ and (3) multiplied by α , we recover (1) for any $\alpha \in [0, 1]$. As a consequence (1) is a straightforward interpolation between (2) and (3) in the *attractive range*.

Now, let us consider the *repulsive range* $\alpha > 1$. It is clear that the inequality is no more the consequence of a simple interpolation. We can also observe that the coefficient $(\alpha - 1)$ in the right-hand side of (1) is now positive. Since

$$G(x) = -\frac{1}{2\pi} \log|x|$$

is the Green function associated with $-\Delta$ on \mathbb{R}^2 , so that we can define

$$(-\Delta)^{-1}f(x) = (G * f)(x) = -\frac{1}{2\pi} \int_{\mathbb{R}^2} \log|x-y| f(y) dy,$$

it is interesting to write (1) as

$$\int_{\mathbb{R}^2} f \log\left(\frac{f}{M}\right) dx + \alpha \int_{\mathbb{R}^2} V f dx + \frac{4\pi}{M}(\alpha-1) \int_{\mathbb{R}^2} f (-\Delta)^{-1}f dx \geq M(\alpha-1)(1+\log\pi). \quad (4)$$

If f has a sufficient decay as $|x| \rightarrow +\infty$, for instance if f is compactly supported, we know that $(-\Delta)^{-1}f(x) \sim -\frac{M}{2\pi} \log|x|$ for large values of $|x|$ and as a consequence,

$$\alpha V + \frac{4\pi}{M}(\alpha-1)(-\Delta)^{-1}f \sim 2(\alpha+1) \log|x| \rightarrow +\infty \quad \text{as } |x| \rightarrow +\infty.$$

In a minimization scheme, this prevents the runaway of the left-hand side in (4). On the other hand, $\int_{\mathbb{R}^2} f \log f dx$ prevents any concentration, and this is why it can be heuristically expected that the left-hand side of (4) indeed admits a minimizer.

Inequality (2) was proved in [8] by E. Carlen and M. Loss (also see [2]). An alternative method based on nonlinear flows was given by E. Carlen, J. Carrillo and M. Loss in [7]: see Section 2 for a sketch of their proof. Our proof of Theorem 1.1 relies on an extension of this approach which takes into account the presence of the external potential V . A remarkable feature of this approach is that it is insensitive to the sign of $\alpha - 1$.

One of the key motivations for studying (4) arises from entropy methods applied to *drift-diffusion-Poisson* models which, after scaling out all physical parameters, are given by

$$\frac{\partial f}{\partial t} = \Delta f + \beta \nabla \cdot (f \nabla V) + \nabla \cdot (f \nabla \phi) \quad (5)$$

with a nonlinear coupling given by the *Poisson* equation

$$-\varepsilon \Delta \phi = f. \quad (6)$$

Here $V = -\log \mu$ is the external *confining potential* and we choose it as in the statement of Theorem 1.1, while $\beta \geq 0$ is a coupling parameter with V , which measures the strength of the external potential. We shall consider more general potentials at the end of this paper. The coefficient ε in (6) is either $\varepsilon = -1$, which corresponds to the *attractive* case, or $\varepsilon = +1$, which corresponds to the *repulsive* case. In terms of applications, when $\varepsilon = -1$, (6) is the equation for the mean field potential obtained from Newton's law of attraction in gravitation, for applications in astrophysics, or for the Keller-Segel concentration of chemo-attractant in chemotaxis. The case $\varepsilon = +1$ is used for repulsive electrostatic forces in semiconductor physics, electrolytes, plasmas and charged particle models.

In view of *entropy methods* applied to PDEs (see for instance [15]), it is natural to consider the *free energy functional*

$$\mathcal{F}_\beta[f] := \int_{\mathbb{R}^2} f \log f \, dx + \beta \int_{\mathbb{R}^2} V f \, dx + \frac{1}{2} \int_{\mathbb{R}^2} \phi f \, dx \quad (7)$$

because, if $f > 0$ solves (5)-(6) and is smooth enough, with sufficient decay properties at infinity, then

$$\frac{d}{dt} \mathcal{F}_\beta[f(t, \cdot)] = - \int_{\mathbb{R}^2} f |\nabla \log f + \beta \nabla V + \nabla \phi|^2 \, dx \quad (8)$$

so that \mathcal{F}_β is a Lyapunov functional. Of course, a preliminary question is to establish under which conditions \mathcal{F}_β is bounded from below. The answer is given by the following result.

Corollary 1.2. Let $M > 0$. If $\varepsilon = +1$, the functional \mathcal{F}_β is bounded from below and admits a minimizer on the set of the functions $f \in L^1_+(\mathbb{R}^2)$ such that $\int_{\mathbb{R}^2} f \, dx = M$ if $\beta \geq 1 + \frac{M}{8\pi}$. It is bounded from below if $\varepsilon = -1$, $\beta \geq 1 - \frac{M}{8\pi}$ and $M \leq 8\pi$. If $\varepsilon = +1$, the minimizer is unique. \square

As we shall see in Section 3.1, Corollary 1.2 is a simple consequence of Theorem 1.1. In the case of the parabolic-elliptic Keller-Segel model, that is, with $\varepsilon = -1$ and $\beta = 0$, this has been used in [12, 4] to provide a sharp range of existence of the solutions to the evolution problem. In [6], the case $\varepsilon = -1$ with a potential V with quadratic growth at infinity was also considered, in the study of intermediate asymptotics of the parabolic-elliptic Keller-Segel model.

Concerning the *drift-diffusion-Poisson* model (5)-(6) and considerations on the *free energy*, in the electrostatic case, we can quote, among many others, [14, 13] and subsequent papers. In the Euclidean space with confining potentials, we shall refer to [10, 11, 3, 1]. However, as far as we know, these papers are primarily devoted to dimensions $d \geq 3$ and the sharp growth condition on V when $d = 2$ has not been studied so far. The goal of this paper is to fill this gap. The specific choice of V has been made to obtain explicit constants and optimal inequalities, but the confining potential plays a role only at infinity if we are interested in the boundedness from below of the free energy. In Section 3.3, we shall give a result for general potentials on \mathbb{R}^2 : see Theorem 3.4 for a statement.

2 Proof of the main result

As an introduction to the key method, we briefly sketch the proof of (2) given by E. Carlen, J. Carrillo and M. Loss in [7]. The main idea is to use the nonlinear diffusion equation

$$\frac{\partial f}{\partial t} = \Delta \sqrt{f}$$

with a nonnegative initial datum f_0 . The equation preserves the mass $M = \int_{\mathbb{R}^2} f dx$ and is such that

$$\frac{d}{dt} \left(\int_{\mathbb{R}^2} f \log f dx - \frac{4\pi}{M} \int_{\mathbb{R}^2} f ((-\Delta)^{-1} f) dx \right) = - \frac{8}{M} \left(\int_{\mathbb{R}^2} |\nabla f^{1/4}|^2 dx \int_{\mathbb{R}^2} f dx - \pi \int_{\mathbb{R}^2} f^{3/2} dx \right).$$

According to [9], the Gagliardo-Nirenberg inequality

$$\|\nabla g\|_2^2 \|g\|_4^4 \geq \pi \|g\|_6^6 \quad (9)$$

applied to $g = f^{1/4}$ guarantees that the right-hand side is nonpositive. By the general theory of fast diffusion equations (we refer for instance to [17]), we know that the solution behaves for large values of t like a self-similar solution, the so-called Barenblatt solution, which is given by $B(t, x) := t^{-2} f_\star(x/t)$. As a consequence, we find that

$$\begin{aligned} & \int_{\mathbb{R}^2} f_0 \log f_0 dx - \frac{4\pi}{M} \int_{\mathbb{R}^2} f_0 ((-\Delta)^{-1} f_0) dx \\ & \geq \lim_{t \rightarrow +\infty} \int_{\mathbb{R}^2} B \log B dx - \frac{4\pi}{M} \int_{\mathbb{R}^2} B ((-\Delta)^{-1} B) dx = \int_{\mathbb{R}^2} f_\star \log f_\star dx - \frac{4\pi}{M} \int_{\mathbb{R}^2} f_\star ((-\Delta)^{-1} f_\star) dx \end{aligned}$$

After an elementary computation, we observe that the above inequality is exactly (2) written for $f = f_0$.

The point is now to adapt this strategy to the case with an external potential. This justifies why we have to introduce a nonlinear diffusion equation with a drift. As we shall see below, the method is insensitive to α and applies when $\alpha > 1$ exactly as in the case $\alpha \in (0, 1)$. A natural question is whether solutions are regular enough to perform the computations below and in particular if they have a sufficient decay at infinity to allow all kinds of integrations by parts needed by the method. The answer is twofold. First, we can take an initial datum f_0 that is as smooth and decaying as $|x| \rightarrow +\infty$ as needed, prove the inequality and argue by density. Second, integrations by parts can be justified by an approximation scheme consisting in a truncation of the problem in larger and larger balls. We refer to [17] for regularity issues and to [15] for the truncation method. In the proof, we will therefore leave these issues aside, as they are purely technical.

Proof of Theorem 1.1. By homogeneity, we can assume that $M = 1$ without loss of generality and consider the evolution equation

$$\frac{\partial f}{\partial t} = \Delta \sqrt{f} + 2\sqrt{\pi} \nabla \cdot (x f).$$

1) Using simple integrations by parts, we compute

$$\int_{\mathbb{R}^2} (1 + \log f) \Delta \sqrt{f} dx = -8 \int_{\mathbb{R}^2} |\nabla f^{1/4}|^2 dx$$

and

$$\int_{\mathbb{R}^2} (1 + \log f) \nabla \cdot (x f) dx = - \int_{\mathbb{R}^2} \frac{\nabla f}{f} \cdot (x f) dx = - \int_{\mathbb{R}^2} x \cdot \nabla f dx = 2 \int_{\mathbb{R}^2} f dx = 2.$$

As a consequence, we obtain that

$$\frac{d}{dt} \int_{\mathbb{R}^2} f \log f dx = -8 \int_{\mathbb{R}^2} |\nabla f^{1/4}|^2 dx + 8\pi \int_{\mathbb{R}^2} \mu^{3/2} dx \quad (10)$$

using

$$\int_{\mathbb{R}^2} \mu^{3/2} dx = \frac{1}{2\sqrt{\pi}}.$$

2) By elementary considerations again, we find that

$$4\pi \int_{\mathbb{R}^2} f(-\Delta)^{-1} (\Delta \sqrt{f}) dx = -4\pi \int_{\mathbb{R}^2} f^{3/2} dx$$

and

$$\begin{aligned} 4\pi \int_{\mathbb{R}^2} \nabla \cdot (xf) (-\Delta)^{-1} f dx &= -4\pi \int_{\mathbb{R}^2} xf \cdot \nabla (-\Delta)^{-1} f dx \\ &= 2 \iint_{\mathbb{R}^2 \times \mathbb{R}^2} f(x) f(y) x \cdot \frac{x-y}{|x-y|^2} dx dy \\ &= \iint_{\mathbb{R}^2 \times \mathbb{R}^2} f(x) f(y) (x-y) \cdot \frac{x-y}{|x-y|^2} dx dy = 1 \end{aligned}$$

where, in the last line, we exchanged the variables x and y and took the half sum of the two expressions. This proves that

$$\frac{d}{dt} \left(4\pi \int_{\mathbb{R}^2} f ((-\Delta)^{-1} f) dx \right) = -8\pi \int_{\mathbb{R}^2} (f^{3/2} - \mu^{3/2}) dx. \quad (11)$$

3) We observe that

$$\mu(x) = \frac{1}{\pi(1+|x|^2)^2} = e^{-V(x)}$$

solves

$$\Delta V = -\Delta \log \mu = 8\pi \mu \quad (12)$$

and, as a consequence,

$$\int_{\mathbb{R}^2} V \Delta \sqrt{f} dx = \int_{\mathbb{R}^2} \Delta V \sqrt{f} dx = 8\pi \int_{\mathbb{R}^2} \mu \sqrt{f} dx.$$

Since

$$\begin{aligned} 2\sqrt{\pi} \int_{\mathbb{R}^2} V \nabla \cdot (xf) dx &= -2\sqrt{\pi} \int_{\mathbb{R}^2} fx \cdot \nabla V dx = -8\sqrt{\pi} \int_{\mathbb{R}^2} \frac{|x|^2}{1+|x|^2} f dx \\ &= -8\sqrt{\pi} + 8\sqrt{\pi} \int_{\mathbb{R}^2} \frac{f}{1+|x|^2} dx = -8\sqrt{\pi} + 8\pi \int_{\mathbb{R}^2} \sqrt{\mu} f dx, \end{aligned}$$

we conclude that

$$\frac{d}{dt} \int_{\mathbb{R}^2} f V dx = 8\pi \int_{\mathbb{R}^2} (\mu \sqrt{f} + \sqrt{\mu} f - 2\mu^{3/2}) dx. \quad (13)$$

Let us define

$$\mathcal{F}[f] := \int_{\mathbb{R}^2} f \log f dx + \alpha \int_{\mathbb{R}^2} V f dx + (1-\alpha)(1+\log \pi) + 2(1-\alpha) \iint_{\mathbb{R}^2 \times \mathbb{R}^2} f(x) f(y) \log |x-y| dx dy.$$

Collecting (10), (11) and (13), we find that

$$\frac{d}{dt} \mathcal{F}[f(t, \cdot)] = -8 \left(\int_{\mathbb{R}^2} |\nabla f^{1/4}|^2 dx - \pi \int_{\mathbb{R}^2} f^{3/2} dx \right) - 8\pi \alpha \int_{\mathbb{R}^2} (f^{3/2} - \mu \sqrt{f} - \sqrt{\mu} f + \mu^{3/2}) dx.$$

Notice that

$$\int_{\mathbb{R}^2} (f^{3/2} - \mu \sqrt{f} - \sqrt{\mu} f + \mu^{3/2}) dx = \int_{\mathbb{R}^2} \varphi \left(\frac{f}{\mu} \right) \mu^{3/2} dx \quad \text{with} \quad \varphi(t) := t^{3/2} - t - \sqrt{t} + 1$$

and that φ is a strictly convex function on \mathbb{R}^+ such that $\varphi(1) = \varphi'(1) = 0$, so that φ is nonnegative. On the other hand, by (9), we know that

$$\int_{\mathbb{R}^2} |\nabla f^{1/4}|^2 dx - \pi \int_{\mathbb{R}^2} f^{3/2} dx \geq 0$$

as in the proof of [7]. Altogether, this proves that $t \mapsto \mathcal{F}[f(t, \cdot)]$ is monotone nonincreasing. Hence

$$\mathcal{F}[f_0] \geq \mathcal{F}[f(t, \cdot)] \geq \lim_{t \rightarrow +\infty} \mathcal{F}[f(t, \cdot)] = \mathcal{F}[f_\star] = 0.$$

This completes the proof of (1). ■

3 Consequences

3.1 Proof of Corollary 1.2

To prove the result of Corollary 1.2, we have to establish first that the *free energy* functional \mathcal{F}_β is bounded from below. Instead of using standard variational methods to prove that a minimizer is achieved, we can rely on the flow associated with (5)-(6).

• **Repulsive case.** Let us consider the *free energy* functional defined in (7) where ϕ is given by (6) with $\varepsilon = +1$, i.e., $\phi = -\frac{1}{2\pi} \log|\cdot| * f$.

Lemma 3.1. Let $M > 0$ and $\varepsilon = +1$. Then \mathcal{F}_β is bounded from below on the set of the functions $f \in L^1_+(\mathbb{R}^2)$ such that $\int_{\mathbb{R}^2} f dx = M$ if $\beta \geq 1 + \frac{M}{8\pi}$. □

Proof. With $g = \frac{f}{M}$ and $\alpha = 1 + \frac{M}{8\pi}$, this means that

$$\begin{aligned} \frac{1}{M} \mathcal{F}_\beta[f] - \log M &= \int_{\mathbb{R}^2} g \log g dx + \beta \int_{\mathbb{R}^2} V g dx - \frac{M}{4\pi} \iint_{\mathbb{R}^2 \times \mathbb{R}^2} g(x) g(y) \log|x-y| dx dy \\ &= (\beta - \alpha) \int_{\mathbb{R}^2} V g dx + \int_{\mathbb{R}^2} g \log g dx + \alpha \int_{\mathbb{R}^2} V g dx - 2(\alpha - 1) \iint_{\mathbb{R}^2 \times \mathbb{R}^2} g(x) g(y) \log|x-y| dx dy \\ &\geq (\beta - \alpha) \int_{\mathbb{R}^2} V g dx - (1 - \alpha) (1 + \log \pi) \end{aligned}$$

according to Theorem 1.1; the condition $\beta \geq \alpha$ is enough to prove that $\mathcal{F}_\beta[f]$ is bounded from below. ■

Proof of Corollary 1.2 with $\varepsilon = +1$. Let us consider a smooth solution of (5)-(6). We refer to [16] for details and to [1] for similar arguments in dimension $d \geq 3$. According to (8), f converges as $t \rightarrow +\infty$ to a solution of

$$\nabla \log f + \beta \nabla V + \nabla \phi = 0.$$

Notice that this already proves the existence of a stationary solution. The equation can be solved as

$$f = M \frac{e^{-\beta V - \phi}}{\int_{\mathbb{R}^2} e^{-\beta V - \phi} dx}$$

after taking into account the conservation of the mass. With (6), the problem is reduced to solving

$$-\Delta \psi = M \left(\frac{e^{-\gamma V - \psi}}{\int_{\mathbb{R}^2} e^{-\gamma V - \psi} dx} - \mu \right), \quad \psi = (\beta - \gamma) V + \phi, \quad \gamma = \beta - \frac{M}{8\pi}$$

using (12). It is a critical point of the functional $\psi \mapsto \mathcal{J}_{M,\gamma}[\psi] := \frac{1}{2} \int_{\mathbb{R}^2} |\nabla \psi|^2 dx + M \int_{\mathbb{R}^2} \psi \mu dx + M \log \left(\int_{\mathbb{R}^2} e^{-\gamma V - \psi} dx \right)$. Such a functional is strictly convex as, for instance, in [10, 11]. We conclude that ψ is unique up to an additional constant. ■

• **Attractive case.** Let us consider the *free energy* functional (7) \mathcal{F}_β where ϕ is given by (6) with $\varepsilon = -1$, i.e., $\phi = \frac{1}{2\pi} \log |\cdot| * f$. Inspired by [12], we have the following estimate.

Lemma 3.2. Let $\varepsilon = -1$. Then \mathcal{F}_β is bounded from below on the set of the functions $f \in L^1_+(\mathbb{R}^2)$ such that $\int_{\mathbb{R}^2} f dx = M$ if $M \leq 8\pi$ and $\beta \geq 1 - \frac{M}{8\pi}$. It is not bounded from below if $M > 8\pi$. □

Proof. With $g = \frac{f}{M}$ and $\alpha = 1 - \frac{M}{8\pi}$, Theorem 1.1 applied to

$$\begin{aligned} & \frac{1}{M} \mathcal{F}_\beta[f] - \log M \\ &= (\beta - \alpha) \int_{\mathbb{R}^2} V g dx + \int_{\mathbb{R}^2} g \log g dx + \alpha \int_{\mathbb{R}^2} V g dx + 2(1 - \alpha) \iint_{\mathbb{R}^2 \times \mathbb{R}^2} g(x) g(y) \log |x - y| dx dy \\ & \geq (\beta - \alpha) \int_{\mathbb{R}^2} V g dx - (1 - \alpha) (1 + \log \pi) \end{aligned}$$

proves that the free energy is bounded from below if $M \leq 8\pi$ and $\beta \geq \alpha$. On the other hand, if $f_\lambda(x) := \lambda^{-2} f(\lambda^{-1} x)$ and $M > 8\pi$, then

$$\mathcal{F}_\beta[f_\lambda] \sim 2M \left(\frac{M}{8\pi} - 1 \right) \log \lambda \rightarrow -\infty \quad \text{as } \lambda \rightarrow 0_+,$$

which proves that \mathcal{F}_β is not bounded from below. ■

Proof of Corollary 1.2 with $\varepsilon = -1$. The proof goes as in the case $\beta = 0$. We refer to [4] and leave details to the reader. ■

Remark 3.3. Let us notice that \mathcal{F}_β is unbounded from below if $\beta < 0$. This follows from the observation that $\lim_{|y| \rightarrow \infty} \mathcal{F}_\beta[f_y] = -\infty$ where $f_y(x) = f(x + y)$ for any admissible f . □

3.2 Duality

When $\alpha > 1$, we can write a first inequality by considering the *repulsive case* in the proof of Corollary 1.2 and observing that

$$\mathcal{J}_{M,\gamma}[\psi] \geq \min \mathcal{J}_{M,\gamma}$$

where $\psi \in W_{\text{loc}}^{2,1}(\mathbb{R}^2)$ is such that $\int_{\mathbb{R}^2} (\Delta \psi) dx = 0$ and the minimum is taken on the same set of functions.

When $\alpha \in [0, 1)$, it is possible to argue by duality as in [5, Section 2]. Since f_\star realizes the equality case in (1), we know that

$$\int_{\mathbb{R}^2} f_\star \log \left(\frac{f_\star}{M} \right) dx + \alpha \int_{\mathbb{R}^2} V f_\star dx + M(1 - \alpha) (1 + \log \pi) = \frac{2}{M} (\alpha - 1) \iint_{\mathbb{R}^2 \times \mathbb{R}^2} f_\star(x) f_\star(y) \log |x - y| dx dy$$

and, using the fact that f_\star is a critical point of the difference of the two sides of (1), we also have that

$$\int_{\mathbb{R}^2} \log \left(\frac{f}{f_\star} \right) (f - f_\star) dx + \alpha \int_{\mathbb{R}^2} V (f - f_\star) dx = \frac{4}{M} (\alpha - 1) \iint_{\mathbb{R}^2 \times \mathbb{R}^2} (f(x) - f_\star(x)) f_\star(y) \log |x - y| dx dy.$$

By subtracting the first identity to (1) and adding the second identity, we can rephrase (1) as

$$\mathcal{F}_{(1)}[f] := \int_{\mathbb{R}^2} f \log \left(\frac{f}{f_\star} \right) dx \geq \frac{4\pi}{M} (1-\alpha) \int_{\mathbb{R}^2} (f - f_\star) (-\Delta)^{-1} (f - f_\star) dx := \mathcal{F}_{(2)}[f].$$

Let us consider the Legendre transform

$$\mathcal{F}_{(i)}^*[g] := \sup_f \left(\int_{\mathbb{R}^2} g f dx - \mathcal{F}_{(i)}[f] \right)$$

where the supremum is restricted to the set of the functions $f \in L^1_+(\mathbb{R}^2)$ such that $M = \int_{\mathbb{R}^2} f dx$. After taking into account the Lagrange multipliers associated with the mass constraint, we obtain that

$$M \log \left(\int_{\mathbb{R}^2} e^{g-V} dx \right) = \mathcal{F}_{(1)}^*[g] \leq \frac{M}{16\pi(1-\alpha)} \int_{\mathbb{R}^2} |\nabla g|^2 dx + M \int_{\mathbb{R}^2} g e^{-V} dx = \mathcal{F}_{(2)}^*[g].$$

We can get rid of M by homogeneity and recover the standard Euclidean form of the Onofri inequality in the limit case as $\alpha \rightarrow 0_+$, which is clearly the sharpest one for all possible $\alpha \in [0, 1)$.

3.3 Extension to general confining potentials with critical asymptotic growth

As a concluding observation, let us consider a general potential W on \mathbb{R}^2 such that

$$W \in C(\mathbb{R}^2) \quad \text{and} \quad \lim_{|x| \rightarrow +\infty} \frac{W(x)}{V(x)} = \beta \quad (\mathcal{H}_W)$$

and the associated *free energy functional*

$$\mathcal{F}_{\beta,W}[f] := \int_{\mathbb{R}^2} f \log f dx + \beta \int_{\mathbb{R}^2} W f dx + \frac{1}{2} \int_{\mathbb{R}^2} \phi f dx$$

where ϕ is given in terms of $f > 0$ by (6). With previous notations, $\mathcal{F}_\beta = \mathcal{F}_{\beta,V}$. Our last result is that the asymptotic behaviour obtained from (\mathcal{H}_W) is enough to decide whether $\mathcal{F}_{\beta,W}$ is bounded from below or not. The precise result goes as follows.

Theorem 3.4. Under Assumption (\mathcal{H}_W) , $\mathcal{F}_{\beta,W}$ defined as above is bounded from below if either $\varepsilon = +1$ and $\beta > 1 + \frac{M}{8\pi}$, or $\varepsilon = -1$, $\beta > 1 - \frac{M}{8\pi}$ and $M \leq 8\pi$. The result is also true in the limit case if $(W - \beta V) \in L^\infty(\mathbb{R}^2)$ and either $\varepsilon = +1$ and $\beta = 1 + \frac{M}{8\pi}$, or $\varepsilon = -1$, $\beta = 1 - \frac{M}{8\pi}$ and $M \leq 8\pi$. \square

Proof. If $(W - \beta V) \in L^\infty(\mathbb{R}^2)$, we can write that

$$\mathcal{F}_{\beta,W}[f] \geq \mathcal{F}_\beta[f] - M \|W - \beta V\|_{L^\infty(\mathbb{R}^2)}.$$

This completes the proof in the limit case. Otherwise, we redo the argument using $\tilde{\beta} V - (\tilde{\beta} V - W)_+$ for some $\tilde{\beta} \in (0, \beta)$ if $\varepsilon = -1$, and for some $\tilde{\beta} \in (1 + \frac{M}{8\pi}, \beta)$ if $\varepsilon = +1$. \blacksquare

Acknowledgment: This work has been partially supported by the Project EFI (ANR-17-CE40-0030) of the French National Research Agency (ANR). The authors thank L. Jeanjean who pointed them some typographical errors. © 2019 by the authors. This paper may be reproduced, in its entirety, for non-commercial purposes.

References

- [1] Arnold, A., Markowich, P., Toscani, G., and Unterreiter, A. (2001). On convex Sobolev inequalities and the rate of convergence to equilibrium for Fokker-Planck type equations. *Comm. Partial Differential Equations*, 26(1-2):43–100.
- [2] Beckner, W. (1993). Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality. *Ann. of Math. (2)*, 138(1):213–242.
- [3] Biler, P. and Dolbeault, J. (2000). Long time behavior of solutions of Nernst-Planck and Debye-Hückel drift-diffusion systems. *Ann. Henri Poincaré*, 1(3):461–472.
- [4] Blanchet, A., Dolbeault, J., and Perthame, B. (2006). Two-dimensional Keller-Segel model: optimal critical mass and qualitative properties of the solutions. *Electron. J. Differential Equations*, 44:1–33.
- [5] Campos, J. and Dolbeault, J. (2012). A functional framework for the Keller-Segel system: logarithmic Hardy-Littlewood-Sobolev and related spectral gap inequalities. *C. R. Math. Acad. Sci. Paris*, 350(21-22):949–954.
- [6] Campos, J. F. and Dolbeault, J. (2014). Asymptotic Estimates for the Parabolic-Elliptic Keller-Segel Model in the Plane. *Comm. Partial Differential Equations*, 39(5):806–841.
- [7] Carlen, E. A., Carrillo, J. A., and Loss, M. (2010). Hardy-Littlewood-Sobolev inequalities via fast diffusion flows. *Proc. Natl. Acad. Sci. USA*, 107(46):19696–19701.
- [8] Carlen, E. A. and Loss, M. (1992). Competing symmetries, the logarithmic HLS inequality and Onofri’s inequality on \mathbb{S}^n . *Geom. Funct. Anal.*, 2(1):90–104.
- [9] Del Pino, M. and Dolbeault, J. (2002). Best constants for Gagliardo-Nirenberg inequalities and applications to nonlinear diffusions. *J. Math. Pures Appl. (9)*, 81(9):847–875.
- [10] Dolbeault, J. (1991). Stationary states in plasma physics: Maxwellian solutions of the Vlasov-Poisson system. *Math. Models Methods Appl. Sci.*, 1(2):183–208.
- [11] Dolbeault, J. (1999). Free energy and solutions of the Vlasov-Poisson-Fokker-Planck system: external potential and confinement (large time behavior and steady states). *J. Math. Pures Appl. (9)*, 78(2):121–157.
- [12] Dolbeault, J. and Perthame, B. (2004). Optimal critical mass in the two-dimensional Keller-Segel model in \mathbb{R}^2 . *C. R. Math. Acad. Sci. Paris*, 339(9):611–616.
- [13] Dressler, K. (1990). Steady states in plasma physics – the Vlasov-Fokker-Planck equation. *Math. Methods Appl. Sci.*, 12(6):471–487.
- [14] Gogny, D. and Lions, P.-L. (1989). Sur les états d’équilibre pour les densités électroniques dans les plasmas. *RAIRO Modél. Math. Anal. Numér.*, 23(1):137–153.
- [15] Jüngel, A. (2016). *Entropy methods for diffusive partial differential equations*. SpringerBriefs in Mathematics. Springer, [Cham].
- [16] Li, X. Asymptotic behavior of Nernst-Planck equation. Preprint Hal: [hal-02310654](https://hal.archives-ouvertes.fr/hal-02310654) and [arXiv:1910.04477](https://arxiv.org/abs/1910.04477), 2019.
- [17] Vázquez, J. L. (2006). *Smoothing and decay estimates for nonlinear diffusion equations*, volume 33 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford.