



Are life-history traits equally affected by global warming? A case study combining a multi-trait approach with fine-grain climate modeling.

Corentin Iltis, Philippe Louâpre, Karolina Pecharová, Denis Thiéry, Sébastien Zito, Benjamin Bois, Jérôme Moreau

► To cite this version:

Corentin Iltis, Philippe Louâpre, Karolina Pecharová, Denis Thiéry, Sébastien Zito, et al.. Are life-history traits equally affected by global warming? A case study combining a multi-trait approach with fine-grain climate modeling.. Journal of Insect Physiology, 2019, 117, pp.103916. 10.1016/j.jinsphys.2019.103916 . hal-02281531

HAL Id: hal-02281531

<https://hal.science/hal-02281531>

Submitted on 25 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 Are life-history traits equally affected by global warming? A case study
2 combining a multi-trait approach with fine-grain climate modeling

3 Corentin Iltis^{*a}, Philippe Louâpre^a, Karolina Pecharová^a, Denis Thiéry^b, Sébastien
4 Zito^a, Benjamin Bois^a, and Jérôme Moreau^a

5 ^a UMR CNRS 6282 Biogéosciences, Université Bourgogne Franche-Comté,
6 6 Boulevard Gabriel, 21000 Dijon, France.

7 ^b UMR INRA 1065 Santé et Agroécologie du Vignoble, Institut des Sciences de la Vigne et du
8 Vin, 71 Avenue Edouard Bourlaux, 33882 Villenave-d'Ornon, France.

9

10 * Corresponding author. E-mail: corentin.iltis@u-bourgogne.fr. ORCID: 0000-0002-1379-

11 8029

Abstract

Predicting species responses to climate change requires tracking the variation in individual performance following exposure to warming conditions. One ecologically relevant approach consists of examining the thermal responses of a large number of traits, both related with population dynamics and trophic interactions (i.e. a multi-trait approach). Based on *in situ* climatic data and projections from climate models, we here designed two daily fluctuating thermal regimes realistically reflecting current and future conditions in Eastern France. These models detected an increase in mean temperature and in the range of daily thermal fluctuations as two local facets of global warming likely to occur in our study area by the end of this century. We then examined the responses of several fitness-related traits in caterpillars of the moth *Lobesia botrana* – including development, pupal mass, survival rates, energetic reserves, behavioral and immune traits expressed against parasitoids – to this experimental imitation of global warming. Increasing temperatures positively affected development (leading to a 31% reduction in the time needed to complete larval stage), survival rates (+19%), and movement speed as a surrogate for larval escape ability to natural enemies (+60%). Conversely, warming elicited detrimental effects on lipid reserves (-26%) and immunity (total phenoloxidase activity: -34%). These findings confirm that traits should differ in their sensitivity to global warming, underlying complex consequences for population dynamics and trophic interactions. Our study strengthens the importance of combining a multi-trait approach with the use of realistic fluctuating regimes to forecast the consequences of global warming for individuals, species and species assemblages.

Keywords

Climate models, defensive behaviors, energetic reserves, global warming, immunity, multi-trait approach.

1. Introduction

Temperature is undeniably one of the most important factors determining individual performance, species evolution and distribution (Angilletta, 2009; Clarke, 2003). In ectothermic animals like insects, whose body temperature and performance are closely related with environmental temperature, the link between a surrogate for individual performance (e.g. development and growth rates, reproductive output) and environmental temperature is typically modeled by an asymmetric and unimodal curve called a thermal performance curve (Martin and Huey, 2008; Sinclair et al., 2016). As temperature increases, individual performance gently rises to a single maximum value associated with a thermal optimum, and thereafter performance declines rapidly.

Thermal performance curves may help to predict how organisms will cope with changes occurring in their thermal environment, such as those related with anthropogenic climate disturbance (Deutsch et al., 2008; Estay et al., 2014; Sinclair et al., 2016). Indeed, one commonly used way to assess the biological consequences of global warming consists of experimentally tracking the variation in organism's performance following exposure to warming conditions designed to simulate future temperatures (for examples, see Bauerfeind and Fischer, 2014; Fischer et al., 2014; Klockmann et al., 2016). In this regard, it is now widely accepted that comparing fluctuating thermal regimes (instead of constant temperatures) significantly improves the ecological realism of global warming experiments, considering the daily variations of temperatures experienced by organisms in their natural settings (Bozinovic et al., 2016, 2011; Colinet et al., 2015; Paaijmans et al., 2013; Zeh et al.,

2015). The existing literature suggests complex patterns of biological responses to warmer conditions because of the combined effects of mean temperature and thermal fluctuations on individual performance (Colinet et al., 2015; Estay et al., 2014; Kingsolver et al., 2015). Part of this variability in thermal responses might be explained by Jensen's inequality, which applies to non-linear functions such as those inherent to thermal performance curves (Martin and Huey, 2008). Jensen's inequality posits that thermal fluctuations around a mean temperature below the thermal optimum (i.e. in the concave part of the thermal performance curve) will enhance performance, while the reverse trend should be observed once the mean temperature of the thermal regime exceeds the thermal optimum, in the convex part of the thermal performance curve (Bozinovic et al., 2011; Foray et al., 2014; Kingsolver et al., 2015; Paaijmans et al., 2013; Terblanche et al., 2010). In a context of fluctuating thermal environments, an increase in mean temperature attributable to global warming could be expected to depress organism performance through transient or permanent exposure to stressful thermal conditions during the warmest hours of the cycle (Colinet et al., 2015; Paaijmans et al., 2013; Vasseur et al., 2014). This is especially true if the increase in mean temperature is accompanied with a widening of thermal fluctuations (Paaijmans et al., 2013; Terblanche et al., 2010; Vasseur et al., 2014). Indeed, high levels of temperature variance have been shown to decrease an organism's ability to tolerate a thermal stress by limiting the expression of plastic responses and reducing the possibility for appropriate acclimation process to take place (Terblanche et al., 2010).

Responses to warming conditions might, however, greatly vary depending on the biological traits measured to infer individual performance, thereby highlighting the need for a multi-trait approach – involving measurements on a large number of traits – in global warming experiments (Fischer et al., 2011; Karl et al., 2011; Laughton et al., 2017). Indeed, fitness-related traits in a given species might differ regarding their thermal optima and

82 associated thermal performance curves (Laughton et al., 2017; Seehausen et al., 2017).
83 Experimental warming might thus yield different effects on the expression of these traits
84 depending on whether thermal conditions applied encompass supraoptimal temperatures for
85 the expression of the traits considered. For instance, in the Indian meal moth *Plodia*
86 *interpunctella* (Lepidoptera: Pyralidae), a rise in mean constant temperature from 20°C to
87 30°C led to an almost twofold decrease in developmental time from egg to adult, while
88 reducing adult lifespan after mating by 32% and increasing the count of immune cells
89 (hemocytes) in larvae hemolymph by more than 50% (Laughton et al., 2017). Furthermore,
90 biological traits are linked to each other by energetic trade-offs and thermal conditions are
91 known to influence resource allocation towards different functions. Convincing examples
92 entail trade-offs between immunity and reproduction (Adamo and Lovett, 2011) as well as
93 between immunity and heat tolerance (Fischer et al., 2011; Karl et al., 2011). Experimental
94 studies employing a multi-trait approach usually incorporate both traits linked with population
95 dynamics (e.g. development and growth rates, survival rates, egg production) and
96 interspecific relationships (e.g. defensive traits such as defensive behaviors and immunity)
97 (Iltis et al., 2018; Karl et al., 2011; Laughton et al., 2017). Indeed, the former help to predict
98 the future population dynamics of the focal species, but species will not respond in isolation
99 to climate disturbance as they are tightly connected by ecological linkages, more specifically
100 trophic interactions (Jeffs and Lewis, 2013; Van der Putten et al., 2010). Integrating defensive
101 traits involved in interspecific relationships – such as host-parasitoid associations – is thus
102 crucial to provide significant insight into how a given species might respond to global
103 warming with respect to its associated ecological context (Iltis et al., 2018). In spite of a fairly
104 robust literature about the effects of thermal fluctuations on performance in a climate change
105 perspective, still few studies have to date associated the ecological realism provided by the
106 use of realistic fluctuating regimes with the multi-trait approach, measuring different traits

related with both population dynamics and species interaction (for examples, see Fischer et al., 2014; Klockmann et al., 2016). Consequently, little is known about the effects of realistic daily thermal cycles on the expression of a broad series of traits within the framework of global warming. Filling this knowledge gap could improve our mechanistic understanding of the responses of traits, individuals and populations to facets of climate change.

In this context, we here explored the consequences of global warming simulated at a local scale for several aspects of an insect's performance. To realistically simulate global warming under controlled conditions, we built two daily fluctuating regimes: one current regime according to local climate observations and one future regime according to the fine-grain predictions provided by climate models in our focal area (Eastern France). We focused on caterpillars of the European grapevine moth *Lobesia botrana* (Lepidoptera: Tortricidae), one of the most severe pests for wine-making activity in Palearctic region (Thiéry et al., 2018). We adopted a global view of the insect's performance encompassing a large number of fitness-related traits, such as development time, pupal mass, survival rates, and the storage patterns of four energetic compounds. In a wider ecological context, we also focused on the abilities of the caterpillars to survive in agroecosystems when faced with natural enemies, especially parasitoids, by examining larval defensive strategies that involve behavioral defenses (aimed at avoiding parasitoid oviposition) and immune defenses (aimed at killing parasitoid eggs) (Iltis et al., 2018; Vogelweith et al., 2014).

We hypothesized that biological traits included in this study should respond in different ways to our simulation of future climates that might be experienced in the same location. Given its tight dependence on environmental temperature as a pure physiological rate (Rebaudo and Rabhi, 2018), we expected larval development to be greatly reduced by warming. The thermal response of larval development should in turn determine the time allowed for feeding in this capital breeding species (i.e. only caterpillars feed), thereby

negatively impacting pupal mass and energetic reserves. We expected defensive behaviors to display more complex patterns of responses to warming, as the thermal modulation of insect behavior is not purely metabolic, that is, does not only arise from short-term variation in metabolic rates or long-term variation in individual physiological state (e.g. amounts and allocation of energetic reserves) ('kinetic effects', Abram et al., 2017). It also involves intentional behavioral adjustments in response to the perception and the integration of thermal information received by thermosensory organs ('integrated effects', Abram et al., 2017). All of these possible effects were included in our experimental design which sought to investigate the overall impacts of warming on the expression of defensive behaviors to infer the future ability of caterpillars to fend off parasitoids. Finally, we postulated that immunity should be negatively affected by high temperatures because immune function in *L. botrana* caterpillars is known to peak at a relatively low thermal optimum (22°C or less) (Iltis et al., 2018).

2. Material and methods

2.1 Thermal regimes

We compared the performance of *L. botrana* caterpillars experiencing two daily fluctuating thermal regimes representative of current and future conditions occurring in Burgundy, Eastern France (Longvic-Dijon weather station, 47,27°N; 5,09°E; altitude = 219 m) (www.meteofrance.com). We focused on the summer period extending from 15th July to 15th August because this one-month period is particularly favorable for *L. botrana*, as indicated by one peak of activity (adult moths emergence) over this date range (Barnay et al., 2001; Martín-Vertedor et al., 2010). A curve of current thermal conditions was designed according to a 20-year recording of *in situ* hourly temperatures (from 1995 to 2014). Based on this curve, we built a mean daily cycle divided into six segments, each lasting 4 h and calculated

as the mean segment over the 30 days (15th July - 15th August) and the 20-year period studied (Table 1). Hereafter, this thermal regime will be referred to as the current regime.

Alongside the current regime, we built the future regime simulating the local conditions that might be found in Burgundy by the end of this century (2081-2100), during the summer period extending from 15th July to 15th August. We selected the most pessimistic of the Representative Concentration Pathways scenarios, called RCP 8.5, which predicts the highest positive radiative forcing (i.e. a high absorbance of solar and infrared radiation) associated with massive and growing emissions of greenhouse gas (www.drias-climat.fr). This scenario was combined with six simulations of high-resolution mesh (12.5 km) applied to the closest point to Dijon (Burgundy). Each simulation was obtained by downscaling a General Circulation Model (GCM) with a Regional Climate Model (RCM) to increase the spatial accuracy of climate predictions. Both GCMs and RCMs are climate tools that enable to follow the evolution of a climate system over time, at different spatial scales: global scale for GCMs, regional or local scale for RCMs nested within GCMs (for more details about the use of climate models in ecology, see Beaumont et al., 2008; Ziter et al., 2012). The six combinations of GCM/RCM used were: CERFACS-CNRM/RCA4, ICHEC-EC-EARTH/RCA4, MPI-ESM-LR/CCLM4, MPI-ESM-LR/REMO019, MPI-ESM-LR/RCA4, MetEir-ECEARTH/RACMO22E. These combinations have been recently used to generate simulations of climate change at small spatial scales across Europe (Alfieri et al., 2015; Dosio, 2016; Kotlarski et al., 2014). We used GCMs originating from the Coupled Model Intercomparison Project Phase 5 (CMIP5), a recent multi-model ensemble widely used to produce climate projections in the context of anthropogenic climate change (Taylor et al., 2012). All the RCMs used to generate downscaled simulations came from the Coordinated Regional Climate Downscaling Experiment across Europe (EURO-CORDEX) and were adapted to downscale the outputs of the GCMs used (Dosio, 2016; Jacob et al., 2014). All

180 data are available on the Drias portal (www.drias-climat.fr). For each simulation, daily
181 maximum and minimum air temperatures were extracted. We then generated six future mean
182 daily cycles (one for each simulation) based on simulated maximum and minimum
183 temperatures and the present daily cycle. The future regime, as the current regime, was
184 composed of six segments of 4 h, each calculated as the mean segment over the six
185 simulations performed (Table 1). Comparison between current and future regimes reveals that
186 global warming should increase both the mean temperature (+ 5.3°C) and the daily thermal
187 range (+ 0.6°C) in our focal area (Table 1).

188 **Table 1** Temperature conditions (°C) for the two daily fluctuating regimes used in this study:
 189 the current regime and the future regime, each divided into 6 segments of 4 h. Daily thermal
 190 range is defined as the difference between maximum and minimum daily temperatures.

	Current regime	Future regime
0h – 4h	16.4	21.4
4h – 8h	17.8	22.9
8h – 12h	22.3	27.8
12h – 16h	24.9	30.5
16h – 20h	22.9	28.3
20h – 0h	19.0	24.2
Mean temperature	20.5	25.8
Daily thermal range	8.5	9.1

191

2.2 Insect stock and larvae collection

We used larvae originating from an inbred and diapause free strain of *L. botrana* maintained at the French National Institute for Agricultural Research (INRA, Villenave d'Ornon, France) for several years under controlled conditions. This stock consists of a large number of caged adults (several thousand per week), to which wild individuals were regularly added to preserve genetic diversity. Hence, larvae coming from this artificial stock display similar patterns of defensive behaviors and basal immunity compared with wild individuals collected in French vineyards (Vogelweith et al., 2014). Moths were reared under standard conditions ($20 \pm 0.5^{\circ}\text{C}$, $60 \pm 5\%$ relative humidity, photoperiod of L17: D6 and 1 h of dusk). Luminosity was 600 lx during photophase and 100 lx at dusk. Two bands of waxed paper were provided as a support for oviposition. Every 2 days, these papers were renewed and transferred to a plastic box covered by a moist piece of paper towel to avoid egg desiccation until hatching.

A total of 2246 freshly-hatched larvae (aged < 24 h) were collected throughout the experiments with a fine brush and individually reared in Eppendorf tubes filled with 1.5 ml of artificial medium (composition for 1000 ml: 1000 ml water, 15 g agar, 86.6 g maize flour, 41.3 g wheat germ, 45.5 g beer yeast, 6 g ascorbic acid, 3.4 g mineral salt, 0.32 g Scala[®], 2.7 g benzoic acid, 2.8 g Nipagin[®] and 5 ml 95% ethanol) (Thiéry and Moreau, 2005). Neonate caterpillars were isolated in tubes so that no larval competition could occur and introduce an experimental bias (Thiéry et al., 2014). Tube caps were pierced and covered by a piece of fine mesh fabric to allow air circulation while preventing larvae from escaping. Individuals were randomly assigned to two incubators (ST 2/2 BASIC, Pol-Eko Aparatura), each simulating one of the two thermal regimes described above (temperature $\pm 0.1^{\circ}\text{C}$, $50 \pm 10\%$ relative humidity, L18: D6, 650 lx). In order to avoid a potential incubator effect, larvae and their associated thermal regime were switched between climate chambers two times a week, as well as the location of individuals inside the new incubator. Temperatures within the climate

chambers were tracked weekly with an independent data logger (Hobo, Onset Computer Corporation), and did not deviate from the thermal regimes programmed during the whole course of the experiments. Larval development was surveyed when caterpillars were swapped between incubators (i.e. two times a week) to minimize disturbance. Once they reached their fifth – and final – instar stage, larvae were randomly assigned to one of three experimental groups all along the experiments, because the different measurements of biological traits involved in this study cannot be taken on a single individual. We focused on the fifth instar because the energetic resources accumulated during the whole larval stage can be quantified at this late larval instar that directly precedes pupation. In addition, measurements of behavioral and immune defenses against parasitoids are classically performed at the end of larval development in this species (Iltis et al., 2018; Vogelweith et al., 2014). In Group 1, the fifth instar larvae were left until pupation in order to measure development time, pupal mass, survival from first instar larva to adult stage, and sex ratio. In Group 2, the fifth instar larvae were frozen and stored to quantify later their energy budgets. In Group 3, the behavioral and immune traits involved in defense of fifth instar caterpillars against larval parasitoids were measured.

2.3 Group 1: development time, pupal mass, survival and sex ratio

A total of 2043 freshly hatched larvae were initially assigned to Group 1 in order to develop until pupation ($n = 981$ individuals for current regime, $n = 1062$ individuals for future regime). We recorded the time (in days) elapsed between hatching and pupation (i.e. the development time of the larval stage), which corresponds to the time during which caterpillars are able to feed. The pupae were collected carefully with fine forceps and weighed (± 0.1 mg) with a balance (Pioneer PA214C, OHAUS). Development time and pupal mass were only recorded for individuals that reached the adult stage to discriminate the results between males and females, which were sexed upon emergence. We also calculated survival (number of

emerged adults relative to the number of freshly hatched larvae initially assigned to each thermal regime), and sex ratio (expressed as the proportion of males in the adult population).

2.4 Group 2: energetic assessments

These measurements initially involved 59 fifth instar larvae ($n = 29$ for current regime, $n = 30$ for future regime). Caterpillars were cleaned with a fine brush to remove any residual of artificial medium from the surface of their body, transferred to Eppendorf tubes, frozen in liquid nitrogen and kept at -80°C for latter quantification of their energy budgets. They were subsequently weighed to obtain fresh body mass (± 0.01 mg) with a microbalance (Quintix 35-15, Sartorius), before being crushed with stainless steel beads (tubes shaken for 90 s at 25 Hz) in 180 μl of aqueous lysis buffer solution (composition: 100 mM KH_2PO_4 , 1 mM dithiothreitol DTT, 1 mM ethylenediaminetetraacetic EDTA, pH = 7.4). Biochemical assays were run on four metabolites potentially involved in energy production: proteins, lipids, soluble carbohydrates and glycogen. The total amount of soluble proteins available was estimated using a DC Protein Assay kit (Bio-Rad). We used the spectrophotometry method described by Foray et al. (2012) for the quantification of lipids, soluble carbohydrates and glycogen.

Crushed samples were placed on ice for 30 min to ensure proteins solubilization, centrifuged (4000 g, 5 min, 4°C) and 5 μl of supernatant from each sample were pipetted to microplate wells together with 35 μl of lysis buffer to obtain diluted samples. Particular attention was paid to avoid collecting lipids (which often form a thin surface layer on the top of the sample) when pipetting the 5 μl of supernatant for protein assays, because lipids might disturb the absorbance readings in the Bradford approach used to quantify proteins (Foray et al., 2012). Then, 5 μl of these diluted samples were transferred to new microplate wells and supplemented with 225 μl of Bradford micro-assay reagent (DC Protein Assay Kit, Bio-Rad).

266 Reaction was allowed to proceed for 15 min at room temperature, after which absorbance
267 readings were taken at 750 nm. Protein concentration in the samples was estimated with a
268 standard curve based on a dilution-series of bovine gamma globulin standard (Pierce, Thermo
269 Fisher Scientific) dissolved in lysis buffer. This dilution-series was constituted by the
270 following protein concentration values: 0, 0.062, 0.125, 0.25, 0.5, 1 and 2 mg.ml⁻¹. Finally, 5
271 µl of lysis buffer, 20 µl of 20 % sodium sulphate solution and 1.5 ml of chloroform-methanol
272 mixture (1:2 v/v) were added to each crushed sample to solubilize total lipids and water-
273 soluble carbohydrates. These samples were then conserved at room temperature for 24h
274 before biochemical assays of the amounts of lipids, soluble carbohydrates and glycogen.

275 At the end of this 24h duration, homogenates were centrifuged at room temperature
276 (4800 g, 6 min). Centrifugation enabled to isolate lipids and soluble carbohydrates
277 (supernatant) from glycogen (pellet). Determination of lipid body amount was performed with
278 a colorimetric method involving vanillin reagent (Alfa Aesar, Thermo Fischer Scientific). For
279 each centrifuged sample, 75 µl of supernatant were transferred to an Eppendorf tube and
280 heated at 90°C until complete solvent evaporation. Then, each tube was supplemented with 40
281 µl of 95-98% sulphuric acid and incubated at 90°C for 10 min. For lipids revelation, 960 µl of
282 vanillin solution (composition: 100 ml ultrapure water, 400 ml 85% orthophosphoric acid,
283 600 mg vanillin) were then deposited in each tube. After vigorous vortexing, 200 µl of each
284 tube were pipetted to microplate wells for absorbance measurements, which were taken at 525
285 nm. The dilution-series used to assess lipid concentration in the samples involved glyceryl
286 trioleate as a standard (Sigma-Aldrich) dissolved in chloroform, and was comprised by the
287 following concentration values: 0, 2.5, 5, 10, 20, 40 and 80 µg.ml⁻¹. The body amounts of
288 both soluble carbohydrates and glycogen were determined with a colorimetric method based
289 on anthrone reagent (Alfa Aesar, Thermo Fisher Scientific). For soluble carbohydrates, 750 µl
290 of supernatant from each centrifuged sample were collected and deposited in an Eppendorf

tube, before being allowed to fully evaporate at 95°C. For glycogen, the remaining supernatant in the centrifuged samples was removed and the pellet was left at room temperature during 30 min for complete solvent evaporation. Then, 2 ml of anthrone solution (composition: 150 ml ultrapure water, 380 ml 95-98% sulfuric acid, 750 mg anthrone) were added to each tube (containing either soluble carbohydrates or glycogen), and all these tubes were heated at 95°C for 17 min. Tubes were subsequently vortexed and 200 µl of each were withdrawn and deposited in microplate wells before readings of optical density, which were performed at 625 nm. The concentration of these two energetic compounds in the samples was estimated through comparison with a dilution-series constituted by the following concentration values of glucose (Sigma-Aldrich): 0, 6.25, 12.5, 25, 50, 100, 200 and 400 µg.ml⁻¹.

For proteins and lipids, two replicates were performed and individuals for which the coefficient of variation exceeded 30% were excluded from further analysis. Repeatability was relatively high, as indicated by the low percentages of individuals removed from the protein (8.5%) and the lipid datasets (5.1%). At the end of this selection procedure, we retained 54 larvae from the protein dataset ($n = 27$ for current regime, $n = 27$ for future regime) and 56 larvae from the lipid dataset ($n = 29$ for current regime, $n = 27$ for future regime).

2.5 Group 3: behavioral and immune defenses

A total of 144 fifth instar larvae were assigned to measurements of defensive traits ($n = 73$ individuals for current regime, $n = 71$ individuals for future regime). Three successive behaviors were quantified, always in the same order ('flee', 'twisting' and 'dropping'), in accordance with the sequence involved in defense against larval parasitoids in natural conditions (Vogelweith et al., 2014). Additionally, these behavioral tests were performed at an ambient temperature strictly identical to the mean temperature of the regime considered:

315 20.5°C if the tested individual was reared in current regime, or 25.8°C if it was reared in
316 future regime (Table 1). Temperature in the testing rooms was controlled with Hobo data
317 loggers. After the completion of each behavioral test, 20 min of rest were given to each larva
318 in a clean Eppendorf tube drilled to allow air circulation before the next test started.

319 The first behavior studied, called ‘flee’, refers to larvae escape ability by moving away
320 from a generic threat without leaving the bunch they feed on. Here, we estimated the
321 locomotor activity (movement speed) of caterpillars as a proxy of ‘flee’ behavior, to infer
322 their ability to evade parasitoids in natural conditions. For this purpose, caterpillars were
323 installed on a plastic squared sheet (84 x 116 cm, squares size: 1 x 1 cm) and acclimated for
324 30 s under the lid of a Petri dish. The lid was then removed and each larva was observed to
325 count the total number of lines crossed during 60 s (the minimum duration required by a larva
326 to exit the sheet). As this behavior occurs before physical contact between the caterpillar and
327 the parasitoid, no simulation of a parasitoid attack was needed to cause the larvae to flee.
328 Locomotor activity was naturally induced by the stress associated with the manipulation of
329 caterpillars with a fine paintbrush during the extraction from their rearing tube, and their
330 exposure to a new environment during behavioral tests.

331 The second behavior, ‘twisting’, consists of multiple and rapid torsions in response to
332 contact between a larva’s body and a parasitoid sting. We mimicked a parasitoid attack by
333 touching the dorsal part of the caterpillar with a fine brush, after 30 s of acclimation under the
334 lid of a Petri dish. Each individual was touched four times in 1 min. The entire sequence was
335 video recorded (HDR-CX220E, Sony) and further analyzed in slow motion with Kinovea
336 0.8.15 software to score the total number of torsions over the four touches.

337 Finally, we examined ‘dropping’ behavior, which describes how caterpillars faced
338 with a foraging parasitoid can drop and weave a silk thread to return to their living patch once

the parasitoid has left it. Larvae were installed at the top of a 150 cm high bracket. After an acclimation of 30 s under a reversed Eppendorf tube, each larva was touched with a fine brush (as described in the previous paragraph) until it fell. For caterpillars weaving a thread, we measured its length with a ruler (precision ± 3 mm) after 5 s of stabilization. Thread length is supposed to be positively linked to larval escape ability (Vogelweith et al., 2014).

Once the three behavioral tests were carried out, larvae were returned to their rearing tube and associated thermal regime for 3 h before the immune measurements were taken. The ability of the larvae to kill parasitoid eggs was estimated by measuring hemocyte concentration and activity of the phenoloxidase (PO). Hemocytes are immune cells that freely circulate in hemolymph of insects and bind to parasitoid eggs. Hemocytes release the PO enzyme that catalyzes melanization reactions occurring around parasitoid eggs as part of the immune response (Lavine and Strand, 2002). Caterpillars were not challenged prior to these immune tests, because they naturally display high levels of basal immunity and do not significantly respond to an infection (Vogelweith et al., 2014). Larvae were chilled on ice for 30 min and then photographed using a stereomicroscope at 12.5x magnification (Stemi 508, Zeiss) to measure the distance between the most distant margins of the head capsule (precision: ± 0.1 μ m) as a reliable estimator of larval body size (Delbac et al., 2010). For this experimental group, we measured larval body size rather than body mass since body size is known to correlate with some of the larval defenses in this species (Iltis et al., 2018) and because caterpillars cannot be accurately weighed while alive (Muller et al., 2015). One 2 μ l sample of hemolymph was then collected from a prick in the dorsal part of the abdomen with a cold sterile glass capillary (Hirschmann Laborgeräte, Eberstadt). This sample was diluted in 20 μ l of cold phosphate-buffered saline (PBS, 8.74 g NaCl, 1.78 g Na₂HPO₄, 1000 ml distilled water, pH 6.5). A volume of 10 μ l of this solution was immediately withdrawn to estimate hemocyte concentration with a Neubauer improved hemocytometer under a phase

contrast microscope at 400x magnification (Primo Star, Zeiss). The remaining 12 μ l of diluted hemolymph were supplemented with 10 μ l of PBS. These samples were frozen in liquid nitrogen and stored at -80°C for the latter estimation of PO activity.

We distinguished between functional PO naturally activated in hemolymph (PO activity) and total PO including functional PO and proenzymes (total PO activity). These measurements were based on the spectrophotometry method described by Vogelweith et al. (2013). Hemolymph samples were allowed to thaw very slowly on ice and centrifuged (4000 g, 15 min, 4°C) before 5 μ l of supernatant were transferred to microplate wells. These samples were supplemented with either 160 μ l of diluted PBS solution (35 ml ultrapure water, 5 ml filtered PBS) for PO activity measurement, or with 160 μ l of chymotrypsin solution (35 ml ultrapure water, 5 ml filtered PBS, 2.45 mg trypsin) for total PO activity measurement. Finally, 20 μ l of L-Dopa solution (40 ml ultrapure water, 160 mg L-Dopamine) were deposited in each well as a colorimetric substrate. An enzymatic reaction proceeded for 40 min at 30°C in a microplate reader (Versamax, Molecular Devices), during which the readings were taken every 15 s at 490 nm. Data were subsequently analyzed with the SOFT-Max Pro 4.0 software. We recorded maximum enzyme activity (V_{\max}), corresponding to the slope of the absorbance curve during its linear phase. All immune assessments were reported to 1 μ l of pure hemolymph.

2.6 Statistical analyses

Development time and pupal mass were normally distributed and displayed homogenous variances (homoscedasticity) between thermal regimes and sexes. Consequently, they were analyzed by means of two-way Analyses of Variance (ANOVAs) integrating thermal regime, sex and their interaction. Survival and sex ratio were expressed in proportions and therefore compared among thermal regimes with Generalized Linear Models (GLMs)-binomial error.

To control for allometric effects on larval physiological and behavioral traits, we included either larval body mass (for energetic reserves) or larval body size (for defensive behaviors and immunity) as a covariate in the statistical models, alongside thermal regime as an explanatory variable. Most of the data were normally distributed (the reserves of the four energetic compounds, hemocyte concentration, total PO activity) or required a logarithm transformation to satisfy normality assumption (thread length and PO activity), and their variances were homogenous between thermal regimes. Accordingly, all these data were studied with analyses of covariance (ANCOVAs) testing for homogeneity of slopes and intercepts, as these models integrated the effects of the explanatory variable (thermal regime), the associated covariate (larval body mass or larval body size) and their interaction. As overdispersed count data, the number of lines caterpillars crossed and the number of twists were analyzed with GLM-negative binomial error. Normality and homoscedasticity were assessed with a Shapiro-Wilk and a Levene test, respectively. For count data, overdispersion (observed variance greater than the theoretical one predicted by the fitted model) was detected with a postregression likelihood ratio test (Cameron and Trivedi, 1990). All statistical analyses were carried out using R 3.5.2 software.

3. Results

3.1 Group 1: development time, pupal mass, survival and sex ratio

Neither development time (two-way ANOVA: $F_{1,832} = 0.04$, $P = 0.84$) nor pupal mass (two-way ANOVA: $F_{1,832} = 0.39$, $P = 0.53$) was impacted by the interaction between thermal regime and sex. Development time was modulated by thermal regime ($F_{1,832} = 639$, $P < 0.001$) and sex ($F_{1,832} = 26.5$, $P < 0.001$) in an additive way: individuals reared in future, warmer conditions needed less time to complete their larval development than those reared in current, cooler conditions, and females developed slower than males (Table 2). Pupal mass was

412 affected by sex ($F_{1,832} = 1308$, $P < 0.001$) but not by thermal regime ($F_{1,832} = 0.67$, $P = 0.41$),
413 as females being significantly heavier than males (Table 2). Survival was influenced by
414 thermal regime (GLM-binomial error: $\chi^2_1 = 10.6$, $P < 0.001$): larvae reared in warmer
415 conditions were more prone to survive until the adult stage than those held in cooler
416 conditions (Table 2). Adult sex ratio did not differ between thermal regimes (GLM-binomial
417 error: $\chi^2_1 = 0.02$, $P = 0.88$) (Table 2).

418 **Table 2** Mean values and 95% confidence intervals of development time (time between egg hatching and pupation), pupal mass, survival (from
 419 larval to adult stage), and sex ratio (proportion of males in the adult population) for the two thermal regimes. Different letters indicate significant
 420 differences for a given trait ($P < 0.05$).

		Current regime	Future regime
Development time (days)	Males	24.6 [24.0; 25.2] ^a	16.9 [16.4; 17.3] ^b
	Females	26.3 [25.5; 27.0] ^c	18.4 [17.8; 19.0] ^d
Pupal mass (mg)	Males	9.00 [8.9; 9.2] ^a	9.00 [8.9; 9.1] ^a
	Females	12.8 [12.5; 13.1] ^b	12.7 [12.4; 12.9] ^b
Survival (%)		43.5 [38.4; 48.6] ^a	48.4 [43.9; 52.9] ^b
Sex ratio (%)		59.3 [54.2; 64.4]	58.5 [54.0; 63.0]

421 3.2 Group 2: energetic assessments

422 None of the four energetic compounds was affected by the interaction between thermal
423 regime and larval body mass (Table 3). Among these compounds, only the body amount of
424 lipids was influenced by thermal regime, while the amounts of proteins, soluble carbohydrates
425 and glycogen remained the same across thermal regimes (Table 3, Fig. 1A, B, C, D).
426 Caterpillars reared in warmer conditions stocked less lipid reserves at the end of their larval
427 development relative to their counterparts reared in cooler conditions (Fig. 1B). Protein and
428 lipid body amounts were both affected positively by larval body mass (Table 3, Fig. 1A, B):
429 heaviest larvae accumulated more reserves than lightest ones (proteins: slope = 44.1; lipids:
430 slope = 32.1).

431 **Table 3** Effects of thermal regime, larval body mass and their interaction on the total body amounts of the four energetic compounds. All data
 432 were studied with analyses of covariance testing for homogeneity of slopes and intercepts. Bold values indicate significant effects ($P < 0.05$).

	Thermal regime		Larval body mass		Regime * Body mass	
	<i>Test value</i>	<i>P</i>	<i>Test value</i>	<i>P</i>	<i>Test value</i>	<i>P</i>
Proteins	$F_{1,50} = 1.72$	0.20	$F_{1,50} = \mathbf{50.0}$	$< \mathbf{0.001}$	$F_{1,50} = 0.001$	0.98
Lipids	$F_{1,52} = \mathbf{5.69}$	$\mathbf{0.02}$	$F_{1,52} = \mathbf{5.34}$	$\mathbf{0.02}$	$F_{1,52} = 1.63$	0.21
Soluble carbohydrates	$F_{1,55} = 0.03$	0.86	$F_{1,55} = 1.50$	0.23	$F_{1,55} = 1.64$	0.20
Glycogen	$F_{1,55} = 2.36$	0.13	$F_{1,55} = 1.98$	0.16	$F_{1,55} = 0.52$	0.47




Fig. 1 Effect of thermal regime (circles and solid lines: current regime, triangles and dashed lines: future regime) in relation with larval fresh body mass on the total body amounts of proteins (A), lipids (B), soluble carbohydrates (C), and glycogen (D). Shaded area around each line corresponds to the 95% confidence interval for the predicted values of the linear regression. Reported *P* values refer to the simple effect of thermal regime on each energetic compound (testing homogeneity of intercepts between solid and dashed lines).

3.3 Group 3: behavioral and immune defenses

None of the behavioral or immune traits was impacted by the interaction between thermal regime and larval body size (Table 4). Among the behavioral traits examined, only the number of lines crossed by caterpillars during ‘flee’ tests was influenced by thermal regime: larvae reared and tested at warmer conditions crossed significantly more lines relative to individuals originating from cooler conditions (Table 4, Fig. 2A). Thermal regime did not affect ‘twisting’ (number of twists) and ‘dropping’ (length of the silk thread woven) behaviors (Table 4, Fig. 2B, C). From an immunological perspective, hemocyte concentration and PO activity were not impacted by thermal regime (Table 4, Fig. 3A, B). However, total PO activity was significantly modulated by thermal regime: larvae reared in warmer conditions displayed lower levels of total PO activity compared with those held in colder conditions (Table 4, Fig. 3C). None of the behavioral and immune traits was modulated by larval body size (Table 4).

453 **Table 4** Effects of thermal regime, larval body size (head capsule width) and their interaction on different defensive behaviors and immune traits
 454 involved in defenses of caterpillars against larval parasitoids. Bold values indicate significant effects ($P < 0.05$).

	Thermal regime		Larval body size		Regime * Body size	
	<i>Test value</i>	<i>P</i>	<i>Test value</i>	<i>P</i>	<i>Test value</i>	<i>P</i>
<i>Defensive behaviors</i>						
Number of lines ^a	$\chi^2_1 = \mathbf{6.88}$	0.02	$\chi^2_1 = 0.24$	0.67	$\chi^2_1 = 0.50$	0.54
Number of twists ^a	$\chi^2_1 = 0.64$	0.35	$\chi^2_1 = 4.10^{-5}$	0.99	$\chi^2_1 = 0.80$	0.30
Thread length ^b	$F_{1,96} = 1.58$	0.21	$F_{1,96} = 1.59$	0.21	$F_{1,96} = 0.05$	0.82
<i>Immune traits</i>						
Hemocyte concentration ^b	$F_{1,72} = 0.64$	0.43	$F_{1,72} = 0.08$	0.78	$F_{1,72} = 0.67$	0.42
PO activity ^b	$F_{1,72} = 3.71$	0.06	$F_{1,72} = 0.89$	0.35	$F_{1,72} = 0.70$	0.41
Total PO activity ^b	$F_{1,72} = \mathbf{9.66}$	< 0.001	$F_{1,72} = 1.28$	0.26	$F_{1,72} = 0.39$	0.53

455 ^a Generalized linear model-negative binomial error

456 ^b Analysis of covariance




Fig. 2 Effect of thermal regime on the mean (\pm 95% c.i.) of the number of lines crossed by larvae during ‘flee’ tests (A), the number of twists over four touches (B), and the length of the silk thread spun (C). Asterisks highlight significant differences (*** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$, n.s. non-significant) and numbers refer to sample sizes.




Fig. 3 Effect of thermal regime on the mean (\pm 95% c.i.) of hemocyte concentration (A), phenoloxidase activity (B), and total phenoloxidase activity (C). All these immune measurements were reported to 1 μl of pure hemolymph collected from fifth instar caterpillars. Asterisks highlight significant differences (*** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$, n.s. non-significant) and numbers refer to sample sizes.

4. Discussion

Within the framework of global warming, we here examined the responses of different fitness-related traits in caterpillars of the moth *L. botrana* to realistic daily fluctuating regimes reflecting current and future conditions in Eastern France. Many of the traits inspected were affected by our experimental imitation of climate change. Warming positively influenced caterpillars' development (leading to a 31% reduction in the time needed to complete all five larval instars), survival from first instar larva to adult stage (+19%) and larval ability to escape natural enemies through the expression of 'flee' behavior (+60%). Conversely, rising temperatures led to depletion of lipid reserves (-26%) and reduced total PO activity (-34%). Such a diversity of thermal responses among traits strengthens the importance of an integrative approach involving measurements on a wide array of traits to accurately forecast the consequences of global warming for an organism's performance, and the way it will interact with adjacent trophic levels.

Some traits related with population dynamics were positively affected by rising temperatures, presumably indicating an overall increase in caterpillars' performance under a simulated global warming scenario. Indeed, caterpillars experiencing higher, future temperatures developed faster and showed increased survival rates compared with individuals kept under thermal conditions similar to those currently recorded in Eastern France. These findings remain in accordance with previous studies showing that the thermal optimum for the development of *L. botrana* larvae – including the laboratory strain used in this study – ranges from 28°C to 30°C (Briere and Pracros, 1998; Iltis et al., 2018). Hence, the experimental warming applied here (mean temperature increasing from 20.5 to 25.8°C) positively impacted development rates by bringing environmental temperatures closer to the thermal optimum of *L. botrana*. Similarly to a previous work conducted on the same species (Thiéry and Moreau, 2005), we found that faster development was accompanied with higher survival rates. One

plausible explanation to these results could involve a phenomenon known as ‘developmental resistance’, whereby disease resistance usually increases throughout larval stages in some butterfly species (McNeil et al., 2010; Shikano et al., 2018). Hence, caterpillars may benefit from a faster development under warmer conditions by spending shorter in the early ontogenetic stages, which are the most susceptible to mortality driven by pathogenic infections (Valadez-Lira et al., 2012).

As a capital breeding species, we would have expected *L. botrana* individuals to incur costs arising from a shrunk larval stage (i.e. a reduced time allowed for feeding) in warmer conditions, especially regarding pupal mass. Contrary to these initial expectations, pupal mass did not differ between thermal regimes. These results confirm that *L. botrana* caterpillars could compensate for a shortened feeding period under hotter conditions through an increase in daily gain of body mass (i.e. growth rate) (Iltis et al., 2018). The underlying mechanisms to support a high growth rate in warm conditions could involve an increased food intake rate and/or an enhanced efficiency at converting food into body matter when exposed to hotter conditions (Bauerfeind and Fischer, 2013; Lee and Roh, 2010; Lemoine et al., 2014). Nonetheless, the 26% decrease in the body amounts of lipids observed in warmer conditions seems to indicate that larvae cannot compensate for the effects of warming on energetic budgets, with costs being specifically apparent on fat reserves (the three other energetic compounds remaining unaffected by temperature). Interestingly, previous work also recorded a reduced lipid storage efficiency for fast developing caterpillars in *Spodoptera exigua* (Lepidoptera: Noctuidae), suggesting that high development rates induced by warming may be accompanied with a decline in lipid reserves (Lee and Roh, 2010). These findings indicate that the lipid energetic compartment could be particularly susceptible to the detrimental effects of a reduced feeding duration caused by an accelerated development under warmer conditions.

Aside from traits related with population dynamics, thermal conditions modulated the expression of several defensive traits involved in trophic interactions between *L. botrana* caterpillars and their natural enemies, especially the guild of parasitoids. Concerning behavioral traits, it is not possible to identify the mechanisms underlying the thermal response of ‘flee’ behavior at this stage. It may arise from long-term effects, whereby temperature met throughout larval development might have impacted the physiological state of caterpillars and their resource partitioning between competing defensive traits (Abram et al., 2017). It could also be attributable to short-term effects related with temperature prevailing during behavioral tests, which might have influenced caterpillars’ metabolic rates and/or triggered thermosensory processes leading to behavioral adjustments following perception of heat conditions (Abram et al., 2017; Soto-Padilla et al., 2018). In terms of trophic dynamics, our findings indicate that increasing temperatures should enhance the ability of *L. botrana* caterpillars to escape from natural enemies through upregulation of ‘flee’ behavior. This does not, however, necessarily imply a lower larval susceptibility to parasitism, provided that the behaviors specifically expressed in case of an attempt of parasitoid oviposition (‘twisting’ and ‘dropping’) remained unaffected by thermal conditions.

Concerning immune traits, total PO activity was the only immune trait affected (negatively) by temperature, congruently with a previous work on *L. botrana* (Iltis et al., 2018) and our hypothesis that global warming could induce immunosuppressive effects in larvae of this species. Intriguingly, the decrease in total PO activity was not explained by a concomitant lowering of hemocyte concentration, which remained constant across thermal treatments in our study. Hemocyte populations are composed by different cell types (four types have been identified in larvae of the moth *Eupoecilia ambiguella*, a relative of *L. botrana*), each having a particular role in immunity (Lavine and Strand, 2002; Vogelweith et al., 2016). In this regard, we could postulate that thermal conditions might influence the

relative abundances of hemocyte types in *L. botrana* larvae, tending to decrease the proportion of cells involved in synthesis of PO zymogens (e.g. oenocytoids). Our findings also indicate that, despite no effect on basal immunity (PO activity), warming could significantly reduce the ability of *L. botrana* caterpillars to respond to a parasitic infection because hotter rearing conditions decreased total PO activity. Indeed, the activation of PO cascade system is required to catalyze melanization of parasitoid eggs (González-Santoyo and Córdoba-Aguilar, 2012; Kanost and Gorman, 2008). However, it is worth noting that, in our immunological assays, immune reactions were allowed to proceed at standard temperature. Considering that, in lepidopteran species, the kinetics of PO enzyme will likely increase alongside larval body temperature up to a certain limit (Xue et al., 2006), it remains possible that larvae reared in warmer conditions invested less in the production of the active form of PO as an adaptive choice to keep the overall enzymatic activity constant across temperatures. Further insights into the functional consequences of this variation in larval immunity – for instance by measuring the *in vivo* ability of caterpillars to encapsulate and melanize foreign bodies at different temperatures – would be highly valuable to predict more thoroughly the impacts of global warming on caterpillars' survival in context of high parasitism pressure as recorded in vineyards.

5. Conclusion

In summary, our study brings new lines of evidence to suggest that multiple traits in a given species may differ in their sensitivity to global warming, underlying complex consequences for population dynamics and trophic interactions. Here, we found that, in terms of population dynamics, warming brought *L. botrana* closer to its thermal optimum for larval development. This should lead to a shorter generation time and therefore a higher future voltinism (number of generations per growing season) of *L. botrana* in Eastern France, as recently observed in meridional areas (Martín-Vertedor et al., 2010). To what extent these benefits might be

mitigated by the reduction of lipid reserves observed under warmer conditions remains not well appreciated for now. A decline in lipids stock might, for instance, negatively impact the survival of the naturally diapausing generation of *L. botrana*, because fat reserves are crucial to survive extended non-feeding periods (Sinclair and Marshall, 2018). It may also persist into the adult stage and adversely impact adult dispersal ability and reproduction (Vande Velde and Van Dyck, 2013). Finally, we found that key traits involved in the trophic interactions between *L. botrana* caterpillars and their larval parasitoids were altered by warming. At this stage, predicting the consequences for future larval mortality induced by natural enemies in agroecosystems is somewhat challenging, provided that the direction of the effect of warming differed among defensive traits, and that parasitoids will also be impacted by rising temperatures, from both physiological and behavioral perspectives (Hance et al., 2007).

Several considerations should, however, be carefully examined to provide a more integrated picture of the impacts of warming for *L. botrana*, with respect to its evolutionary and ecological context. First, our study specifically investigated the plastic response occurring within one generation of *L. botrana* larvae exposed to an abrupt environmental warming. In natural conditions climate disturbance will more likely translate into a gradual shift in thermal conditions whose effects will spread across insect life stages and generations. For this reason, it is now widely acknowledged that the response of insect species to global warming will not solely be determined by intragenerational plasticity, but also by transgenerational plasticity (whereby temperatures faced by one generation will influence the expression of a phenotype in the subsequent generations) in concert with evolutionary processes (Donelson et al., 2018; Sgrò et al., 2016). It would therefore be worth investigating whether these different types of response might interact to shape the insect phenotype when experiencing gradual warming over several successive generations. Second, to what extent the mosaic of microclimatic conditions existing at the vine plant and grape bunch levels might create opportunities for

larval behavioral thermoregulation (through microsite selection) to alleviate the effects of warming must be taken into consideration. Apparently homogenous landscapes are actually composed of an important diversity of thermal conditions at much smaller, biologically relevant spatial scale matching the body size of the insect (Potter et al., 2013; Terblanche et al., 2015). Thus, variation in air temperature recorded in a given site at a macroclimatic scale may bear little relationship to the variation in the operative body temperature of the insect experiencing a set of contrasting microclimatic conditions. In vineyards, grape berry temperature displays an important variability both between grape bunches and within a single bunch, in relation with factors determining direct solar radiation (e.g. bunch exposure, shading provided by leaf canopy), bunch compactness and architecture (Haselgrove et al., 2000; Kiaeian Moosavi et al., 2018; Tello and Añez, 2017). Small scale variation in berry temperature due to sunlight exposure has been shown to deeply impact *L. botrana* performance, particularly in the early ontogenetic stages (egg hatching, larval settlement) (Kiaeian Moosavi et al., 2018). Incorporating this natural thermal complexity at microhabitat scales and the way it would be altered by a rapidly changing climatic context could therefore become a next step towards a more thorough understanding of the consequences of global warming for this insect pest.

Competing interests

The authors declare no conflict of interests.

Acknowledgements

We gratefully acknowledge L. Delbac who managed the insect stock and E. Parfitt for English revision. We also want to thank L. Chevallier, P. Hübner, R. Kuzmin, M. Pêcheur, R. Quinaux and A. Rathery for their practical help in the experiments.

Funding

This work was supported by the Conseil Régional de Bourgogne Franche-Comté (PARI, FABER LOUAPRE-AGREE-BGS and VALEACLIM-BOIS), and the European Union (PO FEDER-FSE Bourgogne 2014/2020 programs).

References

- Abram, P.K., Boivin, G., Moiroux, J., Brodeur, J., 2017. Behavioural effects of temperature on ectothermic animals: unifying thermal physiology and behavioural plasticity. *Biol. Rev.* 92, 1859–1876. <https://doi.org/10.1111/brv.12312>
- Adamo, S.A., Lovett, M.M.E., 2011. Some like it hot: the effects of climate change on reproduction, immune function and disease resistance in the cricket *Gryllus texensis*. *J. Exp. Biol.* 214, 1997–2004. <https://doi.org/10.1242/jeb.056531>
- Alfieri, L., Feyen, L., Dottori, F., Bianchi, A., 2015. Ensemble flood risk assessment in Europe under high end climate scenarios. *Glob. Environ. Chang.* 35, 199–212. <https://doi.org/10.1016/J.GLOENVCHA.2015.09.004>
- Angilletta, M.J., 2009. Thermal adaptation: a theoretical and empirical synthesis, Oxford University Press, Oxford.
- Barnay, O., Hommay, G., Gertz, C., Kienlen, J.C., Schubert, G., Marro, J.P., Pizzol, J., Chavigny, P., 2001. Survey of natural populations of *Trichogramma* (Hym., Trichogrammatidae) in the vineyards of Alsace (France). *J. Appl. Entomol.* 125, 469–477. <https://doi.org/10.1046/j.1439-0418.2001.00575.x>
- Bauerfeind, S.S., Fischer, K., 2014. Simulating climate change: temperature extremes but not means diminish performance in a widespread butterfly. *Popul. Ecol.* 56, 239–250. <https://doi.org/10.1007/s10144-013-0409-y>
- Bauerfeind, S.S., Fischer, K., 2013. Increased temperature reduces herbivore host-plant

640 quality. Glob. Chang. Biol. 19, 3272–3282. <https://doi.org/10.1111/gcb.12297>

641 Beaumont, L.J., Hughes, L., Pitman, A.J., 2008. Why is the choice of future climate scenarios
 642 for species distribution modelling important? Ecol. Lett. 11, 1135–1146.
 643 <https://doi.org/10.1111/j.1461-0248.2008.01231.x>

644 Bozinovic, F., Bastías, D.A., Boher, F., Clavijo-Baquet, S., Estay, S.A., Angilletta, M.J.,
 645 2011. The mean and variance of environmental temperature interact to determine
 646 physiological tolerance and fitness. Physiol. Biochem. Zool. 84, 543–552.
 647 <https://doi.org/10.1086/662551>

648 Bozinovic, F., Medina, N.R., Alruiz, J.M., Cavieres, G., Sabat, P., 2016. Thermal tolerance
 649 and survival responses to scenarios of experimental climatic change: changing thermal
 650 variability reduces the heat and cold tolerance in a fly. J Comp Physiol B 186, 581–587.
 651 <https://doi.org/10.1007/s00360-016-0980-6>

652 Briere, J.-F., Pracros, P., 1998. Comparison of temperature-dependent growth models with the
 653 development of *Lobesia botrana* (Lepidoptera: Tortricidae). Environ. Entomol. 27, 94–
 654 101. <https://doi.org/10.1093/ee/27.1.94>

655 Cameron, A.C., Trivedi, P.K., 1990. Regression-based tests for overdispersion in the Poisson
 656 model. J. Econom. 46, 347–364. [https://doi.org/10.1016/0304-4076\(90\)90014-K](https://doi.org/10.1016/0304-4076(90)90014-K)

657 Clarke, A., 2003. Costs and consequences of evolutionary temperature adaptation. Trends
 658 Ecol. Evol. 18, 573–581. <https://doi.org/10.1016/J.TREE.2003.08.007>

659 Colinet, H., Sinclair, B.J., Vernon, P., Renault, D., 2015. Insects in fluctuating thermal
 660 environments. Annu. Rev. Entomol. 60, 123–140. <https://doi.org/10.1146/annurev-ento-010814-021017>

662 Delbac, L., Lecharpentier, P., Thiery, D., 2010. Larval instars determination for the European

663 grapevine moth (Lepidoptera: Tortricidae) based on the frequency distribution of head-
664 capsule widths. *Crop Prot.* 29, 623–630. <https://doi.org/10.1016/j.cropro.2010.01.009>

665 Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C.,
666 Martin, P.R., 2008. Impacts of climate warming on terrestrial ectotherms across latitude.
667 *Proc. Natl. Acad. Sci. USA.* 105, 6668–6672. <https://doi.org/10.1073/pnas.0709472105>

668 Donelson, J.M., Salinas, S., Munday, P.L., Shama, L.N.S., 2018. Transgenerational plasticity
669 and climate change experiments: where do we go from here? *Glob. Chang. Biol.* 24, 13–
670 34. <https://doi.org/10.1111/gcb.13903>

671 Dosio, A., 2016. Projections of climate change indices of temperature and precipitation from
672 an ensemble of bias-adjusted high-resolution EURO-CORDEX regional climate models.
673 *J. Geophys. Res. Atmos.* 121, 5488–5511. <https://doi.org/10.1002/2015JD024411>

674 Estay, S.A., Lima, M., Bozinovic, F., 2014. The role of temperature variability on insect
675 performance and population dynamics in a warming world. *Oikos* 123, 131–140.
676 <https://doi.org/10.1111/j.1600-0706.2013.00607.x>

677 Fischer, K., Klockmann, M., Reim, E., 2014. Strong negative effects of simulated heat waves
678 in a tropical butterfly. *J. Exp. Biol.* 217, 2892–2898. <https://doi.org/10.1242/jeb.106245>

679 Fischer, K., Kölzow, N., Höltje, H., Karl, I., 2011. Assay conditions in laboratory
680 experiments: is the use of constant rather than fluctuating temperatures justified when
681 investigating temperature-induced plasticity? *Oecologia* 166, 23–33.
682 <https://doi.org/10.1007/s00442-011-1917-0>

683 Foray, V., Desouhant, E., Gibert, P., 2014. The impact of thermal fluctuations on reaction
684 norms in specialist and generalist parasitic wasps. *Funct. Ecol.* 28, 411–423.
685 <https://doi.org/10.1111/1365-2435.12171>

686 Foray, V., Pelisson, P.-F., Bel-Venner, M.-C., Desouhant, E., Venner, S., Menu, F., Giron, D.,
687 Rey, B., 2012. A handbook for uncovering the complete energetic budget in insects: the
688 van Handel's method (1985) revisited. *Physiol. Entomol.* 37, 295–302.
689 <https://doi.org/10.1111/j.1365-3032.2012.00831.x>

690 González-Santoyo, I., Córdoba-Aguilar, A., 2012. Phenoloxidase: a key component of the
691 insect immune system. *Entomol. Exp. Appl.* 142, 1–16. [https://doi.org/10.1111/j.1570-](https://doi.org/10.1111/j.1570-7458.2011.01187.x)
692 [7458.2011.01187.x](https://doi.org/10.1111/j.1570-7458.2011.01187.x)

693 Hance, T., van Baaren, J., Vernon, P., Boivin, G., 2007. Impact of extreme temperatures on
694 parasitoids in a climate change perspective. *Annu. Rev. Entomol.* 52, 107–126.
695 <https://doi.org/10.1146/annurev.ento.52.110405.091333>

696 Haselgrove, L., Botting, D., Heesjwick, R., Høj, P.B., Dry, P.R., Ford, C., Land, P.G.I., 2000.
697 Canopy microclimate and berry composition: the effect of bunch exposure on the
698 phenolic composition of *Vitis vinifera* L cv. Shiraz grape berries. *Aust. J. Grape Wine*
699 *Res.* 6, 141–149. <https://doi.org/10.1111/j.1755-0238.2000.tb00173.x>

700 Iltis, C., Martel, G., Thiéry, D., Moreau, J., Louâpre, P., 2018. When warmer means weaker:
701 high temperatures reduce behavioural and immune defences of the larvae of a major
702 grapevine pest. *J. Pest Sci.* 91, 1315–1326. <https://doi.org/10.1007/s10340-018-0992-y>

703 Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O.B., Bouwer, L.M., Braun, A.,
704 Colette, A., Déqué, M., Georgievski, G., Georgopoulou, E., Gobiet, A., Menut, L.,
705 Nikulin, G., Haensler, A., Hempelmann, N., Jones, C., Keuler, K., Kovats, S., Kröner,
706 N., Kotlarski, S., Kriegsmann, A., Martin, E., van Meijgaard, E., Moseley, C., Pfeifer, S.,
707 Preuschmann, S., Radermacher, C., Radtke, K., Rechid, D., Rounsevell, M., Samuelsson,
708 P., Somot, S., Soussana, J.-F., Teichmann, C., Valentini, R., Vautard, R., Weber, B.,
709 Yiou, P., 2014. EURO-CORDEX: new high-resolution climate change projections for

European impact research. Reg. Environ. Chang. 14, 563–578.
<https://doi.org/10.1007/s10113-013-0499-2>

Jeffs, C.T., Lewis, O.T., 2013. Effects of climate warming on host-parasitoid interactions. Ecol. Entomol. 38, 209–218. <https://doi.org/10.1111/een.12026>

Kanost, M.R., Gorman, M.J., 2008. Phenoloxidases in insect immunity, in: Beckage, N.E. (Eds.), Insect immunology. Academic Press, New York, pp. 69–96.

Karl, I., Stoks, R., De Block, M., Janowitz, S.A., Fischer, K., 2011. Temperature extremes and butterfly fitness: conflicting evidence from life history and immune function. Glob. Chang. Biol. 17, 676–687. <https://doi.org/10.1111/j.1365-2486.2010.02277.x>

Kiaieian Moosavi, F., Cargnus, E., Pavan, F., Zandigiacomo, P., 2018. Effects of grapevine bunch exposure to sunlight on berry surface temperature and *Lobesia botrana* (Lepidoptera: Tortricidae) egg laying, hatching and larval settlement. Agric. For. Entomol. 20, 420–432. <https://doi.org/10.1111/afe.12274>

Kingsolver, J.G., Higgins, J.K., Augustine, K.E., 2015. Fluctuating temperatures and ectotherm growth: distinguishing non-linear and time-dependent effects. J. Exp. Biol. 218, 2218–2225. <https://doi.org/10.1242/jeb.120733>

Klockmann, M., Karajoli, F., Kuczyk, J., Reimer, S., Fischer, K., 2016. Fitness implications of simulated climate change in three species of copper butterflies (Lepidoptera: Lycaenidae). Biol. J. Linn. Soc. 120, 125–143. <https://doi.org/10.1111/bij.12846>

Kotlarski, S., Keuler, K., Christensen, O.B., Colette, A., Déqué, M., Gobiet, A., Goergen, K., Jacob, D., Lüthi, D., van Meijgaard, E., Nikulin, G., Schär, C., Teichmann, C., Vautard, R., Warrach-Sagi, K., Wulfmeyer, V., 2014. Regional climate modeling on European scales: a joint standard evaluation of the EURO-CORDEX RCM ensemble. Geosci. Model Dev. 7, 1297–1333. <https://doi.org/10.5194/gmd-7-1297-2014>

734 Laughton, A.M., O'Connor, C.O., Knell, R.J., 2017. Responses to a warming world:
 735 Integrating life history, immune investment, and pathogen resistance in a model insect
 736 species. *Ecol. Evol.* 7, 9699–9710. <https://doi.org/10.1002/ece3.3506>

737 Lavine, M.D., Strand, M.R., 2002. Insect hemocytes and their role in immunity. *Insect*
 738 *Biochem. Mol. Biol.* 32, 1295–1309. [https://doi.org/10.1016/S0965-1748\(02\)00092-9](https://doi.org/10.1016/S0965-1748(02)00092-9)

739 Lee, K.P., Roh, C., 2010. Temperature-by-nutrient interactions affecting growth rate in an
 740 insect ectotherm. *Entomol. Exp. Appl.* 136, 151–163. [https://doi.org/10.1111/j.1570-](https://doi.org/10.1111/j.1570-7458.2010.01018.x)
 741 [7458.2010.01018.x](https://doi.org/10.1111/j.1570-7458.2010.01018.x)

742 Lemoine, N.P., Burkepile, D.E., Parker, J.D., 2014. Variable effects of temperature on insect
 743 herbivory. *PeerJ* 2, e376. <https://doi.org/10.7717/peerj.376>

744 Martín-Vertedor, D., Ferrero-García, J.J., Torres-Vila, L.M., 2010. Global warming affects
 745 phenology and voltinism of *Lobesia botrana* in Spain. *Agric. For. Entomol.* 12, 169–176.
 746 <https://doi.org/10.1111/j.1461-9563.2009.00465.x>

747 Martin, T.L., Huey, R.B., 2008. Why 'suboptimal' is optimal: Jensen's inequality and
 748 ectotherm thermal preferences. *Am. Nat.* 171, E102–E118.
 749 <https://doi.org/10.1086/527502>

750 McNeil, J., Cox-Foster, D., Slavicek, J., Hoover, K., 2010. Contributions of immune
 751 responses to developmental resistance in *Lymantria dispar* challenged with baculovirus.
 752 *J. Insect Physiol.* 56, 1167–1177. <https://doi.org/10.1016/J.JINSPHYS.2010.03.020>

753 Muller, K., Vogelweith, F., Thiéry, D., Moret, Y., Moreau, J., 2015. Immune benefits from
 754 alternative host plants could maintain polyphagy in a phytopagous insect. *Oecologia*
 755 177:467–475. <https://doi.org/10.1007/s00442-014-3097-1>

756 Paaijmans, K.P., Heinig, R.L., Seliga, R.A., Blanford, J.I., Blanford, S., Murdock, C.C.,

757 Thomas, M.B., 2013. Temperature variation makes ectotherms more sensitive to climate
758 change. *Glob. Chang. Biol.* 19, 2373–2380. <https://doi.org/10.1111/gcb.12240>

759 Potter, K.A., Woods, A.H., Pincebourde, S., 2013. Microclimatic challenges in global change
760 biology. *Glob. Chang. Biol.* 19, 2932–2939. <https://doi.org/10.1111/gcb.12257>

761 Rebaudo, F., Rabhi, V.-B., 2018. Modeling temperature-dependent development rate and
762 phenology in insects: review of major developments, challenges, and future directions.
763 *Entomol. Exp. Appl.* 166, 607–617. <https://doi.org/10.1111/eea.12693>

764 Seehausen, M.L., Régnière, J., Martel, V., Smith, S.M., 2017. Developmental and
765 reproductive responses of the spruce budworm (Lepidoptera: Tortricidae) parasitoid
766 *Tranosema rostrale* (Hymenoptera: Ichneumonidae) to temperature. *J. Insect Physiol.* 98,
767 38–46. <https://doi.org/10.1016/J.JINSPHYS.2016.11.008>

768 Sgrò, C.M., Terblanche, J.S., Hoffmann, A.A., 2016. What can plasticity contribute to insect
769 responses to climate change? *Annu. Rev. Entomol.* 61, 433–451.
770 <https://doi.org/10.1146/annurev-ento-010715-023859>

771 Shikano, I., McCarthy, E., Hayes-Plazolles, N., Slavicek, J.M., Hoover, K., 2018. Jasmonic
772 acid-induced plant defenses delay caterpillar developmental resistance to a baculovirus:
773 slow-growth, high-mortality hypothesis in plant–insect–pathogen interactions. *J.*
774 *Invertebr. Pathol.* 158, 16–23. <https://doi.org/10.1016/J.JIP.2018.09.001>

775 Sinclair, B.J., Marshall, K.E., 2018. The many roles of fats in overwintering insects. *J. Exp.*
776 *Biol.* 221, jeb161836. <https://doi.org/10.1242/jeb.161836>

777 Sinclair, B.J., Marshall, K.E., Sewell, M.A., Levesque, D.L., Willett, C.S., Slotsbo, S., Dong,
778 Y., Harley, C.D.G., Marshall, D.J., Helmuth, B.S., Huey, R.B., 2016. Can we predict
779 ectotherm responses to climate change using thermal performance curves and body
780 temperatures? *Ecol. Lett.* 19, 1372–1385. <https://doi.org/10.1111/ele.12686>

781 Soto-Padilla, A., Ruijsink, R., Sibon, O.C.M., van Rijn, H., Billeter, J.-C., 2018.
 782 Thermosensory perception regulates speed of movement in response to temperature
 783 changes in *Drosophila melanogaster*. J. Exp. Biol. 221, jeb174151.
 784 <https://doi.org/10.1242/jeb.174151>

785 Taylor, K.E., Stouffer, R.J., Meehl, G.A., 2012. An overview of CMIP5 and the experiment
 786 design. Bull. Am. Meteorol. Soc. 93, 485–498. [https://doi.org/10.1175/BAMS-D-11-](https://doi.org/10.1175/BAMS-D-11-00094.1)
 787 00094.1

788 Tello, J., Añez, J.I., 2017. What do we know about grapevine bunch compactness? A state-of-
 789 the-art review. Aust. J. Grape Wine Res. 24, 6–23. <https://doi.org/10.1111/ajgw.12310>

790 Terblanche, J.S., Karsten, M., Mitchell, K.A., Barton, M.G., Gibert, P., 2015. Physiological
 791 variation of insects in agricultural landscapes: potential impacts of climate change, in:
 792 Bjorkman, C., Niemela, P. (Eds.), Climate Change and Insect Pests. CABI, Wallingford,
 793 pp. 92–118.

794 Terblanche, J.S., Nyamukondiwa, C., Kleynhans, E., 2010. Thermal variability alters climatic
 795 stress resistance and plastic responses in a globally invasive pest, the Mediterranean fruit
 796 fly (*Ceratitis capitata*). Entomol. Exp. Appl. 137, 304–315.
 797 <https://doi.org/10.1111/j.1570-7458.2010.01067.x>

798 Thiéry, D., Louâpre, P., Muneret, L., Rusch, A., Sentenac, G., Vogelweith, F., Iltis, C.,
 799 Moreau, J., 2018. Biological protection against grape berry moths. A review. Agron.
 800 Sustain. Dev. 38, 15. <https://doi.org/10.1007/s13593-018-0493-7>

801 Thiéry, D., Monceau, K., Moreau, J., 2014. Larval intraspecific competition for food in the
 802 European grapevine moth *Lobesia botrana*. Bull. Entomol. Res. 104, 517–524.
 803 <https://doi.org/10.1017/S0007485314000273>

804 Thiéry, D., Moreau, J., 2005. Relative performance of European grapevine moth (*Lobesia*

805 *botrana*) on grapes and other hosts. *Oecologia* 143, 548–557.
 806 <https://doi.org/10.1007/s00442-005-0022-7>

807 Valadez-Lira, J.A., Alcocer-Gonzalez, J.M., Damas, G., Nuñez-Mejía, G., Oppert, B.,
 808 Rodriguez-Padilla, C., Tamez-Guerra, P., 2012. Comparative evaluation of
 809 phenoloxidase activity in different larval stages of four lepidopteran pests after exposure
 810 to *Bacillus thuringiensis*. *J. Insect Sci.* 12, 1–11. <https://doi.org/10.1673/031.012.8001>

811 Van der Putten, W.H., Macel, M., Visser, M.E., 2010. Predicting species distribution and
 812 abundance responses to climate change: why it is essential to include biotic interactions
 813 across trophic levels. *Philos. Trans. R. Soc. B Biol. Sci.* 365, 2025–2034.
 814 <https://doi.org/10.1098/rstb.2010.0037>

815 Vande Velde, L., Van Dyck, H., 2013. Lipid economy, flight activity and reproductive
 816 behaviour in the speckled wood butterfly: on the energetic cost of territory holding.
 817 *Oikos* 122, 555–562. <https://doi.org/10.1111/j.1600-0706.2012.20747.x>

818 Vasseur, D.A., DeLong, J.P., Gilbert, B., Greig, H.S., Harley, C.D.G., McCann, K.S., Savage,
 819 V., Tunney, T.D., O'Connor, M.I., 2014. Increased temperature variation poses a greater
 820 risk to species than climate warming. *Proceedings. Biol. Sci.* 281, 20132612.
 821 <https://doi.org/10.1098/rspb.2013.2612>

822 Vogelweith, F., Moret, Y., Monceau, K., Thiéry, D., Moreau, J., 2016. The relative abundance
 823 of hemocyte types in a polyphagous moth larva depends on diet. *J. Insect Physiol.* 88,
 824 33–39. <https://doi.org/10.1016/J.JINSPHYS.2016.02.010>

825 Vogelweith, F., Thiéry, D., Moret, Y., Colin, E., Motreuil, S., Moreau, J., 2014. Defense
 826 strategies used by two sympatric vineyard moth pests. *J. Insect Physiol.* 64, 54–61.
 827 <https://doi.org/10.1016/j.jinsphys.2014.03.009>

828 Vogelweith, F., Thiery, D., Moret, Y., Moreau, J., 2013. Immunocompetence increases with

829 larval body size in a phytophagous moth. *Physiol. Entomol.* 38, 219–225.
830 <https://doi.org/10.1111/phen.12025>

831 Xue, C.-B., Luo, W.-C., Chen, Q.-X., Wang, Q., Ke, L.-N., 2006. Enzymatic properties of
832 phenoloxidase from *Pieris rapae* (Lepidoptera) larvae. *Insect Sci.* 13, 251–256.
833 <https://doi.org/10.1111/j.1744-7917.2006.00091.x>

834 Zeh, J.A., Bonilla, M.M., Su, E.J., Padua, M. V., Anderson, R. V., Zeh, D.W., 2015. Constant
835 diurnal temperature regime alters the impact of simulated climate warming on a tropical
836 pseudoscorpion. *Sci. Rep.* 4, 3706. <https://doi.org/10.1038/srep03706>

837 Ziter, C., Robinson, E.A., Newman, J.A., 2012. Climate change and voltinism in Californian
838 insect pest species: sensitivity to location, scenario and climate model choice. *Glob.*
839 *Chang. Biol.* 18, 2771–2780. <https://doi.org/10.1111/j.1365-2486.2012.02748.x>

840



Global warming



Dijon



Vineyards


+ 5.3°C in mean temperature

+ 0.6°C in daily thermal range

Positive effect



Lobesia botrana



Negative effect

Development




Survival



Movement speed



Lipid reserves



Immunity

