

Very high specific activity erbium 169Er production for potential receptor-targeted radiotherapy

R. Formento-Cavaier, U. Köster, B. Crepieux, V.M. Gadelshin, Ferid Haddad, T. Stora, K. Ulrich Wendt

► To cite this version:

R. Formento-Cavaier, U. Köster, B. Crepieux, V.M. Gadelshin, Ferid Haddad, et al.. Very high specific activity erbium 169Er production for potential receptor-targeted radiotherapy. Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms, 2019, 10.1016/j.nimb.2019.04.022 . hal-02291338

HAL Id: hal-02291338 https://hal.science/hal-02291338

Submitted on 7 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Very high specific activity erbium ¹⁶⁹Er production for potential receptor-targeted radiotherapy

R. Formento-Cavaier^{1,2,7}, U. Köster³, B. Crepieux⁴, V. M. Gadelshin^{5,6}, F. Haddad^{2,7}, T. Stora⁴ and K. Wendt⁵

¹ Advanced Accelerator Applications, a Novartis company, 20 rue Diesel, 01630 Saint-Genis-Pouilly, France

⁷ Subatech, CNRS/IN2P3, IMT Atlantique, Université de Nantes, CS 20722 44307 Nantes cedex, France

ABSTRACT

Erbium ¹⁶⁹Er is one of the most interesting radiolanthanides for new potential receptor-targeted β^{2} therapy applications due to its low energy β^{2} emissions, very low intensity γ rays and the possibility to use ⁶⁸Ga or ⁴⁴Sc as companion for diagnostic in a theranostics approach. Currently it can be produced in reactors through the neutron activation of highly enriched ¹⁶⁸Er. The low specific activity of the produced carrier-added ¹⁶⁹Er is limiting its use for receptor-targeted therapy. Nonetheless it is used for radiosynoviorthesis of small joints. The aim of this work is to develop a new large-scale production method for the supply of very high specific activity ¹⁶⁹Er. Highly enriched ¹⁶⁸Er target has been irradiated at ILL nuclear reactor and shipped to CERN-MEDICIS. There, the irradiated sample has been mass separated in order to isolate ¹⁶⁹Er from the high amount of remaining stable ¹⁶⁸Er. The proof of principle for a preclinical dose production has been demonstrated with a collection of ≈ 17 MBq. The specific activity obtained was ≈ 240 GBq/mg (≈ 200 times higher than the product obtained at End of Bombardment - EOB) and the overall separation efficiency was $\approx 0.2\%$. Several improvements for the future have been identified and are promising. One of them is the installation of the new laser laboratory at CERN-MEDICIS that will allow to improve the selective ionization of erbium atoms leading to an increase of the efficiency of the method. This method can provide the supply of high specific activity ¹⁶⁸Er, first for preclinical studies, and opens also the potential for future large-scale supply.

1. Introduction

Erbium ¹⁶⁹Er is one of the most interesting radiolanthanides for new potential receptor-targeted β^{-} therapy applications. Its main physical characteristics (see Table 1) are low energy β^{-} emissions and the near absence of γ rays. This is an advantage for dosimetry and radioprotection as well as for limiting damage to healthy tissues further away. The low energy β^{-} emissions can diversify the applications allowing ¹⁶⁹Er to be suitable for tumors not efficiently treatable with higher energies β^{-} emissions. In Table 1 are resumed the main physical properties of ¹⁶⁹Er compared to the actual β^{-} emitter of reference ¹⁷⁷Lu, currently used for commercially available radiopharmaceuticals for treatment of neuroendocrine tumors [1].

Table 1.

Main physical properties comparison between $^{169}\mathrm{Er}$ and $^{177}\mathrm{Lu}$ (data are extracted from NuDat 2 [2]

Radioisotope	¹⁶⁹ Er	¹⁷⁷ Lu
Half-life (days)	9.39	6.65
Mean/maximum β ⁻ emissions (keV)	98/343 (45%) 101/351 (55%)	48/177 (11.6%) 112/385 (9%) 149/498 (79.4%)
γ emissions (keV)	109.8 (1.3E-3%) 118.2 (1.4E-4%)	112.9 (6.2%) 208.4 (10.4%)
Mean/maximum penetration in soft tissues	0.2/1 mm [3]	0.23/1.7 mm [4]

As it is a trivalent element, it can be used with the same chelators as those used for coupling the diagnostic counterparts such as ^{68}Ga or $^{44}\text{Sc},$ and thus easily be used for theranostics applications. Due to its characteristics, it has already been used in medicine for radiosynovectomy (also called radiosynoviorthesis) of small joints such as finger joints [3]. Radiosynoviorthesis has been used since 1952 with the injection of radiopharmaceuticals based on pure or dominant β emitters. Different isotopes are currently used for different sized joints depending on their physical characteristics, in particular the electron energy emission (higher β - energy for larger sizes of the treated joint): ⁹⁰Y is used for knee joints, while ¹⁸⁶Re is used for elbow and ankle joints. For those applications, the radionuclides are used as a colloid in citrate form [3] and low specific activity is sufficient, thus carrier-added ¹⁶⁹Er can be used. The radionuclide batches are currently produced in nuclear reactors irradiating highly enriched (up to 98.2%) ¹⁶⁸Er targets. The neutron capture reaction $^{168}{\rm Er}(n,\gamma)^{169}{\rm Er}$ has a rather small cross section, namely 2.3 barns for thermal neutrons. This is limiting the specific activity. Considering a neutron flux of 1.2·10¹⁵ n/cm⁻²s⁻¹, available in a high-flux nuclear reactor such as that of Institut Laue-Langevin (ILL) in Grenoble, and ten days of irradiation, a specific activity of ≈ 5 GBq/mg can be achieved. Still, this corresponds to a ratio between ¹⁶⁸Er and ¹⁶⁹Er nuclides of 600:1 at end of irradiation and even higher at the time of use. The dilution of ¹⁶⁹Er in stable erbium limits the benefits in receptor-targeted therapies as stable erbium isotopes will compete with the radioactive one for the radiolabeling. Consequently, only a small fraction of the targeting vectors would carry "useful" radioactive 169Er while all others would just carry "cold" stable Er that is not therapeutically active. Therefore, higher specific activities are required. The theoretical specific activity of pure non-carrier-added 169Er would correspond to 3 TBq/mg. It is

² GIP Arronax, 1 rue Aronnax, 44800 Saint Herblain, France

³ Institut Laue-Langevin, 71 Avenue des Martyrs, 38042 Grenoble, France

⁴ CERN, Esplanade des Particules 1, 1211 Genève, Switzerland

⁵ Institute of Physics, Johannes Gutenberg University Mainz, Staudingerweg 7, 55128 Mainz, Germany

⁶ Ural Federal University, Mira st. 19, 620002 Ekaterinburg, Russia

defined as the activity of ¹⁶⁹Er over its mass, corresponding to the ideal scenario where only pure ¹⁶⁹Er is present.

For improving the specific activity, ¹⁶⁹Er should be separated from stable erbium ¹⁶⁸Er. As they are isotopes of the same element, conventional chemical separation processes are not beneficial. In the present work the use of EMIS technology (ElectroMagnetic Isotope Separation) for a reactor irradiated enriched ¹⁶⁸Er target is studied. This will allow producing very high specific activity batches of ¹⁶⁹Er for the potential application in vectorized radiotherapy. The experiments have been performed combining irradiations at ILL's high flux nuclear reactor in Grenoble with the mass separation of the irradiated samples at CERN-MEDICIS [5]. The experimental campaign results are shown and future improvements of the system are discussed.

2. Experimental method

2.1. Off-line separation of stable erbium

The first step was to determine the experimental overall efficiency of stable erbium extraction, ionization and mass separation. This allows determining extrapolated yield values for production estimations. The separation of a known mass of stable erbium was performed in the off-line laboratory at CERN. For this experiment MEDICIS target unit #631-M, the one dedicated to the following separation of 169Er at CERN-MEDICIS, was used. It consists in a conventional ISOLDE type target unit, with inside an empty target container and a tungsten surface ion source. Usually a tantalum plug is used to close the tantalum target container, but with the high operation temperatures this risk getting diffusion-bonded. This behavior is fine when the target is irradiated with CERN's proton beam because the target container has not to be opened after use. In our case as the target container has to be filled several times, the plug should be easily removable. Therefore, for guaranteeing the retrieval of the inserted samples a graphite plug has been used.

The inserted sample was prepared from erbium plasma standard solution (1 mg/ml in 5% HNO₃. Specpure, Alfa Aesar). Natural Erbium is composed of 6 stable isotopes among which ¹⁶⁶Er is the most abundant, 33.5%. This isotope was followed during our "cold" tests. An aliquot of 100 μ l, corresponding to 3.6 * 10¹⁷ atoms of natural erbium, was poured on a graphite disc placed on a molybdenum boat. The boat prevents any direct contact between the sample and the tantalum container to avoid reactions of graphite and tantalum at high temperature. By means of a lamp, the sample was heated to evaporate the acid solution and deposit the erbium salt Er(NO₃)₃. In the Fig. 1 the preparation of the sample is shown.



Fig.1. Erbium sample used for the offline erbium separation efficiency measurements.

Then the sample was loaded into the target container. After vacuum pumping the target container was heated to ≈ 2100 °C. The extraction was performed over 7.8 h collecting 3.15×10^{14} positively charged ions of 166 Er. As the initial number of 166 Er atoms in the sample was $3.6 \times 10^{17} \times 0.335 = 1.21 \times 10^{17}$, the overall efficiency of the mass separation was $\frac{3.15 \times 10^{14}}{1.21 \times 10^{17}} \times 100 = 0.26\%$. When reaching the maximum temperature in the target, the Er⁺ current on mass 166 peaked at 1.7 nA. Then as the temperature remained constant until the end of the measurement the release of 166 Er from the target exponentially decreased over time as expected. Nevertheless, once the target was removed from the oven, we noticed molybdenum boat damages on the contact points between the boat and the graphite. For this reason, following theoretical and experimental considerations, rhenium was chosen as future boat material.

This test showed the feasibility of erbium extraction and laid the basis for the radioactive experiment.

2.2. Off-line separation of radioactive erbium

A quartz ampoule was filled with (7 ± 0.1) mg of Er₂O₃ powder enriched to 98.2% in ¹⁶⁸Er and irradiated for 6.5 days at the high-flux reactor of ILL in a neutron flux of 1.2×10^{15} n.cm⁻²s⁻¹ leading to a calculated specific activity of 3.7 GBq/mg. Then the radioactive sample was shipped to CERN where it was put inside one of MEDICIS fume hoods. The ampoule was opened and the erbium oxide was dissolved within 3 ml of 1M HNO₃. The solution was then transferred from the ampoule to the support consisting in a graphite holder inserted in a rhenium boat (Fig. 2).



Fig. 2. Graphite holder used for the ¹⁶⁹Er separation. The erbium solution has been poured inside and evaporated to have a deposit on the graphite.

Then as in the test experiment, the solution was heated to around 40 °C to evaporate the solution to dryness on the holder. The sample was then inserted inside the MEDICIS target unit (#631-M), which was later positioned on the mass separator front end (Fig. 3). This operation did not lead to a significant radiation dose of the operators since ¹⁶⁹Er has very low γ emission. As before the target and ion source unit was then pumped to vacuum. After bringing the ion source to the operation temperature, the target was slowly heated to release the Er. The Er ions were mass-separated and the ¹⁶⁹Er was implanted into zinc coated gold foils (Fig. 4).



Fig. 3. MEDICIS target unit positioned on the mass separator front end for the $^{169}\mathrm{Er}$ collection



Fig. 4. Collection foils used. Backside view of the foils facing each a 10 mm diameter collimator. The Zn coating is on the opposite side.

3. Results

The collection lasted 24 hours. The operating temperature was around 2000-2100 °C both in the target container and the surface ion source. The readout of the current from the collection foils was in agreement with the Faraday cup measurements upstream and the collimator readout. The losses in the collimator were almost constant all along the collection oscillating around 7%. The total ion current of ¹⁶⁹Er and its isobaric contaminants peaked at 3.8 nA when the target container reached the maximum temperature, then the signal decreased exponentially as expected. The number of atoms collected was $2.56 * 10^{14}$. The y spectrometry analysis showed that 17(6) MBq of 169 Er 1 and 10(4) kBq of 169 Yb had been collected on the zinc coated gold foil. This corresponds to $\approx 2 * 10^{13}$ ¹⁶⁹Er atoms and $4 * 10^{10}$ ¹⁶⁹Yb atoms. The difference between the integrated current and the detected activities is assumed to correspond to a tail of stable ¹⁶⁸Er ions. At the time of the separation the quantity of ¹⁶⁸Er was $\approx 2 * 10^3$ times higher compared to ¹⁶⁹Er (calculated specific activity of ≈ 1.3 GBq/mg). Therefore, a small part of the tail of the beam profile at mass 168 can be found on one side of the Gaussian profile of the mass 169 beam. The mass separation improved the ratio of ¹⁶⁸Er to ¹⁶⁹Er from \approx 2000:1 to \approx 10:1, meaning a gain of a factor \approx 200. Correspondingly the specific activity was increased from ≈ 1.3 GBq/mg to \approx 240 GBq/mg, closer to the desired specific activity for a potential clinical receptor-targeted radiotherapy application. These results should be confirmed through an ICP-MS measurement after complete decay of the radionuclides. However, only ≈ 17 MBq of ^{169}Er were extracted from the \approx 9 GBq of ^{169}Er present in the MEDICIS target container before mass separation. The overall efficiency was consequently $\approx 0.2\%$ close to the value obtained during the off-line test with stable erbium.

¹ The large uncertainty in the activity determination results from the interference correction in the γ ray analysis. All γ rays populated in ¹⁶⁹Er decay are also populated with much higher intensity in the ¹⁶⁹Yb decay. Thus, despite the much lower ¹⁶⁹Yb activity, it actually dominates the γ ray spectra and ¹⁶⁹Er is only observed as increment at certain energies.

The described experiment represents the first production of high specific activity ¹⁶⁹Er. The proof of principle of the production of a usable dose was performed. Nevertheless, the production yield and the quality of the product can be improved. The simplest improvement is the reduction of the residual ¹⁶⁸Er content, which can be achieved optimizing the position of the mass separator slit. The optimal position for the maximum reduction of stable erbium minimizing the ¹⁶⁹Er should be found. A more important improvement is needed to increase the overall efficiency, as 0.2% is too low for practical purposes. A much higher ionization efficiency can be expected when replacing surface ionization by resonant laser ionization.

4. Resonant laser ionization

The installation of lasers has not yet been completed at the MEDICIS facility [6]. Therefore, the experiments have been performed at Johannes Gutenberg University of Mainz, using the RISIKO mass separator [7]. The tests have been performed with stable erbium.

An erbium plasma standard solution (1 mg/ml in 5% of HNO₃, Specpure, Alfa Aesar), was diluted by a factor of ten. 3 μ l of the diluted erbium solution were deposited on a zirconium foil and dried by a heat lamp. The 3 μ l of erbium correspond to 1.08 * 10¹⁵ erbium atoms. For this experiment a two-step scheme has been selected that had been previously developed by the LARISSA group [8]. The full analysis of this experiment will be presented in a separate paper.



Fig. 5. Mass scan performed on the mass range of atomic erbium ions and erbium oxide ions. The red curve represents the ion signal with lasers switched on, whereas the dashed curve shows the ion signal with lasers switched off.

In Fig. 5 an example of a mass spectrum is presented. The laser On / Off enhancement ratio ranged from \approx 1000 to \approx 10000 during the main part of the ion extraction. As expected all the laser ionized erbium was observed as atomic ions, no laser enhancement was observed for erbium oxide ions. The significant presence of atomic gadolinium and gadolinium oxide did not interfere with the measurement of the erbium separation efficiency. Indeed, its atomic masses range from 152 to 160 and the oxides from 168 to 176, while the erbium measurement was performed on mass 166.

5. Discussion and future perspectives

¹⁶⁹Er is an interesting isotope for receptor-targeted radiotherapy due to its physical and chemical characteristics. However, it is not currently available with a sufficient specific activity because of the quite small cross section (2.3 b) for the neutron activation reaction of ¹⁶⁸Er. This leads to a strong dilution of radioactive ¹⁶⁹Er by stable ¹⁶⁸Er. The combination of irradiation in a high-flux nuclear reactor with electromagnetic mass separation is proposed to overcome this issue. The proof-of-principle experiments performed at the ILL reactor and the CERN-MEDICIS facility showed the feasibility of this production method. The specific activity could be improved from \approx 1.3 GBq/mg to \approx 240 GBq/mg. This was the first production of high specific activity ¹⁶⁹Er and \approx 17 MBq of mass-separated ¹⁶⁹Er were obtained. However, the overall efficiency was only 0.2 %. Boosting this number by resonant laser ionization was studied at JGU Mainz using the RISIKO mass separator. The finalization of the MELISSA laser laboratory at CERN-MEDICIS in the first half of 2019 will allow verifying the experimental gain factor in the production yields of ¹⁶⁹Er.

This work has demonstrated that ¹⁶⁹Er can be produced at high specific activities and be available for preclinical studies. Preclinical trials could be performed starting from 2019. In parallel, the abovementioned improvements will allow soon to get much higher activity making ¹⁶⁹Er available for receptor-targeted radiotherapy clinical trials.

Acknowledgments

This research project has been supported by a Marie Skłodowska-Curie innovative training network fellowship of the European commission's horizon 2020 program under contract number 642889 MEDICIS-PROMED.

This work has been, in part, supported by a grant from the French National Agency for Research called "Investissements d'Avenir", Equipex Arronax-Plus noANR-11-EQPX-0004 and Labex IRON noANR-11-LABX-18-01.

References

- J. Strosberg *et. al.*, Phase 3 Trial of ¹⁷⁷Lu-Dotatate for Midgut Neuroendocrine Tumors, The New England Journal of Medicine, 376 (2017) 125-135.
- [2]. National Nuclear Data Center (NNDC), Brookhaven National Laboratory, visited on the 30th of November 2018, https://www.nndc.bnl.gov/nudat2/
- [3]. J. Farahati *et al.*, Post-radiosynovectomy imaging of Er-169 using scintigraphy and autoradiography, Clinical Case Reports, 5 (2017):1048– 1050.
- [4]. J.J. Zaknun *et al.*, The joint IAEA, EANM, and SNMMI practical guidance on peptide receptor radionuclide therapy (PRRNT) in neuroendocrine tumours. Eur J Nucl Med Mol Imaging 40 (2013) 800– 816.

- [5]. R. M. Augusto *et al.*, CERN-MEDICIS (Medical Isotopes Collected from ISOLDE): a new facility. Applied Sciences, 4 (2014) 265.
- [6]. V. M. Gadelshin et al., MELISSA: Laser Ion Source Setup at CERN-MEDICIS facility. Blueprint, Proc. of EMIS 2018, this volume.
- [7] C. Mattolat *et. al.*, An all-solid-state high repetition rate titanium:sapphire laser system for resonance ionization laser ion sources, AIP conference proceeding, 1104 (2009) 114.
- [8] D Studer *et al.*, RIS in dysprosium and erbium for scheme development and determination of the first ionization potential, LA3NET Laser Ion Sources Workshop (2016).