



HAL
open science

Characteristic early membrane effects induced by tryptophan in pulvinar motor cells

Fabienne Dédaldéchamp, Christelle Moyen, Pierrette Fleurat-Lessard, Gabriel
Roblin

► **To cite this version:**

Fabienne Dédaldéchamp, Christelle Moyen, Pierrette Fleurat-Lessard, Gabriel Roblin. Characteristic early membrane effects induced by tryptophan in pulvinar motor cells. *Plant Physiology and Biochemistry*, 2019, 142, pp.528-535. 10.1016/j.plaphy.2019.08.006 . hal-02296344

HAL Id: hal-02296344

<https://hal.science/hal-02296344>

Submitted on 20 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **PLAPHY-D-19-00294 > revised final version**

2 **Characteristic early membrane effects induced by tryptophan in**
3 **pulvinar motor cells**

4

5 **Fabienne Dédaldéchamp^{1*}, Christelle Moyen², Pierrette Fleurat-Lessard¹ and**
6 **Gabriel Roblin¹**

7

8

9

10 *¹ University of Poitiers, UFR Sciences Fondamentales et Appliquées, UMR CNRS*
11 *7267 Laboratoire EBI Ecologie et Biologie des Interactions, Equipe SEVE Sucres &*
12 *Echanges Végétaux Environnement, 3, rue Jacques Fort, Bât. B31, TSA 51106,*
13 *86073 POITIERS CEDEX 9, France*

14 *² University of Bourgogne Franche-Comté, Laboratoire de Chrono-Environnement,*
15 *UMR CNRS 6249 UsC INRA, 16 route de Gray, 25000 Besançon, France*

16

17

18 *Corresponding author

19 *E-mail address:* fabienne.dedaldechamp@univ-poitiers.fr

20 *Tel:* 0(33)5 49 45 37 60

21

22 Resubmitted 2 July 2019

23 With 2 tables and 2 figures

24 Supplementary data S1 and S2 with 2 figures

25 **Running title:** Tryptophan as a signal molecule in plants

26

27 **Highlight** Tryptophan modifies transmembrane ionic equilibrium in plant cells as
28 indicated by variations induced on transmembrane electrical potential, proton
29 movement and osmocontractile ion-driven pulvinar cell reaction.

30

31 **ABSTRACT (197 words)**

32 Tryptophan at concentrations higher than 0.1 mM, triggered characteristic early
33 physiological effects such as rapid (within 5 min) dose-dependent membrane
34 hyperpolarization in *Mimosa pudica* motor cells and modification of the time course
35 of the spontaneous proton efflux monitored in the incubation medium of pulvinar
36 tissues. The rapid modifications of the leaf turgor-mediated movements seen on the
37 primary pulvini of *M. pudica* following a shock and on *Cassia fasciculata* leaflets
38 during a transition from light to darkness indicate that tryptophan disturbed the ionic
39 migrations involved in the electrophysiological events and in the osmocontractile
40 reaction of the motor cells. These reactions were specific to tryptophan compared to
41 those induced by serine and 5-hydroxytryptophan. The tryptophan mode of action
42 cannot be linked to a direct modification of the plasma membrane H⁺-ATPase
43 activity as monitored on purified pulvinar plasma membrane vesicles. The
44 tryptophan metabolism-linked products tryptamine and indole also inhibited the
45 motile reactions, activated in a continuous manner the H⁺ secretion of pulvinar
46 tissues and showed properties of a protonophore and an ATPase activity inhibitor on
47 plasma membrane vesicles, respectively. The specific behavior of tryptophan in the
48 reaction studies here is discussed in light of the previously reported action of
49 phytohormones.

50

51 **Keywords:** amino acids, *Cassia fasciculata*, membrane potential, *Mimosa pudica*,
52 nasties, osmocontraction, pulvinus, tryptamine, tryptophan.

53 Abbreviations: Ind, indole; OHTrp, 5-hydroxytryptophan; PMVs, plasma membrane
54 vesicles; Ser, serine; Try, tryptamine, Trp, tryptophan.

55

56 1. Introduction

57 The amino acid tryptophan (Trp) plays a primary role in all living organisms,
58 from bacteria and fungi to plants and animals, by being an essential amino acid
59 required for protein synthesis. In addition, Trp intervenes in the synthesis of many
60 pivotal compounds. In animals which are incapable of synthesizing this amino acid,
61 Trp after being ingested as residues within proteins is used for the synthesis of
62 serotonin, melatonin and vitamin nicotinic acid (Moffett and Namboodiri, 2003). In
63 plants, Trp plays a direct role in regulating plant development by providing
64 precursors for the multiple pathways of synthesis of the hormone indole-3-acetic acid
65 (IAA) that controls numerous physiological processes (Woodward and Bartel, 2005).
66 Furthermore, Trp has been shown to assist in plant tolerance of heavy metals. Thus,
67 the Trp content increased in *Noccaea caerulea* treated with increasing
68 concentrations of Cd (Zemanova et al., 2014), resulting in reduced Cd availability to
69 the plants, decreased Cd transport and reduced Cd accumulation (Sanjaya et al.,
70 2008). In addition, enzymes involved in Trp synthesis were induced under different
71 kinds of stresses such as amino acid starvation, the oxidative stress-inducing
72 herbicide acifluorfen and the abiotic elicitor α -amino butyric acid (Zhao et al., 1998).
73 Plants also utilize Trp as a precursor in the synthesis of a large variety of secondary
74 metabolites, known to intervene in pathogen defense responses such as serotonin
75 (Ishihara et al., 2008), phytoalexins (particularly camalexin), glucosinolates and
76 indole-derived alkaloids (Radwanski and Last, 1995; Frerigmann et al., 2016).
77 However, Trp has been shown to affect *per se* various long-term physiological
78 processes in plants as shown by its effects observed on leaf growth and cell
79 expansion in *Arabidopsis* (Jing et al., 2009).

80 In addition to its action on these types of processes involving gene regulation,
81 Trp, similar to other amino acids, induces short-term cellular events, as evidenced by
82 its triggering of rapid ionic fluxes during their uptake through the plasma membrane
83 mediated by proton-coupled symports (Delrot et al., 2001 and references therein).

84 The aim of this work was to show whether, compared to other natural amino
85 acids, Trp triggers characteristic early events at the membrane level, and in
86 particular, noteworthy modifications of the bioelectrical transmembrane potential and
87 disturbances of proton fluxes. For this purpose, the motor cells of pulvini of *Mimosa*
88 *pudica* and *Cassia fasciculata* were chosen because the kinetics, energetics and some

89 of the reaction steps of the stress-induced turgor driven leaf movements are now well
90 documented (Samejima and Sibaoka, 1980; Moran 2007). The reaction of the
91 primary pulvini of *M. pudica* following a shock consists of a rapid spectacular
92 drooping movement of the leaf in approximately 2 s, and the recovery takes 15-20
93 min (Volkov et al., 2010). Leaflet movements can be induced in *M. pudica* and *C.*
94 *fasciculata* by transferring plants from light to darkness or from darkness to light. As
95 the result of the coordinated contraction of the pulvinar cells, the motor reactions are
96 driven by rapid turgor changes mainly associated with K⁺ and Cl⁻ fluxes. In plant
97 cells, K⁺ fluxes have been shown to be coupled to reverse H⁺ fluxes driven by the
98 regulation of the activity of a proton-pumping ATPase located at the plasma
99 membrane (Serrano, 1989). Thus, attention has been paid on the way Trp may have a
100 direct effect on the H⁺-ATPase using purified plasma membrane vesicles (PMVs)
101 isolated from pulvinar tissues of *M. pudica*, which allow such an experimental
102 approach.

103 The results obtained regarding Trp action on these different biological events
104 are discussed in light of the comparative effects observed following treatments with
105 5-hydroxytryptophan (OHTrp) and by compounds linked either to Trp synthesis,
106 namely, serine (Ser) and indole (Ind) (Kriechbaumer et al., 2008) or to Trp
107 catabolism, namely, tryptamine (Try) (Kang et al., 2008). It can be concluded that,
108 among amino acids, Trp triggers specific actions on early ionic exchanges at the
109 plasma membrane level and may serve as a signal modulating more long-lasting
110 physiological processes triggered in plants by environmental changes.

111

112

113

114 2. Materials and methods

115

116 2.1. Plant growth conditions

117

118 Seeds of *Mimosa pudica* L. and *Cassia fasciculata* Michx. were germinated in
119 an organic compost. Seedlings and older plants were grown in the compost watered
120 daily and were kept in climate-controlled chambers at 27 ± 0.5 °C and $65 \pm 5\%$
121 relative humidity. Illumination was regulated to give 16 h of light (photophase 06:00-
122 22:00 hours) provided by fluorescent tubes (mixing Osram fluora and Osram day-
123 light types) giving a photon flux density (400-800 nm) of $80 \mu\text{mol m}^{-2} \text{s}^{-1}$ at the plant
124 apex.

125

126 2.2. Electrophysiological measurements

127

128 The particular anatomy of the primary pulvinus of *M. pudica* characterized by
129 many layers of parenchyma cells surrounding the central cylinder allows impalement
130 of a microelectrode in a well-defined kind of cell. The transmembrane potential was
131 measured by the classical electrophysiological method using glass microelectrodes
132 (tip diameter $< 1 \mu\text{m}$, tip resistance from 5 to 30 M Ω) drawn from capillaries
133 provided with an internal microfibre (GC 150F15; Clark Electromedical
134 Instruments). Briefly, a leaf was excised from the stem of 2-month-old plants and the
135 pulvinus fixed to the bottom of a 4 ml plexiglass chamber filled with a buffered
136 medium (10 mM MES, pH 5.5) containing 1 mM NaCl, 0.1 mM KCl, and 0.1 mM
137 CaCl₂. The microelectrode was impaled into a motor cell of the abaxial (“extensor”)
138 half of the organ. Under these conditions, the calculated value of the resting
139 transmembrane potential (Ψ_0) from 33 assays was -118 ± 4 mV (SEM). For details,
140 see [Saeedi et al. \(2013\)](#) and references therein.

141

142 2.3. Measurement of pH variations

143

144 To record H⁺ excretion, primary pulvini (400 mg) of 2-month-old *M. pudica*
145 plants generally bearing 10 leaves were excised at 10:00 hours. The organs were
146 divided in transverse sections and treated following the same procedure previously
147 described ([Saeedi et al., 2013](#)): an incubation medium composed of 0.50 mM CaCl₂,

148 0.25 mM MgCl₂, and variations of pH read on a pH-meter provided with combined
149 electrodes (Futura micro-combination, Beckman Coulter) and linked to a
150 potentiometric recorder. Compounds were added 3 hours after the beginning of the
151 experiments when the pH decrease in the bathing medium reached a plateau. To
152 quantify the amount of mobilized protons, titration was performed with NaOH or
153 HCl at 5×10^{-3} M onto a 2 ml incubation medium aliquot collected 3 h after
154 application of the various tested compounds. The experiments were repeated at least
155 3 times.

156

157 *2.4. Preparation and use of plasma membrane vesicles (PMVs)*

158

159 Purified PMVs were prepared by phase partitioning of microsomal fractions
160 from *M. pudica* primary pulvini. PMVs were frozen in liquid nitrogen and stored at -
161 80 °C. The PMVs were put in the inside-out configuration by adding 0.05% Brij in
162 the assay medium. In the 4 batches of vesicles used in this work, treatment with the
163 plasma membrane H⁺-ATPase inhibitor sodium orthovanadate at 0.25 mM showed
164 that 60% of the enzyme activity can be attributed to plasma membrane functioning.
165 Vanadate-sensitive ATPase activity of the PMVs was measured in a medium
166 buffered with 50 mM MES/Tris at pH 6.5. Proton pumping was followed by the
167 decrease of 9-aminoacridine absorbance at 495 nm in a medium buffered with 10
168 mM MES/Tris at pH 6.5. The reaction was started by the addition of 3 mM MgSO₄
169 into the medium. For details, see [Saeedi et al. \(2013\)](#).

170

171

172 *2.5. Experimental procedure for observation of pulvinar movements*

173

174 Plantlets of *M. pudica* bearing the first fully developed leaf were excised at
175 10:00 hours by cutting the hypocotyl 3 cm below the cotyledons and then were
176 dipped in distilled water contained in small plastic tubes to allow for recovery from
177 harvesting. Leaf position determined by the angle formed by the petiole and the line
178 prolonging the hypocotyl was measured using a transparent protractor provided with
179 a movable pointer, allowing angle measurement by homothety preventing therefore
180 leaves from mechanical touch. At 14:00 hours (middle of the photophase), the
181 distilled water was replaced by the tested solution buffered at pH 5.5 with 2.5 mM

182 MES. After measurement of the initial angle (α_i), the pulvini were stimulated by a
183 shock at 1-h intervals for 4 h. Motor reactions were monitored by measuring the
184 variation of the angle ($\Delta\alpha$) within 10 s after the stimulation. The experiments were
185 repeated 3 times on ten seedlings.

186 Fully expanded leaves of *C. fasciculata* were excised at 09:00 hours from 2-
187 month-old plants bearing 12 leaves. The cut petioles were dipped in distilled water to
188 recover. Leaves were then transferred to test solutions buffered with 2.5 mM MES at
189 pH 5.5 (except in particular assays dealing with pH action) for durations extending
190 from 5 to 180 min and were exposed to darkness in the middle of the photophase of
191 the photoperiodic cycle or in the light in the middle of the nyctophase (in this case,
192 see [additional data S2](#)). At the end of the incubation period, the petiole tips were
193 rinsed with buffer solution and the leaves were transferred to 2.5 mM MES adjusted
194 at pH 5.5. The pulvinar movements induced by the variation in the light regime were
195 monitored by measuring the distance between the leaflet tips with a calliper square
196 which prevents disturbing of the leaf position by a mechanical touch. This linear
197 measurement was then converted into angular values according to the formula $d/2 =$
198 $L/2 \sin \sigma/2$ in which σ was the calculated angle, d was the distance measured
199 between the leaflet tips and L the maximum distance measured when the leaflets
200 were fully opened ($\sigma = 180$ degrees). The curves represent the mean values obtained
201 from the sum of at least three separate experiments on 8 individual leaves.

202

203 *2.6. Uptake and metabolism of tryptophan*

204

205 The experiments dealing with 1 mM labelled Trp uptake were carried out on
206 the excised plantlets of *M. pudica* as described above. The uptake of [^3H] Trp (final
207 specific activity of the solution 8 kBq ml $^{-1}$) was measured by sectioning primary
208 pulvini from the leaves with a razor blade at regular intervals. The excised organs
209 were weighed and digested in a cocktail of 30% hydrogen peroxide, 70% perchloric
210 acid and Triton X100 (2:1:1, v/v/v) for 24h at 55 °C. Radioactivity was then counted
211 after addition of a scintillation medium (Ecolite, ICN) by liquid scintillation
212 spectrometry (Packard Instruments). The experiments were repeated at least three
213 times on 10 seedlings, so that each experimental point represents observations on 30
214 motor organs.

215 In a specific assay, a set of pulvini was harvested after 2 h of treatment in the
216 labelled Trp, before being washed twice in 30 ml cold Trp-free buffered medium.
217 The samples were ground by mortar and pestle with 1 g quartz in 1 ml distilled
218 water. After centrifugation, the residue was extracted again with medium and the
219 combined extracts were vacuum evaporated. Thin-layer chromatography was
220 performed to identify the Trp on Merck silica gel 60F254 plates (2 mm thick)
221 developed with butanol / acetic acid / H₂O (60:20:20, v/v/v). In these conditions, the
222 R_F value of the Trp visualized under UV light (254 nm) was 0.66. A 50- μ l volume of
223 the labelled extract was deposited onto a plate. After migration, the plate was dried
224 and 0.5-cm-high silica gel areas were scraped off along the lane and placed into
225 scintillation vials with 4 ml Ecolyte for counting.

226 227 *2.7. Data analysis*

228
229 Data were statistically analyzed according to Student-Fisher t test. Depending
230 whether the number of sample (n) was above or below 20, data were respectively
231 given as mean \pm standard deviation (SD) and mean \pm standard error (SE) in the
232 figures and tables. A significant level of $P \leq 0.05$ was selected for determination of
233 statistical significance.

234 235 *2.8. Chemicals*

236
237 All chemicals were purchased from Sigma-Aldrich Chimie (St-Quentin-
238 Fallavier, France). Solutions of Trp, OHTrp, Try₂ and Ser were prepared at 10 mM
239 final concentration and diluted in 2.5 mM HEPES (pKa 3.5 and 7.5) for experiments
240 conducted at pH 4.5 and 7.5, and in 2.5 mM MES (pKa 6.1) for experiments
241 conducted at pH 5.5 and 6.5. The pH of the solutions was adjusted with either 0.1 M
242 KOH or 0.1 M HCl. Control assays have shown that buffering the media did not
243 affect the pulvinar movements. Ind at 1 mM final concentration was dissolved in
244 absolute ethanol, the final concentration of solvent being 0.3%. A stock solution of
245 Fusicoccin (FC) prepared at 1 mM was dissolved in 7% final concentration of
246 methanol. It has been verified that the respective solvent concentration assayed in
247 controls did not affect the studied processes.

248

3. Results

3.1. Effect of tryptophan on the transmembrane potential of motor cells

The addition of Trp to the bathing medium of *M. pudica* motor cells at pH 5.5 induced a long-lasting hyperpolarization of the motor cell membrane in a concentration-dependent manner from 0.05 (threshold value) to 1 mM. **Fig. 1A** presents typical recordings of the time course of the bioelectrical modifications: the lag period lasted some minutes and the subsequent rise of potential reached a plateau within 20-30 min after application of the compound. Additional quantitative data are given in **Table 1**. The effect of Trp completely differed from that of OHTrp (**Fig. 1A**) and serine (**Table 1**) both of which induced a depolarization indicating that the effect-of Trp is specific compared to that of the other tested amino acids. FC at 10 μ M strongly hyperpolarized the cell membrane, ensuring the bioelectrical responsiveness of the impaled cell.

3.2. Effect of Tryptophan on the H^+ -ATPase activity

In a previous study, it was shown that slices of primary pulvini spontaneously acidified their incubation medium as a result of the FC-activated proton pump activity (Saeedi et al. 2013). **Fig. 1B** shows the time course of the pH variation monitored in the bathing medium of the motor organ tissues after addition of the tested compounds. In the presence of Trp, the time course monitored in the controls was modified and displayed a biphasic pattern. First, Trp promoted a transient concentration-dependent acidification of the medium after a latency of approximately 5 min in the range 0.1 - 1 mM, lasting from 0.5 h for 0.1 mM to 1.5 h for 1 mM. Then, in the next phase, it induced a dose-dependent rise in pH lasting for 2-3 h. By contrast, 1 mM OHTrp and 1 mM Ser both induced a sharp rise of pH observed some min after application which reached an optimum level after 2.5 h for OHTrp and 1 h for Ser, before decreasing back to the level of the control in about 5 h for OHTrp and 2 h for Ser. Another pattern was seen with 1 mM Try and 1 mM Ind as they both induced a continuous pH decrease in the bathing medium starting approximately 15 min after application of the products and lasting 3 h for Try and 5 h for Ind (**Fig. 1B**).

282 Additional data quantified the amount of mobilized proton following the various
283 treatments (**Table 1**).

284 We further investigated whether a close relationship could be drawn between
285 the H⁺ fluxes monitored previously and a direct effect on the functioning of the
286 plasma membrane H⁺-ATPase activity. In this goal, additional experiments were
287 conducted on PMVs purified from pulvinar tissues in which proton pump assays and
288 determination of ATPase catalytic activity were carried out. As seen in **Table 2**, Trp
289 did not significantly modify both components of the proton motive force showing the
290 same behavior as Ser and OHTrp. By contrast, Try strongly inhibited the proton
291 pumping but did not modify the catalytic activity, presenting, therefore, the
292 characteristics of a protonophore. Ind showed a vanadate-like behavior with
293 particular characteristics. Indeed, Ind inhibited more strongly proton pumping (86%
294 vs. 60%) but inhibited ATPase activity with a lower efficiency (37% vs. 66%).

295

296 *3.3.Characteristics of the inhibition of the osmocontractile reaction by tryptophan*

297

298 The seismonastic reaction of the *M. pudica* primary pulvinus was gradually
299 inhibited in a dose-dependent manner for Trp concentrations higher than 1 mM (**Fig.**
300 **1C**). The effect was clearly seen one hour after the Trp addition and developed over
301 the course of the experiment. After 2 h of Trp action, the inhibition was 6% at 1 mM,
302 18% at 5 mM and 35% at 10 mM. The time course of Trp effect can be closely
303 paralleled with the amount of the labelled product measured in the motor organs
304 (**Fig. 1C**). The Trp uptake inhibition (77%) obtained following a treatment with 10
305 μM of the protonophore CCCP argues for an active Trp uptake mechanism driven by
306 a H⁺ gradient-dependent process and supports the hypothesis of a Trp-H⁺ symport in
307 motor cells. TLC analysis using [³H] labelled Trp indicated that after 2 h of
308 treatment, 97% of the radioactivity was recovered at the R_f of the Trp migration spot,
309 indicating that Trp metabolism occurred at a very low extent in these experimental
310 conditions. Furthermore, the fact that the Trp-induced decrease of angle variation
311 (Δα) resulted from an incomplete leaf sinking and not from an incomplete leaf
312 recovery (since the pulvini returned to their initial position after the stimulation)
313 argues for a restriction of the K⁺ efflux, while the K⁺ influx was not affected. OHTrp
314 showed no significant effect whereas Try and Ind both strongly inhibited the reaction

315 (25 and 64% after a 2 h treatment, respectively). Of note, Ind drastically hindered the
316 recovery as measured by the increasing α_i (inset in **Fig. 1D**).

317 As shown in **Fig. 2A**, Trp disturbed the dark-induced motor reactions of *C.*
318 *fasciculata* leaflets. The handling of this model is simpler than that of *M. pudica*
319 because of its low seismonastic sensitivity. The induced closure was inhibited in a
320 dose-dependent manner at concentrations higher than 0.1 mM. After 2 h of Trp
321 action, the inhibition was 6% at 0.1 mM, 26% at 0.5 mM, 50% at 1 mM and 78% at
322 5 mM. It should be stressed that the *Cassia* model is more sensitive (approximately 5
323 times) than the primary pulvini of *M. pudica*. To determine the influence of treatment
324 duration, 1 mM Trp was applied for 5, 15, 30, 60, and 180 min before the onset of
325 darkness. **Fig. 2B** shows that a treatment as short as 5 min induced a significant
326 hindering of the reaction. This inhibitory effect increased with treatment duration
327 being complete after 30 min. Applied at different pH in buffered media, 1 mM Trp
328 inhibited the dark-induced reaction at all the pH values tested but the inhibition was
329 well-marked towards alkaline pH values (**Fig. 2C**). Nevertheless, the choice of pH
330 5.5 in 2.5 mM MES was thought to be the most suitable in our experiments as it was
331 near the physiological apoplastic pH. Data in **Fig. 2D** show that the L- or D-
332 configuration of the molecule did not influence the reaction. By contrast, a
333 modification of the structure by the introduction of an OH-group abolished the
334 efficiency. Try and Ind both showed an inhibitory effect in the same range of
335 concentrations (from 0.1 to 1 mM).

336 As shown in the histograms of **Fig. 2E**, the Trp inhibition induced on dark-
337 induced reactions was noteworthy compared to other amino acids. While most of the
338 amino acids tested did not influence the leaflet closure, it must be noted, however,
339 that arginine, cysteine, phenylalanine and serine induced a low but significant
340 inhibition when applied for 3h in this assay.

341
342
343
344
345
346
347
348

349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380

4. Discussion

4.1. Tryptophan as a multifunctional component in the plant cell life

Trp is an essential aromatic amino acid since it is one of the building blocks for protein synthesis and the source of many important compounds, particularly melatonin and nicotinamide adenine dinucleotide in animals (Moffett and Namboodiri, 2003) and IAA and phytoalexins in plants (Zhao et al., 1998). In plants, Trp has been shown to affect various long-term physiological processes such as organ growth (Jing et al., 2009), responses to stress and pathogen elicitors (Zhao et al., 1998; Bednarek et al., 2009) and tolerance of heavy metals (Sanjaya et al., 2008). Nevertheless, little has been reported about the mechanism by which Trp affects growth and development at the cellular and molecular level.

In this work, we focused our investigations on the early events triggered by Trp on plant cell functioning. Thus, we determined that Trp is rapidly recognized at the cell membrane level and induced characteristic and specific short-term cellular events by disturbing processes involved in rapid changes of membrane permeability to water and ions which are the early processes allowing osmocontractile pulvinar reactions.

4.2. Tryptophan as a putative signal molecule

The proposal that Trp is rapidly recognized at the plasma membrane level is greatly supported by the observation of the large concentration-dependent membrane hyperpolarization triggered a few minutes after application of the compound, indicating early modifications in ionic fluxes. Ion fluxes through the plasma membrane are common and early events reported as a prerequisite for subsequent cell reaction (Colcombet et al., 2009; and references therein). The early recognition of Trp at the membrane level can be analyzed according to three phases.

381 (a) The occurrence of a lag period indicates that the recognition of the
382 compound at the membrane triggers a chain reaction requiring a lapse of time to
383 express the bioelectrical response and suggests therefore that Trp does not act
384 directly on particular membrane effectors. A complementary study should now be
385 carried out to identify the steps triggered before the onset of the hyperpolarization
386 with a particular focus on the possible involvement of a Ca^{2+} signaling process.
387 Indeed, Ca^{2+} is considered to be a ubiquitous internal second messenger that
388 regulates various cellular processes acting through intracellular $[\text{Ca}^{2+}]$ increase in
389 response to various environmental stimuli (Knight, 2000) or following recognition of
390 elicitors (Lecourieux et al., 2002).

391 (b) The membrane hyperpolarization could result either from cation effluxes or
392 anion influxes. However, close lag times in $\Delta\Psi$ and ΔpH variations (Table 1) and
393 similar time courses of the bioelectrical and the pH variations noted in the bathing
394 medium of pulvini strongly suggest that proton efflux may be the main process
395 sustaining the hyperpolarization (Fig. 1). However, it can be concluded from
396 experiments performed on PMVs that Trp did not increase the activity of the H^+ -
397 ATPase by directly acting on the pump (Table 2) when compared to FC that also
398 hyperpolarized the pulvinar cells and activated the H^+ -ATPase activity in PMVs
399 (Moyen et al., 2007). This outcome clearly demonstrated different modes of action
400 for these two compounds. Obviously, the membrane potential variations triggered by
401 Trp strongly differed from those evoked by OHTrp (Fig. 1) and glycine (Roblin et al.
402 2018) on *M. pudica* motor cells and by various amino acids on *Lemna gibba* cells
403 (Fisher and Lüttge, 1980) and on barley mesophyll cells (Felle and Zimmermann,
404 2007). In these latter cases, potential variations appear as depolarization spikes after
405 a latency period reduced to seconds and followed by the return to the basal potential
406 in approximately 30 min. Felle and Zimmermann (2007) monitored the temporal
407 sequence of the ionic migrations showing that Ca^{2+} was the first ion to move and was
408 required to trigger the subsequent Cl^- and K^+ fluxes. As shown in Fig. 1B, the time
409 course of the proton translocation appears complex possibly because of a dual role of
410 Trp. In the first minutes following its application, Trp induced a continuous H^+ efflux
411 from cells reminding of an auxin-like effect, as observed with IAA (Bourbouloux et
412 al., 1994) and, in a second step, the observed H^+ influx may be related to its uptake
413 (Fig. 1C) via a H^+ /substrate cotransport mechanism evidenced in amino acid

414 transport by plant cells (Delrot et al., 2001). The recorded curve represents the
415 resultant of the bidirectional proton migration (See additional data S1).

416 (c) By comparing the time course of bioelectrical variations and proton fluxes,
417 it should be stressed that both the plateau and the repolarization phase did not
418 coincide with the beginning of the pH increase recorded in the bathing medium,
419 suggesting that H⁺ influx into the cells may not be the only driving component
420 involved in this latter process. Considering the ionic species (Cl⁻ and K⁺) underlying
421 the action potential (Samejima and Sibaoka, 1980), it can be expected that these
422 pivotal ions may also contribute to the early bioelectrical modifications triggered by
423 Trp. The time course of pH variations also evidenced that H⁺ fluxes were disturbed
424 in a long-lasting manner (for hours) and, could therefore be considered to be
425 involved in the regulation of long-term processes. In this direction, H⁺ fluxes were
426 suggested to be involved in elicitation processes (Schaller and Oecking, 1999) and
427 cytosolic acidification to act as a second messenger, mediating MAP kinase
428 activation in the response of plant cells to various stresses (Tena and Renaudin,
429 1998).

430 By considering its electrogenic properties shown here, Trp may constitute an
431 intermediary in long distance signaling by inducing an electrical stimulus through its
432 transport over long distance in plant as other amino acids. Thus, Trp may create a
433 direct cell-to-cell electrical conduction following its accumulation and therefore may
434 be implicated in plant resistance processes. Indeed, action potentials or wound-
435 induced potential variations are able to transmit information over long distances
436 (Fromm and Lautner, 2007) and to induce the expression of pathogen response genes
437 (Fisahn et al., 2004). However, Trp is unlikely to be directly involved in the
438 Hypersensitive Response as revealed by the assay of Ueno et al. (2003) showing that
439 following infection with *Magnaporthe grisea*, H₂O₂ formation and subsequent
440 lesions on the leaves were due to an accumulation of Trp resulting from increased
441 Trp synthesis.

442

443 *4.3. Tryptophan as an osmotic agent*

444

445 The data obtained on the osmocontractile reaction of pulvinar cells (Fig. 1, 2)
446 indicated that the application of Trp modified bulk ionic migration and the associated
447 water movement by considering the well documented mechanism driving motor cell

448 shrinkage following the application of shock and darkness. Large K^+ and Cl^- effluxes
449 (Samejima and Sibaoka, 1980; Moran, 2007) induced a rapid decrease in cell turgor.
450 The ionic migrations are likely regulated by the activity of K^+ and Cl^- channels and
451 the subsequent water movement allowed by the activity of aquaporins present in
452 motor cell membranes (Fleurat-Lessard et al., 1997; Temmei et al., 2005; Moran,
453 2007). As seen in Fig. 1C, there is a close relationship between the extent of
454 inhibition of the motor reaction of *M. pudica* pulvini and the amount of Trp available
455 in the vicinity of the motor cells. The fact that the sole drooping movement was
456 affected but not the recovery argues for a process involving a vectorial water influx,
457 resulting from the hindering of the K^+ efflux. This hypothesis is corroborated by the
458 Trp-induced inhibitory effect on the dark-induced leaflet movement of *C.*
459 *fasciculata*, (Fig. 2A) and by the Trp-promoting effect observed on the light-induced
460 leaflet movement (See Supplementary data 2). However, it should be stressed that
461 there is no direct link between motile reaction (and the underlying K^+ and water
462 fluxes) and the process of Trp uptake, since the uptake of amino acids is stimulated
463 at an acidic pH (Otsiogo-Oyabi and Roblin, 1985; and reference therein) whereas
464 leaflet movement was promoted at neutral/alkaline pH levels (Fig. 2C). These
465 remarks concerning an increase in water availability in the cells are in accordance
466 with some data showing that Trp increased the water content of wheat and maize
467 cells and reduced the effect of drought in maize (Rao et al., 2012; and references
468 therein).

469

470 *4.4. The effects induced by tryptophan are specific in several aspects*

471

472 Many previous studies have claimed that the physiological responses seen after
473 an exogenous application of Trp may result from the conversion of Trp into IAA
474 (Abdel-Monem et al., 2010; and references therein). However, other works have also
475 shown that Trp exerts a direct effect on plant cell physiology in processes as different
476 as embryo germination in *Triticum aestivum* (Morris et al., 1988) and on cell
477 expansion in *Arabidopsis* mutants (Jing et al., 2009). A major point to ascertain in
478 this study was that Trp acted itself and that the effects may not be attributed to
479 products of its metabolism, in particular to the possible synthesis of IAA. On the
480 motor cell model, Trp induced similar physiological effects as IAA
481 molecules regarding the induction of membrane hyperpolarization, the inhibitory

482 effect on the shock-induced reaction (unpublished result), the inhibitory effect on the
483 dark-induced reaction and the promoting effect on the light-induced reaction (Everat-
484 Bourbouloux et al., 1990). It should be emphasized that Trp is less effective than
485 IAA, since 1 mM of the amino acid was required to trigger reactions of similar
486 amplitude as those obtained with 10 μ M IAA. However, Trp triggers a complex
487 biphasic change in H⁺ fluxes as previously described (Fig. 1B) in contrast with IAA
488 and FC which continuously acidified the bathing medium of motor organs
489 (Bourbouloux et al., 1994). This pattern of Trp responses looks similar to those
490 obtained with 2,4-D. Indeed, the herbicide also induced at first a pH acidification
491 followed by a pH rise in relation to its phenolic structure (Moyen et al., 2007),
492 indicating that the proton exchanges are modulated according to the effect triggered
493 by each moiety of the molecular complex. The fact that Trp is the active component
494 was first supported by the short latency (within min) occurring between the Trp
495 application and the observation of the cell responses which tended to rule out a
496 putative effect of a compound resulting from the metabolism of Trp by the motor
497 cells. Such eventuality was disproven with the use of [³H] labelled Trp combined
498 with TLC analysis, which evidenced a very low rate of metabolism during the first 2
499 h of Trp uptake by the tissues.

500 We have also characterized the specific effect of Trp by including in our study
501 the effects of compounds involved in its synthesis (Ind and Ser) and its metabolism
502 (Try). Clearly, we have shown that these compounds acted through mechanisms
503 which are fundamentally different from those underlying Trp effects, although they
504 can result in the same final osmoregulated reaction of the pulvinar cells. Thus, we
505 showed that Try acted on membrane functioning through a protonophore effect and
506 that Ind acted directly on both processes of the H⁺-ATPase activity as observed on
507 PMVs. In previous reports, we showed that disturbing the H⁺-ATPase functioning
508 resulted in modification of the motile reactions (Saeedi et al., 2013). However, an
509 additional effect has to be considered, since a sole protonophore action would induce
510 a pH alkalisation in the bathing medium of motor tissues as observed with
511 dinitrophenol (Saeedi et al., 2013). In contrast, Try induced a continuous
512 acidification of the medium, thus possibly meaning that Try induced a modification
513 of the membrane permeability leading to a leakage of protons. This continuous H⁺
514 efflux may be one step in a chain reaction leading to a necrotic effect linked to H₂O₂
515 formation as observed in rice infected with the fungus *Magnaporthe grisea* (Ueno et

516 [al., 2003](#)). In the case of Ind, stronger ionic disorders have to be considered as
517 indicated by the also unexpected medium acidification and by the observation of the
518 large gradual increase in initial angle in course of the assay (see inset in [Fig. 1D](#)).
519 These last data may indicate that Ind drastically disturbed cell functioning and led to
520 a necrosis of the pulvinar cells upon increased treatment duration that requires
521 further microscopic observations to identify the wounded cell compartments.

522 It can further be highlighted that, compared to many other amino acids, Trp
523 affects in a specific way the electrophysiological events, the proton exchange status
524 and more generally the ionic migrations as indicated by the data on the stimulus-
525 induced motile reactions. Of note, the Trp-induced membrane potential changes
526 strongly differed from those induced by the structurally related compound OHTrp
527 which triggered a spiky depolarization of the motor cell membrane identical to that
528 evoked by glycine application ([Roblin et al; 2018](#)). Furthermore, differences were
529 also apparent in the shock- and dark-induced reactions: Trp strongly inhibited these
530 reactions while glycine promoted them ([Otsiogo-Oyabi and Roblin, 1984](#)).
531 Interestingly, arginine, cysteine, phenylalanine and Ser also inhibited the dark-
532 induced reactions in *C. fasciculata* leaflets when observed after 3 hours of action
533 ([Fig. 2E](#)). The case of Ser is interesting since this amino acid combines with Ind to
534 form Trp ([Kriechbaumer et al., 2008](#)), so that we can suppose that the effect observed
535 4 h after the application on the leaves could result from Trp synthesis. Altogether,
536 these data prompted us to investigate further whether these particular amino acids
537 may play particular roles in the plant cell functioning. In this way, a recent study
538 concerning cysteine has confirmed that this amino acid also induced characteristic
539 effects on the processes studied here ([Roblin et al., 2018](#)).

540

541 **5. Conclusions**

542 The data highlight that some amino acids may have specific roles outside of a
543 simply metabolic role as demonstrated by the characteristic properties evoked by Trp
544 in plant cell physiology. Further experiments should be devoted to the underlying
545 mechanisms of these non-metabolic effects, since some actions of Trp did not
546 correspond to a classical mode of action as compared to other amino acids. In
547 particular, the mechanism sustaining the variation of membrane permeability to
548 water and ions should be investigated further, as it does not result from an activation
549 of the proton pump activity, raising the hypothesis of a direct action on ionic

550 channels (K^+ channels in particular). Another pivotal line of research would be
551 focused on the processes linked to the membrane sensing of Trp. The perception at
552 the membrane level may be achieved through a specific receptor as shown for IAA
553 (Woodward and Bartel, 2005) or may be induced through a transceptor (Dinkeloo et
554 al., 2018) linking transport function and signaling in the cells.

555

556

557

558

559

560 **Supplementary material**

561 Supplementary data S1. Putative explanation of the complex time course of pH
562 variations monitored in the bathing medium of pulvinar tissues of *M. pudica*
563 following application of tryptophan.

564 Supplementary data S2. Effect of 1 mM tryptophan and 1 mM serine on the light-
565 induced opening of *Cassia fasciculata* leaflets.

566

567 **Author contributions**

568 Christelle Moyen performed the electrophysiological studies, Fabienne
569 Dédaldéchamp participated to the assays on the metabolism experiments, Pierrette
570 Fleurat-Lessard and Gabriel Roblin carried out the observations on leaf reactions and
571 the analysis carried out on PMVs. All the authors participated in the writing of the
572 manuscript and approved the final version.

573

574 **Acknowledgments**

575 This work was supported by UMR CNRS 7267 Laboratoire EBI Ecologie et Biologie
576 des Interactions, Equipe SEVE Sucres & Echanges Végétaux Environnement,

577

578

References

- Abdel-Monem, A.A., El-Bassiouny, H.M.S., Rady, M.M., Gaballah, M.S., 2010. The role of tryptophan and Prozac (5-hydroxytryptophan) on the growth, some biochemical aspects and yield of two sunflower cultivars grown in saline soil. *Int. J. Acad. Res.* 2, 254-262.
- Bednarek, P., Piślewska-Bednarek, M., Svatoš, A., Schneider, B., Doubský, J., Mansurova, M., Humphry, M., Consonni, C., Panstruga, R., Sanchez-Vallet, A., Molina, A., Schulze-Lefert, P., 2009. A glucosinolate metabolism pathway in living plant cells mediates broad-spectrum antifungal defense. *Science* 323, 101-106.
- Bourbouloux, A., Fleurat-Lessard, P., Roblin, G., 1994. Effects of jasmonic acid on motor cell physiology in *Mimosa pudica* L. and *Cassia fasciculata* Michx. *Plant Cell Physiol.* 35, 389-396.
- Colcombet, J., Mathieu, Y., Peyronnet, R., Agier, N., Lelièvre, F., Barbier-Brygoo, H., Frachisse, J.M., 2009. R-type anion channel activation is an essential step for ROS-dependent innate immune response in *Arabidopsis* suspension cells. *Funct. Plant Biol.* 36, 832-843.
- Delrot, S., Atanassova, R., Gomès, S., Coutos-Thévenot, P., 2001. Plasma membrane transporters: a machinery for uptake of organic solutes and stress resistance. *Plant Sci.* 161, 391-404.
- Dinkeloo, K., Boyd, S., Pilot, G., 2018. Update on amino acid transporter functions and on possible amino acid sensing mechanisms in plants. *Seminars in Cell & Developmental Biol.* 74, 105-113.
- Everat-Bourbouloux, A., Fleurat-Lessard, P., Roblin, G., 1990. Comparative effects of indole-3-acetic acid, gibberellic acid and benzylaminopurine on the dark- and light-induced pulvinar movements in *Cassia fasciculata*. *J. Exp. Bot.* 41, 315-324.
- Felle, H.H., Zimmermann, M.R., 2007. Systemic signalling in barley through action potentials. *Planta* 226, 203-214.
- Fisahn, J., Herde, O., Willmitzer, L., Pena-Cortès, H., 2004. Analysis of the transient increase in cytosolic Ca²⁺ during the action potential of higher plants with high temporal resolution: requirement of Ca²⁺ transients for induction of jasmonic acid biosynthesis and PINII gene expression. *Plant Cell Physiol.* 45, 456-459.

- Fisher, E., Lüttge, U., 1980. Membrane potential changes related to glycine transport in *Lemna gibba* G1. *Plant Physiol.* 65, 1004-1008.
- Fleurat-Lessard, P., Bouché-Pillon, S., Leloup, C., Bonnemain, J-L., 1997. Distribution and activity of the plasma membrane H⁺-ATPase in *Mimosa pudica* L. in relation to ionic fluxes and leaf movements. *Plant Physiol.* 113, 747-753.
- Frerigmann, H., Pislewska-Bedarek, M., Sánchez-Vallet, A., Molina, A., Glawischnig, E., Gigolashvili, T., Bednarek, P., 2016. Regulation of pathogen-triggered tryptophan metabolism in *Arabidopsis thaliana* by MYB transcription factors and indole glucosinolate conversion products. *Molec. Plant* 9, 682-695.
- Fromm, J., Lautner, S., 2007. Electrical signals and their physiological significance in plants. *Plant Cell Envir.* 30, 249-257.
- Ishihara, A., Hashimoto, Y., Tanaka, C., Dubouzet, J.G., Nakao, T., Matsuda, T., Miyagawa, H., Wasaka, K., 2008. The tryptophan pathway is involved in the defense responses of rice against pathogenic infection via serotonin production. *Plant J.* 54, 481-495.
- Jing, Y., Cui, D., Bao, F., Hu, Z., Qin, Z., Hu, Y., 2009. Tryptophan deficiency affects organ growth by retarding cell expansion in *Arabidopsis*. *Plant J.* 57, 511-521.
- Kang, K., Kang, S., Lee, K., Park, M., Back, K., 2008. Enzymatic features of serotonin biosynthesis enzymes and serotonin biosynthesis in plants. *Plant Signal. Behav.* 3, 389-390.
- Knight, H., 2000. Calcium signaling during abiotic stress in plants. *Internat. Rev. Cytol.* 195, 269-324.
- Kriechbaumer, V., Weigang, L., Fießelmann, A., Letzel, T., Frey, M., Gierl, A., Glawischnig, E., 2008. Characterisation of the tryptophan synthase alpha subunit in maize. *BMC Plant Biol.* 8, 44.
- Lecourieux, D., Mazars, C., Pauly, N., Ranjeva, R., Pugin, A., 2002. Analysis and effects of cytosolic free calcium elevation in response to elicitors in *Nicotiana plumbaginifolia* cells. *Plant Cell* 14, 2627-2641.
- Moffett, J.R., Namboodiri, M.A., 2003. Tryptophan and the immune response. *Immunol. Cell Biol.* 81, 247-265.
- Moran, N., 2007. Osmoregulation of leaf motor cells. *FEBS Lett.* 581, 2337-2347.

- Morris, C.F., Mueller, D.D., Faubion, J.M., Paulsen, G.M., 1988. Identification of L-tryptophan as an endogenous inhibitor of embryo germination in white wheat. *Plant Physiol.* 88, 435-440.
- Moyen, C., Bonmort, J., Roblin, G., 2007. Membrane effect of 2,4-dichlorophenoxyacetic acid in motor cells of *Mimosa pudica* L. *Plant Physiol. Biochem.* 45, 420-426.
- Otsiogo-Oyabi, H., Roblin, G., 1984. Effects of glycine on dark- and light-induced pulvinal movements and modification of proton fluxes in the pulvinus of *Mimosa pudica* during glycine uptake. *Planta* 161, 404-408.
- Otsiogo-Oyabi, H., Roblin, G., 1985. Uptake of glycine by excised pulvini of *Mimosa pudica* in relation to proton pump activity. *Physiol. Plant.* 65, 40-44.
- Radwanski, E.R., Last, R.L., 1995. Tryptophan biosynthesis and metabolism: biochemical and molecular genetics. *Plant Cell* 7, 921-934.
- Rao, S.R., Qayyum, A., Razzaq, A., Ahmad, M., Mahmood, I., Sher, A., 2012. Role of foliar application of salicylic acid and L-tryptophan in drought tolerance of maize. *J. Anim. Plant Sci.* 22, 768-772.
- Roblin, G., Octave, S., Faucher, M., Fleurat-Lessard, P., Berjeaud, J-M., 2018. Cysteine: a multifaceted amino acid involved in signalling, plant resistance and antifungal development. *Plant Physiol. Biochem.* 129, 77-89.
- Saeedi, S., Rocher, F., Bonmort, J., Fleurat-Lessard, P., Roblin, G., 2013. Early events induced by salicylic acid in motor cells of *Mimosa pudica* pulvinus. *J. Exp. Bot.* 64, 1829-1836.
- Samejima, M., Sibaoka, T., 1980. Changes in the extracellular ion concentration in the main pulvinus of *Mimosa pudica* during rapid movement and recovery. *Plant Cell Physiol.* 21, 467-479.
- Sanjaya, Hsiao, P-Y., Su, R-C., Ko, S-S., Tong, C-G., Yang, R-Y., Chan, M-T., 2008. Overexpression of *Arabidopsis thaliana* tryptophan synthase beta 1 (*AtTSB1*) in *Arabidopsis* and tomato confers tolerance to cadmium stress. *Plant Cell Environ.* 31, 1074-1085.
- Schaller, A., Oecking, C., 1999. Modulation of plasma membrane H⁺-ATPase activity differentially activates wound and pathogen defense responses in tomato plants. *Plant Cell* 11, 263-272.
- Serrano, R., 1989. Structure and function of plasma membrane ATPase. *Annu. Rev. Plant Physiol. Plant Molec. Biol.* 40, 61-94.


- Temmei, Y., Uchida, S., Hoshino, D., Kanzawa, N., Kuwahara, M., Sasaki, S., Tsuchiya, T., 2005. Water channel activities in *Mimosa pudica* plasma membrane intrinsic proteins are regulated by direct interaction and phosphorylation. *FEBS Lett.* 579, 4417-4422.
- Tena, G., Renaudin, J., 1998. Cytosolic acidification but not auxin at physiological concentration is an activator of MAP kinases in tobacco cells. *Plant J.* 16, 173-182.
- Ueno, M., Shibata, H., Kihara, J., Honda, Y., Arase, S., 2003. Increase tryptophan decarboxylase and monoamine oxidase activities induce Sekiguchi lesion formation in rice infected with *Magnaporthe grisea*. *Plant J.* 36, 215-228.
- Volkov, A.G., Foster, J.C., Ashby, T.A., Walker, R.K., Johnson, J.A., Markin, V.S., 2010. *Mimosa pudica*: electrical and mechanical stimulation of plant movements. *Plant Cell Environ.* 33, 163-173.
- Woodward, A.W., Bartel, B., 2005. A receptor for auxin. *Plant Cell* 17, 2425-2429.
- Zemanova, V., Pavlik, M., Pavliková, D., Tlustos, P., 2014. The significance of methionine, histidine and tryptophan in plant response and adaptation to cadmium stress. *Plant Soil Environ.* 60, 426-432.
- Zhao, J., Williams, C.C., Last, R.L., 1998. Induction of Arabidopsis tryptophan pathway enzymes and camalexin by amino acid starvation, oxidative stress, and an abiotic elicitor. *Plant Cell* 10, 359-370.

Table 1. Effect of tryptophan (Trp) at various concentrations, 5-hydroxytryptophan (OHTrp) at 1 mM and serine (Ser) at 1 mM on the characteristics of the transmembrane potential of pulvinar motor cells and on the characteristics of the proton fluxes measured in the bathing medium of excised pulvini of *Mimosa pudica*. Calculations of $\Delta\Psi$ and of the amount of mobilized protons were respectively made 10 min and 90 min after application of the compounds. n: number of assays. Mean \pm SD. Superscript letters indicate significant differences at the 5% probability by Student-Fisher *t* test (*b-f*) or values not significantly different from the control (*a*). +, Hyperpolarization; -, depolarization.

Treatment	Membrane potential changes				Proton fluxes					
	[Conc]	Latency	$\Delta\Psi$	n	Latency	Acidification duration	H ⁺ excreted	pH rise duration	H ⁺ absorbed	n
	(mM)	(s)	(mV)		(s)	(min)	(nmole)	(min)	(nmole)	
Control	0	-	+1 \pm 1 ^a	(5)	-	-	-	-	-	(3)
Trp	0.05	480 \pm 80 ^b	+7 \pm 7 ^b	(5)	-	-	-	-	-	(3)
Trp	0.1	360 \pm 60 ^c	+24 \pm 3 ^c	(4)	370 \pm 30 ^b	60 \pm 5 ^b	15 \pm 5 ^b	80 \pm 8 ^b	24 \pm 6 ^b	(3)
Trp	0.5	200 \pm 48 ^d	+35 \pm 8 ^d	(4)	180 \pm 12 ^c	80 \pm 5 ^c	54 \pm 8 ^c	100 \pm 10 ^c	66 \pm 8 ^c	(3)
Trp	1	120 \pm 36 ^e	+48 \pm 8 ^e	(5)	138 \pm 24 ^d	96 \pm 4 ^d	90 \pm 8 ^d	150 \pm 10 ^d	105 \pm 10 ^d	(4)
OH-Trp	1	15 \pm 1 ^f	-28 \pm 6 ^e	(6)	240 \pm 72 ^e	-	-	180 \pm 10 ^e	135 \pm 10 ^e	(4)
Ser	1	19 \pm 6 ^f	-14 \pm 5 ^f	(4)	120 \pm 20 ^f	-	-	60 \pm 5 ^e	105 \pm 8 ^d	(3)

Table 2. Modifications of net proton movement (measured by absorbance variation from the control absorbance at 495 nm) and vanadate-sensitive ATPase activity in plasma membrane vesicles purified from *Mimosa pudica* pulvini treated with tryptophan, 5-hydroxytryptophan, tryptamine, indole, serine and vanadate. Values are mean \pm SD. Into brackets, percentage of inhibition (-) or activation (+). Letters following bracket indicate significant differences at the 5% probability by Student-Fisher *t* test (*b*, *c*) or values not significantly different from the control (*a*).

Product	Concentration (mM)	Absorbance variation (unit mg.prot ⁻¹ .min ⁻¹)	n	Vanadate-sensitive activity (nmol Pi.mg prot ⁻¹ .min ⁻¹)	n
Control	0	0.35 \pm 0.03 <i>a</i>	8	433 \pm 17 <i>a</i>	8
Tryptophan	1	0.34 \pm 0.01 [-3] <i>a</i>	8	452 \pm 28 [+4] <i>a</i>	8
Tryptamine	1	0.05 \pm 0.03 [-86] <i>b</i>	6	444 \pm 23 [+3] <i>a</i>	8
Indole	1	0.05 \pm 0.02 [-86] <i>b</i>	8	272 \pm 43 [-37] <i>b</i>	8
Vanadate	0.25	0.14 \pm 0.06 [-60] <i>c</i>	8	147 \pm 25 [-66] <i>c</i>	8
5OH Tryptophan	1	0.31 \pm 0.02 [-11] <i>a</i>	4	421 \pm 15 [-3] <i>a</i>	4
Serine	1	0.36 \pm 0.05 [+3] <i>a</i>	3	441 \pm 13 [+2] <i>a</i>	4



579

Fig. 1. Characteristics of the early events induced by tryptophan (Trp) in primary pulvini of *Mimosa pudica*. (A) Typical recordings of transmembrane potential variations in pulvinar parenchyma motor cells induced by tryptophan at various concentrations (0.05, 0.1, 0.5, 1 mM) and by 1 mM 5OH-tryptophan (OHTrp). Compounds were added at time 0 (black arrow) and 10 μ M fusicoccin (FC) was added at 60 min (black arrowheads). The number of assays is indicated in Table 1. (B) Representative time courses of pH variations monitored in the bathing medium of pulvinar tissues induced by various compounds. In the upper part, effect of tryptophan at various concentrations (0.1, 0.5, 1 mM) and of 1 mM 5OH-tryptophan (OHTrp). In the lower part, effect of serine (Ser), tryptamine (Try) and indole (Ind) applied at 1 mM. The experiments were conducted at least three times with the same general result. (C) Time courses of the inhibition of the seismonastic pulvinar osmocontractile reaction following treatment with tryptophan at various concentrations expressed in mM (mean \pm SE, $n = 30$) and time course of the amount of 1 mM labelled tryptophan absorbed by the pulvinar tissues from excised leaves (white symbols, dotted line) and effect of CCCP (Δ) at 10 μ M on the uptake (mean \pm SD, $n = 4$). (D) Comparative time courses of the inhibition of the seismonastic pulvinar osmocontractile reaction observed after application of serine, 5OH-tryptophan, tryptamine and indole (at 1 mM) (mean \pm SE, $n = 30$). In inset, initial angle in degrees measured at the beginning of the treatment (α_{i0}) and after 4 h (α_{i4}) (mean \pm SE, $n = 30$). Co: control.

580




Fig. 2. Characteristics of the inhibitory effect induced by tryptophan on the dark-induced closure of *Cassia fasciculata* leaflets. (A) Modification in the time course of the closure as a function of concentration applied (from 0.1 to 5 mM) for 3 h before the darkness (black arrow). (B) Effect of 1 mM tryptophan treatment duration. (C) Effect of the pH of the incubation medium as indicated in controls (black columns) and in 1 mM tryptophan-treated leaves (white columns). (D) Effects of 1 mM, tryptophan in the L- and D-form, 1 mM 5OH-tryptophan (OH), and of indole (Ind) and tryptamine (Try) at various concentrations (a, b, c respectively 0.1, 0.5 and 1 mM). (E) Comparative effect of tryptophan with other natural amino acids (Co: control; a: tryptophan, b: alanine; c: arginine; d: asparagine; e: aspartic acid; f: cysteine; g: glutamine; h: glutamic acid; i: glycine; j: histidine; k: hydroxyproline, l: isoleucine; m: leucine; n: lysine; o: methionine; p: phenylalanine; q: proline; r: serine; s: threonine; t: tyrosine; u: valine). Otherwise indicated, the calculations were made 1 h after the start of darkness on sets of leaflets treated for 3 h with the compounds prepared in buffer at pH 5.5. Mean \pm SE; n = 24. Superscript letters and asterisks in (E) indicate significant differences at 5% probability level by Student-Fisher *t* test (*b-d*) or values not significantly different from the control (*a*).

581