

# Towards a Better Understanding of Emotion Blends: The Case of Anger-Related Emotion Blends Elicited Via Film Clips

Alex Lafont, Joceline Roge, Daniel Ndiaye, Jean-Michel Boucheix

### ▶ To cite this version:

Alex Lafont, Joceline Roge, Daniel Ndiaye, Jean-Michel Boucheix. Towards a Better Understanding of Emotion Blends: The Case of Anger-Related Emotion Blends Elicited Via Film Clips. Cognition, Brain, Behavior. An Interdisciplinary Journal, 2019, 2 (23), pp.77-99. 10.24193/cbb.2019.23.05. hal-02298345

## HAL Id: hal-02298345 https://hal.science/hal-02298345v1

Submitted on 26 Sep 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	
2	Towards a Better Understanding of Emotion Blends: The Case of Anger-Related Emotion
3	Blends Elicited Via Film Clips
4	
5	Abstract
6	We sought to better understand the effect of film clips supposed to elicit different intensities
7	of anger on feelings and cardiac activity among our participants. In experiment 1, we tested in
8	21 volunteers the ability of two film clips to elicit two intensities of anger: a slight one and
9	strong one. In experiment 2, we used the same film clips in order to investigate, in 36
10	volunteers, how cardiovascular patterns may differ between the emotional states we elicited.
11	Results suggest that the film clips produced a physiological reappraisal pattern, with a lower
12	heart rate and higher heart rate variability values (RMSSD and SD1) during exposure to film
13	clips that were selected to elicit anger. Results of both experiments also suggest that film clips
14	selected to elicit anger, not only triggered anger, but also other negative emotions.
15	Consequently, we propose a method of assessing emotions which takes emotion blends into
16	account.
17	Keywords: emotion; reappraisal; heart rate variability; film clips
18	
19	
20	
	Alex Lafont 1, Joceline Rogé 1, Daniel Ndiaye 2, Jean-Michel Boucheix 3
	1 Université Lyon, IFSTTAR, TS2, LESCOT, F-69675 Lyon, France 2 Université Paris-Est, COSYS, LEPSIS, IFSTTAR, F-77447 Marne-la-Vallée, France

3 Université de Bourgogne Franche-Comté, LEAD-CNRS, F-21065 Dijon, France

LAFONT, Alex, ROGE, Joceline, NDIAYE, Daniel, BOUCHEIX, Jean-Michel, 2019, Towards a Better Understanding of Emotion Blends: The Case of Anger-Related Emotion Blends Elicited Via Film Clips, Cognition, Brain, Behavior. An Interdisciplinary Journal, 2, XXIII, 77-99, DOI: 10.24193/ cbb.2019.23.05

"There are more than a thousand definitions of what emotion is and the only point on 21 which everyone agrees, is that the concept is hard... to define" (Belzung, 2007). According to 22 Kleinginna and Kleinginna, (1981) who analyzed and classified 92 definitions of emotion, it 23 can be argued that emotion consists of four main components: a) cognitive b) subjective 24 c) physiological d) behavioral. Although different ordering of these components could be 25 practically all the time defended, cognitive, subjective and physiological features appear to play 26 an upstream role in emotion emergence (Niedenthal, Krauth-Gruber, & Ric, 2006, p. 10-11; 27 Plutchik, 2001). The present research coupled subjective and cardiac measures in order to 28 investigate these early processes. The aim was to answer the question of what is happening 29 throughout the processing of emotional stimuli supposed to elicit anger with different 30 intensities. Does physiology solely vary according to the intensity of the emotion felt, or 31 physiological variations observed during emotion elicitation could reflect other processes (e.g., 32 33 emotion regulation)? With the aim of addressing this issue, let us firstly discuss about emotion, especially how can it be assessed. 34

35

#### **Subjective Emotional Assessment**

Earliest studies in psychology generally used subjective data to assess individual 36 emotional states. Different approaches to emotion have been conceptualized in order to 37 categorize feelings. Some are based on the view that there is a finite number of emotions 38 (Ekman, 1984, 1992; Izard, 1977). Others are based on multidimensional models in which 39 emotion can be defined as a function of broader features (e.g., Russell, 1980; Watson & 40 Tellegen, 1985, Fontaine, Scherer, Roesch, & Ellsworth, 2007). Both discrete and dimensional 41 approaches to emotion are widely used by psychologists in experimental contexts (Scherer, 42 2005). 43

44 Alternatively, other models have emerged from cognitive appraisal approaches of 45 emotion (e.g., Frijda, 1986; Schachter & Singer, 1962), which assume that there are cognitive

#### TOWARDS A BETTER UNDERSTANDING OF EMOTION BLENDS

processes that evaluate the emotional event, understand what caused it and assess ways of 46 coping with it. In other words, "appraisals are transformations of raw sensory input into 47 psychological representations of emotional significance" (Clore & Ortony, 2000). Scherer 48 (2005) created a tool for subjective emotional assessment, placing appraisal as a key feature of 49 his model. This tool has a specific arrangement of 16 discrete emotions placed around a circle 50 divided along two axes: valence (positive versus negative) along the horizontal line, and coping 51 potential (control/power) along the vertical line. Thus, individuals can rate the intensity of their 52 feelings on the basis of the distance from the hub of the wheel and the size of the circles. In this 53 way, the intensity of emotion can be expressed as the degree to which individuals are affected 54 by emotion (Scherer, Shuman, Fontaine, & Soriano, 2013). The present study is thus based on 55 the appraisal approaches of emotion, especially Scherer's model (2005). However, besides 56 subjective feelings, emotion also affects physiology. 57

58

#### **Emotional Assessment through Cardiac Measures**

In accordance with the neurovisceral integration model (Thayer & Lane, 2000; Thayer 59 & Lane, 2009) there is some evidence supporting brain connections, implicated in affective 60 processes like emotion regulation, and cardiac activity controlled by both the sympathetic and 61 parasympathetic branches of the autonomic nervous system (ANS). According to Kreibig 62 (2010), cardiac measurements, particularly heart rate (HR), defined as the number of 63 contractions of the heart per minute, and heart rate variability (HRV) which refers to the 64 variations between consecutive heartbeats are widely used to investigate the impact of emotion 65 on ANS activity. While sympathetic activity tends to increase HR in a few seconds, 66 parasympathetic activity tends to decrease HR and its response is faster (0.2-0.6 s) (Berntson et 67 al., 1997). 68

#### 69 Cardiac Measures as an Indicator of Anger

According to many authors, the ANS produces specific activations in response to 70 specific emotions (Cacioppo, Klein, Berntson, & Hatfield, 1993; Levenson, Ekman, & Friesen, 71 1990; Stemmler, Aue, & Wacker, 2007). According to Kreibig's review, anger is one of the 72 most widely investigated emotions. Several studies have revealed typical variations in metrics 73 related to cardiac activity while experiencing emotions with high arousal such as anger. For 74 75 instance, many authors have found a significant increase in HR due to anger elicitation (Fairclough & Spiridon, 2012; Fernández et al., 2012; Lobbestael, Arntz, & Wiers, 2008; Marci, 76 Glick, Loh, & Dougherty, 2007). 77

Some studies have assessed HRV to explore emotion using *time-domain* methods. 78 Among the most commonly used indexes suitable for short-term recordings, the root mean 79 square of successive NN interval<sup>1</sup> differences (RMSSD) appears particularly relevant, notably 80 81 indicating short-term variability in cardiac rhythm (Task Force of The European Society of Cardiology and The North American Society of Pacing and Electrophysiology, 1996). Indeed, 82 several studies have shown that the RMSSD value decreases when participants have to carry 83 out an anger-related imagery task (Kreibig, 2010; Pauls & Stemmler, 2003). Consequently, 84 lower RMSSD values are expected when anger is experienced in the current study. 85

Furthermore, the mediation of cardiac activity by complex control systems assumes that non-linear mechanisms are involved in cardiac rhythm variations (Task Force of The European Society of Cardiology and The North American Society of Pacing and Electrophysiology, 1996). Among *non-linear* methods used to study HRV, the Poincaré plot is a popular tool because of the relative ease of interpreting one of the parameters. More specifically, the SD1 index, describes short-term variability in cardiac rhythm (Goshvarpour, Abbasi, &

<sup>&</sup>lt;sup>1</sup> (i.e., the interval from one R peak to the subsequent R peak)

Goshvarpour, 2017; Tarvainen, 2014). Consequently, SD1 should vary similarly to RMSSD
leading to lower values when an individual experiences anger.

Another interesting non-linear measure is related to entropy, which is used to quantify 94 signal complexity and irregularity. Two entropy indexes can be calculated: Approximate 95 entropy (ApEn) and sample entropy (SampEn). Low values for both indexes indicate greater 96 signal regularity. However, SampEn index is more accurate than ApEn. Additionally it is 97 preferred for short-term recordings (See Riganello, Cortese, Arcuri, Quintieri, & Dolce, 2015; 98 Tarvainen, 2014; Valenza, Allegrini, Lanatà, & Scilingo, 2012 for further information about 99 ApEn and SampEn). According to Valenza et al. (2012), entropy tends to decrease significantly 100 during arousal. It is well known that anger is characterized by high arousal (see Russell, 1980). 101 Consequently, lower values of SampEn are expected when anger is experienced. 102

Furthermore, cardiac measures do not only reflect a direct expression of anger.
According to several authors, cardiac activity vary is affected by emotion regulation (León,
Hernández, Rodríguez, & Vila, 2009; Porges, Doussard-Roosevelt, & Maiti, 1994).

#### 106 Cardiac Measures as an Indicator of Cognitive Reappraisal

107 As briefly introduced previously, appraisal theories of emotion focus on immediate cognitive processes of evaluation of meaning, causal attribution and coping capabilities 108 (Niedenthal et al., 2006, p. 13). Regarding the last point (i.e., coping capabilities), it suggests 109 that a stimulus can be reappraised in order to regulate emotions felt by individuals and thus face 110 a psychological stress. According to Gross (1998), emotion regulation depends on the 111 individuals' ability to adjust physiological arousal on a momentary basis (see also Appelhans 112 & Luecken, 2006). As a consequence, cognitive reappraisal could be identified through cardiac 113 114 measures.

115 According to Wranik and Scherer (2010), anger often occurs when one's goals are 116 obstructed. This highlights the importance of considering the cognitive appraisal step,

determining the impact of a specific event on an individual. For instance, Denson, Grisham, 117 and Moulds (2011) asked three groups of participants to watch an anger-inducing video. In one 118 group (control), participants just had to watch the video. In the second group (suppression), 119 they had to try to control their facial expression and to behave in such a way that a person 120 watching them would not know what they were feeling. In the last group (reappraisal), 121 participants had to manage their emotional reactions; they were asked to try to maintain a 122 neutral mood and not to be overwhelmed by the anger elicited by the video. The authors 123 observed a significant increase in HRV (RMSSD) and decreased HR among participants in the 124 reappraisal condition. Their results reflect a physiological correlate of adaptive emotion 125 regulation. This physiological pattern is consistent with the notion of reappraisal as an active 126 and adaptive means of cognitively changing our response to emotional situations. This process 127 allows greater autonomic flexibility, and improves the ability to view anger-inducing events as 128 129 non-threatening. The same kinds of pattern were observed when using pictures to induce a negative emotional state (Kuoppa, Tarvainen, Karhunen, & Narvainen, 2016; Sarlo, Palomba, 130 Buodo, Minghetti, & Stegagno, 2005; Stemmler et al., 2007). 131

Furthermore, as links have been found between Poincaré's SD1 index and time-domain measures (i.e., RMSSD), higher values for SD1 can be expected when emotion-regulation strategies emerge. In addition, as parasympathetic activity is predominant when individuals reappraise emotional events (Denson et al., 2011) and entropy is higher when that branch of ANS is in control (Valenza et al., 2012), it can be assumed that SampEn values will be higher when individuals regulate their emotion.

Hence, a decrease in HR together with an increase in time and non-linear HRV indexes
(RMSSD, SD1 and SampEn) could be expected when emotion-regulation strategies are
implemented.

#### 141 Existence of an Anger Intensity Threshold?

As previously seen, cardiac patterns may vary according to the way that individuals 142 manage their emotions. Several studies investigating different intensities of the same negative 143 emotion have effectively shown that emotion-regulation strategies could depend on the 144 intensity of the elicited emotion (Luce, Payne, & Bettman, 1999; Yao & Lin, 2015). More 145 precisely, Luce et al. (1999) found that high levels of negative emotion induce regulation 146 strategies such as avoidance coping. This raises the question of whether the level of anger 147 intensity produces a specific cardiac pattern, in other words an anger threshold pattern. Indeed, 148 it could be expected that a cardiac pattern reflecting reappraisal strategies would only occur 149 when a high intensity of anger is elicited. By contrast, a cardiac pattern reflecting direct 150 processing of anger should emerge when a moderate intensity of anger is induced. 151

To our knowledge, no study has investigated different levels of anger using both HR and time and non-linear aspects of HRV. In order to see how emotional processing differs as a function of the intensity of the anger experienced, the present study combined subjective and physiological measurements to examine the specific cardiac patterns associated with different intensity levels of anger.

157

#### **Emotion Elicitation**

According to Baldaro et al. (2001), film clips are commonly used to induce emotions 158 because of their high ecological validity and their ease of presentation. Several researchers have 159 developed databases of film clips suitable for mood induction procedures (Gross & Levenson, 160 1995; Schaefer, Nils, Sanchez, & Philippot, 2010). Prior research used the most recent database 161 to study emotion. For instance, Fernández et al. (2012) investigated the objective physiological 162 responses (skin conductance level and heart rate) and subjective emotional responses elicited 163 through a set of emotional film clips capable to induce anger, fear, sadness, disgust amusement 164 tenderness and a neutral state. In the same vein, the Schaefer's database (2010) has been used 165

#### TOWARDS A BETTER UNDERSTANDING OF EMOTION BLENDS

to compare different types of real-life emotional elicitations: film clips eliciting anger, fear or
sadness and a hostile social interaction (Cabral, Tavares, Weydmann, das Neves, & de Almeida,
2018). However; no study focused on the ability for the film clips belonging to this database to
elicit different intensities of a same emotion. Moreover, no study investigated the propensity
for these film clips to elicit emotion blends. In the current study, the Schaefer's database (2010)
thus enabled us to select relevant film clips for the study's objective.

In line with all the studies above, we expected that the chosen film clips (see below) would elicit two intensities of anger (slight and strong). In addition, we expected that the physiological patterns observed would reflect either a classic anger pattern, or could reflect a cognitive reappraisal pattern depending on the intensity of anger elicited.

176

#### **Experiment 1**

The goal of Experiment 1 was to study the capacity of specific film clips to elicit 177 178 different intensities of anger. In addition, we sought to explore the potential presence of other emotions felt while exposed to the film clips. In this context, emotional state was assessed 179 through the ratings of the intensities felt by the participants for eight negative emotions while 180 exposed to specific film clips. Given the classification of Schaefer and colleagues (2010), we 181 assumed that intensity rating of anger would be lower with the slight anger film clip than with 182 one expected to elicit strong anger. This assumption was called the *subjective feeling* 183 hypothesis. 184

185 Method

Participants and design. Twenty-one participants (volunteers) took part in the study, 12 males and 9 females, aged from 17 to 51 years (M = 29; SD = 11.5). All participants reported normal or corrected-to-normal vision and audition. The experimental design consisted of a counterbalanced within-subjects film-clip factor (neutral, slight anger, and strong anger). The dependent variable was the subjective intensity of eight negative emotions (i.e., anger, 191 frustration, contempt, disgust, guilt, shame, fear and sadness). Intensity was rated using a score192 varying from 0 to 100 for each emotion.

Material. We used the Schaefer's validated emotional film clip database in which we 193 selected three film clips on the basis of their propensity to elicit two different levels of anger 194 and a neutral state. For the anger-inducing film clips, the selection was based first on a set of 195 10 film clips with high anger discreetness scores (i.e., mean score obtained on the anger item 196 of the Differential Emotional Scale (DES) (Izard, Dougherty, Bloxom, & Kotsch, 1974), minus 197 mean scores obtained on all remaining items of the DES). Note that DES scale allows, through 198 the rating of groups of emotional adjectives, to assess the ability of stimuli to trigger 199 differentiated emotional feeling states. Then, two of the film clips with the highest anger 200 discreetness scores (i.e., film clips with the highest propensity to elicit anger) were selected on 201 the basis of their mean anger scores on the DES. In this way, we chose two film clips that 202 203 potentially elicited anger at different levels: slight (no. 30 with a score of 3.16) and strong (no. 2 with a score of 5.04). For neutral film clip, which was used as a control condition, we selected 204 205 a clip with a very low level on the anger item of DES (score of 1). Precautions were also taken 206 regarding this clip, for which no other item of DES exceeded a score of 1, thus ensuring that the film clip could be considered as neutral (no. 49). 207

The neutral film clip showed a woman walking along the street, the one inducing slight 208 209 anger showed a man undergoing intense interrogation, and the one inducing strong anger showed a man randomly shooting people from his balcony. Unfortunately, the film clips were 210 of different length. Hence, it was necessary to make some changes to obtain three comparable 211 videos. First, we shortened the film clip inducing slight anger by cutting some scenes that did 212 not affect the narrative structure. Secondly, we extended the neutral film clip using a video 213 214 scene that matched the rest of the clip, including color and sound, as closely as possible. In this way, we obtained three film clips of comparable length (mean length =  $1 \min 55$  s; SD = 10 s). 215

To measure the dependent variable, we used the Emotional Wheel (EW) (Rogé, El 216 Zufari, Vienne, & Ndiaye, 2015) inspired by the Geneva Emotion Wheel (Scherer, 2005), which 217 was specifically designed to combine both discrete and dimensional approaches in self-report 218 assessments of emotion. The EW provides both valence and intensity information about a set 219 of 16 emotions: eight negative (i.e., anger, frustration, contempt, disgust, guilt, shame, fear and 220 sadness) and eight positive (i.e., pride, elation, happiness, satisfaction, relief, hope interest and 221 surprise). As two film clips were selected in order to mainly elicit anger (i.e., negative emotion), 222 positive emotions were seen as epiphenomena. Therefore, only negative emotions were 223

analyzed in this article (also see Rogé et al., 2015 who did the same).

Procedure. Participants were seated comfortably in a quiet room, and completed the 225 experiment individually. They watched the three film clips, for which the presentation order 226 has been fully counterbalanced, on a computer screen (visual angle of 32°x21°) and were asked 227 228 to rate their emotional state after each clip. To this end, they had to draw a cross, on each segment (item) of the EW indicating the highest emotional intensities they felt for each emotion 229 230 while watching the film clip. Data collected from this EW were the lengths (in mm), expressed as a percentage, between the beginning of each segment and the mark drawn on it by the 231 participant. Each length corresponded to an intensity rating of the experienced emotion. 232

Intensity was assessed for each emotion and each film clip on the same wheel. This allowed participants to compare their own assessments while watching the clips and thus offering them the opportunity to make an accurate emotional appraisal. The total experiment lasted about 15 minutes.


237 **Results of Experiment 1** 

The assumptions underlying the ANOVAs were checked using the Kolmogorov– Smirnov test for normal distribution. Then, we conducted a repeated measure analysis of variance on the intensity scores, with two within-subjects factors, film clip (neutral vs. slight anger vs. strong anger) and Emotional Wheel item (anger vs. frustration vs. contempt vs. disgust vs. guilt vs. shame vs. fear vs. sadness), to study how the participants felt after viewing the film clips. Comparisons of means were performed using the post-hoc Newman-Keuls test, and means were considered as significantly different when the probability of a Type 1 error was less than or equal to 0.05. Pearson correlation coefficients between intensity of anger and other negative emotions were also calculated. Correlation coefficients were considered as significant when the probability of a Type 1 error was less than 0.05

Results of the ANOVAs showed significant differences between emotional assessments of the three film clips (F(2, 40) = 62.8, p = .001,  $\eta p^2 = 0.76$ ) as well as significant differences between assessment the items of the Emotional Wheel (F(7, 140) = 21.16, p = .001,  $\eta p^2 = 0.51$ ). The interaction between film clips and items of the Emotional Wheel was also significant (F(14, 280) = 10.23, p = .001,  $\eta p^2 = 0.34$ ).

253 Regarding anger, comparisons of means showed that the intensities of anger differed significantly between the three film clips (p < .001). Participants rated a very low level of anger 254 255 (M = 1.96, SD = 1.37) for the neutral film clip, a moderate level (M = 39.16, SD = 5.39) for the slight anger clip, and the highest level (M = 69.05, SD = 5.45) for the strong anger clip as 256 illustrated by the average emotional assessment for the three film clips presented in Figure 1. 257 Therefore, it can be concluded that the anger clips elicited two distinct levels of anger as 258 compared to our control condition (i.e., neutral film clip) supporting the subjective feelings 259 hypothesis. This finding is also congruent with the study conducted by Schaefer et al. (2010). 260

261





263

Figure 1. Average emotional assessment for the three film clips

264

Indeed, as can be observed in Figure 1, negative emotions other than anger also received high levels of subjective intensity ratings. To gain a better understanding of the links between anger and the set of negative emotions, correlation analyses were performed in order to highlight emotions that could vary in the same way as anger. The results presented in Table 1 show that subjective intensity of anger correlated only with sadness for the neutral film clip, with contempt, guilt and disgust for the slight anger film clip, and with frustration, contempt, disgust, sadness, fear and shame for the strong film clip.

272

- 273 Table 1.
- 274 *Pearson correlation coefficients between anger intensity for each film clip and the intensities*

Film clips	Sadness	Shame	Fear	Frustration	Contempt	Guilt	Disgust
Neutral	0.73*	0.10	0.02	0.32	0.05	-0.06	0.12
Slight	0.13	0.25	0.35	0.37	0.67*	0.49*	0.44*
Anger							
Strong	0.92*	<b>0.48</b> *	0.44*	0.58*	0.82*	0.11	<b>0.81</b> *
Anger							

275 *of other negative emotions* 

- 276
- 277

Surprisingly, several significant correlations between intensity of anger and intensity of other negative emotions were observed depending on the propensity of the film clips to elicit anger. It should be noted that anger is a particular emotion which is difficult to induce in laboratories. Indeed, Harmon-Jones, Amodio and Zinner (2007) argued in a book chapter that "whereas other emotions can be induced using film clips or photographs, anger is difficult to induce using such stimuli".

On the basis of the constructed approach of emotion (Lindquist, Siegel, Quigley, & 284 Barrett, 2013; Barrett, 2017) which suggests that mixes of emotions are possible, we proposed 285 a new method of calculation to assess emotion blends. To this end, we developed a global 286 emotional indicator we named global indicator (GI). It allows investigating the strength of the 287 relationship between a main emotion elicited and others related to it. In the current study, two 288 289 of the film clips used (i.e., slight and strong anger film clips) were previously validated as mainly eliciting anger (see Schaefer et al., 2010). As a large number of correlations between 290 anger and other negative emotions was also observed, we decided to build an indicator in order 291

to take into account the relationship between anger and other negative emotions. To this end, 292 the strength of the relationship between a given emotion and anger was weighted using the 293 value of the correlation coefficient between the intensity ratings for both emotions. Therefore, 294 we firstly calculated the product between the correlation coefficient with anger for the specific 295 emotion considered, the probability of a  $\beta$ -type error for this correlation and the intensity 296 collected on the EW for this emotion. Then, we made the sum of this product for all investigated 297 emotions (i.e., eight EW's emotions). Then, we divided the result by eight. As the value of this 298 index increases, the intensity of negative emotions felt related to anger increases too. 299

300 
$$GI = \frac{\sum_{i=anger}^{sadness} (Correlation Coefficient_i \times (1 - p_i) \times Intensity_i)}{8}$$

The assumptions underlying the ANOVAs were checked using the Kolmogorov-301 Smirnov test for normal distribution. A repeated measure ANOVA was then performed on the 302 GI values, with film clip (neutral vs. slight anger vs. strong anger) as a within subject factor. 303 Results showed a significant effect of the three film clips on GI values (F(2, 40) = 68.55), 304 p = .001,  $\eta p^2 = 0.77$ ). Comparisons of means showed that GI values differed significantly 305 between the three film clips (p < .001) with the lowest values (M = 0.92, SD = 1.69) for the 306 neutral film clip, moderate values (M = 8.81, SD = 4.89) for the slight anger clip, and the 307 308 highest values (M = 16.68, SD = 6.49) for the strong anger clip.

The analyses from Experiment 1 showed that the film clips were able to elicit two different intensities of anger and different blends of other specific negative emotions. The GI used varied together with the intensities of anger across the conditions, suggesting the following new categorization: low level emotion blend (LLEB) while exposed to the neutral film clip, moderate level emotion blend (MLEB) while exposed to the slight anger film clip and high level emotion blend (HLEB) while exposed to the strong anger film clip. 315

#### **Experiment 2**

Assuming the film clips used in Experiment 1 elicited two specific intensities of anger, and three different emotion blends (LLEB, MLEB and HLEB), the aim of this second experiment was to examine how cardiovascular patterns might differ between these three emotional states. In this context several hypotheses were examined.

In line with the results of Experiment 1, it was expected that subjective intensity of anger would correlate with sadness for the neutral film clip, with contempt, guilt and disgust for the slight anger clip, and with frustration, contempt, disgust, sadness, fear and shame for the strong anger clip. We named this the *comparable correlations hypothesis*.

Based on studies previously mentioned in the introduction, the specific set of negative emotions that could emerge when watching the film clips should provide an indication of the effect on cardiac measures. One possibility is that cardiac data would directly reflect how negative emotions, especially anger were processed. In this case, there would be an increase in HR and a decrease in HRV (e.g., RMSSD, SD1 and SampEn). We named this the *direct anger pattern hypothesis*.

Another possibility is that cardiac data variations reflect emotion regulation strategies. In that case, there would be a decrease in HR and an increase in HRV (e.g., RMSSD, SD1 and SampEn). We called this the *reappraisal pattern hypothesis*.

Furthermore, in view of the evidence for a link between strong negative feelings and emotion regulation strategies notably raised by Luce et al. (1999), we expected that the cardiac pattern described above, reflecting these emotion regulation strategies, would occur when the intensity of anger experienced was the highest, whereas a classic anger pattern (i.e., HR increase and a decrease RMSSD, SD1 and SampEn values) should occur for our slight anger elicitation. Consequently, it is assumed that an intensity threshold could exist between the MLEB and HLEB. We called this the *threshold hypothesis*. 340 Method

Participants and design. Thirty-six participants (18 males, 18 females) took part in this second experiment, aged between 20 and 43 years (M = 28.5; SD = 6.6). All participants reported normal or corrected-to-normal vision and audition. A between-subjects design was used, with film clip as between-subjects factor (neutral, slight anger, and strong anger). The dependent variables were intensity scores on each segment of the EW (i.e., eight emotions) and cardiac indices (HR, RMSSD, SD1, and SampEn).

Material and procedure. We used the same three film clips as in Experiment 1, 347 together with the EW (see above). Three Ag-AgCl pre-gelled electrodes were attached to the 348 participants, following a modified lead II configuration, to record their cardiac activity. 349 Electrodes were connected to Bionomadix transmitter (BIOPAC Systems Inc.) allowing 350 wireless connections with MP150 data recording system (BIOPAC Systems Inc.). After a few 351 352 seconds of recording to ensure correct equipment functioning, participants seated comfortably and were asked to relax during five minutes in order to record cardiac baseline. This five-minute 353 354 rest session was divided in two: a first rest session with eyes closed (2 min 30 s) and a second with eyes open (2 min 30 s). Closed-eye period allowed individuals to calm down, then opened-355 eye period was considered as cardiac baseline standard. 356

They then watched one of the three film clips on a large screen (visual angle of 357 31° x 17°). The sound track was played through speakers positioned behind the participant in 358 order to create an immersive watching situation. Participants were randomly assigned to one of 359 the three experimental conditions. Before and after viewing the film clip, participants were 360 asked to assess the highest emotional intensities they felt throughout the baseline and the film 361 clip watching stages respectively using the EW. As for Experiment 1, they had to draw a cross, 362 on each segment of the EW indicating the highest emotional intensities they felt for each 363 emotion while watching the film clip. Participants were alone in the room, except for a short 364

interview at the end of the experiment. The total test lasted about 30 minutes. Electrocardiogram
(ECG) was continuously recorded and sampled at 1000 Hz from the beginning of the first
baseline until the end of the film clip. The ECG was analog low-pass (0.05 Hz) and high-pass
(35 Hz) filtered at acquisition, as approximately 5–30 Hz range covers most of the frequency
content of QRS complex (Pahlm & Sörnmo, 1984). Triggers were manually positioned during
the recording to define the beginning and the end of each baseline and film clip session.

Measures and data processing. Subjective emotional state was assessed using the EW 371 as in Experiment 1. More precisely, data collected from this questionnaire were the lengths (in 372 mm), expressed as a percentage, between the beginning of each segment and the mark drawn 373 on it by the participant. Each length corresponded to an intensity rating of the experienced 374 emotion. As for Experiment 1, only negative emotions were analyzed. For the physiological 375 emotional state, the recorded cardiac signal was processed in order to calculate different 376 377 metrics. First, each participant's filtered cardiac signal was visually checked to correct any artifacts (see Berntson et al., 1997; Berntson & Stowell, 1998 for artifact correction methods 378 379 used). R peaks were detected on cardiac signal in order to calculate R-R intervals corresponding 380 to the time between two R peaks.

Several indexes were then calculated with RR interval series using Kubios HRV 381 software v.2.2. First, HR was computed in beats per minute (b.p.m.). HRV metrics were also 382 computed. For HRV time domains, the square root of the mean squared differences of 383 successive NN intervals (RMSSD) was calculated. Indeed, "in contrast to other time-domain 384 (i.e., pNN50) and time-frequency (i.e., HF-HRV) measures, changes in breathing rate have been 385 found to not significantly affect RMSSD" (Penttilä et al., 2001). Regarding HRV non-linear 386 domains, the Poincaré plot provided a graphic display of the correlations between successive 387 RR intervals (i.e., plot of RRj+1 as a function of RRj). Standard deviation of the points 388 perpendicular to the line of identity denoted by SD1 was analyzed in order to parameterize the 389

shape's ellipse formed by cloud points describing short-term variability in cardiac rhythm.
Finally, SampEn was computed by computing and comparing vectors derived from RR
intervals series (See Tarvainen (2014) for more information about how ApEn and SampEn are
calculated).

#### **Results of Experiment 2**


**Subjective emotional assessment.** The assumptions underlying the ANOVAs were checked using the Kolmogorov–Smirnov test for normal distribution and Levene's test for variance homogeneity for all analyses. The similarity of the three film clip groups (neutral, slight anger and strong anger) in terms of subjective emotional state at the beginning of the experiment (i.e. before watching the film clip) was checked using the Kruskal-Wallis test due to non-normal distribution of emotional intensity as illustrated by the average emotional assessment before viewing the film clip presented in Figure 2.


To study whether the film clips elicited two subjectively felt intensities of anger, we conducted an ANOVA with film clip (neutral vs. slight anger vs. strong anger) as betweensubjects factor and Emotional Wheel item (anger vs. frustration vs. contempt vs. disgust vs. guilt vs. shame vs. fear vs. sadness) as within-subjects factor. Comparisons of means were conducted using a post-hoc Newman-Keuls test, and means were considered as significantly different when the probability of a Type 1 error was less than or equal to 0.05.

Pearson correlation coefficients were calculated between subjective intensity of anger and other negative emotions as in Experiment 1 in order to investigate their potential relationships (note that correlation coefficients were again considered as significant when the probability of a Type 1 error was less than 0.05).

To study whether the film clips elicited three distinct emotion blends, we conducted an
ANOVA on global indicator values, with the film clips (neutral vs. slight anger vs. strong anger)

- 414 as between-subjects factor. Comparisons of means were conducted using a post-hoc Newman-
- 415 Keuls test.
- 416





419

First, results did not show any significant differences between film clip groups regarding assessment of negative emotions experienced (Emotional Wheel items) before watching the film clip: anger (p = .72), frustration (p = .60), contempt (p = .47), disgust (p = .94), guilt (p = .37), shame (p = .74), fear (p = .59) and sadness (p = .54) as illustrated in Figure 2. Groups were comparable for each emotion (Fs < 1).

Secondly, significant differences were found between emotional assessments of the three film clips (F(2, 33) = 34.73, p = .001,  $\eta p^2 = 0.678$ ) and between assessment of negative emotions (Emotional Wheel items) (F(7, 231) = 16.23, p = .001,  $\eta p^2 = 0.33$ ). The interaction between film clips and Emotional Wheel item was also significant (F(14, 231) = 5.52, p = .001,  $\eta p^2 = 0.25$ ). In addition, Newman-Keuls post hoc tests revealed that intensities of anger differed significantly between each film clip (p < .001). The neutral film clip triggered a very low level of anger (M = 2.72, SD = 2.85), the slight anger clip a moderate level of anger (M = 42.92, SD = 19.17), and the strong anger clip was rated with the highest intensity (M = 70.45, SD = 14.22). This result is consistent with the anger intensity pattern observed in Experiment 1.

As shown in Experiment 1, exposure to film clips elicited other negative emotions (see Figure 2, after film clips). Correlation results, which are reported in Table 2 indicated that intensity of anger correlated with sadness, shame and guilt for the neutral film clip; with shame, contempt and disgust for the slight anger film clip; and with sadness, shame, frustration, contempt, guilt and disgust for the strong anger clip.

- 440
- 441 Table 2.

442 Pearson correlation coefficients between anger intensity for each film clip and intensities other
443 negative emotions

Film clips	Sadness	Shame	Fear	Frustration	Contempt	Guilt	Disgust
Neutral	0.89**	0.74**	0.55	0.04	0.27	0.84**	0.26
Slight Anger	0.02	0.84**	0.46	0.36	0.70*	0.26	0.90**
Strong Anger	0.60*	0.78**	0.47	0.82**	0.67*	0.60*	0.85**

444 \* 
$$p < .05$$
. \*\*  $p < .01$ .

445

446 As in Experiment 1, the number of correlations between intensity of anger and intensity 447 of other negative emotions increased with the propensity of the film clips to elicit anger. Once 448 again, this suggests that watching the film clip elicited not only a varying intensity of anger, but also a mixture of other emotions. Nevertheless, the results did not support our *comparable correlations hypothesis* since correlations between the intensity of anger and intensities of other
negative emotions were not the same as Experiment 1.

Results regarding emotion blends highlighted significant differences between emotional assessments of the three film clips (F(2, 33) = 38.58, p = .001,  $\eta p^2 = 0.70$ ). Post hoc tests revealed that GI values differed significantly following the film clip (p < .001) with the lowest values (M = 1.53, SD = 1.50) for the neutral film clip, moderate values (M = 14.41, SD = 8.92) for the slight anger clip, and the highest values (M = 30.58, SD = 10.76) for the strong anger clip. Emotion blend categorization provided results consistent with those observed for anger assessment.

Emotional assessment using cardiac data. The assumptions underlying the ANOVAs were checked using the Kolmogorov–Smirnov test for normal distribution and Levene's test for variance homogeneity for all analyses. The similarity of the three film clip groups (neutral, slight anger and strong anger) for all cardiac metrics investigated (i.e., HR, RMSSD, SD1 and SampEn) at baseline (i.e. open eyes) was checked using ANOVA due to the normal distribution of each cardiac metric.

In order to study whether film clips elicited different physiological emotional states, 465 statistical analyses were conducted on the cardiac data collected while watching the film clips. 466 For metric distributions, data normality and homogeneity were checked as previously and 467 suitable statistical tests were then applied. Once normality had been checked (i.e., HR and 468 SampEn), we conducted ANOVAs with film clip as between-subjects factor (neutral vs. slight 469 anger vs. strong anger). Comparisons of means were also conducted using a post-hoc Newman-470 Keuls test, and means were considered as significantly different when the probability of a Type 471 1 error was less than or equal to 0.05. By contrast, when cardiac metric values did not follow a 472 normal distribution (i.e., RMSSD, SD1), a Kruskal-Wallis test was performed using film clip 473

474 as between-subjects factor (neutral vs. slight anger vs. strong anger). Mann-Whitney tests were475 then conducted to compare groups.

476 Pearson correlation coefficients were also calculated between subjective intensity of
477 anger, GI values, and cardiac indexes in order to investigate the links between subjective
478 emotional assessment and cardiac data.

First, results at baseline did not show any significant differences between groups for any cardiac metric values (p > .05). These results indicate that groups were comparable regarding cardiac metrics at baseline as illustrated by the means and standard deviations for each cardiac metric at baseline in Table 3.

483

484 Table 3.

485 *Means and standard deviations for each cardiac metric at baseline according to the* 486 *experimental conditions* 

Film clips	HR	RMSSD	SD1	SampEn
Neutral	<i>M</i> = 74.24, SD = 12.	<i>M</i> = 39.51, SD = 19.	M = 28.05, SD = 14.	M = 1.48, SD = .37
	43	73	02	
Slight Anger	<i>M</i> = 66.05, SD = 7.6	M = 80.24, SD = 48.	<i>M</i> = 56.99, SD = 34.	<i>M</i> = 1.46, SD = .24
	4	75	64	
Strong	<i>M</i> = 65.48, SD = 9.9	M = 62.62, SD = 45.	<i>M</i> = 44.46, SD = 32.	<i>M</i> = 1.77, SD = .46
Anger	3	15	08	

487

Results showed a significant effect of film clip (F(2, 33) = 5.73, p = .007,  $\eta p^2 = 0.26$ , CI = [64.04, 71.75]) for HR, with significant differences between the neutral (M = 75.59, SD = 12.95) and slight anger (M = 66.12, SD = 7.19) clips, and differences between the neutral and the strong anger clips (M = 61.97, SD = 9.32), indicating that anger elicited lower HR than a neutral situation.

23

Film clip had a significant effect on RMSSD values (H(2, 36) = 7.59, p = .02, CI = [41.68, 72.64]), with significant differences between the neutral (M = 36.72, SD = 21.89) and slight anger (M = 65.27, SD = 33.26) clips, and differences between neutral and strong anger clips (M = 69.48, SD = 66.20).

For non-linear measures, a significant effect of film clip was found on SD1 values (H(2, 36) = 7.47, p = .02, CI = [29.6, 51.61]), with significant differences between neutral (M = 26.08, SD = 15.56) and slight anger (M = 46.36, SD = 23.64) groups. Trends (p = .051) were also observed between neutral and strong anger groups (M = 49.38, SD = 47.08), and between slight anger and strong anger groups (p = .055).

The final indicator was *SampEn*. Results showed a significant effect of film clip ( $F(2, 32) = 6.67, p = .004, \eta p^2 = .29, CI = [-1.22, 9.99]$ ), with significant differences between neutral (M = 1.41, SD = .31) and slight anger groups (M = 1.89, SD = .37), and between slight anger and strong anger groups (M = 1.57, SD = .30).

506 Considering only cardiac data, results are consistent with the *reappraisal pattern* 507 *hypothesis* but not with the *direct anger pattern hypothesis*. This suggests that anger-eliciting 508 film clips likely lead individuals to establish emotion-regulation strategies. Unexpectedly, our 509 results did not show a cardiac reappraisal pattern only for the highest induced intensity of anger. 510 Hence, they are not consistent with the *threshold hypothesis*. This point will be discussed in the 511 general discussion.

512

#### **General Discussion**

The aim of the present research was to study the effects of specific intensities of anger on subjective and cardiac features of an individual's emotional state. The first experiment showed the ability of a set of film clips to elicit two subjective intensities of anger: a slight one and a strong one. Results suggested that film clips elicited, not only anger, but also other negative emotions. A global indicator (GI) of emotions felt allows us to highlight that films clips elicited three specific negative emotion blends (i.e., low, moderate and high). Experiment
2 provided comparable results for the subjective emotional assessment since the anger intensity
ratings and the GI values both showed similar ranges to the first experiment (i.e., a low level, a
moderate one and high one).

Interestingly, the results of both experiments showed that the number of correlations 522 between intensity of anger and intensity of other negative emotions increased with the 523 propensity of the film clips to elicit anger. This suggests that film clips used to elicit anger have 524 525 the ability to converge negative emotions toward overall negative states depending on the level of anger intensity they elicit. As the intensity of anger and other negative emotions (e.g., the 526 emotion blends) increase, individuals may have a less accurate assessment of their own 527 emotions. Indeed, as intensity is a concept related to the way that individuals are affected by 528 stimuli (Scherer et al., 2013), it is possible that the overall negative states triggered by anger-529 530 inducing film clips emotionally overwhelm individuals. Consequently, they likely fail to assess their emotional state accurately. 531

Furthermore, cardiac data are consistent with a reappraisal pattern for both anger 532 conditions. Indeed, anger-eliciting film clips led to lower HR than the neutral clip. This result 533 is in line with other studies that found HR deceleration due to negative emotion elicitations 534 (Kuoppa et al., 2016; Sarlo et al., 2005; Stemmler et al., 2007) and emotion-regulation strategies 535 (Denson et al., 2011). In addition, higher values for RMSSD and SD1 were observed when 536 individuals watched the anger film clips. This increase in short-term variability is associated 537 with high frequency variations in cardiac rhythm. This testifies to the dominance of the 538 parasympathetic branch of the ANS. It also suggests the establishment of emotion-regulation 539 strategies (Berntson, Cacioppo, & Quigley, 1993; Butler, Wilhelm, & Gross, 2006; Denson et 540 al., 2011; León, Hernández, Rodríguez, & Vila, 2009). Furthermore, according to Valenza et 541 al. (2012), entropy is higher when the parasympathetic branch of ANS is in control. 542

#### TOWARDS A BETTER UNDERSTANDING OF EMOTION BLENDS

543 Consequently, the significantly higher value for SampEn when participants were watching the 544 slight anger film clip is congruent with an emotion-regulation strategy. Nevertheless, significant 545 differences only appeared between the neutral and slight anger groups. An explanation could 546 be that SampEn was influenced by specific emotions only contained in the mixture of negative 547 emotions elicited by film clip associated with slight anger as illustrated in Table 2.

The next question is why participants seemed reappraise their feelings when watching 548 both anger-eliciting film clips. As described above, anger intensities correlated with other 549 550 negative emotions. Since the intensity of subjective anger and the number of converging negative emotions was higher following the two anger-eliciting film clips than the neutral clip, 551 we can assume that highly negative states resulted from watching these clips. According to 552 Luce et al. (1999), emotion-regulation strategies are established when highly intense negative 553 emotions are experienced. Hence, it can be assumed that the highly negative states elicited by 554 555 the anger-eliciting film clips triggered a greater need to reappraise the situation.

However, why did the appraisal pattern not only appear for the film clip supposed to 556 557 elicit strong anger? According to Luce et al. (1999) and Yao and Lin (2015), high intensities of negative emotions could be involved in the emergence of emotion-regulation strategies. In the 558 present study, the possible anger threshold predicting which cardiac pattern would be expressed 559 does not appear to be located between slight anger and strong anger intensity as expected, but 560 before our slight anger condition. Moreover, one of the key features of the reappraisal approach 561 is the immediate emergence of cognitive processes of coping strategies when faced with an 562 emotional event (Niedenthal, Krauth-Gruber, & Ric, 2006). Consequently, it is possible that 563 both anger film clips were emotionally appraised as eliciting a sufficient intensity of anger for 564 the establishment of emotion-regulation strategies. This regulatory effort interpretation of the 565 566 results is in line with the polyvagal theory (Porges, 1995; Porges, Doussard-Roosevelt, & Maiti, 1994; Porges, Doussard-Roosevelt, Portales, & Greenspan, 1996). This model postulates that 567

the vagus nerve, which is involved in ANS parasympathetic activity, reflects active engagement with the environment in line with emotion-regulation strategies. Unfortunately, our protocol did not enable us to identify this threshold. Future studies should investigate more intensities of negative emotion to address this issue.

Our research also raises issues about how to precisely assess emotional state. As a complement to basic approaches of emotions which distinguish emotions according to specific labels and "ANS fingerprints"(e.g., Ekman, 1992; Saarimäki et al., 2016) or dimensional approaches only accounting for some dimensions like valence and arousal (e.g., Russell, 1980), our results are particularly congruent with appraisal theories of emotion (e.g., Scherer, 2009). Indeed, as attested by the cognitive reappraisal pattern we found, emotions seem to emerge after a cognitive appraisal where a stimulus is evaluated for its meaning.

Furthermore, according to Siegel et al. (2018), the classical view of emotion that 579 580 consider some emotion categories having a specific autonomic nervous system (ANS) "fingerprint", has some weaknesses and need to be questioned. Indeed, for a given emotion 581 582 category, ANS variations are the result of random error or epiphenomenon (e.g., the method used to elicit emotion). The work carried out by Feldman Barrett describing a new approach of 583 emotion, called the theory of constructed emotion (Barrett, 2017), provides interesting outputs 584 for a better understanding of the nature of emotion. It hypothesizes that emotion is a complex 585 entity and that "an emotion category is a population of context-specific, highly variable 586 instances that need not share an ANS fingerprint" (Siegel et al., 2018). The same authors also 587 suggest that "ANS patterns are expected to be highly variable within an emotion category and 588 to overlap with other categories". Therefore, we can talk about emotion as a global phenomenon 589 consisting of many instances. In accordance with Barrett (2017); Siegel et al., (2018) we 590 591 consider emotions as a complex entities and we argue that emotion would be formed by a mix of feelings related to a main emotion. The number of emerging emotions for each film clip and 592

the correlations observed between the intensity of anger felt and other negative emotions in this study effectively shows the difficulty to elicit an emotion in isolation. GI could offer a promising global emotional assessment of negative emotions taking into account the relationships between a main emotion and others related.

597

598

599	References
600	Appelhans, B. M., & Luecken, L. J. (2006). Heart rate variability as an index of regulated
601	emotional responding. Review of General Psychology, 10(3), 229–240.
602	https://doi.org/10.1037/1089-2680.10.3.229
603	Baldaro, B., Mazzetti, M., Codispoti, M., Tuozzi, G., Bolzani, R., & Trombini, G. (2001).
604	Autonomic Reactivity during Viewing of an Unpleasant Film. Perceptual and Motor
605	Skills, 93(3), 797-805. https://doi.org/10.2466/pms.2001.93.3.797
606	Barrett, L. F. (2017). The theory of constructed emotion: an active inference account of
607	interoception and categorization. Social Cognitive and Affective Neuroscience, 12(1),
608	1-23. https://doi.org/10.1093/scan/nsw154
609	Belzung, C. (2007). Biologie des émotions (de Boeck).
610	Berntson, G. G., Cacioppo, J. T., & Quigley, K. S. (1993). Respiratory sinus arrhythmia:
611	Autonomic origins, physiological mechanisms, and psychophysiological implications.
612	Psychophysiology, 30(2), 183–196.
613	Berntson, G. G., & Stowell, J. R. (1998). ECG artifacts and heart period variability: don't miss
614	a beat! Psychophysiology, 35(1), 127–132.
615	Berntson, G. G., Thomas Bigger, J., Eckberg, D. L., Grossman, P., Kaufmann, P. G., Malik,
616	M., Van Der Molen, M. W. (1997). Heart rate variability: Origins, methods, and
617	interpretive caveats. Psychophysiology, 34(6), 623-648.
618	Butler, E. A., Wilhelm, F. H., & Gross, J. J. (2006). Respiratory sinus arrhythmia, emotion, and
619	emotion regulation during social interaction. Psychophysiology, 43(6), 612–622.
620	Cabral, J. C. C., Tavares, P. de S., Weydmann, G. J., das Neves, V. T., & de Almeida, R. M.
621	M. (2018). Eliciting Negative Affects Using Film Clips and Real-Life Methods.
622	Psychological Reports, 121(3), 527-547. https://doi.org/10.1177/0033294117730844

- 623 Cacioppo, J. T., Klein, D. J., Berntson, G. C., & Hatfield, E. (1993). The Psychophysiology of
- Emotion. In The handbook of emotion (M. Lewis, J.M. Haviland (Eds.), pp. 119–142).
  New York: Guilford Press.
- Clore, G. L., & Ortony, A. (2000). Cognitive Neuroscience of Emotion. In R. D. R. Lane, L.
  Nadel, G. L. Ahern, J. Allen, & A. W. Kaszniak (Eds.), Cognitive Neuroscience of
  Emotion (pp. 24–61). Oxford University Press.
- Denson, T. F., Grisham, J. R., & Moulds, M. L. (2011). Cognitive reappraisal increases heart
  rate variability in response to an anger provocation. Motivation and Emotion, 35(1), 14–
  22.
- Ekman, P. (1984). Expression and the nature of emotion. In Approaches to emotion (P. Ekman,
  K. Scherer (Eds.), pp. 319–343). Hilsdale, NJ: Lawrence Erlbaum Associates, Inc.
- Ekman, P. (1992). An argument for basic emotions. Cognition and Emotion, 6(3–4), 169–200.
  https://doi.org/10.1080/02699939208411068
- Fairclough, S. H., & Spiridon, E. (2012). Cardiovascular and electrocortical markers of anger
  and motivation during a simulated driving task. International Journal of
  Psychophysiology, 84(2), 188–193. https://doi.org/10.1016/j.ijpsycho.2012.02.005
- 639 Fernández, C., Pascual, J. C., Soler, J., Elices, M., Portella, M. J., & Fernández-Abascal, E.
- 640 (2012). Physiological Responses Induced by Emotion-Eliciting Films. Applied
  641 Psychophysiology and Biofeedback, 37(2), 73–79. https://doi.org/10.1007/s10484-012642 9180-7
- 643 Frijda, N. H. (1986). The Emotions. Cambridge University Press.
- Fontaine, J. R. J., Scherer, K. R., Roesch, E. B., & Ellsworth, P. C. (2007). The world of
  emotions is not two-dimensional. *Psychological Science*, *18*(12), 1050–1057.
  https://doi.org/10.1111/j.1467-9280.2007.02024.x

647

- Goshvarpour, A., Abbasi, A., & Goshvarpour, A. (2017). Indices from lagged poincare plots of
  heart rate variability: an efficient nonlinear tool for emotion discrimination.
  Australasian Physical & Engineering Sciences in Medicine, 40(2), 277–287.
  https://doi.org/10.1007/s13246-017-0530-x
- Gross, J. J. (1998). The emerging field of emotion regulation: An integrative review. *Review of General Psychology*, 2(3), 271–299. https://doi.org/10.1037/1089-2680.2.3.271
- Gross, J. J., & Levenson, R. W. (1995). Emotion elicitation using films. Cognition & Emotion,
  9(1), 87–108. <u>https://doi.org/10.1080/02699939508408966</u>
- Harmon-Jones, E. (2007). Trait anger predicts relative left frontal cortical activation to anger-
- 657 inducing stimuli. International Journal of Psychophysiology, 66(2), 154–160.
  658 https://doi.org/10.1016/j.ijpsycho.2007.03.020
- Izard, C. E. (1977). Human Emotions. New York: Plenum Press. Retrieved from
  http://www.springer.com/us/book/9780306309861
- Izard, C. E., Dougherty, F. E., Bloxom, B. M., & Kotsch, N. E. (1974). The Differential
  Emotions Scale: A method of measuring the subjective experience of discrete emotions.
  Nashville: Vanderbilt University, Department of Psychology.
- Kleinginna, P. R., & Kleinginna, A. M. (1981). A categorized list of emotion definitions, with
  a suggestion for a consensual definition. *Motivation and Emotion*, 5(3), 263–291.
  https://doi.org/10.1007/BF00993889
- Kreibig, S. D. (2010). Autonomic nervous system activity in emotion: A review. Biological
  Psychology, 84(3), 394–421. https://doi.org/10.1016/j.biopsycho.2010.03.010
- 669 Kuoppa, P., Tarvainen, M. P., Karhunen, L., & Narvainen, J. (2016). Heart rate reactivity
- associated to positive and negative food and non-food visual stimuli (pp. 5279–5282).
- 671 IEEE. https://doi.org/10.1109/EMBC.2016.7591918

León, I., Hernández, J. A., Rodríguez, S., & Vila, J. (2009). When head is tempered by heart:
Heart rate variability modulates perception of other-blame reducing anger. Motivation

and Emotion, 33(1), 1–9. https://doi.org/10.1007/s11031-008-9112-2

- Levenson, R. W., Ekman, P., & Friesen, W. V. (1990). Voluntary facial action generates
  emotion-specific autonomic nervous system activity. Psychophysiology, 27(4), 363–
  384.
- Lindquist, K. A., Barrett, L. F., Bliss-Moreau, E., & Russell, J. A. (2006). Language and the
  perception of emotion. *Emotion*, 6(1), 125–138. https://doi.org/10.1037/15283542.6.1.125
- Lobbestael, J., Arntz, A., & Wiers, R. W. (2008). How to push someone's buttons: A
  comparison of four anger-induction methods. Cognition & Emotion, 22(2), 353–373.
  https://doi.org/10.1080/02699930701438285
- Luce, M. F., Payne, J. W., & Bettman, J. R. (1999). Emotional Trade-Off Difficulty and Choice.
  Journal of Marketing Research, 36(2), 143. https://doi.org/10.2307/3152089
- 686 Macatee, R. J., Albanese, B. J., Schmidt, N. B., & Cougle, J. R. (2017). The moderating
- 687 influence of heart rate variability on stressor-elicited change in pupillary and attentional
- 688 indices of emotional processing: An eye-Tracking study. Biological Psychology, 123,
- 689 83–93. https://doi.org/10.1016/j.biopsycho.2016.11.013
- Marci, C. D., Glick, D. M., Loh, R., & Dougherty, D. D. (2007). Autonomic and prefrontal
  cortex responses to autobiographical recall of emotions. Cognitive, Affective, &
- 692 Behavioral Neuroscience, 7(3), 243–250. https://doi.org/10.3758/CABN.7.3.243
- 693 Niedenthal, P. M., Krauth-Gruber, S., & Ric, F. (2006). Psychology of emotion: interpersonal,
- 694 experiential, and cognitive approaches. New York: Psychology Press.

- Pahlm, O., & Sörnmo, L. (1984). Software QRS detection in ambulatory monitoring a
  review. Medical & Biological Engineering & Computing, 22(4), 289–297.
  https://doi.org/10.1007/BF02442095
- Pauls, C. A., & Stemmler, G. (2003). Repressive and defensive coping during fear and anger.
  Emotion, 3(3), 284–302. https://doi.org/10.1037/1528-3542.3.3.284
- Penttilä, J., Helminen, A., Jartti, T., Kuusela, T., Huikuri, H. V., Tulppo, M. P., ... Scheinin, H.
- (2001). Time domain, geometrical and frequency domain analysis of cardiac vagal
  outflow: effects of various respiratory patterns. Clinical Physiology (Oxford, England),
  21(3), 365–376.
- Plutchik, R. (2001). The Nature of Emotions. *American Scientist*, 89(4), 344.
   https://doi.org/10.1511/2001.4.344
- Porges, S. W. (1995). Orienting in a defensive world: Mammalian modifications of our
  evolutionary heritage. A Polyvagal Theory. Psychophysiology, 32(4), 301–318.
  https://doi.org/10.1111/j.1469-8986.1995.tb01213.x
- Porges, S. W., Doussard-Roosevelt, J. A., & Maiti, A. K. (1994). Vagal tone and the
  physiological regulation of emotion. Monographs of the Society for Research in Child
  Development, 59(2–3), 167–186. https://doi.org/10.1111/j.1540-5834.1994.tb01283.x
- Porges, S. W., Doussard-Roosevelt, J. A., Portales, A. L., & Greenspan, S. I. (1996). Infant
  regulation of the vagal ?brake? predicts child behavior problems: A psychobiological
  model of social behavior. Developmental Psychobiology, 29(8), 697–712.
  https://doi.org/10.1002/(SICI)1098-2302(199612)29:8<697::AID-DEV5>3.0.CO;2-O
- Ravaja, N. (2004). Contributions of Psychophysiology to Media Research: Review and
  Recommendations. Media Psychology, 6(2), 193–235.
  https://doi.org/10.1207/s1532785xmep0602\_4

- Riganello, F., Cortese, M. D., Arcuri, F., Quintieri, M., & Dolce, G. (2015). How Can Music
  Influence the Autonomic Nervous System Response in Patients with Severe Disorder
  of Consciousness? Frontiers in Neuroscience, 9.
  https://doi.org/10.3389/fnins.2015.00461
- Rogé, J., El Zufari, V., Vienne, F., & Ndiaye, D. (2015). Safety messages and visibility of
  vulnerable road users for drivers. Safety Science, 79, 29–38.
  https://doi.org/10.1016/j.ssci.2015.05.002
- Russell, J. A. (1980). A circumplex model of affect. Journal of Personality and Social
  Psychology, 39(6), 1161–1178. <u>https://doi.org/10.1037/h0077714</u>
- Saarimäki, H., Gotsopoulos, A., Jääskeläinen, I. P., Lampinen, J., Vuilleumier, P., Hari, R., ...
  Nummenmaa, L. (2016). Discrete Neural Signatures of Basic Emotions. *Cerebral Cortex* (*New York, N.Y.: 1991*), 26(6), 2563–2573.
  https://doi.org/10.1093/cercor/bhv086
- 732Sarlo, M., Palomba, D., Buodo, G., Minghetti, R., & Stegagno, L. (2005). Blood pressure733changes highlight gender differences in emotional reactivity to arousing pictures.734BiologicalPsychology,70(3),188–196.
- 735 https://doi.org/10.1016/j.biopsycho.2005.01.005
- Schachter, S., & Singer, J. (1962). Cognitive, social, and physiological determinants of
  emotional state. Psychological Review, 69(5), 379–399.
  https://doi.org/10.1037/h0046234
- Schaefer, A., Nils, F., Sanchez, X., & Philippot, P. (2010). Assessing the effectiveness of a
  large database of emotion-eliciting films: A new tool for emotion researchers. Cognition
- 741 & Emotion, 24(7), 1153–1172. https://doi.org/10.1080/02699930903274322
- 742 Scherer, K. R. (2005). What are emotions? And how can they be measured? Social Science
- 743 Information, 44(4), 695–729. https://doi.org/10.1177/0539018405058216

- Scherer, K. R., Shuman, V., Fontaine, J. R. J., & Soriano, C. (2013). The GRID meets the 744 745 Wheel: Assessing emotional feeling via self-report. https://doi.org/10.13140/RG.2.1.2694.6406 746
- Siegel, E. H., Sands, M. K., Van den Noortgate, W., Condon, P., Chang, Y., Dy, J., ... Barrett, 747 L. F. (2018). Emotion fingerprints or emotion populations? A meta-analytic 748 investigation of autonomic features of emotion categories. Psychological Bulletin, 749 144(4), 343–393. https://doi.org/10.1037/bul0000128 750
- Stemmler, G., Aue, T., & Wacker, J. (2007). Anger and fear: Separable effects of emotion and 751 motivational direction on somatovisceral responses. International Journal of 752 Psychophysiology, 66(2), 141–153. https://doi.org/10.1016/j.ijpsycho.2007.03.019 753

Tarvainen, M. (2014). Kubios HRV user's guide version 2.2. 754

- Task Force of The European Society of Cardiology and The North American Society of Pacing 755 756 and Electrophysiology. (1996). Heart rate variability: standards of measurement, physiological interpretation and clinical use. Circulation, 93(5), 1043–1065. 757
- 758 Thayer, J. F., & Lane, R. D. (2000). A model of neurovisceral integration in emotion regulation and dysregulation. Journal of Affective Disorders, 61(3), 201–216. 759
- Thayer, Julian F., & Lane, R. D. (2009). Claude Bernard and the heart-brain connection: 760 Further elaboration of a model of neurovisceral integration. Neuroscience & 761 Biobehavioral Reviews, 33(2), 81–88. https://doi.org/10.1016/j.neubiorev.2008.08.004
- 762
- Valenza, G., Allegrini, P., Lanatà, A., & Scilingo, E. P. (2012). Dominant Lyapunov exponent 763
- and approximate entropy in heart rate variability during emotional visual elicitation. 764 Frontiers in Neuroengineering, 5. https://doi.org/10.3389/fneng.2012.00003 765
- Watson, D., & Tellegen, A. (1985). Toward a consensual structure of mood. Psychological 766 767 Bulletin, 98(2), 219–235. https://doi.org/10.1037/0033-2909.98.2.219

768	Wranik, T., & Scherer, K. R. (2010). Why Do I Get Angry? A Componential Appraisal
769	Approach. In M. Potegal, G. Stemmler, & C. Spielberger (Eds.), International
770	Handbook of Anger: Constituent and concomitant biological, psychological, and social
771	processes (pp. 243-266). Springer New York. https://doi.org/10.1007/978-0-387-
772	89676-2_15
773	Yao, C. W., & Lin, T. Y. (2015). Consumer behaviour with negative emotion in e-tailing service

- Yao, C. W., & Lin, T. Y. (2015). Consumer behaviour with negative emotion in e-tailing service
   environment. International Journal of Information and Communication Technology,
- 775 7(1), 73. https://doi.org/10.1504/IJICT.2015.065990
- 776

777