



HAL
open science

Omalizumab Therapy for Mast Cell-Mediator Symptoms in Patients with ISM, CM, MMAS, and MCAS

Richard Lemal, Guillemette Fouquet, Louis Terriou, Mélanie Vaes, Cristina Bulai Livideanu, Laurent Frenzel, Stéphane Barete, Danielle Canioni, Ludovic Lhermitte, Julien Rossignol, et al.

► **To cite this version:**

Richard Lemal, Guillemette Fouquet, Louis Terriou, Mélanie Vaes, Cristina Bulai Livideanu, et al.. Omalizumab Therapy for Mast Cell-Mediator Symptoms in Patients with ISM, CM, MMAS, and MCAS. *Journal of Allergy and Clinical Immunology: In Practice*, 2019, 7 (7), pp.2387-2395.e3. 10.1016/j.jaip.2019.03.039 . hal-02299382

HAL Id: hal-02299382

<https://hal.science/hal-02299382>

Submitted on 20 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Omalizumab therapy for mast cell-mediator symptoms**
2 **in patients with ISM, CM, MMAS and MCAS**
3

4 Richard Lemal^{1*}, MD, Guillemette Fouquet^{2*}, MD, Louis Terriou³, MD, Mélanie Vaes^{4,5}, MD, Cristina
5 Bulai Livideanu⁶, MD, Laurent Frenzel⁷, MD, PhD, Stéphane Barete⁸, MD, PhD, Danielle Canioni⁹,
6 MD, Ludovic Lhermitte¹⁰, MD, PhD, Julien Rossignol^{11,12}, MD, PhD, Michel Arock^{13,14}, PharmD, PhD,
7 Patrice Dubreuil¹⁵, PhD, Olivier Lortholary¹⁶, MD, PhD, and Olivier Hermine^{2,7,11}, MD, PhD
8

9 **Authors' affiliations**

- 10 1. Histocompatibility laboratory, EA 7453 - Université Clermont Auvergne, CHU de Clermont-Ferrand,
11 France
12 2. Imagine Institute, INSERM U1163 and CNRS ERL 8654, Paris Descartes University, Sorbonne
13 Paris Cité, Paris, France
14 3. Department of Internal Medicine and Clinical Immunology, Hôpital Huriez, CHRU Lille, France
15 4. Department of Hematology, Université Libre de Bruxelles, Hôpital Erasme, Brussels, Belgium
16 5. Department of Hematology, Université Libre de Bruxelles, CHU Tivoli, La Louvière, Belgium
17 6. French Reference Center for Mastocytosis (CEREMAST), Department of dermatology, hôpital
18 Larrey, CHU Toulouse, France
19 7. Department of Hematology, Necker Children's Hospital, APHP, Paris, France
20 8. Dermatology Department, Pitié-Salpêtrière Hospital, AP-HP, Paris, France
21 9. Pathology Department, Necker Children's Hospital, APHP, Paris, France
22 10. Hematology, Necker Children's Hospital, APHP, Paris, France
23 11. French Reference Center for Mastocytosis (CEREMAST), Necker Children's Hospital, APHP,
24 Paris, France
25 12. Hematology department, Gustave Roussy, Université Paris-Saclay, F-94805, Villejuif, France
26 13. LBPA CNRS UMR8113, Ecole Normale Supérieure de Paris-Saclay, Cachan, France
27 14. Hematology, Pitié-Salpêtrière Hospital, APHP, Paris, France
28 15. Centre de Recherche en Cancérologie de Marseille, Inserm U1068, Marseille, France
29 16. Necker Pasteur Center for Infectious Diseases and Tropical Medicine, Necker Children's Hospital,
30 APHP, Paris, France

31 *These authors equally contributed to this work.
32

33 **Corresponding author**

34 Dr Richard Lemal
35 CHU de Clermont-Ferrand
36 Unité d'histocompatibilité – Centre de Biologie
37 58, rue Montalembert
38 63000 CLERMONT-FERRAND
39 Phone: +33 4 73 75 19 84
40 Fax: +33 4 73 75 19 89
41 Email: rlemal@chu-clermontferrand.fr
42

43 **Running title:** Omalizumab in systemic mastocytosis

44 **Abstract:** 249 words – **Manuscript:** 3662 words – **Figures:** 2 – **Supplemental Figures:** 5 – **Tables:** 2
45 – **References:** 33
46

47 **Abbreviations**

48 CEREMAST: French National Reference Center for Mastocytosis

49 CM: cutaneous mastocytosis

50 CR: complete response

51 IgE: immunoglobulin E

52 ISM: indolent systemic mastocytosis

53 MCA: mast cell activation

54 MCAS: Mast Cell Activation Syndrome

55 MMAS: Monoclonal MCAS

56 MR: major response

57 PR: partial response

58 SM: systemic mastocytosis

59 SM-AHN: systemic mastocytosis with associated hematologic neoplasm

60 **Key words:** omalizumab, mast cell disorders, mastocytosis, IgE

61

62

63 **Highlights boxes**

64 1. What is already known about this topic?

65 Clinical experience of omalizumab in mast cell disorders is limited, with mainly case-reports and a 14-
66 patients retrospective series testifying for its efficacy in recurrent anaphylaxis, venom immunotherapy
67 and mast cell activation symptoms.

68

69 2. What does this article add to our knowledge?

70 We prospectively surveyed 56 patients with a systemic mast cell disorder receiving omalizumab,
71 following the same administration scheme. The overall response rate was 76.8% with a very favorable
72 safety profile.

73

74 3. How does this study impact current management guidelines?

75 This real-life study describes the symptoms which encompass the best response, the median time to
76 response and it provides guidelines about optimal modalities of omalizumab administration in mast cell
77 disorders.

78

79

80

81

82

83

84

85 **Abstract**

86
87 **Background.** Patients with mast cell diseases may suffer from various distressing symptoms, which
88 can be insufficiently controlled with available therapies, severely affecting their quality of life. There is
89 a need for new and safe treatment options for these patients.

90 **Objectives.** We aimed to evaluate safety and efficacy of omalizumab administration in patients with a
91 symptomatic mast cell disorder.

92 **Methods.** We included 55 patients with a mast cell disorder associated with debilitating symptoms
93 who received omalizumab treatment between January 2015 and December 2017, after a
94 multidisciplinary team meeting at the French National Reference Center for Mastocytosis
95 (CEREMAST).

96 **Results.** A complete response was achieved for 1 patient (1.8%), a major response for 30 patients
97 (54.5%) and a partial response for 12 patients (21.8%), resulting in an overall best response rate of
98 78.2% (43/55 patients). The response was persistent at least 3 months in 33/43 (76.7%) responding
99 patients. At last follow-up, the final overall response rate was 58.2% (32/55 patients). Median time to
100 first response was 2 months and median time to best response was 6 months. Omalizumab was
101 dramatically effective on all superficial and general vasomotor symptoms and on most gastrointestinal
102 or urinary symptoms, and partially effective on most neuropsychiatric symptoms. Safety profile was
103 acceptable, except for one severe adverse event (edema of the larynx and dyspnea after the first
104 injection of omalizumab). Side effects were reported in 16 patients (28.6%), mainly of low to mild
105 intensity, yet causing interruption of treatment in 6 patients (10.7%).

106 **Conclusion.** Omalizumab seems to be a useful therapeutic option to control mast cell-mediator
107 symptoms and displays a favorable safety profile.

108

109 **Introduction**

110 Mast cells are effector cells involved in allergic and inflammatory reactions (1-3). They derive from
111 myeloid stem cells, then migrate in tissues and organs (1). Mast cells are a well-known source for
112 many proinflammatory mediators (such as histamine, tryptase, prostaglandin D₂, platelet-activating
113 factor, leukotrienes, various cytokines and chemokines) and express the high-affinity IgE receptor
114 FcεRI (1-3). Once activated by an allergen that aggregates IgE and FcεRI complexes, or by non-IgE
115 related stimuli, mast cells self-activate and release their mediators, which can cause many local and
116 systemic symptoms.

117 Systemic mastocytosis (SM) is a heterogeneous hematological neoplasm characterized by clonal
118 expansion and accumulation of mast cells in different tissues, including the bone marrow, the skin, the
119 gastrointestinal tract, the liver and the spleen (1). SM is opposed to cutaneous mastocytosis where
120 only the skin is involved, even though most adult patients with cutaneous mastocytosis have SM. SM
121 is further divided in several subcategories of increasing severity: indolent SM (ISM), smoldering SM
122 (SSM), SM with associated hematologic neoplasm (SM-AHN), aggressive SM (ASM) and mast cell
123 leukemia (MCL) (4). Some patients display typical clinical symptoms of mast cell activation (MCA), but
124 do not meet the criteria for mastocytosis: these patients are diagnosed with another mast cell disorder
125 called Mast Cell Activation Syndrome (MCAS), which can be clonal (if *KIT* mutation (mandatory) +/-
126 aberrant phenotype (CD2 and/or CD25) is proven) (MMAS) or non clonal (5).

127 In ISM, SSM, some SM-AHN, MMAS or non clonal MCAS, clinical symptoms caused by MCA are
128 often in the foreground, and can be very diverse ranging from chronic mild pruritus, musculoskeletal
129 pain, fluctuating abdominal cramping and diarrhea, to acute episodes of potentially life-threatening
130 anaphylaxis (1, 5). Patients with mast cell disorders can also present neuropsychiatric symptoms with
131 unknown pathophysiological mechanisms.

132 Several pharmacologic agents can target MCA symptoms such as antihistamines, corticosteroids,
133 cromolyn sodium or leukotriene inhibitors. When those agents are insufficient to control symptoms,
134 targeting mast cells with tyrosine kinase inhibitors, immunomodulatory or cyto-reductive therapies such
135 as imatinib, masitinib, alpha-interferon or cladribine can be necessary (4, 6, 7) and some patients can
136 relapse, experience adverse events, or become refractory to these therapies. However, these
137 therapies should only be used in patients with mastocytosis and not in other mast cell disorders.
138 Uncontrolled MCA symptoms can severely affect the quality of life of patients, and there is a great
139 need for alternative and safe treatment options (8).

140 Omalizumab is a humanized anti-IgE monoclonal antibody already approved for the treatment of
141 allergic asthma and chronic spontaneous urticaria (9-11). In these diseases, omalizumab binds to free
142 IgE, decreases the expression of FcεRI and therefore increases the threshold for degranulation of
143 mast cells and basophils (10).

144 Clinical experience of omalizumab in mast cell disorders is limited, with mainly case-reports testifying
145 for its efficacy in recurrent anaphylaxis (12-19). It has also been shown to help control reactions to
146 venom immunotherapy (17). In some cases, a “flare” effect has been observed after injection of
147 omalizumab (20), which pleads for a first in-hospital administration. More recently, a 14-patients cohort
148 study evaluated the efficacy and safety of omalizumab in SM (21). This study reported a promising
149 efficacy of omalizumab in SM, especially in anaphylaxis but with improvement of other MCA
150 symptoms and quality of life as well, along with a favorable safety profile.

151 We aimed to confirm these results by analyzing our national cohort of 55 patients who received
152 omalizumab since January 2015, within the institutional framework of the French National Reference
153 Center for Mastocytosis (CEREMAST).

154

155 **Methods**

156 **Patients.** From January 2015 to December 2017, the national multidisciplinary expert team from the
157 CEREMAST recommended treatment with omalizumab for 65 patients with symptomatic mast cell
158 disorder, treated in 11 different centers in France and Belgium. Patients had severe or debilitating
159 symptoms refractory to classical treatments of mast cell disorders, including antihistamines, cromolyn
160 sodium or leukotriene inhibitors. According to CEREMAST rules of functioning, patients with such
161 treatment decision were prospectively surveyed in the CEREMAST national database, in order to
162 record efficacy and safety data. The use of this computerized database (22) to store personal
163 information was authorized by the French National Data Protection Commission (CEREMAST
164 authorization: CNIL: No1445939 (October 29, 2010)).

165 In January 2018, we screened these 65 eligible patients, and the different centers were contacted in
166 order to update data with last follow-up. We included patients who received at least one injection of
167 omalizumab. Nine patients who did not receive omalizumab, who were lost from follow-up or for whom
168 data was deemed insufficient were excluded from this study. One patient with SM-AHN was also
169 excluded for this study. We retained 55/65 patients with a confirmed diagnosis of mast cell disorder
170 according to the World Health Organization (WHO) criteria (23) (n=40 patients) and with clonal and
171 non clonal MCAS (n= 1 and n=14 patients). MCAS was diagnosed according to recommendations, by
172 the association of typical clinical symptoms, increase in serum total tryptase by at least 20% above
173 baseline plus 2 ng/ml during or within 4 hours after a symptomatic period, and response of clinical
174 symptoms to histamine receptor blockers (HR1 +/- HR2 inverse agonists) or ‘MC-targeting’ agents like
175 cromolyn 1 (5).

176 All 55 patients were informed that they received a drug non registered neither for mastocytosis nor
177 MCAS, but that could be prescribed after recommendation from a multidisciplinary team meeting at
178 the CEREMAST. All patients gave their consent for this prospective survey.

179 Clinical and biological data were collected from the CEREMAST database for all 55 patients, and
180 completed with medical files and with a descriptive questionnaire sent to the clinician in charge of the
181 treated patient(s). The study was conducted according to the Declaration of Helsinki.

182 **Omalizumab therapy.** The recommended starting dose of omalizumab was 150 mg every 2 weeks
183 subcutaneously, without taking into account patient weight nor total IgE level. Patients were strongly
184 encouraged to continue their conventional therapy during omalizumab therapy. First injection was
185 always performed at the hospital, and the following doses could be injected at home. The rhythm of
186 physician's supervision was every four to six weeks during the first three months then every three to
187 six months depending on tolerance and efficacy. A dose escalation to a maximum of 300 mg every
188 two weeks was allowed in order to improve a partial efficacy. A weekly administration of omalizumab
189 was permitted (150 mg/week), in case of reoccurrence of symptoms prior to the following dose of
190 omalizumab, in a patient responding to therapy. This reappearance of symptoms a few days before
191 the following dose of omalizumab was assessed by the patient and its physician. It should be noted
192 that there is no proof of the efficacy of these regimens (dose escalation or weekly dosing) and that
193 they are not recommended outside of a clinical trial. Treatment was given as long as it was effective
194 and well-tolerated.

195 **Efficacy assessment: symptoms improvement.** All symptoms were assessed before and during
196 treatment, from medical files and/or via an assessment questionnaire sent to the clinician in charge of
197 the treated patients. We collected all symptoms present before treatment, and attributed the number 1
198 for all existing symptoms at baseline. Then, for each symptom, we determined the evolution on
199 treatment, including time to first response and time to best response.

200 To semi-quantitatively evaluate the symptoms evolution, we established a "symptom score" summing
201 up the response for each symptom under omalizumab therapy:

- 202 • 1.5: Worsening of the symptom.
- 203 • 1: No change.
- 204 • 0.75: Slight improvement.
- 205 • 0.5: Substantial improvement.
- 206 • 0.25: Dramatic improvement.
- 207 • 0: Complete disappearance of the symptom.

208 Quality of life assessment with classical forms was not mandatory in this real-life study; thus effect of
209 omalizumab on quality of life was not assessable.

210 **Efficacy assessment: depth of response.** Responses were assigned at time of best response, from
211 the "symptom score". A complete response (CR) was defined as the resolution of all preexisting
212 symptoms. A major response (MR) was defined as a decrease superior to 50% of preexisting
213 symptoms, according to symptom score. A partial response (PR) was defined as a decrease of 10% to

214 50% of preexisting symptoms, according to symptom score. Absence of response was defined as a
215 decrease of less than 10% of preexisting symptoms, according to symptom score, or worsening of
216 preexisting symptoms or apparition of new symptoms.

217 **Efficacy assessment: biological effects of omalizumab.** Blood parameters such as tryptase and
218 total IgE levels were analysed as in daily practice in the different centres. They were collected at
219 diagnosis, and in some cases before and during treatment with omalizumab.

220 **Safety assessment.** All physicians following patients under omalizumab therapy were requested to
221 report all adverse events occurring under therapy, following the in-law daily practice reporting system.

222 **Statistical analyses.** All analyses were performed with the GraphPad Prism 7.00 software.
223 Differences in the symptom score between baseline and follow-up under omalizumab therapy were
224 calculated using Wilcoxon signed rank test. Tests were performed two-tailed. Mean tryptase and total
225 IgE levels were compared with paired Student T tests. Reported p-values were considered statistically
226 significant if below 0.05.

227

228 **Results**

229 **Patients.** 55 patients with a confirmed diagnosis of mast cell disorder according to the World Health
230 Organization (WHO) criteria and treated with omalizumab were included in this study. Median age at
231 diagnosis was 41 years [2-87] and median age at treatment with omalizumab was 48 years [17-93].

232 The diagnoses were: indolent systemic mastocytosis (ISM; 29 patients, 52.7%), mast cell activation
233 syndrome (MCAS; 15 patients, 27.3%) and cutaneous mastocytosis (CM; 11 patients, 20%). Forty-
234 nine patients/55 were evaluated for *KIT* mutation; *KIT* D816V mutation was found in 27/49 patients
235 (55.1%): 23/28 (82.1%) patients with ISM, 3/9 (33.3%) patients with CM and 1/12(8.3%) patient with
236 MCAS.

237 A large majority of these patients already received all classical therapies against symptoms related to
238 mast cell activation, including antihistamines, cromolyn sodium or leukotriene inhibitors. 45/55 patients
239 (82%) continued at least one of the classical therapies that they were receiving before initiation of
240 omalizumab. Some patients already received multiple tyrosine kinase inhibitors, immunomodulatory or
241 cytoreductive treatments. Patients' characteristics at inclusion, including previous treatments, are
242 detailed in Table 1.

243 **Best response rates.** Among the 55 patients, we observed a CR for 1 patient (1.8%), a MR for 30
244 patients (54,5%) and a PR for 12 patients (21.8%), resulting in symptom improvement for 43 patients
245 (overall response rate of 78,2%).

246 The overall response rate was 82.9% among the 41 patients with primary MCAS, and 68.7 among the
247 14 patients with idiopathic MCAS. Detailed response rates are presented in Table 2.

248 **Time to first response and duration of response.**

249 Median time to response was 2 months and median time to best response was 6 months. Regarding
250 specific symptoms improvement, median time to first symptom improvement was 1 month [1 – 3
251 months] for general vasomotor symptoms (anaphylaxis, malaise, palpitations) and improvement
252 required 2 to 3 months for all the others symptoms [1 – 10 months].

253 At last follow-up, 32 patients (58.2%) remained on omalizumab therapy. Median duration of
254 omalizumab therapy was 9 months for the whole cohort, and 11 months for responding patients, i.e.
255 reaching at least a PR [2 – 36 months]. Median duration of therapy was 2 months for non responding
256 patients, 10 months for patients in PR, 11 months in patients with MR and 17 months for the patient in
257 complete response.

258 Response was persistent in most responding patients, with a sustained response of at least 3 months
259 in 33/43 patients (76.7%). Only 4 patients lost their acquired response under omalizumab, after 2 to 13
260 months of treatment.

261 Among the 43 responding patients, 11 have stopped therapy because of: loss of response (4
262 patients), intolerance (3 patients), loss from follow-up (2 patients), pregnancy intent (2 patients).

263 The 12 patients who did not respond to omalizumab all stopped treatment, after 1 to 6 months: 9
264 patients (75%) stopped for lack of efficacy while 3 patients (25%) stopped for intolerance. Follow-up
265 data are summarized in Figure 2.

266 Separated follow-up data for patients with primary MCAS (ISM + CM + MMAS) and non clonal
267 idiopathic MCAS patients are presented in supplemental Figures E3 and E4.

268 **Response rate at last follow-up.** At last follow-up, 32 patients were still on therapy. Among these 32
269 patients, 1 patient was still in CR (1.8%), 23 in MR (41.8%) and 8 in PR (14.5%), resulting in a
270 persistent overall response rate of 58.2%. Detailed final response rates are presented in Table 2.

271 **Dose modifications.** In 15/55 patients (27.3%), a dose escalation was prescribed in order to try to
272 improve symptom control with omalizumab, according to CEREMAST initial recommendations (up to
273 300 mg every 15 days for 14 patients, and up to 600 mg every 15 days for 1 patient, which was not a
274 recommended schedule and without objective measures to substantiate efficacy). Among these 15
275 patients, 14 patients (93%) improved their response, with 6 patients (40%) reaching a MR and 8
276 patients (53.3%) improving their PR.

277 In our cohort, 3 patients (5.5%) had a reoccurrence of symptoms prior to the following dose of
278 omalizumab administered every two weeks. In these patients, omalizumab's therapy was modified to a
279 weekly schedule, to reduce the dose interval to one week – this weekly schedule is however not FDA-
280 approved and should not be prescribed outside of a clinical trial. Among these 3 patients, 2 patients
281 improved their response, reaching 1 MR and 1 PR, and 1 patient was not improved with closer
282 administration and stopped omalizumab.

283 **Symptoms improvement.** Symptoms and numbers of patients with each symptom are indicated in
284 Figure 1 for ISM and MCAS patients. The mean number of symptoms per patient was 11.

285 Omalizumab was dramatically effective on all general and superficial vasomotor symptoms, and on
286 most gastrointestinal or urinary symptoms. Omalizumab was also, but often partially, effective on most
287 neuropsychiatric symptoms. Some patients had an improvement of their musculoskeletal pains or
288 headaches. Clinical efficacy data are presented in Figure 1.

289 We performed separate analyses for symptoms improvement in patients with primary MCAS (ISM +
290 CM + MMAS) and non clonal idiopathic MCAS patients (supplemental Figures E1 and E2). The
291 response profile to omalizumab therapy does not seem to be different between these entities, except
292 for some symptoms with only low number of patients leading to lower statistical power.

293 **Safety.** Overall, safety profile was acceptable. Sixteen patients (29%) presented side effects, among
294 whom 6 patients interrupted omalizumab therapy because of these side effects. One patient (1.8%)
295 developed a severe adverse event with edema of the larynx and dyspnea 30 minutes after the first
296 injection of omalizumab. Symptoms were controlled after corticosteroids, antihistamine and β 2
297 mimetics therapy, but persisted moderately around 3 weeks according to the patient. Three patients
298 (5.5%) presented a significant “flare” effect with accentuation of their symptoms on therapy, requiring
299 omalizumab discontinuation. Other side effects included: fatigue (3 patients), fever (3 patients),
300 headache (2 patients), malaise (2 patients), nausea (2 patients), abdominal pain (2 patients), bleeding
301 (increased menstruations and epistaxis for 1 patient, ecchymosis for 1 patient) pruritus (1 patient),
302 local rash (1 patient), sweating (1 patient).

303 **Biological effects.** Blood tests during therapy were not mandatory in this survey, but 15 patients were
304 tested for serum tryptase before and on omalizumab therapy and 15 patients were tested for serum
305 total IgE level before and on omalizumab therapy.

306 We did not observe any significant modification of the tryptase level on therapy with omalizumab, with
307 a mean value of 30.3 ng/mL before omalizumab and 32 ng/mL on therapy (after at least 3 months of
308 treatment).

309 We observed an increase in total IgE levels on therapy, with a mean value of 235.1 UI/mL [3.5 – 1616]
310 before omalizumab and 772.8 UI/mL [6.6 – 4356] on therapy. Among the 15 tested patients, 11
311 (73,3%) presented an increase of total IgE level while the 4 remaining patients (26,7%) presented
312 stable total IgE levels. The evolution of total IgE levels was not correlated with treatment efficacy or
313 tolerance in this small tested sub-group of patients.

314 Tryptase and total IgE levels evolutions on therapy are presented in supplemental Figure E5.

315

316 **Discussion**

317 This study presents data on the efficacy and safety of omalizumab in mast cell disorders, mainly
318 primary MCAS (ISM, CM and MMAS) and non clonal idiopathic MCAS. We included 55 patients in this
319 analysis, which is to our knowledge the largest cohort to date since only 29 patients with SM treated
320 omalizumab have been reported so far, in 10 different series (12-21).

321 We observed an overall response rate of 78.2% in our series, which is in line with the overall response
322 rate of 78.6% recently published by Broesby-Olsen et al. (21) in their 14-patients cohort.

323 In this series and when compiling single-cases observations, omalizumab seemed especially efficient
324 on recurrent anaphylaxis. In our study, we observed that omalizumab had a very positive effect on all
325 general and superficial vasomotor symptoms, including but not restricted to anaphylaxis, and on most
326 gastrointestinal or urinary symptoms. Regarding anaphylaxis, 13/14 patients had an improvement of
327 their symptoms, reaching independence of epinephrine after beginning of omalizumab; but
328 anaphylaxis worsened in one patient, which remains unexplained.

329 Efficacy seems more modest for musculoskeletal pains, headaches or neuropsychiatric symptoms, yet
330 a few patients significantly improved their symptoms on therapy. These results were similar to those
331 reported by Broesby-Olsen et al.: in their series, the lowest response rate was observed for
332 musculoskeletal pains and neuropsychiatric symptoms (21).

333 In our study, omalizumab's efficacy was rapidly observed with a median time to first response of 2
334 months and a median time to best response of 6 months. Time to response nevertheless ranged from
335 1 to 10 months, depending on the patient and on the symptom considered. Consistently with available
336 data, the most rapid effect of omalizumab was observed on general vasomotor symptoms, including
337 anaphylaxis. We also observed persistent responses, with an ongoing maximum duration of 36
338 months at last follow-up. Only 4 patients lost their response, after 2 to 13 months of treatment.

339 Interestingly, we observed a possible improvement of symptoms with dose adjustment and/or closer
340 administration. Even if we did not observe a clear relationship between omalizumab dosing and depth
341 of response in the whole cohort, these observations claim for a dosing effect, which seems variable
342 over time and which remains to be explored.

343 With respect to omalizumab's safety, a "flare" effect has previously been described at the beginning of
344 therapy in some patients with mastocytosis (20). Among our 55 patients, 3 patients presented a similar
345 "flare" effect, which has rapidly resolved without sequelae. However, in one patient, the flare effect
346 was severe with edema of the larynx and dyspnea occurring 30 minutes after the first injection. Apart
347 from this case, safety profile was acceptable, with mainly mild side effects yet leading to termination of
348 the treatment in 6 patients (10.9%). Broesby-Olsen et al. also reported an overall favorable tolerance
349 of omalizumab, with subjective side-effects of moderate severity in 3 patients, leading to termination of
350 the treatment in 2 out of 14 patients (14.3%) (21). Although omalizumab's safety profile appears
351 manageable in these series, the rate of side effects seems higher than expected as compared to other
352 diseases such as chronic idiopathic/spontaneous urticaria (24). However, most of severe adverse
353 events occurred after the first injection, which justifies the need for at least day care hospitalization at
354 the beginning of therapy.

355 Concerning omalizumab's biological effects, similarly to Broesby-Olsen et al., we did not find any
356 improvement in tryptase levels under therapy (21), demonstrating as expected that anti-IgE therapy is

357 not cytoreductive. We observed an increase in total IgE levels in 11 patients among the 15 tested
358 patients. This effect might be due to stabilization of total IgE (which includes free and omalizumab-
359 bound fractions), as previously reported in allergy patients (25) and in chronic spontaneous urticaria
360 (26). Total IgE levels did not seem to correlate to clinical efficacy nor tolerance in our series. In allergic
361 diseases already routinely treated with omalizumab, a reduction of inflammatory mediator release by
362 omalizumab requires a 95% reduction in serum free IgE levels to modify allergen responses via
363 receptor downregulation (25, 27, 28). Therefore, in allergic asthma, doses of omalizumab are adapted
364 to serum total free IgE level and body weight to achieve this reduction in free IgE (29, 30). In
365 symptomatic chronic idiopathic/spontaneous urticaria, which is not a classic allergen-driven disease,
366 dose of omalizumab is fixed and measurement of total or free IgE level under therapy is not necessary
367 (9, 24, 31, 32). Omalizumab's mechanism of action in symptomatic chronic idiopathic/spontaneous
368 urticaria is however not entirely understood (26, 33). Further studies remain necessary to better
369 understand how omalizumab therapy results in improvement of mast cell disorders symptoms, and
370 whether and how free or total IgE levels are correlated with clinical efficacy.

371 There are some limitations to our study, mainly due to the absence of use of validated scoring test as
372 well as quality of life assessment in this real-life evaluation. Moreover, as this was not a randomized
373 study, patients were all aware of the treatment that they were given. Some of the responses, and
374 especially temporary responses, may thus be due to a placebo effect. It should also be noted that
375 dosing of omalizumab was not adapted on serum total IgE level nor on body weight, and that some
376 schedules of treatment used in our study (dose escalation and weekly schedule) are not
377 recommended in FDA approval for asthma and chronic urticaria.". Despite these limitations, this study
378 provides valuable information on omalizumab therapy in patients with mast cell disorders, in which
379 clinical experience is very limited. A few recommendations could directly derive from our real-life
380 experience : 1/ Omalizumab can improve all symptoms but is most effective on general and superficial
381 vasomotor and gastrointestinal symptoms; 2/ Efficacy is usually quick and persistent in responding
382 patients; 3/ Even if rare, the risk of severe side effects exists at therapy initiation and thus the first(s)
383 administration(s) should be performed in-hospital; 4/ In case of insufficient efficacy, increasing the
384 dose up to 300 mg every 15 days can improve the response in majority of the cases.

385

386 **Author contributions**

387 **Conception and design:** RL, GF, LF, SB, DC, LL, JR, MA, OL, OH

388 **Collection and assembly of data:** RL, GF, LT, MV, CBL, LL

389 **Data analysis and interpretation:** RL, GF, JR, MA, OL, OH

390 **Manuscript writing – original draft:** RL, GF, JR, MA, OL, OH

391 **Manuscript writing – review, editing, and final validation of the manuscript:** RL, GF, OH

392

Table 1. Patients' characteristics.

		N = 55
General characteristics		
Age at diagnosis, years, median (range)		41 (2 – 87)
Age at omalizumab treatment, years, median (range)		48 (17 – 93)
Time from diagnosis to omalizumab treatment, median (range)		3 years (2 months - 37 years)
Sex ratio (M/F)		0.44
Disease classification in the whole cohort		
Primary MCAS (N=41)	Indolent Systemic Mastocytosis (ISM), N (%)	29 (52.7)
	- <i>KIT</i> D816V mutation positive, N	23
	- <i>KIT</i> D816V mutation negative, N	5
	- <i>KIT</i> D816V mutation not evaluated	1
	Cutaneous Mastocytosis (CM), N (%)	11 (20)
	- <i>KIT</i> D816V mutation positive, N	3
- <i>KIT</i> D816V mutation negative, N	6	
- <i>KIT</i> D816V mutation not evaluated	2	
Idiopathic MCAS (N=14)	Mast Cell Activation Syndrome (MCAS), N (%)	15 (27.3)
	- <i>KIT</i> D816V mutation positive (=MMAS), N	1
	- <i>KIT</i> D816V mutation negative, N	11
	- <i>KIT</i> D816V mutation not evaluated	3
WHO 2016 criteria		
Multifocal dense infiltrates of MCs (≥15 MCs in aggregates) in BM biopsies and/or in sections of other extracutaneous organ(s) (n=23 evaluated)		13 (56%)
25% of all MCs are atypical cells (type I or type II) on BM smears or are spindle-shaped in MC infiltrates detected on sections of visceral organs (n=39 evaluated)		39 (44%)
<i>KIT</i> point mutation at codon 816 in the BM or another extracutaneous organ (n=49 evaluated)		27 (55.1%)
MCs in BM or blood or another extracutaneous organ exhibit CD2 and/or CD25 (n=34 evaluated)		18 (53%)
Baseline serum tryptase level >20 ng/mL (n=55 evaluated)		21 (38%)
Disease presentation		
Involved skin, N (%)		35 (63.6)
	Maculopapular cutaneous mastocytosis, N (%)	31 (56.4)
	Histological criteria, N (%)	28 (50.9)
	<i>KIT</i> D816V mutation, N (%) (n= 22 evaluated)	14 (63.6)

Involved bone marrow, N (%) (n=49 evaluated)	28 (57.1)
Histological criteria, N (%)	13 (26)
Cytological criteria, N (%)	16 (32.7)
Phenotypical criteria, N (%)	17 (34.7)
KIT D816V mutation, N (%)	22 (44.9)
Decreased bone density, N (%)	17 (30.9)
Tryptase and total IgE levels at diagnosis (N=15)	
Tryptase (ng/mL), mean (range)	30.1 (2.5 – 145)
Total IgE (UI/mL), mean (range)	151.2 (2 – 1616)
Previous lines of therapy	
Symptomatic therapies	55 (100)
H2 antihistamine, N (%)	50 (90.9)
H1 antihistamine, N (%)	49 (89)
Montelukast, N (%)	39 (70.9)
Sodium cromoglycate, N (%)	27 (49)
Cytoreductive or targeted therapies (restricted to mastocytosis patients)	17 (30.9)
Cladribine, N (%)	11 (20)
Masitinib, N (%)	8 (14.3)
Imatinib, N (%)	5 (8.9)
Rapamycine, N (%)	3 (5.4)
Interferon, N (%)	2 (3.6)

Table 2. Response rates.

	All patients (N=55)	Primary MCAS (N=41)				Idiopathic MCAS (N=14)
		All (N=41)	ISM (N=29)	CM (N=11)	MMAS (N=1)	
Best response						
ORR, N (%)	43 (78.2)	34 (82.9)	23 (79.3)	10 (90.9)	1 (100)	9 (64.3)
CR, N (%)	1 (1.8)	1 (2.4)	1 (3.4)	0	0	0
MR, N (%)	30 (54.5)	22 (53.7)	16 (55.2)	5 (45.45)	1	8 (57.1)
PR, N (%)	12 (21.8)	11 (26.8)	6 (20.7)	5 (45.45)	0	1 (7.1)
No response, N (%)	12 (21.8)	7 (17.1)	6 (20.7)	1 (9.1)	0	5 (35.7)
Response at last follow up						
Persistent ORR, N (%)	32 (58.2)	25 (61)	17 (58.6)	7 (63.6)	1 (100)	7 (50)
CR, N (%)	1 (1.8)	1 (2.4)	1 (3.4)	0	0	0
MR, N (%)	23 (41.8)	17 (41.5)	12 (41.4)	4 (36.3)	1	6 (42.9)
PR, N (%)	8 (14.5)	7 (17.1)	4 (13.8)	3 (27.3)	0	1 (7.1)
No persistent response, N (%)	23 (41.8)	16 (39)	12 (41.4)	4 (36.3)	0	7 (50)
No response, N (%)	12 (21.8)	7 (17.1)	6 (20.7)	1 (9.1)	0	5 (35.7)
Loss of response, N (%)	4 (7.3)	3 (7.3)	2 (6.9)	1 (9.1)	0	1 (7.1)
Intolerance, N (%)	3 (5.4)	3 (7.3)	1 (3.4)	2 (18.2)	0	0
Lost from follow-up, N (%)	4 (7.3)	3 (7.3)	3 (10.3)	0	0	1 (7.1)

ORR: overall response rate, CR: complete response, MR: major response, PR: partial response

References

1. Metcalfe DD. Mast cells and mastocytosis. *Blood*. 2008;112(4):946-56.
2. Rivera J, Gilfillan AM. Molecular regulation of mast cell activation. *The Journal of allergy and clinical immunology*. 2006;117(6):1214-25; quiz 26.
3. Galli SJ, Grimaldeston M, Tsai M. Immunomodulatory mast cells: negative, as well as positive, regulators of immunity. *Nature reviews Immunology*. 2008;8(6):478-86.
4. Valent P, Akin C, Hartmann K, Nilsson G, Reiter A, Hermine O, et al. Advances in the Classification and Treatment of Mastocytosis: Current Status and Outlook toward the Future. *Cancer research*. 2017;77(6):1261-70.
5. Valent P, Akin C, Arock M, Brockow K, Butterfield JH, Carter MC, et al. Definitions, criteria and global classification of mast cell disorders with special reference to mast cell activation syndromes: a consensus proposal. *International archives of allergy and immunology*. 2012;157(3):215-25.
6. Siebenhaar F, Akin C, Bindslev-Jensen C, Maurer M, Broesby-Olsen S. Treatment strategies in mastocytosis. *Immunology and allergy clinics of North America*. 2014;34(2):433-47.
7. Lortholary O, Chandesris MO, Bulai Livideanu C, Paul C, Guillet G, Jassem E, et al. Masitinib for treatment of severely symptomatic indolent systemic mastocytosis: a randomised, placebo-controlled, phase 3 study. *Lancet*. 2017;389(10069):612-20.
8. Hermine O, Lortholary O, Leventhal PS, Catteau A, Soppelsa F, Baude C, et al. Case-control cohort study of patients' perceptions of disability in mastocytosis. *PloS one*. 2008;3(5):e2266.
9. Maurer M, Rosen K, Hsieh HJ, Saini S, Grattan C, Gimenez-Arnau A, et al. Omalizumab for the treatment of chronic idiopathic or spontaneous urticaria. *The New England journal of medicine*. 2013;368(10):924-35.
10. Kawakami T, Blank U. From IgE to Omalizumab. *Journal of immunology (Baltimore, Md : 1950)*. 2016;197(11):4187-92.
11. Kaplan AP, Gimenez-Arnau AM, Saini SS. Mechanisms of action that contribute to efficacy of omalizumab in chronic spontaneous urticaria. *Allergy*. 2017;72(4):519-33.
12. Lieberoth S, Thomsen SF. Cutaneous and gastrointestinal symptoms in two patients with systemic mastocytosis successfully treated with omalizumab. *Case reports in medicine*. 2015;2015:903541.
13. Carter MC, Robyn JA, Bressler PB, Walker JC, Shapiro GG, Metcalfe DD. Omalizumab for the treatment of unprovoked anaphylaxis in patients with systemic mastocytosis. *The Journal of allergy and clinical immunology*. 2007;119(6):1550-1.
14. Kibsgaard L, Skjold T, Deleuran M, Vestergaard C. Omalizumab induced remission of idiopathic anaphylaxis in a patient suffering from indolent systemic mastocytosis. *Acta dermato-venereologica*. 2014;94(3):363-4.
15. Douglass JA, Carroll K, Voskamp A, Bourke P, Wei A, O'Hehir RE. Omalizumab is effective in treating systemic mastocytosis in a nonatopic patient. *Allergy*. 2010;65(7):926-7.
16. Molderings GJ, Raithel M, Kratz F, Azemar M, Haenisch B, Harzer S, et al. Omalizumab treatment of systemic mast cell activation disease: experiences from four cases. *Internal medicine (Tokyo, Japan)*. 2011;50(6):611-5.
17. Kontou-Fili K, Filis CI. Prolonged high-dose omalizumab is required to control reactions to venom immunotherapy in mastocytosis. *Allergy*. 2009;64(9):1384-5.
18. Siebenhaar F, Kuhn W, Zuberbier T, Maurer M. Successful treatment of cutaneous mastocytosis and Meniere disease with anti-IgE therapy. *The Journal of allergy and clinical immunology*. 2007;120(1):213-5.
19. Pitt TJ, Cisneros N, Kalicinsky C, Becker AB. Successful treatment of idiopathic anaphylaxis in an adolescent. *The Journal of allergy and clinical immunology*. 2010;126(2):415-6; author reply 6.
20. Jandus P, Hausmann O, Haeberli G, Gentinetta T, Mueller U, Helbling A. Unpredicted adverse reaction to omalizumab. *Journal of investigational allergology & clinical immunology*. 2011;21(7):563-6.
21. Broesby-Olsen S, Vestergaard H, Mortz CG. Omalizumab prevents anaphylaxis and improves symptoms in systemic mastocytosis: Efficacy and safety observations. 2018;73(1):230-8.
22. Garcelon N, Neuraz A, Salomon R, Faour H, Benoit V, Delapalme A, et al. A clinician friendly data warehouse oriented toward narrative reports: Dr. Warehouse. *Journal of biomedical informatics*. 2018;80:52-63.
23. Arber DA, Orazi A, Hasserjian R, Thiele J, Borowitz MJ, Le Beau MM, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016;127(20):2391-405.
24. Saini SS, Bindslev-Jensen C, Maurer M, Grob JJ, Bulbul Baskan E, Bradley MS, et al. Efficacy and safety of omalizumab in patients with chronic idiopathic/spontaneous urticaria who remain symptomatic on H1 antihistamines: a randomized, placebo-controlled study. *The Journal of investigative dermatology*. 2015;135(1):67-75.

25. Hamilton RG, Marcotte GV, Saini SS. Immunological methods for quantifying free and total serum IgE levels in allergy patients receiving omalizumab (Xolair) therapy. *Journal of immunological methods*. 2005;303(1-2):81-91.
26. Metz M, Staubach P, Bauer A, Brehler R, Gericke J, Kangas M, et al. Clinical efficacy of omalizumab in chronic spontaneous urticaria is associated with a reduction of FcepsilonRI-positive cells in the skin. *Theranostics*. 2017;7(5):1266-76.
27. MacGlashan DW, Jr., Bochner BS, Adelman DC, Jardieu PM, Togias A, McKenzie-White J, et al. Down-regulation of Fc(epsilon)RI expression on human basophils during in vivo treatment of atopic patients with anti-IgE antibody. *Journal of immunology (Baltimore, Md : 1950)*. 1997;158(3):1438-45.
28. Saini SS, MacGlashan DW, Jr., Sterbinsky SA, Togias A, Adelman DC, Lichtenstein LM, et al. Down-regulation of human basophil IgE and FC epsilon RI alpha surface densities and mediator release by anti-IgE-infusions is reversible in vitro and in vivo. *Journal of immunology (Baltimore, Md : 1950)*. 1999;162(9):5624-30.
29. Nezos A, Papageorgiou A, Fragoulis G, Ioakeimidis D, Koutsilieris M, Tzioufas AG, et al. B-cell activating factor genetic variants in lymphomagenesis associated with primary Sjogren's syndrome. *Journal of autoimmunity*. 2014;51:89-98.
30. Steiss JO, Schmidt A, Rudloff S. Monitoring of omalizumab therapy by measuring free IgE using a bedside immunoassay. *Clinical laboratory*. 2015;61(1-2):31-7.
31. Kaplan A, Ledford D, Ashby M, Canvin J, Zazzali JL, Conner E, et al. Omalizumab in patients with symptomatic chronic idiopathic/spontaneous urticaria despite standard combination therapy. *The Journal of allergy and clinical immunology*. 2013;132(1):101-9.
32. Saini S, Rosen KE, Hsieh HJ, Wong DA, Conner E, Kaplan A, et al. A randomized, placebo-controlled, dose-ranging study of single-dose omalizumab in patients with H1-antihistamine-refractory chronic idiopathic urticaria. *The Journal of allergy and clinical immunology*. 2011;128(3):567-73.e1.
33. Chang TW, Chen C, Lin CJ, Metz M, Church MK, Maurer M. The potential pharmacologic mechanisms of omalizumab in patients with chronic spontaneous urticaria. *The Journal of allergy and clinical immunology*. 2015;135(2):337-42.

Figure legends

Figure 1. Symptoms improvement in the whole cohort (N=55).

*Comparison of the symptom score before and under omalizumab therapy: p-value: ns: > 0.05; *: p < 0.05; **:p < 0.01; ***:p < 0.001*

Figure 2. Follow-up of responses in the whole cohort (N=55).



