



HAL
open science

Monitoring Vegetation Fraction Cover of French Mediterranean Forests for Evapotranspiration and Water Stress Mapping

Albert Oliosio, Chloé Ollivier, Nicolas Martin-StPaul, Guillaume Simioni, Marie Weiss, P.C. Guillevic, Olivier Marloie, Simon Carriere, Hendrik Davi, Frederic Huard

► **To cite this version:**

Albert Oliosio, Chloé Ollivier, Nicolas Martin-StPaul, Guillaume Simioni, Marie Weiss, et al.. Monitoring Vegetation Fraction Cover of French Mediterranean Forests for Evapotranspiration and Water Stress Mapping. ESA Living Planet Symposium 2019, May 2019, Milan, Italy. 2019. <hal-02299668>

HAL Id: hal-02299668

<https://hal.science/hal-02299668v1>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Monitoring Vegetation Fraction Cover of French Mediterranean Forests for Evapotranspiration and Water Stress Mapping



Oliosio A.¹, Ollivier C.¹, Martin-StPaul N.², Simioni G.²,
Weiss M.¹, Marloie O.², Guillevic P.³, Carrière S.¹,
Davi H.², Huard F.⁴

1 – EMMAH Environnement Méditerranéen et Modélisation des Agro-Hydrosystèmes, INRA-Avignon Université, Avignon, France
2 – Unité de Recherche sur les Forêt Méditerranéenne, INRA, Avignon, France
3 – Dep. of Geographical Sciences, Univ. of Maryland, College Park, USA
4 – Terrestrial Information Systems Laboratory, NASA GSFC, Greenbelt, MD, USA;
5 – Agroclim, INRA, Avignon, France


Evapotranspiration (ET), which is a fundamental variable of the hydrological cycle, is still not known with precision over forested area in Mediterranean areas. Remote sensing data, that inform on vegetation coverage and meteorological reanalysis can be used to improve our knowledge on the distribution of ET in space and time.

A simple model using remote sensed fraction cover, meteorological data and soil AWC from various source is presented. When considering spatially available information, uncertainties in model inputs (fcover, ETo, precipitation and soil information) strongly impact the simulation of evapotranspiration

Fraction of Vegetation Cover fcover


fcover was obtained from remote sensing data:

1. Copernicus fcover products at 1 km (1999-2018) and 300 m (2014-2018). [fcover is estimated from SPOT-VEGETATION and PROBA-V reflectances using neural networks trained on a large amount of ground data (BELMANIP databases)]
2. fcover calculated from MODIS VI thanks to a linear relation established on PROSAIL simulations (varying LAI, leaf inclination, leaf chlorophyll, leaf dry matter and leaf water content).



ET Model

SimpKcET was developed as a simplified form of the crop coefficient – reference evapotranspiration approach that is classically used for monitoring evapotranspiration and water requirements for agricultural covers (ex: FAO56)



Precipitation and reference evapotranspiration ETo

ETo calculated with the Penman-Monteith model (assuming albedo 0.2 and $r_s = 60 \text{ m.s}^{-1}$). Meteorological data from SAFRAN reanalysis (MétéoFrance), ERA5 reanalysis (ECMWF) and MSWEP precipitation product.



Fontblanche experimental site:

Pinus halepensis – Q. ilex mixed forest, 20 km east of Marseille, France. 50 % rocks in top soil layers. Maximum available water content around 150 mm.

Puechabon experimental site:

Quercus ilex forest, 35 km north-west Montpellier, France. Soil is characteristics of calcareous epikarst: shallow with a high proportion of stone and rocks in the top layers (75%). Maximum available water content is around 120 mm.



Simulations: from random values of **Kmax**, **b**, **frocks** and **SAWC** around local values (+/- 20% to +/- 100% depending on the level of knowledge), all different **fcover** estimations, ERA5, SAFRAN and MSWEP reanalysis data

Results

- mean simulated ET are in good agreement with measurements;
- variability in simulations is mainly driven by rain products followed by **Kmax**, **fcover** and soil maximum available water content **SAWC**
- **frocks** has a strong impact on soil evaporation, whis is particularly visible in winter

