



HAL
open science

The molecular structure of gauche-1,3-butadiene: Experimental proof of non-planarity

Joshua Baraban, M. A. Martin-Drumel, P. Bryan Changala, Sandra Eibenberger, Matthew Nava, David Patterson, John F. Stanton, G. Barney Ellison, Michael Mccarthy

► To cite this version:

Joshua Baraban, M. A. Martin-Drumel, P. Bryan Changala, Sandra Eibenberger, Matthew Nava, et al.. The molecular structure of gauche-1,3-butadiene: Experimental proof of non-planarity. *Angewandte Chemie International Edition*, 2018, 57 (7), pp.1821-1825. 10.1002/anie.201709966 . hal-02305403

HAL Id: hal-02305403

<https://hal.science/hal-02305403>

Submitted on 4 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The molecular structure of *gauche*-1,3-butadiene: Experimental proof of non-planarity

Joshua H. Baraban^{1†}, Marie-Aline Martin-Drumel^{2‡}, P. Bryan Changala³,
Sandra Eibenberger⁴, Matthew Nava^{5§}, David Patterson^{4¶}, John F. Stanton⁶,
G. Barney Ellison¹, Michael C. McCarthy^{2*}

¹*Department of Chemistry and Biochemistry, University of Colorado, Boulder, CO 80309, USA*

²*Harvard-Smithsonian Center for Astrophysics and School of Engineering & Applied Sciences, Harvard University, Cambridge, Massachusetts 02138, United States*

³*JILA, National Institute of Standards and Technology and Department of Physics, University of Colorado, Boulder, CO 80309, USA*

⁴*Department of Physics, Harvard University, Cambridge, MA 02138*

⁵*Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*

⁶*Quantum Theory Project, Depts. of Chemistry and Physics, Univ. of Florida, Gainesville, FL 32611*

Abstract: The planarity of the second stable conformer of 1,3-butadiene – the archetypal diene for the Diels-Alder reaction, in which a planar conjugated diene and a dienophile combine to form a ring – is not established. The most recent high level calculations predict the species to adopt a twisted, *gauche* structure due to steric interactions between the inner terminal hydrogens rather than a planar, *cis* structure favored by the conjugation of the doubled bonds. Here we unambiguously prove experimentally that the structure *cis*-1,3-butadiene is indeed *gauche* with a substantial dihedral angle of 34°, in excellent agreement with theory. Observation of two tunneling components indicates that the molecule undergoes facile interconversion between two equivalent enantiomeric forms. Comparison of experimentally determined structures for *gauche*- and *trans*-butadiene provides an opportunity to examine the effects of conjugation and steric interactions.

[†]Current address: Dept. of Chemistry, Ben-Gurion University of the Negev, Beer-Sheva, 84105, Israel


[‡]Current address: Institut des Sciences Moléculaires d'Orsay, CNRS, Univ. Paris-Sud, Université Paris-Saclay, Orsay, France

[§]Current address: Department of Chemistry, Harvard University, Cambridge, MA 02138, USA

[¶]Current address: Department of Physics, University of California, Santa Barbara, CA 93106, USA

*Corresponding author. Email: mmccarthy@cfa.harvard.edu

Table of contents



The second stable conformer of 1,3-butadiene is non-planar:

The structure of *gauche*-butadiene has been determined using a combination of sensitive microwave experiments, unambiguously proving its non-planarity with a twist angle of 34° . High level theoretical calculations show that the *cis* structure is a transition state between the two *gauche minima* on the potential energy surface (see figure).

J. H. Baraban, M.-A. Martin-Drumel, P. B. Changala, S. Eibenberger, M. Nava, D. Patterson, J. F. Stanton, G. B. Ellison, M. C. McCarthy

Keywords: Rotational spectroscopy, Molecular dynamics, Quantum Chemistry, Ab initio calculations, Conjugation, Structure elucidation

Cyclic structures play essential roles in organic and biological chemistry. Perhaps the most important ring forming reaction is the stereospecific Diels-Alder cycloaddition,^[1,2] where a conjugated diene and a dienophile combine; the classic example is the reaction of 1,3-butadiene and ethylene to form cyclohexene. While this clearly requires that butadiene contort from its most stable *trans* form^[3-5] to allow ring closure, there was substantial debate as to the nature of the lowest reaction pathway,^[6] which is notoriously opaque to the usual experimental tools of mechanistic analysis.^[7] The eventual consensus is that the reaction proceeds via a synchronous transition state within a concerted mechanism.^[8] For all the diversity of Diels-Alder transition state possibilities,^[9,10] butadiene is believed to adopt a planar structure in the lowest, aromatically stabilized, pericyclic transition state,^[11] no doubt because orbital considerations suggest a planar *cis* conformation for optimal overlap with the π -system of the dienophile.^[12]

It is therefore surprising that the planarity of *cis*-butadiene itself remains a topic of interest^[13] and even controversy. Although the planarity of *trans*-butadiene is well established and attributed to conjugation between the two double bonds, steric interaction between the inner terminal hydrogens of *cis*-butadiene is a competing factor that favors a non-planar *gauche* conformation. After the existence of a second conformer was first established in 1946,^[14] numerous studies were unable to determine its structure or even its symmetry definitively.^[15-27] The advent of modern electronic structure methods^[28-30] and careful examination of gas-phase Raman spectra^[31] have led to the prevailing view that *cis*-butadiene is in fact *gauche*.^[32-35] Despite this progress, conclusive detection and structural characterization of this elementary molecule have remained elusive for sixty years.^[36,37]

Here we report the first microwave spectrum of “*cis*” butadiene, which proves unambiguously that it possesses a non-planar *gauche* structure. The experimental evidence for non-planarity, which includes not only a nearly complete equilibrium structure but also vivid observation of tunneling between equivalent *gauche* minima, was acquired with a combination of sensitive microwave techniques and high accuracy *ab initio* calculations. Examination of the structures and energetics obtained from both experiment and theory reveals detailed information about how the interplay of steric and conjugation effects determines the structure of this fundamental reaction intermediate.

We have recorded the gas phase spectrum of *gauche*-1,3-butadiene using Fourier-transform microwave (FTMW) spectroscopy, a powerful technique capable of distinguishing isomers, conformers, and even enantiomers.^[38] A particular advantage of microwave spectroscopy in this case is its immunity to spectral interference from the *trans* conformer, which has no dipole moment. Prior attempts to detect *gauche*-butadiene by microwave techniques were unsuccessful presumably owing to a combination of several factors: this conformer

lies 2.93 kcal/mol above the ground state *trans* conformer,^[39] it possesses a very small dipole moment^[40] (calculated to be 0.09 D in this work), and its rotational transition frequencies are sensitive functions of the torsional $\angle\text{C}=\text{C}-\text{C}=\text{C}$ angle τ (Fig. 1). Observation of the microwave spectrum of *gauche*-butadiene in the present work required the use of two complementary experimental techniques: cavity enhanced FTMW combined with a supersonic expansion^[41] and chirped-pulse FTMW in a cryogenic buffer gas cell.^[42] The population of the minor conformer was increased by using a 1 kV electric discharge or a heated inlet capillary (245 °C) in the two experiments, respectively. In the buffer gas cell, butadiene molecules rapidly thermalize rotationally to the cell temperature (~ 10 K) through collisions with cold helium atoms, while the *gauche-trans* relaxation proceeds much more slowly. The molecules diffuse through the cold gas until they collide with the cell wall, where they freeze; during this time, the sample is repeatedly interrogated. The combination of a high repetition rate (30 kHz) and cryogenic low noise amplifiers (noise temperature of ~ 9 K) yields significantly higher signal-to-noise than in the cavity-enhanced pulsed jet experiment. The spectral resolution in the buffer gas cell is limited by collisional broadening with helium to ~ 80 kHz, while the cavity-enhanced pulsed jet work yields a time-of-flight broadening of ~ 5 kHz. Initial spectral searches were guided by high-level predictions of the rotational constants, and transitions were readily observed near the predicted frequencies. The large inertial defect (-2.5 amu \AA^2 , Table S2) is a clear sign of non-planarity, which was confirmed by the semi-experimental equilibrium structure^[43], r_e^{se} , derived from subsequent spectroscopy of five isotopically substituted species of butadiene ($^{13}\text{CH}_2\text{CHCHCH}_2$, $\text{CH}^{13}\text{CHCHCH}_2$, $\text{CH}_2\text{CDCDCH}_2$, $\text{CD}_2\text{CHCHCD}_2$, $\text{CD}_2\text{CDCDCD}_2$). Unfortunately, the lack of commercial asymmetrically deuterated samples precludes the determination of a complete structure at present, but efforts to remedy this are underway.

Conclusive evidence that the equilibrium structure of the species being probed is *gauche*, rather than *cis*, is the doublet structure of the measured transitions (Fig. 2). Such doublets can only arise from tunneling between the two equivalent C_2 *gauche* minima on the potential energy surface, which causes the low-lying vibrational states to split into two components, e.g. $v = 0^+$ and $v = 0^-$ for the ground state, each of which has its own manifold of rotational levels.

Table 1 shows the experimentally determined and calculated structures of *gauche*-butadiene. The *ab initio* geometry was extrapolated from optimizations at the (all-electron) AE-CCSD(T) level of theory with cc-pCVTZ through cc-pCV5Z basis sets, and supplemented by a perturbative quadruples (Q) correction calculated with the ANO0 basis set and the frozen core (FC) approximation. The (Q) correction for the dihedral τ is surprisingly large (almost 1°), presumably because of the complex electron correlation associated with conjugation. Together with zero-point corrections from second-order vibrational Møller-Plesset theory^[44]

(VMP2) using FC-CCSD(T) calculations and the ANO1 basis set,^[45,46] this structure accurately predicted the rotational constants for all isotopologues – the final calculations described here match all observed rotational constants to within 6 MHz. All electronic structure calculations were carried out with CFOUR;^[47] the VMP2 calculations were performed using NITROGEN.^[48] The resulting rovibrational energies were then reduced to spectroscopic constants for the $v = 0^+$ and 0^- *gauche*-butadiene levels.

While the observed transitions do not yield any direct measurement of the $0^+/0^-$ tunneling splitting, Δ_{inv} , a fit of Coriolis interactions between the two tunneling components allows us to determine Δ_{inv} (Fig. 3). The best fit results are obtained for the heaviest isotopologue available, d_6 , where $\Delta_{inv}=16.577(16)$ GHz, within 5% of theory. Based on this agreement and a similar, preliminary fit for the normal isotopologue, we believe the *ab initio* value of 55 GHz is likely to be accurate to within 20%, corresponding to a tunneling lifetime of approximately 18 ps. This result is entirely compatible with the proposal of Michl *et al.*^[20,22] that rapid tunneling could cause the matrix infrared spectrum to appear as that of a planar C_{2v} *cis*-butadiene.

Two major factors influence the structure of *gauche*-butadiene: conjugation of the double bonds and steric interaction between the inner terminal hydrogens. The former favors planarity, as in the *trans* conformer, and the latter is destabilizing at the planar *cis* geometry; a compromise between the two effects leads to the *gauche* minimum. Conjugation of the two π bonds in planar butadiene imparts a stabilization of roughly 4 kcal/mol (Table S4), while twisting about the dihedral angle τ (Fig. 1) reduces the conjugation by a factor of roughly $\cos^2 \tau$, implying that the steric repulsion between the two *endo* hydrogens is decreased by at least 1.3 kcal/mol in *gauche*- relative to *cis* butadiene.

A structural comparison between *gauche*- and *trans*-butadiene is particularly illustrative of the competition between these two effects (Table 1). It has been shown^[49] that conjugation in *trans*-butadiene shortens the C–C bond (relative to propene^[50] or a formal sp^2-sp^2 bond length^[51]) and slightly elongates the C=C bonds (relative to ethylene or propene).^[49,52] To minimize the steric interaction between the inner hydrogens (H_1 , Fig. 1), *gauche*-butadiene adopts a surprisingly large twist angle of 34° for the C=C–C=C skeleton, over a third of the way to where conjugation would be completely destroyed. The C–C bond nearly returns to the canonical sp^2-sp^2 value because this displacement directly increases the H_1-H_1 distance; by contrast, the C=C bond is only slightly shorter than in the *trans* conformer. Other more subtle distortions with similar effects are also evident: for example, the $\angle C-C=C$ angle opens by almost 1° compared to *trans*-butadiene. In general, modes with small force constants experience larger distortions, which relieves strain most efficiently.

The determination of the structure of *gauche*-butadiene advances our experimental knowledge of the

quintessential Diels-Alder reaction pathway. The stereochemistry of this reaction has been established by reacting 1,1,4,4-tetradeutero-1,3-butadiene with both *cis*- and *trans*-1,2-dideuteroethylene, with the results supporting a concerted mechanism.^[53] The activation energy has been determined by kinetics studies to be 27 kcal/mol, both by examining the Diels-Alder reaction itself^[54] and by combining data on the *retro*-Diels-Alder reaction^[55,56] with the known enthalpy of reaction^[57]. Especially since the threshold energy is likely to be several kcal/mol lower^[58], this implies that the *trans-gauche* conformational change in butadiene accounts for a substantial portion of the total energy required to initiate the Diels-Alder reaction with ethylene. We believe that the detection and characterization of the reactive conformation of butadiene points the way towards studying the pre-reaction complex^[59] and transition state of this important reaction.

Acknowledgements

We thank Norman Craig (Oberlin), Ken Wiberg (Yale), and Norman Allinger (Univ. Georgia) for helpful discussions. The work in Cambridge is supported by NSF Grants CHE-1566266 and DBI-1555781. The National Science Foundation (CBET-1403979) has provided support for J.H.B. and G.B.E. P.B.C. is supported by an NSF Graduate Research Fellowship (Award No. DGE1144083). J.F.S. would like to thank the NSF (Grant CHE-1361031), which also provided support for J.H.B. S.E. acknowledges funding through the Schrödinger Fellowship of the Austrian Science Fund (J3796-N36). We thank E.S. Palmer and P. Antonucci for technical assistance and G.T. Buckingham for assistance with early laboratory searches.

References

- [1] O. Diels, K. Alder, *Justus Liebigs Annalen der Chemie* **1928**, *460*, 98–122.
- [2] “The Nobel Prize in Chemistry 1950”. *Nobelprize.org. Nobel Media AB 2014. Web. 28 Jun 2016.* http://www.nobelprize.org/nobel_prizes/chemistry/laureates/1950/.
- [3] V. Schomaker, L. Pauling, *Journal of the American Chemical Society* **1939**, *61*, 1769–1780.
- [4] A. Almenningen, O. Bastiansen, M. Trættestad, *Acta Chemica Scandinavica* **1958**, *12*, 1221.
- [5] D. Marais, N. Sheppard, B. Stoicheff, *Tetrahedron* **1962**, *17*, 163 – 169.
- [6] K. N. Houk, J. Gonzalez, Y. Li, *Accounts of Chemical Research* **1995**, *28*, 81–90.
- [7] W. E. Doering, W. Roth, *Tetrahedron* **1962**, *18*, 67 – 74.
- [8] H. Lischka, E. Ventura, M. Dallos, *ChemPhysChem* **2004**, *5*, 1365–1371.
- [9] Y. Li, K. N. Houk, *Journal of the American Chemical Society* **1993**, *115*, 7478–7485.
- [10] A. Z. Bradley, M. G. Kociolk, R. P. Johnson, *The Journal of Organic Chemistry* **2000**, *65*, 7134–7138, PMID: 11031040.
- [11] T. Sexton, E. Kraka, D. Cremer, *The Journal of Physical Chemistry A* **2016**, *120*, 1097–1111, PMID: 26785172.
- [12] R. B. Woodward, R. Hoffmann, *Angewandte Chemie International Edition in English* **1969**, *8*, 781–853.
- [13] P.-Q. Liao, N.-Y. Huang, W.-X. Zhang, J.-P. Zhang, X.-M. Chen, *Science* **2017**, *356*, 1193–1196.
- [14] J. G. Aston, G. Szasz, H. W. Woolley, F. G. Brickwedde, *The Journal of Chemical Physics* **1946**, *14*, 67–79.
- [15] M. E. Squillacote, R. S. Sheridan, O. L. Chapman, F. A. L. Anet, *Journal of the American Chemical Society* **1979**, *101*, 3657–3659.
- [16] M. E. Squillacote, T. C. Semple, P. W. Mui, *Journal of the American Chemical Society* **1985**, *107*, 6842–6846.
- [17] D. Feller, E. R. Davidson, *Theoretica chimica acta* **1985**, *68*, 57–67.
- [18] J. J. Fisher, J. Michl, *Journal of the American Chemical Society* **1987**, *109*, 1056–1059.
- [19] K. B. Wiberg, R. E. Rosenberg, *Journal of the American Chemical Society* **1990**, *112*, 1509–1519.
- [20] B. R. Arnold, V. Balaji, J. Michl, *Journal of the American Chemical Society* **1990**, *112*, 1808–1812.
- [21] B. R. Arnold, V. Balaji, J. W. Downing, J. G. Radziszewski, J. J. Fisher, J. Michl, *Journal of the American Chemical Society* **1991**, *113*, 2910–2919.
- [22] H. C. Choi, M. Kertesz, S. Dobrin, J. Michl, *Theoretical Chemistry Accounts* **1999**, *102*, 196–206.
- [23] Y. Furukawa, H. Takeuchi, I. Harada, M. Tasumi, *Bulletin of the Chemical Society of Japan* **1983**, *56*, 392–399.
- [24] L. A. Carreira, *The Journal of Chemical Physics* **1975**, *62*, 3851–3854.
- [25] N. C. Craig, R. L. Sams, *The Journal of Physical Chemistry A* **2008**, *112*, 12637–12646, PMID: 19007193.

- [26] J. R. Durig, W. E. Bucy, A. R. H. Cole, *Canadian Journal of Physics* **1975**, *53*, 1832–1837.
- [27] M. Kofranek, A. Karpfen, H. Lischka, *Chemical Physics Letters* **1992**, *189*, 281 – 286.
- [28] J. E. Rice, B. Liu, T. J. Lee, C. M. Rohlfing, *Chemical Physics Letters* **1989**, *161*, 277 – 284.
- [29] I. L. Alberts, H. F. Schaeffer III, *Chemical Physics Letters* **1989**, *161*, 375 – 382.
- [30] H. Guo, M. Karplus, *The Journal of Chemical Physics* **1991**, *94*, 3679–3699.
- [31] R. Engeln, D. Consalvo, J. Reuss, *Chemical Physics* **1992**, *160*, 427 – 433.
- [32] P. Boopalachandran, N. Craig, P. Groner, J. Laane, *The Journal of Physical Chemistry A* **2011**, *115*, 8920–8927.
- [33] P. Boopalachandran, N. C. Craig, J. Laane, *The Journal of Physical Chemistry A* **2012**, *116*, 271–281.
- [34] S. V. Krasnoshchekov, N. C. Craig, P. Boopalachandran, J. Laane, N. F. Stepanov, *The Journal of Physical Chemistry A* **2015**, *119*, 10706–10723, PMID: 26437183.
- [35] D. Feller, N. C. Craig, *The Journal of Physical Chemistry A* **2009**, *113*, 1601–1607, PMID: 19199679.
- [36] D. R. Lide, Jr., *The Journal of Chemical Physics* **1962**, *37*, 2074–2079.
- [37] W. Caminati, G. Grassi, A. Bauder, *Chemical Physics Letters* **1988**, *148*, 13 – 16.
- [38] D. Patterson, M. Schnell, J. M. Doyle, *Nature* **2013**, *497*, 475–477.
- [39] J. Saltiel, D. F. Sears Jr., A. M. Turek, *The Journal of Physical Chemistry A* **2001**, *105*, 7569–7578.
- [40] W. Caminati, B. Vogelsanger, A. Bauder, *Journal of Molecular Spectroscopy* **1988**, *128*, 384 – 398.
- [41] K. N. Crabtree, M. R. Talipov, J. Martinez, Oscar, G. D. O’Connor, S. L. Khursan, M. C. McCarthy, *Science* **2013**, *342*, 1354–1357.
- [42] D. Patterson, J. M. Doyle, *Molecular Physics* **2012**, *110*, 1757–1766.
- [43] P. Pulay, W. Meyer, J. E. Boggs, *The Journal of Chemical Physics* **1978**, *68*, 5077–5085.
- [44] P. B. Changala, J. H. Baraban, *The Journal of Chemical Physics* **2016**, *145*, 174106.
- [45] J. Almlöf, P. R. Taylor, *The Journal of Chemical Physics* **1987**, *86*, 4070–4077.
- [46] L. McCaslin, J. F. Stanton, *Molecular Physics* **2013**, *111*, 1492–1496.
- [47] J. F. Stanton, J. Gauss, M. E. Harding, P. G. Szalay, *CFOUR, Coupled-Cluster techniques for Computational Chemistry, a quantum-chemical program package*, With contributions from A.A. Auer, R.J. Bartlett, U. Benedikt, C. Berger, D.E. Bernholdt, Y.J. Bomble, L. Cheng, O. Christiansen, M. Heckert, O. Heun, C. Huber, T.-C. Jagau, D. Jonsson, J. Jusélius, K. Klein, W.J. Lauderdale, F. Lipparini, D.A. Matthews, T. Metzroth, L.A. Mück, D.P. O’Neill, D.R. Price, E. Prochnow, C. Puzzarini, K. Ruud, F. Schiffmann, W. Schwalbach, C. Simmons, S. Stopkowitz, A. Tajti, J. Vázquez, F. Wang, J.D. Watts and the integral packages MOLECULE (J. Almlöf and P.R. Taylor), PROPS (P.R. Taylor), ABACUS (T. Helgaker, H.J. Aa. Jensen, P. Jørgensen, and J. Olsen), and ECP routines by A. V. Mitin and C. van Wüllen. For the current version, see <http://www.cfour.de>.
- [48] *NITROGEN, Numerical and Iterative Techniques for Rovibronic Energies with General Internal Coordinates, a program written by P. B. Changala.* <http://www.colorado.edu/nitrogen>.
- [49] N. C. Craig, P. Groner, D. C. McKean, *The Journal of Physical Chemistry A* **2006**, *110*, 7461–7469, PMID: 16759136.

- [50] H. S. Tam, M. D. Harmony, *The Journal of Physical Chemistry* **1991**, *95*, 9267–9272.
- [51] M. Dewar, H. Schmeising, *Tetrahedron* **1959**, *5*, 166 – 178.
- [52] M. D. Harmony in *Equilibrium Structural Parameters*, Vol. 24 of *Vibrational Spectra and Structure*, J. R. Durig (Ed.), Elsevier, **1999**, chapter 1, pp. 1 – 83.
- [53] K. N. Houk, Y. T. Lin, F. K. Brown, *Journal of the American Chemical Society* **1986**, *108*, 554–556, PMID: 22175504.
- [54] D. Rowley, H. Steiner, *Discuss. Faraday Soc.* **1951**, *10*, 198–213.
- [55] M. Uchiyama, T. Tomioka, A. Amano, *The Journal of Physical Chemistry* **1964**, *68*, 1878–1881.
- [56] W. Tsang, C. M. Rosado-Reyes, *The Journal of Physical Chemistry A* **2015**, *119*, 7155–7162, PMID: 25584988.
- [57] J. B. Pedley, R. D. Naylor, S. P. Kirby, *Thermochemistry of Organic Compounds 2nd ed.*, Chapman and Hall, New York, **1986**.
- [58] M. Menzinger, R. Wolfgang, *Angewandte Chemie International Edition in English* **1969**, *8*, 438–444.
- [59] S. Novick, J. M. Lehr, W. Klemperer, *Journal of the American Chemical Society* **1973**, *95*, 8189–8191.
- [60] D. Papoušek, M. Aliev, *Molecular Vibrational-rotational Spectra: Theory and Applications of High Resolution Infrared, Microwave, and Raman Spectroscopy of Polyatomic Molecules*, of *Studies in Physical and Theoretical Chemistry*, Elsevier, **1982**.
- [61] C. M. Western, *Journal of Quantitative Spectroscopy and Radiative Transfer* **2017**, *186*, 221 – 242.




Figure 1: The structure of *gauche*-butadiene. The C_2 symmetry b -axis is vertical. The dihedral $\angle C=C-C=C$ is indicated (τ) on the right.




Figure 2: Torsional potential and rovibrational energy levels of butadiene. The potential was calculated at the FC-CCSD(T)/ANO1 level of theory (see also Table S1). The expanded view on the right shows the rotational level structure of the ground state tunneling doublet and the b -type microwave transitions allowed within each tunneling component. Dashed arrows mark transitions in the inset spectrum, which consists of 2×10^6 averages collected in ~ 5 minutes, while flowing 0.7 sccm butadiene through a 245 °C nozzle along with 6 sccm He into the cryogenic buffer gas cell.




Figure 3: Reduced term value plot for d_6 butadiene, illustrating the near-resonant Coriolis interaction between tunneling components that allows the determination of Δ_{inv} from the microwave data. E+/O+/E-/O- rovibrational symmetries^[60,61] are represented by filled/open circles and squares, respectively. 0^+ levels (solid lines) from $K = 1f$ tune into resonance with 0^- levels (dashed lines) of the same 0^+ symmetry from $K = 0f$ at $J = 4$.

Table 1: Butadiene geometries. Bond lengths in Å, angles in degrees. The experimental structures are semi-experimental equilibrium structures, r_e^{se} , obtained by correcting experimental rotational constants for zero-point rotation-vibration effects calculated *ab initio*.

Parameter	<i>gauche</i>		<i>trans</i>		<i>cis</i>
	Calc. ^a	Expt.	Calc. ^a	Expt. ^[49]	Calc. ^a
C–C	1.4673	1.48(2)	1.4542	1.4539(10)	1.4684
C=C	1.3369	1.33(1)	1.3385	1.3376(10)	1.3379
∠C–C=C	124.51	124.1(5)	123.57	123.62(10)	126.16
∠C=C–C=C (τ)	33.87	33.8(13)	planar		planar
r_{CH_1}	1.0817	1.08(1)	1.0819	1.0819(10)	1.0814
∠H ₁ –C=C	121.21	123(2)	120.94	120.97(10)	122.11
∠H ₁ –C=C–C	2.07	2(1)	planar		planar
r_{CH_2}	1.0798	[1.07983] ^b	1.0796	1.0793(10)	1.0795
∠H ₂ –C=C	121.10	118(1)	121.48	121.47(10)	120.75
∠H ₂ –C=C–H ₃	2.41	[2.410] ^b	planar		planar
r_{CH_3}	1.0837	1.083(5)	1.0843	1.0847(10)	1.0829
∠H ₃ –C–C	116.54	116.2(6)	116.69	116.47(10)	115.66
∠H ₃ –C–C=C	-147.26	-146.4(7)	planar		planar

^a Extrapolation of optimized all electron CCSD(T)/pCVTZ–5Z geometries, supplemented by a higher order perturbative quadruples correction calculated with the ANO0 basis set and the frozen core approximation. ^b Numbers in brackets were fixed to the *ab initio* values.