



HAL
open science

Discussing the first velocity dispersion profile of an ultra-diffuse galaxy in MOND

Michal Bílek, Oliver Müller, Benoit Famaey

► **To cite this version:**

Michal Bílek, Oliver Müller, Benoit Famaey. Discussing the first velocity dispersion profile of an ultra-diffuse galaxy in MOND. *Astronomy & Astrophysics - A&A*, 2019, 627, pp.L1. 10.1051/0004-6361/201935840 . hal-02306467v1

HAL Id: hal-02306467

<https://hal.science/hal-02306467v1>

Submitted on 5 Oct 2019 (v1), last revised 23 May 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LETTER TO THE EDITOR

Discussing the first velocity dispersion profile of an ultra-diffuse galaxy in MOND

Michal Bílek¹, Oliver Müller¹, and Benoit Famaey¹

Université de Strasbourg, CNRS, Observatoire astronomique de Strasbourg (ObAS), UMR 7550, 67000 Strasbourg, France
e-mail: bilek@astro.unistra.fr

Received tba; accepted tba

ABSTRACT

Using Jeans modelling, we calculate the velocity dispersion profile of the ultra-diffuse galaxy (UDG) Dragonfly 44 in MOND. For the nominal mass-to-light ratio from the literature and an isotropic profile, the agreement with the data is excellent near the center of the galaxy. However, in modified gravity, close to the cluster core, the gravitational environment should bring the galaxy back towards the Newtonian behavior. This success of the isolated MOND prediction for the central velocity dispersion could then mean that the galaxy is at large distances ($\gg 5$ Mpc) from the cluster core, as hinted by the fact that close-by UDGs share similar velocities with a dispersion well below that of the cluster itself. There is however a two-sigma tension in the outer part of the UDG due to an increase of the observed dispersion profile with respect to the flat MOND prediction. This deviation could simply be a measurement error. Other possibilities could be, for a UDG far from the cluster, a higher-than-nominal baryonic mass with a tangentially anisotropic dispersion profile, or even a sign of a dark baryonic halo. If the UDG is closer to the cluster core, the deviation could be a sign that it is in the process of disruption.

Key words. Galaxies: individual: Dragonfly 44; Galaxies: clusters: individual: Coma cluster; Galaxies: kinematics and dynamics

1. Introduction

Ultra-diffuse galaxies (UDG) are very extended ($r_{\text{eff}} > 1.5$ kpc) low-surface brightness ($\mu_V > 25$ mag arcsec⁻²) objects. They have been identified in different galactic environments for decades (e.g., Sandage & Binggeli 1984; Impey et al. 1988). These galaxies have recently experienced a revival (Crnojević et al. 2014; van Dokkum et al. 2015a; Koda et al. 2015; van der Burg et al. 2016; Martínez-Delgado et al. 2016; Merritt et al. 2016; Román & Trujillo 2017a; Venhola et al. 2017; Wittmann et al. 2017; Müller et al. 2018; Mancera Piña et al. 2019). In galaxy clusters, UDGs do not contain gas while in sparser environments, they can be gas dominated (van Dokkum et al. 2015a; Shi et al. 2017; Leisman et al. 2017; Papastergis et al. 2017; Sardone et al. 2019), following the well-known density-morphology relation (Dressler 1980). Several formation scenarios have been proposed in the Λ CDM context: they might be tidal dwarf galaxies, galaxies formed by collapse of gas in galaxy outflows, dwarf galaxies that experienced strong tidal stripping or repeated episodes of intense star formation and others (e.g. van Dokkum et al. 2015a; Yozin & Bekki 2015; Amorisco & Loeb 2016; Di Cintio et al. 2017; Chan et al. 2018; Toloba et al. 2018). Observations suggest that at least some UDGs form in the field and then are accreted to galaxy clusters where they experience environmental quenching (Román & Trujillo 2017b; Ferré-Mateu et al. 2018; Alabi et al. 2018).

The dynamics of these systems provides exciting insights to the discussion. For instance, van Dokkum et al. (2018a) announced the discovery of an apparently dark matter free UDG in the field of NGC 1052. Soon, in the same field around NGC 1052 a second galaxy lacking dark matter was found (van Dokkum et al. 2019a). These two galaxies – NGC 1052-DF2 and

NGC 1052-DF4 – have sparked a vast discussion in the literature (e.g., van Dokkum et al. 2018b; Trujillo et al. 2019; Martin et al. 2018; Kroupa et al. 2018; Famaey et al. 2018; Emsellem et al. 2018; Fensch et al. 2018; Müller et al. 2019a,b; Nusser 2019; Laporte et al. 2019).

As low surface-brightness objects, UDGs are expected to show complex dynamical behavior in the context of Modified Newtonian Dynamics (MOND, Milgrom 1983; here we only consider MOND modified gravity theories and not modified inertia theories). As a historic note, MOND was proposed to solve the missing mass problem in high surface brightness spiral galaxies (e.g., Rubin et al. 1978; Bosma 1981; Rubin et al. 1982) and has since excelled in reproducing the dynamics of a much larger range of galaxies (e.g., McGaugh & de Blok 1998; McGaugh & Milgrom 2013; Lelli et al. 2017, see also the extensive of review in Famaey & McGaugh 2012). At the time MOND was proposed, the existence of UDGs was unknown. In MOND, a dynamical system appears Newtonian when the gravitational acceleration is greater than about the threshold of $a_0 = 1.2 \times 10^{-13}$ km s⁻² (Milgrom 1983; Begeman et al. 1991). Below, a departure from the Newtonian dynamics occurs such that it appears as if it hosted dark matter. The dark matter behavior is at its full strength only if the galaxy is isolated. In the case the object resides in a strong gravitational field of the environment, the deviations from the Newtonian dynamics can effectively be suppressed (e.g. Milgrom 1983; Bekenstein & Milgrom 1984), which is due to the non-linearity of any MOND theory (Milgrom 2014).¹ In other words, the object will appear dark matter free, even though its internal acceleration is below a_0 , thanks to the so-called external

¹ However, the external field effect can be weak, and even almost negligible, in some modified inertia versions of MOND (Milgrom 2011).

field effect (EFE). To explain the dynamics of two UDGs of the NGC 1052 group in MOND – NGC 1052-DF2 and NGC 1052-DF4 (if these are not in the foreground of their putative host group, see Trujillo et al. 2019) –, it is crucial to take the EFE into account, mostly removing the tension for NGC 1052-DF2 (Kroupa et al. 2018; Famaey et al. 2018), and lessening it for NGC 1052-DF4 (Müller et al. 2019a). Interestingly, the UDG Dragonfly 44 in the Coma cluster behaves very differently from the UDGs of the NGC 1052 group. It exhibits a pronounced deviation from Newtonian dynamics. Early measurements of its stellar velocity dispersion yielded $\sim 47 \text{ km s}^{-1}$ (van Dokkum et al. 2016), well above the nominal isolated MOND prediction. However, all these previous studies relied on the global velocity dispersion estimated at the half light radius of the galaxy, but did not take into account the overall shape and profile of the velocity dispersion due to the difficulty in measuring it.

Now, the first velocity dispersion profile of a UDG, namely Dragonfly 44, was presented recently by van Dokkum et al. (2019b). The data was taken with the Keck Cosmic Web Imager (KCWI) on Mauna Kea during 25.3 hrs of observations. The velocity dispersion profile was studied in the context of Newtonian gravity with several types of dark matter profiles in van Dokkum et al. (2019b). Here we use these data to discuss Dragonfly 44 in MOND.

2. Velocity dispersion profile expected in MOND in isolation

Let us first study the velocity dispersion profile of Dragonfly 44 as expected by MOND, assuming that the galaxy is far away from the Coma cluster center and any other galaxies, so that the EFE is negligible. We calculated it using the spherically symmetric Jeans equation (e.g., Binney & Tremaine 2008)

$$\frac{1}{\rho} \frac{d(\rho \sigma_r^2)}{dr} + 2 \frac{\beta(r)}{r} \sigma_r^2 = a(r), \quad (1)$$

where ρ is the density of tracers, σ_r the radial velocity dispersion, β the anisotropy parameter, and a the of the radial gravitational acceleration. In our case, ρ is the star density of the galaxy that we approximated by a Sérsic sphere whose parameters were obtained by fitting the photometric profile (van Dokkum et al. 2015a, 2016, 2019b), i.e. the I -band luminosity of $L_I = 3.0 \pm 0.6 \times 10^8 L_\odot$, the effective radius of $R_e = 4.7 \text{ kpc}$ and the Sérsic index of $n = 0.94$. We used the analytic approximation for the density of a Sérsic sphere by Lima Neto et al. (1999) with the update by Márquez et al. (2000). The stellar mass-to-light ratio expected from the stellar population synthesis model of Dragonfly 44 is between 1 and 1.5 (van Dokkum et al. 2019b). This density determines the MOND gravitational field $a(r)$ (Milgrom 1983). The Newtonian gravitational field was recalculated to the MOND field assuming the so-called simple interpolation function (see, e.g., Famaey & McGaugh 2012). The parameter β was considered to be zero corresponding to an isotropic profile. We then obtained the line-of-sight velocity dispersion using the formulas given in the Appendix of Mamon & Łokas (2005).

We did not perform any fitting. Rather, we assumed a fixed mass-to-light ratio of 1.3 and the best-estimate of the galaxy luminosity as $3.0 \times 10^8 L_\odot$, which lead to our fiducial stellar mass of $M_{*,0} = 3.9 \times 10^8 M_\odot$. The resulting line-of-sight velocity dispersion profile is shown in Fig. 1. The agreement with observations in the center of the UDG is remarkably good given that the MOND model had no free parameters.




Fig. 1. Comparison of the measured velocity dispersion profile of the ultra diffuse galaxy Dragonfly 44 to the MOND no-fitting model and the Newtonian model. A fixed mass-to-light ratio of 1.3 and the best-estimate of the galaxy luminosity as $3.0 \times 10^8 L_\odot$ were assumed, together with isotropy. We illustrate the effect of relaxing some of those assumptions in Fig. 2.

In the dark matter framework, it is important to realize that this central velocity dispersion could have been very different from the MOND prediction, and there would actually be no obvious reason why this agreement would be so good given the unclear origin of UDGs. Indeed, for comparison, the dotted line is an isotropic model employing Newtonian gravity without dark matter, which clearly underestimates the velocity dispersion by more than 3σ .

On the other hand, while the deviation of the MOND models from observation increases with radius, the data points are still within the 2σ uncertainty limit so that the no-fitting model is consistent with the data. We discuss the implications of our results and possible reasons – beyond the possible measurement error – for the deviation in the outer part of the profile in the following section.

3. Discussion

3.1. Dragonfly 44 far from the cluster core

The fact that the isotropic MOND model calculated with the assumption of isolation was generally successful in Dragonfly 44 and the Newtonian was not, suggests that the UDG has to be far from the Coma cluster core in the framework of MOND. If Dragonfly 44 was sufficiently close to the center of the Coma cluster for a long time, the UDG would be subject to the EFE exerted by the cluster so that the gravity and velocity dispersion would be lower than in isolation – in the most extreme case they would have the Newtonian values, see, e.g., Fig. 1 in Müller et al. (2019a) for the predicted impact of the EFE onto different UDGs near massive galaxies as a function of their 3D separation. To give an approximate scale for how far away Dragonfly 44 has to be from the Coma cluster center in order for the EFE to be negligible, we calculate the separation at which the acceleration caused by the cluster equals the acceleration in the UDG at $1 R_e$ in the isolated case. A dedicated MOND estimate of the mass of the cluster in MOND is $4.6 \times 10^{14} M_\odot$ (Sanders 2003). This includes also the contribution of the dark matter that MOND predicts for galaxy clusters. We conclude that the distance be-

tween Dragonfly 44 and the Coma cluster has to be well beyond 5 Mpc. For comparison, the projected separation is 1.7 Mpc (angular separation of 1° at the distance of 100 Mpc). The wide field map of the UDGs around the Coma cluster by Zaritsky et al. (2019) indeed reveals that UDGs occur at least to the distance of 15 Mpc from the cluster center. The UDGs lie in cosmic filaments of galaxies that seem to flow toward the cluster. Intriguingly, another evidence in favor of Dragonfly 44 being quite far from the Coma cluster itself is the fact noted by van Dokkum et al. (2019b) that the UDGs close to Dragonfly 44 have similar radial velocities, whose differences are much smaller than the average velocity dispersion in the Coma cluster itself. If we require Dragonfly 44 to head toward the cluster, the difference of radial velocities of the cluster and the galaxy (van Dokkum et al. 2015b) implies that the galaxy is in the background of the cluster.

3.2. The effect of anisotropy

The deviation of the predicted vs. the observed velocity dispersion profile of Dragonfly 44 is less than 2 sigmas at all radii, and thus it could simply be a systematic error in this difficult observation. The deviation indeed increases as the surface brightness of the object decreases, and the data seem to imply that the dispersion profile is almost flat then suddenly start rising, which is very unlikely in the Λ CDM framework too. Nevertheless, for completeness, we now discuss in the rest of the paper the possibility that the deviation is – at least partly – real.

It is first mandatory to discuss the effect of the well known downside of the Jeans modeling approach, namely the ignorance of the anisotropy parameter. The effect of changing this parameter is illustrated in Fig. 2. When an anisotropy parameter of 0.5 is assumed, which corresponds to a radial anisotropy as hinted by the observations (van Dokkum et al. 2019b), the expected velocity dispersion profile is decreasing. The choice of an externally radial anisotropy $\beta_{ER} = r/(r + 1.4R_e)$ that is isotropic in the center and becomes radial at larger radii matches the central data points better than both the isotropic and radial anisotropies but still deviates at larger radii. A similar profile is expected to arise if the galaxy was in the external field dominated regime in the past (Wu et al. 2008). Nevertheless, it is observationally difficult to determine the anisotropy even for much brighter galaxies and as van Dokkum et al. (2019b) note, the signs for radial anisotropy have other possible interpretations.

The modeled velocity profile for tangential anisotropy with $\beta = -0.5$ has a shape similar to the observed profile but it has systematically lower values than observed. Assuming a higher M/L ratio and galaxy luminosity, each at their 2σ upper limit, leads to the stellar mass of $1.9M_{*,0}$, which reaches the lower one sigma uncertainty limits of most of the data points. Nevertheless, stellar M/L ratios are generally subjects of high uncertainties caused by the ignorance of the evolution of AGB stars and the initial mass function (e.g., Zibetti et al. 2009; Martinsson et al. 2013) such that the uncertainty in M/L given by van Dokkum et al. (2019b) may be underestimated. A higher M/L together with a larger distance caused by Dragonfly 44 lying in the background of the cluster makes then a stellar mass of $3M_\odot$ plausible. This value together with $\beta = -0.5$ provides an excellent model to the data, see the dash-dotted line in Fig. 2.




Fig. 2. Various models of the MOND velocity dispersion profile of Dragonfly 44 discussed in Sect. 3.2. Top panel: Radially anisotropic models. The isotropic model – as in Fig. 1 – is displayed as a solid line, the externally radial anisotropy model as a dashed line, and the pure radial anisotropy model as a dot-dashed line. Bottom panel: Tangentially anisotropic models. Three models for the tangential anisotropy with different stellar masses, as discussed in Sect. 3.2, are displayed with the solid, dashed, and dot-dashed lines.

3.3. Dragonfly 44 being disrupted?

Now we consider the possibility that Dragonfly 44 is actually closer to the cluster core. In this case, such a UDG cannot be stable, or it needs additional dark matter. Hence Milgrom (2015) suggested that UDGs in the Coma cluster could fall on radial orbits into the Coma cluster for the first time, where they would then be disrupted by the combination of the EFE and tidal forces. Again, this possibility is consistent with the facts that UDGs are observed to flow to the Coma cluster in filaments, that several UDGs close to Dragonfly 44 share a similar radial velocity, and several other observational pieces of evidence that some UDGs form outside clusters (see the references in the introduction). The tidal radius (Zhao 2005) of Dragonfly 44 is equal to $2R_e$ if the three dimensional separation from the cluster is 2.6 Mpc. This is larger than the projected on-sky separation so that the possibility that Dragonfly 44 is being affected by tidal forces should be considered seriously.

If the velocity dispersion of the UDG was set in a region where the EFE is negligible, we do not expect that the velocity dispersion would decrease when the UDG entered in the EFE dominated regime. The fact that the gravitational well becomes

shallower rather means that the object becomes unbound from its own gravity and dissolves. The outer parts of the object would obtain extra acceleration by tidal forces. This could then explain the observed rising velocity dispersion profile of Dragonfly 44. This scenario has a few other appealing features: it would explain why Dragonfly 44 is very elongated for a non-rotating object and why van Dokkum et al. (2019b) noted hints that the stars of the galaxy are on radial orbits. A simulation of a satellite being disturbed by a field of a more massive host is depicted in Fig. 5 of Brada & Milgrom (2000), where we can see that the object is indeed expanding and has an elongated shape.

One could suspect that the downside of this explanation is that we would have to observe the 10 Gyr old UDG (van Dokkum et al. 2019b) in a rather special period of its lifetime. With the expansion velocity estimated as the deviation of the data from the equilibrium isotropic model of around 10 km s^{-1} , the stars pass the effective radius in 460 Myr. If this expansion took a much longer time the object would dissolve. Moreover, the tidal stripping cannot be very effective because the galaxy possesses many (~ 100) globular clusters. On the other hand, in this scenario we do not observe the galaxy at a random instance of its life but when it already entered the cluster so that signs of disruption are more expected based on the closeness of the galaxy to the cluster. All these considerations clearly need support by future numerical simulations.

3.4. Dragonfly 44 with a halo of baryonic dark matter?

The fact that many UDGs in the Coma cluster field lie within the projected radius where the tidal forces are expected to be substantial, might suggest that they need additional dark halos to survive in MOND (Milgrom 2015, the density of UDGs as a function of cluster-centric distance was studied, e.g., in van der Burg et al. 2016).

Indeed, many independent pieces of evidence suggest that MOND does require dark matter in galaxy clusters that could be made of baryons (see, e.g., the review in Milgrom 2008). One possibility, suggested by Milgrom (2015), would be that cluster UDGs could host many of these dark baryons. The observed rising velocity dispersion profile could then be interpreted as a consequence of the presence of such a dark halo around the UDGs. However, a typical dark halo rotation curve would not reproduce the outer spike in velocity dispersions hinted at by the data. But note that the baryonic dark halos of UDGs have likely different density profiles from what we are used for for the non-baryonic dark halos in Newtonian gravity (Navarro et al. 1997). Moreover, the fact that the velocity dispersion in the center of Dragonfly 44 matches the MOND prediction for isolation gives us an important information: if Dragonfly 44 is at a large distance from the cluster core, and that the EFE is weak, then its halo must be extended so that it has no strong gravitational influence near the center of the galaxy. On the other hand, if the EFE acting on the UDG is substantial, then the increase of gravity due to the halo in the central parts must *coincidentally* compensate for the decrease of the MOND boost.

The presence of a dark baryonic halo might be hinted to by the fact that Dragonfly 44 has around ten times more globular clusters than dwarf galaxies of the same luminosity (Amorisco et al. 2018). If we interpret the empirical relation between the galaxy luminosity and number of globular clusters as a relation between the total baryonic mass of the galaxy and the number of globular clusters, then it follows from Fig. 6 of Amorisco et al. (2018) that the baryonic mass of Dragonfly 44 is around

$10^{10} M_{\odot}$, i.e. the amount of dark baryons in this galaxy is a few tens of $M_{*,0}$.

We should note that Dragonfly 44 can be without a dark matter halo while the UDGs near the cluster center have one. Many formation scenarios for UDGs were suggested in the Λ CDM context and therefore the same should be allowed in the MOND context. At least one way to form a UDG without substantial amounts of dark baryons is known: some UDGs seem to be tidal dwarf galaxies (Bennet et al. 2018) and these are not supposed to contain a much larger proportion of dark baryons than their parent galaxies, i.e. the spiral galaxies, and those do not seem to contain much dark baryons (McGaugh et al. 2000; Tiret & Combes 2009).

3.5. How to find the answer?

One of the main uncertainties in interpreting the data for Dragonfly 44 is that we do not know whether the EFE exerted by the cluster is negligible, nor whether the galaxy is in dynamical equilibrium.

In the case of hypothetical baryonic dark matter halos suggested by Milgrom (2015), the easiest way to detect them would be to focus the future observations on UDGs that are in projection far enough from the cluster to be considered isolated. Then a higher than predicted velocity dispersion would be sign of a dark halo. For the Coma cluster and a typical UDG, it means a projected separation of at least 4.5° (assuming the median UDG mass and radius given by van Dokkum et al. 2015a of $6 \times 10^7 M_{\odot}$ and $r_e = 2.8 \text{ kpc}$, respectively). Figure 2 suggest that the outer parts of UDGs are particularly suitable for this task since the velocity dispersion profile of Dragonfly 44 is quite insensitive to the precise value of the anisotropy parameter (the reasons for this are probably related to the results of Wolf et al. 2010).

Concerning UDGs in the central parts of the clusters, in order to distinguish the cases of disrupting UDGs or UDGs with dark halos, we would have to compare them to their analogs in simulations on their supposed orbits. From this perspective, it is most advantageous to observe in the future the UDGs close to cluster centers, but at distances where the internal dynamics of the UDGs is not expected to be influenced by tidal forces by the main competitors of MOND, the theories supposing Newtonian gravity and non-baryonic dark matter.

An interesting result would be if all UDGs in clusters came out to have velocity dispersions corresponding to the isolated case similarly to Dragonfly 44. The chance that all of the central UDGs are just lying there in projection and in reality are in the outskirts is dim (van der Burg et al. 2016). Similarly, the fraction of UDGs having a dark halo whose dynamical effect would precisely compensate for the decrease of MOND gravity due to the EFE should be low. Then, two possibilities would arise: either MOND is excluded as a modification of gravity (but this would leave the question open of why the observations would agree with the isolated MOND prediction), or the UDGs are on their first radial infall and are in the process of disruption as suggested by Milgrom (2015).

Note however that some theories actually expect that the internal dynamics of UDGs is not influenced by the cluster. For instance, MOND could be modified such that it does not require dark matter in galaxy clusters as proposed in Zhao & Famaey (2012). This could help the galaxies survive, but Hodson & Zhao (2017) showed that the high velocity dispersion, now measured at the last data point of Dragonfly 44, could not be explained in this way, which would bring us back to the hypothesis of disruption. Another exotic possibility is that the empiri-

cal MOND effect in galaxies is due to the coupling of baryons to the phonons of a dark matter superfluid core which would encompass their baryonic profile. As extensively discussed in Sect. IX.B of Berezhiani et al. (2018), galaxy clusters would have small superfluid cores of about 100 kpc and no EFE would then act on the UDGs residing outside of this core, while they would still obey the isolated MOND prediction in the central parts, but have an additional effect from dark matter itself in the outskirts. Finally, the EFE can be weak in some modified inertia versions of MOND (Milgrom 2011), but this would leave the question open of why UDGs in groups, such as NGC 1052, seem to be affected by the EFE, apart if they are, in fact, foreground objects.

4. Conclusion

Recently, the first velocity dispersion profile of an ultra-diffuse galaxy, namely Dragonfly 44, has been measured and analyzed in the framework of Newtonian dynamics and dark matter (van Dokkum et al. 2019b). Here we computed Jeans models for this galaxy in the framework of MOND. Assuming the galaxy to reside in isolation we found that an isotropic model reproduces the observed velocity dispersion near the center without any *ad hoc* tuning of free parameters. For the outer parts, the modeled velocity dispersion deviates from the measured data points but it is still within the two sigma uncertainty limit of the measurement. While the modeled velocity dispersion profile is almost flat, the observed one is steeply rising in the outer parts.

We proposed several interpretations of our results:

1. the galaxy is far from the cluster (thus in isolation) and the outer deviation is simply a measurement error
2. the galaxy is far from the cluster, and the outer deviation is mostly an effect of tangential anisotropy for a very high stellar mass of the galaxy, which brings back the discrepancy at less than one sigma.
3. the galaxy is closer to the cluster core, and is in the process of disruption: the inner parts are not yet observably affected by the EFE and the deviation in the outskirts is caused by tidal deformation.
4. the deviation is a manifestation of a baryonic dark halo around the UDG, which raises the question of why it then obeys the central velocity dispersion predicted by MOND in isolation.

We also discussed more exotic possibilities such as modifications of inertia without the EFE or superfluid dark matter, in which case the galaxy does not need to be far away from the cluster core to obey the isolated MOND prediction, but still requires an explanation for the outer data points mismatch. In order to test these scenarios, it is necessary to measure the velocity dispersion of more of these objects as well as to perform simulations of the evolution of UDGs moving through galaxy clusters.

Acknowledgements. O.M. is grateful to the Swiss National Science Foundation for financial support.

References

Alabi, A., Ferré-Mateu, A., Romanowsky, A. J., et al. 2018, MNRAS, 479, 3308
 Amorisco, N. C. & Loeb, A. 2016, MNRAS, 459, L51
 Amorisco, N. C., Monachesi, A., Agnello, A., & White, S. D. M. 2018, MNRAS, 475, 4235
 Begeman, K. G., Broeils, A. H., & Sanders, R. H. 1991, MNRAS, 249, 523
 Bekenstein, J. & Milgrom, M. 1984, ApJ, 286, 7

Bennet, P., Sand, D. J., Zaritsky, D., et al. 2018, ApJ, 866, L11
 Berezhiani, L., Famaey, B., & Khoury, J. 2018, Journal of Cosmology and Astro-Particle Physics, 2018, 021
 Binney, J. & Tremaine, S. 2008, Galactic Dynamics: Second Edition (Princeton University Press)
 Bosma, A. 1981, AJ, 86, 1791
 Brada, R. & Milgrom, M. 2000, ApJ, 541, 556
 Chan, T. K., Kereš, D., Wetzel, A., et al. 2018, MNRAS, 478, 906
 Crnojević, D., Sand, D. J., Caldwell, N., et al. 2014, ApJ, 795, L35
 Di Cintio, A., Brook, C. B., Dutton, A. A., et al. 2017, MNRAS, 466, L1
 Dressler, A. 1980, ApJ, 236, 351
 Emsellem, E., van der Burg, R. F. J., Fensch, J., et al. 2018, arXiv e-prints [arXiv:1812.07345]
 Famaey, B., McGaugh, S., & Milgrom, M. 2018, MNRAS, 480, 473
 Famaey, B. & McGaugh, S. S. 2012, Living Reviews in Relativity, 15, 10
 Fensch, J., van der Burg, R. F. J., Jerabkova, T., et al. 2018, arXiv e-prints [arXiv:1812.07346]
 Ferré-Mateu, A., Alabi, A., Forbes, D. A., et al. 2018, MNRAS, 479, 4891
 Hodson, A. O. & Zhao, H. 2017, A&A, 607, A109
 Impey, C., Bothun, G., & Malin, D. 1988, ApJ, 330, 634
 Koda, J., Yagi, M., Yamanoi, H., & Komiyama, Y. 2015, ApJ, 807, L2
 Kroupa, P., Hāghi, H., Javanmardi, B., et al. 2018, Nature, 561, E4
 Laporte, C. F. P., Agnello, A., & Navarro, J. F. 2019, MNRAS, 484, 245
 Leisman, L., Haynes, M. P., Janowiecki, S., et al. 2017, ApJ, 842, 133
 Lelli, F., McGaugh, S. S., Schombert, J. M., & Pawlowski, M. S. 2017, ApJ, 836, 152
 Lima Neto, G. B., Gerbal, D., & Márquez, I. 1999, MNRAS, 309, 481
 Mamon, G. A. & Łokas, E. L. 2005, MNRAS, 363, 705
 Mancera Piña, P. E., Aguerri, J. A. L., Peletier, R. F., et al. 2019, MNRAS, 485, 1036
 Márquez, I., Lima Neto, G. B., Capelato, H., Durret, F., & Gerbal, D. 2000, A&A, 353, 873
 Martin, N. F., Collins, M. L. M., Longeard, N., & Tollerud, E. 2018, ApJ, 859, L5
 Martínez-Delgado, D., Läsker, R., Sharina, M., et al. 2016, AJ, 151, 96
 Martinsson, T. P. K., Verheijen, M. A. W., Westfall, K. B., et al. 2013, A&A, 557, A131
 McGaugh, S. & Milgrom, M. 2013, ApJ, 775, 139
 McGaugh, S. S. & de Blok, W. J. G. 1998, ApJ, 499, 66
 McGaugh, S. S., Schombert, J. M., Bothun, G. D., & de Blok, W. J. G. 2000, ApJ, 533, L99
 Merritt, A., van Dokkum, P., Danieli, S., et al. 2016, ApJ, 833, 168
 Milgrom, M. 1983, ApJ, 270, 365
 Milgrom, M. 2008, New A Rev., 51, 906
 Milgrom, M. 2011, arXiv e-prints, arXiv:1111.1611
 Milgrom, M. 2014, MNRAS, 437, 2531
 Milgrom, M. 2015, MNRAS, 454, 3810
 Müller, O., Famaey, B., & Zhao, H. 2019a, A&A, 623, A36
 Müller, O., Jerjen, H., & Binggeli, B. 2018, A&A, 615, A105
 Müller, O., Rich, R. M., Román, J., et al. 2019b, A&A, 624, L6
 Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, ApJ, 490, 493
 Nusser, A. 2019, MNRAS, 484, 510
 Papastergis, E., Adams, E. A. K., & Romanowsky, A. J. 2017, A&A, 601, L10
 Román, J. & Trujillo, I. 2017a, MNRAS, 468, 703
 Román, J. & Trujillo, I. 2017b, MNRAS, 468, 4039
 Rubin, V. C., Ford, Jr., W. K., & Thonnard, N. 1978, ApJ, 225, L107
 Rubin, V. C., Ford, Jr., W. K., Thonnard, N., & Burstein, D. 1982, ApJ, 261, 439
 Sandage, A. & Binggeli, B. 1984, AJ, 89, 919
 Sanders, R. H. 2003, MNRAS, 342, 901
 Sardone, A., Pisano, D. J., Burke-Spolaor, S., Mascoop, J. L., & Pol, N. 2019, ApJ, 871, L31
 Shi, D. D., Zheng, X. Z., Zhao, H. B., et al. 2017, ApJ, 846, 26
 Tiret, O. & Combes, F. 2009, A&A, 496, 659
 Toloba, E., Lim, S., Peng, E., et al. 2018, ApJ, 856, L31
 Trujillo, I., Beasley, M. A., Borlaff, A., et al. 2019, MNRAS [arXiv:1806.10141]
 van der Burg, R. F. J., Muzzin, A., & Hoekstra, H. 2016, A&A, 590, A20
 van Dokkum, P., Abraham, R., Brodie, J., et al. 2016, ApJ, 828, L6
 van Dokkum, P., Danieli, S., Abraham, R., Conroy, C., & Romanowsky, A. J. 2019a, arXiv e-prints [arXiv:1901.05973]
 van Dokkum, P., Danieli, S., Cohen, Y., et al. 2018a, Nature, 555, 629
 van Dokkum, P., Danieli, S., Cohen, Y., Romanowsky, A. J., & Conroy, C. 2018b, ApJ, 864, L18
 van Dokkum, P., Wasserman, A., Danieli, S., et al. 2019b, arXiv e-prints [arXiv:1904.04838]
 van Dokkum, P. G., Abraham, R., Merritt, A., et al. 2015a, ApJ, 798, L45
 van Dokkum, P. G., Romanowsky, A. J., Abraham, R., et al. 2015b, ApJ, 804, L26
 Venhola, A., Peletier, R., Laurikainen, E., et al. 2017, A&A, 608, A142
 Wittmann, C., Lisker, T., Ambachew Tilahun, L., et al. 2017, MNRAS, 470, 1512
 Wolf, J., Martinez, G. D., Bullock, J. S., et al. 2010, MNRAS, 406, 1220
 Wu, X., Famaey, B., Gentile, G., Perets, H., & Zhao, H. 2008, MNRAS, 386, 2199
 Yozin, C. & Bekki, K. 2015, MNRAS, 452, 937
 Zaritsky, D., Donnerstein, R., Dey, A., et al. 2019, ApJS, 240, 1
 Zhao, H. & Famaey, B. 2012, Phys. Rev. D, 86, 067301
 Zhao, H. S. 2005, A&A, 444, L25
 Zibetti, S., Charlot, S., & Rix, H.-W. 2009, MNRAS, 400, 1181