



Have conditions under which women undergo abortion improved around the world ?

Agnès Guillaume, Clémentine Rossier

► To cite this version:

Agnès Guillaume, Clémentine Rossier. Have conditions under which women undergo abortion improved around the world ?. N-IUSSP - IUSSP's online news magazine, 2019. <hal-02307012>

HAL Id: hal-02307012

<https://hal.science/hal-02307012v1>

Submitted on 7 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Have conditions under which women undergo abortion improved around the world?

Les conditions de l'avortement dans le monde: évolutions récentes

[Agnès Guillaume, Clémentine Rossier](#)



Agnès Guillaume and Clémentine Rossier review the evidence on abortion in the world since 1990. Few countries have liberalized their laws since then, and regressive movements are frequent. The diffusion of medical abortion and post-abortion care have made abortion safer, but progress is lagging for the poorest women and countries. While better contraception means fewer abortions, other adverse factors are raising the number of unintended pregnancies and the need for abortion.


Have conditions under which women undergo abortion improved around the world? Access to safe and legal abortion informs on a key aspect of women's status, namely their ability to control whether and when to have children. Guillaume and Rossier (2018) examine recent progress in women's use of abortion worldwide, from legal, public health and sociodemographic perspectives.

Legalization of abortion at the global level: one step forward, one step back

Legalization of abortion represents a crucial step in securing women access to safe abortion. While a few countries have recently liberalized their abortion laws (Ireland, Argentina,...), progress at the global level remains slow. The Global Abortion Policies database (WHO, 2017) shows that abortion laws are still closely related to countries' level of development. In developed countries, 95% of women of reproductive age benefit from liberal laws (i.e., abortion is legal on request, or at least on economic or social grounds), compared to 45% of women in less developed countries, and only 8% in least developed countries (Figure 1). In addition, and whatever the region, numerous barriers, such as short gestational limits, compulsory parental or spousal consent, requirements for medical or legal approval, or mandatory waiting periods, can jeopardize access to legal abortion services. Moreover, in

certain liberal countries, such as Uruguay or Italy, providers frequently use their right to conscientious objection, further limiting access.

Figure 1. Distribution (%) of women aged 15-49 years by permitted legal grounds for abortion and level of development in 2013



Sources: authors' calculations on the basis of World Abortion Policies 2013 - World Population Prospects: The 2017 Revision (2015 figures).

Two opposing positions have weighed equally in legal debates around the world in recent decades: those promoting women's sexual and reproductive rights as defined in international conferences, and those seeking to protect the rights of the embryo. As a result, laws tend to oscillate between periods of easing and tightening of restrictions, liberalization is rare, and the right to abortion can be challenged even when it has already been established. In Nicaragua, for example, even therapeutic abortion was banned in 2006, despite the resulting risks for women's health and lives. Active and widespread campaigns at both national and international levels are currently attempting to reverse these measures. Conversely, in late 2016, the Polish government abandoned its plan to limit access to abortion, but a new law has been in preparation since January 2018, aimed at restoring measures that restrict the right to abortion to two situations: cases where the woman's health or life are in danger; and cases of rape or incest. Recently in the United States, where laws are defined at the state level, the right to abortion has been contested in certain states.

Abortions have become safer around the world, but many women are excluded

While legality and safety were closely associated in the 1980s, that is no longer the case thanks to the diffusion of new abortion techniques (aspiration and medical method) since the 1990s and 2000s. Use of medical methods (mainly misoprostol) has increased in both restrictive and liberal countries, decreasing the risk of illegal abortions. In settings where abortion is legal, aspiration has replaced curettage at longer gestational periods, except in a number of Asian and former Soviet countries. As a result, three categories of risk are now recognized for abortions, independently of legal status: safe (recommended techniques *and* providers), less safe (recommended technique *or* provider), and least safe (neither of these safety criteria are met). Estimates of abortion safety in different world regions indicate that just over half (55%) of the 56 million annual induced abortions in 2010–2014 were safe, 31% less safe and 15% least safe (Table 1) (Ganatra et al., 2017).

Table 1. Proportions of abortions by estimated degree of safety and geographical zone, 2010-2014

	Safe		Less safe (1)		Least safe (2)		Unsafe (1+2)	
	%	90% CI	%	90% CI	%	90% CI	%	90% CI
World	54.9	[49.9;59.4]	30.7	[25.5;35.6]	14.4	[11.5;18.1]	45.1	[40.6;50.1]
Developed countries ^a	87.5	[81.9;89.6]	12.4	[10.2;17.9]	0.08	[0.0;1.36]	12.5	[10.4;18.1]
Developing countries	50.5	[45.2;55.9]	33.2	[27.0;38.3]	16.3	[13.1;20.7]	49.5	[44.1;54.9]
North America	99	[97.7;99.8]	0.9	[0.2;2.3]	0	[0.0;0.03]	0.9	[0.2;2.3]
Europe	88.8	[80.3;91.7]	11.2	[7.8;19.3]	0	[0.0;0.02]	11.2	[8.3;19.7]
Asia	62.1	[54.8;67.2]	29.7	[23.5;36.6]	8.3	[4.9;13.3]	37.8	[32.8;45.2]
Latin America	23.6	[8.8;47.0]	59.7	[32.7;72.2]	16.7	[8.8;33.4]	76.4	[53.0;91.3]
Africa	24.4	[18.6;33.6]	27.6	[21.2;37.0]	48	[36.5;52.9]	75.6	[66.4;81.4]
Oceania	66.3	[61.4;77.7]	7.8	[3.5;17.9]	25.9	[11.5;31.1]	33.7	[22.3;38.6]

a United Nations Population Division definition.

Interpretation: The proportion of safe abortions in the world in 2010-2014 is estimated at 54.9%, and there is a 90% chance that the real figure is between 49.9% and 59.4%. “Unsafe” is the sum of “less” and “least safe”.

Source: Ganatra et al. (2017).

Today, misoprostol is used illegally throughout Latin America and in Asian countries with restrictive legislation, such as the Philippines. In Africa, it is less common, but its use is growing in some countries, such as Uganda, Gabon, and Nigeria. Across regions, it is increasingly common for women to obtain these medications through informal markets or online; some international and local NGOs also sell and deliver these products or provide helplines offering guidance to women.

Thanks to the growing use of misoprostol and the implementation of post abortion care programs since the 1990s, the frequency of abortion-related deaths has decreased worldwide. There is today about one complication for every four or five unsafe abortions worldwide (Singh and Maddow-Zimet, 2016). However, abortion complications and deaths are still distributed very unequally, and disproportionately affect the world’s poorest women and less developed countries, especially in sub-Saharan Africa.

Better contraception means fewer abortions, but other factors are involved

It is estimated that today (2010–2014), on average, at global level, 35 out of every 1,000 women aged 15 to 44 years have an induced abortion each year (Sedgh et al., 2016). This corresponds to 56 million annual abortions, or 25% of pregnancies worldwide. While abortion is frequent everywhere, its rate is lower in liberal countries with better sexual and reproductive programs and higher contraceptive prevalence.

However, other important factors determine women’s need for abortions, as shown by a review of abortion seekers’ profiles in different world regions. In Asia, women who have abortions are often older, already mothers, and married. Premarital sex remains rare, and family formation begins early; abortion thus serves mainly at the end of reproductive life to limit the number of children if contraception fails, or to obtain male children in settings where sex-selective abortion is practiced. In Africa, by contrast, the desired family size remains large, which limits the number of abortions within marriage and at the end of reproductive life. And while premarital sexual activity is more common than in Asia, it is nonetheless socially condemned and often poorly protected, which can lead to unwanted pregnancies and abortions at young ages. The profile of Latin America is intermediate, between Asia and

Africa: it is closer to Africa at young ages, but as in Asia, many women also seek to limit births at the end of their reproductive life.

In Europe and North America, where effective contraception is widely used by women both at the beginning and the end of reproductive life, abortions are less frequent but more linked to the phase of family formation, when fertility intentions are ambivalent and changing, and contraceptive failures more frequent. The proportion of repeat abortions among all abortions has increased in countries such as Sweden, New Zealand and France. Paradoxically, this reflects improvements in the contraceptive practices of the population as a whole, with abortions concentrated among women who repeatedly experience contraceptive failure or use methods ill-suited to their situation.

In general terms, abortion has become safer across the world, but inequalities in the risks associated with illegal abortion have increased. Securing abortion as a right is not making much progress, despite the international recognition of rights to equality, health, and free and responsible decision-making on the number and spacing of children. And while effective contraceptive services lower the occurrence of unintended pregnancies and the need for abortion, other factors such as emerging premarital sexuality, delayed parenthood, smaller desired family size, sexual violence and prenatal sex selection, all pull in the opposite direction.

References

Ganatra Bela, Gerdt Caitlin, Rossier Clémentine, et al. (2017) “Global, regional, and subregional classification of abortions by safety, 2010–14: Estimates from a Bayesian hierarchical model”, *The Lancet*, 390 (10110), pp. 2372–2381.

Guillaume Agnès, Rossier Clémentine (2018) Abortion around the world. An overview of legislation, measures, trends, and consequences. *Population*, 73(2), 217-306.

Sedgh Gilda, Bearak Jonathan, Singh Susheela, Akinrinola Bankole, Anna Popinchalk, Bela Ganatra, Clémentine Rossier, et al. (2016) “Abortion incidence between 1990 and 2014: Global, regional, and subregional levels and trends”, *The Lancet*, 388(10041), pp. 258–267.

Singh Susheela, Maddow-Zimet Isaac (2016) “Facility-based treatment for medical complications resulting from unsafe pregnancy termination in the developing world, 2012: A review of evidence from 26 countries”, *BJOG: An International Journal of Obstetrics & Gynaecology*, 123(9), pp. 1489–1498.

WHO – [World Health Organization \(2017\) Global Abortion Policies Database, Geneva, WHO,](#)