



**HAL**  
open science

# Ionization of strontium atoms by ultrashort pulses of a Ti-Sapphire laser

A Dimitriou, Alexandre Marciniak, F. Lépine, C. Bordas, S. Cohen

► **To cite this version:**

A Dimitriou, Alexandre Marciniak, F. Lépine, C. Bordas, S. Cohen. Ionization of strontium atoms by ultrashort pulses of a Ti-Sapphire laser. *Journal of Physics: Conference Series*, 2015, 635 (9), pp.092083. 10.1088/1742-6596/635/9/092083 . hal-02308760

**HAL Id: hal-02308760**

**<https://hal.science/hal-02308760>**

Submitted on 5 Feb 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PAPER • OPEN ACCESS

## Ionization of strontium atoms by ultrashort pulses of a Ti-Sapphire laser

To cite this article: A Dimitriou *et al* 2015 *J. Phys.: Conf. Ser.* **635** 092083

View the [article online](#) for updates and enhancements.

### Related content

- [Various Kinds Waves and Solitons Interaction Solutions of Boussinesq Equation Describing Ultrashort Pulse in Quadratic Nonlinear Medium](#)  
Bang-Xing Guo, Zhan-Jie Gao and Ji Lin
- [Fibre laser with a subterahertz repetition rate of ultrashort pulses in the telecom range](#)  
A V Andrianov, V M Mylnikov, M Yu Koptev *et al.*
- [\(3+1\)-dimensional nonlinear propagation equation for ultrashort pulsed beam in left-handed material](#)  
Hu Yong-Hua, Fu Xi-Quan, Wen Shuang-Chun *et al.*



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

## Ionization of strontium atoms by ultrashort pulses of a Ti-Sapphire laser


A. Dimitriou<sup>†1\*</sup>, A. Marciniak<sup>‡2</sup>, F. Lepine<sup>‡3</sup>, C. Bordas<sup>‡4</sup>, S. Cohen<sup>†5</sup>

<sup>†</sup> Dept. of Physics, University of Ioannina, GR 45110 Ioannina, Greece

<sup>‡</sup>Institute Lumière Matière, Université Lyon 1, UMR 5306, 10 rue Ada Byron 69622, Villeurbanne, France

**Synopsis** We present experimental photoelectron spectra obtained by ionizing two-valence electron Sr atoms with  $\approx 25$  fs Ti-Sapphire laser pulses and we show preliminary evidence that electron-electron correlation effects dominate for laser intensities  $< 0.4 \times 10^{13} \text{ W}\cdot\text{cm}^{-2}$ , while a single electron picture prevails when the laser intensity is increased further.

The two-electron Alkaline Earth atoms present dense manifolds of doubly excited autoionizing states above their first ionization threshold. Hence, their continua are structured and these states are dominated by electron-electron interaction effects. The interaction and subsequent ionization of Alkaline Earth atoms with intense laser fields reveals the absorption of photons within the structured continua. For laser pulses of long duration ( $\sim$ ns), there exist above-threshold-absorption pathways which are dominated by the presence of these doubly excited states and lead to the production of excited ions [1]. On the contrary, for pulses of short duration ( $\sim$ fs) energy level structure issues are of much less importance and one expects a single-active-electron picture to prevail [2,3]. In this case ions should be mostly produced to their ground state. This latter expectation is here experimentally examined by ionizing Sr atoms via  $\sim 25$  fs,  $\sim 800$  nm laser pulses, with intensities ranging from  $I \approx 2 \times 10^{12} \text{ W}\cdot\text{cm}^{-2}$  to  $\approx 5 \times 10^{13} \text{ W}\cdot\text{cm}^{-2}$ . A preliminary analysis of measured photoelectron spectra (recorded via a velocity map imaging (VMI) apparatus [4] at Institute Lumière Matière (ILM), Université Lyon 1) shows that for  $I < 0.4 \times 10^{13} \text{ W}\cdot\text{cm}^{-2}$  electron-electron interaction dominates and population is created at excited states of  $\text{Sr}^+$ . As the laser intensity is further increased, however, the spectra are composed solely by a series of Above Threshold Ionization (ATI [3]) electron peaks (see Fig. 1) and consequently the single-active-electron picture appears to dominate. These findings suggest that the key factor controlling the dominance of one or the other mechanism is laser intensity, rather than pulse duration.



**Figure 1.** High laser intensity electron energy spectra of Sr ionized by 25 fs Ti-Sapphire laser pulses.

This research has been co-financed by the European Union (European Social Fund – ESF) and Greek national funds through the Operational Program "Education and Lifelong Learning" of the National Strategic Reference Framework (NSRF) - Research Funding Program: THALES (ISEPUMA). Investing in knowledge society through the European Social Fund.

\* **Present address:** Dept. of Physics, University of Crete, GR 71003, Heraklion, Greece

### References

- [1] I. Lontos *et al* 2008 *J. Phys. B: At. Mol. Opt. Phys.* **41** 045601
- [2] M. J Nandor *et al* 1999 *Rhys. Rev. A.* **60** R1771
- [3] P. Agostini *et al* 1979 *Phys. Rev. Lett.* **42** 1127
- [4] A. T. J. B. Eppink and D. H. Parker 1997 *Rev. Sci. Instr.* **68** 3477

<sup>1</sup>E-mail: [adimitr@physics.uoi.gr](mailto:adimitr@physics.uoi.gr)

<sup>2</sup>E-mail: [alexandre.marciniak@univ-lyon1.fr](mailto:alexandre.marciniak@univ-lyon1.fr)

<sup>3</sup>E-mail: [franck.lepine@univ-lyon1.fr](mailto:franck.lepine@univ-lyon1.fr)

<sup>4</sup>E-mail: [christian.bordas@univ-lyon1.fr](mailto:christian.bordas@univ-lyon1.fr)

<sup>5</sup>E-mail: [scohen@uoi.gr](mailto:scohen@uoi.gr)

