



**HAL**  
open science

# Asymptotic behavior for the Vlasov-Poisson equations with strong uniform magnetic field and general initial conditions

Mihai Bostan

► **To cite this version:**

Mihai Bostan. Asymptotic behavior for the Vlasov-Poisson equations with strong uniform magnetic field and general initial conditions. 2019. <hal-02310174>

**HAL Id: hal-02310174**

**<https://hal.science/hal-02310174v1>**

Preprint submitted on 9 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 Asymptotic behavior for the Vlasov-Poisson  
2 equations with strong uniform magnetic field and  
3 general initial conditions

4 Mihai BOSTAN \*

5 (June 2, 2019)

6 **Abstract**

7 We investigate the Vlasov-Poisson equations perturbed by a strong external  
8 uniform magnetic field. We study the asymptotic behavior of the solutions, based  
9 on averaging techniques. We analyze the case of general initial conditions. By  
10 filtering out the oscillations, we are led to a profile. We prove strong convergence  
11 results and establish second order estimates.

12 **Keywords:** Vlasov-Poisson system, two-scale analysis, averaging, homogenization.

13 **AMS classification:** 35Q75, 78A35, 82D10.

14 **1 Introduction**

15 We consider a population of charged particles of mass  $m$ , charge  $q$ , whose density is  
16 denoted by  $f = f(t, x, v)$ , depending on time  $t$ , position  $x$  and velocity  $v$ . In order to  
17 study the magnetic confinement, we focus on the asymptotic behavior of the Vlasov-  
18 Poisson equations, with strong external magnetic field [16, 17, 18, 24, 25, 11, 13, 22, 3,  
19 4, 5, 6, 14, 19, 15, 12].

20 Neglecting the curvature of the magnetic lines, that is  $\mathbf{B}^\varepsilon = (0, 0, B^\varepsilon) = (0, 0, B/\varepsilon)$ ,  
21 we are led to

$$\partial_t f^\varepsilon + v \cdot \nabla_x f^\varepsilon + \frac{q}{m} \{E[f^\varepsilon(t)](x) + B^\varepsilon \perp v\} \cdot \nabla_v f^\varepsilon = 0, \quad (t, x, v) \in \mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2. \quad (1)$$

The notation  $\perp(\cdot)$  stands for the rotation of angle  $-\pi/2$  of the velocity  $v = (v_1, v_2)$ ,  
*i.e.*,  $\perp v = \mathcal{R}(-\pi/2)v = (v_2, -v_1)$ ,  $v = (v_1, v_2) \in \mathbb{R}^2$  and  $\varepsilon > 0$  is a small parameter  
related to the ratio between the cyclotronic period and the advection time scale. The

---

\*Aix Marseille Université, CNRS, Centrale Marseille, I2M, Marseille France, Centre de  
Mathématiques et Informatique, UMR 7373, 39 rue Frédéric Joliot Curie, 13453 Marseille Cedex  
13 France. E-mail : [mihai.bostan@univ-amu.fr](mailto:mihai.bostan@univ-amu.fr).

electric field  $E[f^\varepsilon(t)] = -\nabla_x \Phi[f^\varepsilon(t)]$  derives from a potential, satisfying the Poisson equation

$$-\varepsilon_0 \Delta_x \Phi[f^\varepsilon(t)] = q \int_{\mathbb{R}^2} f^\varepsilon(t, x, v) dv, \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^2$$

1 where  $\varepsilon_0$  represents the electric permittivity. Appealing to the fundamental solution  
 2  $z \rightarrow -\frac{1}{2\pi} \ln |z|, z \in \mathbb{R}^2 \setminus \{0\}$ , we have

$$\Phi[f^\varepsilon(t)](x) = -\frac{q}{2\pi\varepsilon_0} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln |x - x'| f^\varepsilon(t, x', v') dv' dx' \quad (2)$$

3 and

$$E[f^\varepsilon](x) = \frac{q}{2\pi\varepsilon_0} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} f^\varepsilon(x', v') \frac{x - x'}{|x - x'|^2} dv' dx'. \quad (3)$$

4 For any particle density  $f = f(x, v)$ , we denote by  $E[f]$  the Poisson electric field  
 5 associated to the charge density  $\rho[f] = q \int_{\mathbb{R}^2} f(\cdot, v) dv$ . We also use the notation  
 6  $j[f] = q \int_{\mathbb{R}^2} f(\cdot, v) v dv$  for the current density of  $f$ . We complete the above system by  
 7 the initial condition

$$f^\varepsilon(0, x, v) = f_{\text{in}}(x, v), \quad (x, v) \in \mathbb{R}^2 \times \mathbb{R}^2. \quad (4)$$

8 Very recently, the Vlasov-Poisson equations with strong external non homogeneous  
 9 magnetic field have been studied, when considering well prepared initial densities [9,  
 10 14]. As usual, we are looking for quantities with small variations over a cyclotronic  
 11 period (the guiding center), and we average with respect to the fast cyclotronic motion  
 12 in order to obtain the effective Vlasov equation. Following this strategy, most of  
 13 the previous studies provided formal or rigorous (based on compactness arguments)  
 14 gyrokinetic approximations for the transport of charged particles under the action of  
 15 strong external magnetic fields. The subject matter in [9] was to derive second order  
 16 regular reformulations of the Vlasov-Poisson equations with strong magnetic field and  
 17 to perform the error analysis. One of the key point was to split the advection field  
 18 of the Vlasov equation into a fast and slow dynamics, such that the guiding center is  
 19 left (exactly) invariant by the fast dynamics. It was shown that in this case the fast  
 20 dynamics becomes periodic (even for a non homogeneous magnetic field), and therefore  
 21 the homogenization procedure simplifies a lot : instead of taking ergodic means, it is  
 22 enough to average over one period.

23 When the magnetic field is uniform, it is possible to go further in our analysis, by  
 24 considering smooth initial particle densities, not necessarily well prepared. We mention  
 25 that most of the studies concentrate only on models with well prepared initial particle  
 26 densities. We intend to extend the analysis in [9] for general initial conditions. The  
 27 asymptotic behavior is more complicated because the particle densities  $(f^\varepsilon)_{\varepsilon>0}$  present  
 28 fast oscillations in time. We appeal to a two scale approach by working in a extended  
 29 phase space supplemented by a fast time variable  $s = t/\varepsilon$ . Up to a second order term,  
 30 the oscillations of the family  $(f^\varepsilon)_{\varepsilon>0}$  can be described in terms of a profile solving a  
 31 regular reformulation of the Vlasov-Poisson equations, see also [25].

### 32 **Theorem 1.1**

33 *Let  $f_{\text{in}} \in C_c^2(\mathbb{R}^2 \times \mathbb{R}^2)$  be a non negative, smooth, compactly supported particle den-*  
 34 *sity. We denote by  $(f^\varepsilon)_{\varepsilon>0}$  the solutions of the Vlasov-Poisson equations (1), (2) with*

1 uniform external magnetic field  $\frac{B}{\varepsilon} \neq 0$ , corresponding to the initial condition  $f_{\text{in}}$ . Then  
 2 for any  $T \in \mathbb{R}_+$ , there is a constant  $C_T > 0$  such that for any  $\varepsilon > 0$

$$\sup_{t \in [0, T]} \left\{ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left[ f^\varepsilon(t, x, v) - \tilde{F} \left( t, x + \varepsilon \frac{\perp v}{\omega_c}, \mathcal{R}(\omega_c \frac{t}{\varepsilon}) \left( v - \varepsilon \frac{\perp E[\tilde{F}(t)]}{B} \right) \right) \right]^2 dv dx \right\}^{1/2} \leq C_T \varepsilon^2$$

3 where  $\tilde{F}$  is the solution of

$$\partial_t \tilde{F} + \varepsilon \frac{\perp E[\tilde{F}(t)]}{B} \cdot \nabla_X \tilde{F} + \varepsilon \frac{\perp (j[\tilde{F}(t)] - \rho[\tilde{F}(t)] \tilde{V})}{2\varepsilon_0 B} \cdot \nabla_{\tilde{V}} \tilde{F} = 0, \quad (t, X, \tilde{V}) \in \mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2$$

4 corresponding to the initial condition

$$\tilde{F}(0, X, \tilde{V}) = f_{\text{in}} \left( X - \varepsilon \frac{\perp \tilde{V}}{\omega_c}, \tilde{V} + \varepsilon \frac{\perp E[f_{\text{in}}](X)}{B} \right), \quad (X, \tilde{V}) \in \mathbb{R}^2 \times \mathbb{R}^2.$$

5 Our paper is organized as follows. In Section 2 we derive formally the effective Vlasov-  
 6 Poisson equations. We introduce the average operators along the fast dynamics and  
 7 discuss their main properties. The effective model is studied in Section 3, and we sum-  
 8 marize its properties. The well posedness of this model is a direct consequence of the  
 9 well posedness of the vorticity formulation for the 2D incompressible Euler equations.  
 10 Section 4 is devoted to the error analysis. We establish second order estimates, by  
 11 constructing a suitable corrector on the extended phase space, supplemented by the  
 12 fast time variable. Some generalizations are indicated in the last section.

## 13 2 Asymptotic analysis by formal arguments

14 The well posedness of the Vlasov-Poisson problem is well known, see [1] for weak  
 15 solutions, and [26, 21, 23, 2] for strong solutions. Essentially the same arguments  
 16 provide the global existence and uniqueness for the strong solution of the Vlasov-  
 17 Poisson problem with external magnetic field cf. Theorem 2.1 [9].

### Theorem 2.1

Consider a non negative, smooth, compactly supported initial particle density  $f_{\text{in}} \in C_c^1(\mathbb{R}^2 \times \mathbb{R}^2)$  and a smooth magnetic field  $B \in C_b^1(\mathbb{R}^2)$ . There is a unique particle density  $f \in C^1(\mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2)$ , whose restriction on  $[0, T] \times \mathbb{R}^2 \times \mathbb{R}^2$  is compactly supported for any  $T \in \mathbb{R}_+$ , whose Poisson electric field is smooth  $E[f] \in C^1(\mathbb{R}_+ \times \mathbb{R}^2)$ , satisfying

$$\partial_t f + v \cdot \nabla_x f + \frac{q}{m} (E[f(t)] + B \perp v) \cdot \nabla_v f = 0, \quad (t, x, v) \in \mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2$$

$$E[f(t)](x) = \frac{q}{2\pi\varepsilon_0} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} f(t, x', v') \frac{x - x'}{|x - x'|^2} dv' dx', \quad (t, x) \in \mathbb{R}_+ \times \mathbb{R}^2$$

$$f(0, x, v) = f_{\text{in}}(x, v), \quad (x, v) \in \mathbb{R}^2 \times \mathbb{R}^2.$$

18 Moreover, if for some integer  $k \geq 2$  we have  $f_{\text{in}} \in C_c^k(\mathbb{R}^2 \times \mathbb{R}^2)$ ,  $B \in C_b^k(\mathbb{R}^2)$ , then  
 19  $f \in C^k(\mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2)$  and  $E[f] \in C^k(\mathbb{R}_+ \times \mathbb{R}^2)$ .

We assume that the initial particle density is smooth

$$f_{\text{in}} \geq 0, \quad f_{\text{in}} \in C_c^2(\mathbb{R}^2 \times \mathbb{R}^2)$$

and that the magnetic field is uniform  $B^\varepsilon = B/\varepsilon \neq 0$ . We know by Theorem 2.1 that for every  $\varepsilon > 0$ , there is a unique strong solution  $f^\varepsilon \in C^2(\mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2)$  (whose restriction on  $[0, T] \times \mathbb{R}^2 \times \mathbb{R}^2$  is compactly supported for any  $T \in \mathbb{R}_+$ ),  $E^\varepsilon := E[f^\varepsilon] \in C^2(\mathbb{R}_+ \times \mathbb{R}^2)$  for the Vlasov-Poisson problem with external magnetic field  $B^\varepsilon = B/\varepsilon$ . By standard arguments, we also have uniform estimates with respect to  $\varepsilon > 0$  for the  $L^\infty$  norm of the electric field  $E^\varepsilon|_{[0, T] \times \mathbb{R}^2}$  and the size of the support of the particle density  $f^\varepsilon|_{[0, T] \times \mathbb{R}^2 \times \mathbb{R}^2}$ . Let us denote by  $(X^\varepsilon, V^\varepsilon)(t; t_0, x, v)$  the characteristics associated to (1)

$$\frac{dX^\varepsilon}{dt} = V^\varepsilon(t; t_0, x, v), \quad \frac{dV^\varepsilon}{dt} = \frac{q}{m} \left[ E^\varepsilon(t, X^\varepsilon(t; t_0, x, v)) + \frac{B}{\varepsilon} \perp V^\varepsilon(t; t_0, x, v) \right]$$

1

$$X^\varepsilon(t_0; t_0, x, v) = x, \quad V^\varepsilon(t_0; t_0, x, v) = v.$$

2 The strong external magnetic field induces a large cyclotronic frequency with respect  
3 to the reciprocal advection time scale, and therefore a fast dynamics. Indeed, by  
4 introducing the characteristic scales  $(\bar{t}, \bar{x}, \bar{v})$  for time, length, velocity, we have  $\bar{t} = \bar{x}/\bar{v}$   
5 and  $\omega_c \bar{t} \sim 1/\varepsilon$ . We use the notations  $\omega_c^\varepsilon = qB^\varepsilon/m = \omega_c/\varepsilon, \omega_c = qB/m \sim 1/\bar{t}$ .  
6 It is well known that the guiding center,  $X^\varepsilon(t) + \varepsilon \perp V^\varepsilon(t)/\omega_c$  has small variations  
7 over one cyclotronic period [14, 22, 25]. Another quantity having small variations is  
8  $\mathcal{R}(\omega_c t/\varepsilon) [V^\varepsilon(t) - \varepsilon \frac{\perp E^\varepsilon(t, X^\varepsilon(t))}{B}]$ , see [20].

9 **Lemma 2.1**

10 *The following quantities*

$$X^\varepsilon(t) + \varepsilon \perp V^\varepsilon(t)/\omega_c, \quad \mathcal{R}(\omega_c t/\varepsilon) \left[ V^\varepsilon(t) - \varepsilon \frac{\perp E^\varepsilon(t, X^\varepsilon(t))}{B} \right]$$

11 *have small variations in time.*

**Proof.**

By direct computations we have

$$\frac{d}{dt} \left[ X^\varepsilon(t) + \varepsilon \frac{\perp V^\varepsilon(t)}{\omega_c} \right] = \varepsilon \frac{\perp E^\varepsilon(t, X^\varepsilon(t))}{B}.$$

Notice also that

$$\frac{d}{dt} \left[ V^\varepsilon(t) - \varepsilon \frac{\perp E^\varepsilon(t, X^\varepsilon(t))}{B} \right] = \frac{\omega_c}{\varepsilon} \perp \left[ V^\varepsilon(t) - \varepsilon \frac{\perp E^\varepsilon(t, X^\varepsilon(t))}{B} \right] - \varepsilon \frac{d}{dt} \left[ \frac{\perp E^\varepsilon(t, X^\varepsilon(t))}{B} \right]$$

and therefore

$$\frac{d}{dt} \left\{ \mathcal{R}(\omega_c t/\varepsilon) \left[ V^\varepsilon(t) - \varepsilon \frac{\perp E^\varepsilon(t, X^\varepsilon(t))}{B} \right] \right\} = -\varepsilon \mathcal{R}(\omega_c t/\varepsilon) \frac{d}{dt} \left[ \frac{\perp E^\varepsilon(t, X^\varepsilon(t))}{B} \right].$$

12

□

1 Motivated by the above calculation, we introduce the relative velocity with respect to  
 2 the electric cross field drift

$$\tilde{v} = v - \varepsilon \frac{{}^\perp E^\varepsilon(t, x)}{B} \quad (5)$$

3 and the new particle density

$$\tilde{f}^\varepsilon(t, x, \tilde{v}) = f^\varepsilon \left( t, x, \tilde{v} + \varepsilon \frac{{}^\perp E[f^\varepsilon(t)](x)}{B} \right), \quad (x, \tilde{v}) \in \mathbb{R}^2 \times \mathbb{R}^2. \quad (6)$$

4 It is easily seen that

$$\rho[\tilde{f}^\varepsilon(t)] = q \int_{\mathbb{R}^2} \tilde{f}^\varepsilon(t, \cdot, \tilde{v}) \, d\tilde{v} = q \int_{\mathbb{R}^2} f^\varepsilon(t, \cdot, v) \, dv = \rho[f^\varepsilon(t)], \quad t \in \mathbb{R}_+$$

5 and

$$\begin{aligned} j[f^\varepsilon(t)] &= q \int_{\mathbb{R}^2} f^\varepsilon(t, \cdot, v) v \, dv = q \int_{\mathbb{R}^2} \tilde{f}^\varepsilon(t, \cdot, \tilde{v}) \left( \tilde{v} + \varepsilon \frac{{}^\perp E^\varepsilon(t)}{B} \right) \, d\tilde{v} \\ &= j[\tilde{f}^\varepsilon(t)] + \varepsilon \frac{{}^\perp E^\varepsilon(t)}{B} \rho[\tilde{f}^\varepsilon(t)], \quad t \in \mathbb{R}_+. \end{aligned}$$

6 Therefore the Poisson electric fields corresponding to the particle densities  $f^\varepsilon, \tilde{f}^\varepsilon$  coin-  
 7 cide

$$E[f^\varepsilon(t)] = E[\tilde{f}^\varepsilon(t)], \quad t \in \mathbb{R}_+$$

and we can use the same notation  $E^\varepsilon(t)$  for denoting them. We introduce the notations

$$B_0 := |B| > 0, \quad \omega_0 := |\omega_c| > 0.$$

Observe that the new particle densities  $(\tilde{f}^\varepsilon)_{\varepsilon>0}$  are smooth,  $\tilde{f}^\varepsilon \in C^2(\mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2)$  and that the restrictions to  $[0, T] \times \mathbb{R}^2 \times \mathbb{R}^2$  are compactly supported, uniformly with respect to  $\varepsilon \in ]0, 1]$ , for any  $T \in \mathbb{R}_+$ . We obtain the following problem in the new coordinates  $(x, \tilde{v})$

$$\begin{aligned} \partial_t \tilde{f}^\varepsilon + \left( \tilde{v} + \varepsilon \frac{{}^\perp E^\varepsilon}{B} \right) \cdot \nabla_x \tilde{f}^\varepsilon - \varepsilon \left[ \partial_t \left( \frac{{}^\perp E^\varepsilon}{B} \right) + \partial_x \left( \frac{{}^\perp E^\varepsilon}{B} \right) \left( \tilde{v} + \varepsilon \frac{{}^\perp E^\varepsilon}{B} \right) \right] \cdot \nabla_{\tilde{v}} \tilde{f}^\varepsilon \\ + \frac{\omega_c}{\varepsilon} {}^\perp \tilde{v} \cdot \nabla_{\tilde{v}} \tilde{f}^\varepsilon = 0, \quad (t, x, \tilde{v}) \in \mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2 \end{aligned}$$

$$\tilde{f}^\varepsilon(0, x, \tilde{v}) = f_{\text{in}} \left( x, \tilde{v} + \varepsilon \frac{{}^\perp E[f_{\text{in}}](x)}{B} \right), \quad (x, \tilde{v}) \in \mathbb{R}^2 \times \mathbb{R}^2.$$

8 Thanks to the continuity equation

$$\partial_t \rho[f^\varepsilon] + \text{div}_x j[f^\varepsilon] = 0$$

the time derivative of the electric field  $E^\varepsilon$  can be written in terms of the particle density  $f^\varepsilon$  (or  $\tilde{f}^\varepsilon$ )

$$\begin{aligned} \partial_t E[f^\varepsilon] &= \frac{1}{2\pi\varepsilon_0} \int_{\mathbb{R}^2} \partial_t \rho[f^\varepsilon(t)](x - x') \frac{x'}{|x'|^2} \, dx' \\ &= -\frac{1}{2\pi\varepsilon_0} \int_{\mathbb{R}^2} \text{div}_x j[f^\varepsilon](x - x') \frac{x'}{|x'|^2} \, dx' \\ &= -\frac{1}{2\pi\varepsilon_0} \text{div}_x \int_{\mathbb{R}^2} \frac{x'}{|x'|^2} \otimes j[f^\varepsilon(t)](x - x') \, dx' \\ &= -\frac{1}{2\pi\varepsilon_0} \text{div}_x \int_{\mathbb{R}^2} \frac{x - x'}{|x - x'|^2} \otimes \left( j[\tilde{f}^\varepsilon(t)](x') + \varepsilon \frac{{}^\perp E^\varepsilon(t, x')}{B} \rho[\tilde{f}^\varepsilon(t)](x') \right) \, dx'. \end{aligned}$$

1 The change of coordinates (5), (6) leads to the problem

$$\partial_t \tilde{f}^\varepsilon + \varepsilon a^\varepsilon[\tilde{f}^\varepsilon(t)] \cdot \nabla_{x, \tilde{v}} \tilde{f}^\varepsilon + \frac{b^\varepsilon(x, \tilde{v})}{\varepsilon} \cdot \nabla_{x, \tilde{v}} \tilde{f}^\varepsilon = 0, \quad (t, x, \tilde{v}) \in \mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2 \quad (7)$$

$$\tilde{f}^\varepsilon(0, x, \tilde{v}) = f_{\text{in}} \left( x, \tilde{v} + \varepsilon \frac{^\perp E[f_{\text{in}}](x)}{B} \right), \quad (x, \tilde{v}) \in \mathbb{R}^2 \times \mathbb{R}^2$$

where  $b^\varepsilon \cdot \nabla_{x, \tilde{v}} = \varepsilon \tilde{v} \cdot \nabla_x + \omega_c \cdot \nabla_{\tilde{v}}$  and for any particle density  $\tilde{f}$ ,  $a^\varepsilon[\tilde{f}] \cdot \nabla_{x, \tilde{v}}$  stands for the vector field

$$\begin{aligned} a^\varepsilon[\tilde{f}] \cdot \nabla_{x, \tilde{v}} &= \frac{^\perp E[\tilde{f}]}{B} \cdot \nabla_x + \frac{1}{2\pi\varepsilon_0 B} \operatorname{div}_x \int_{\mathbb{R}^2} \frac{^\perp(x-x')}{|x-x'|^2} \otimes \left( j[\tilde{f}] + \varepsilon \frac{^\perp E[\tilde{f}]}{B} \rho[\tilde{f}] \right) (x') dx' \cdot \nabla_{\tilde{v}} \\ &\quad - \partial_x \left( \frac{^\perp E[\tilde{f}]}{B} \right) \left( \tilde{v} + \varepsilon \frac{^\perp E[\tilde{f}]}{B} \right) \cdot \nabla_{\tilde{v}}. \end{aligned}$$

2 Notice that the vector fields  $a^\varepsilon[\tilde{f}] \cdot \nabla_{x, \tilde{v}}$  and  $b^\varepsilon \cdot \nabla_{x, \tilde{v}}$  are divergence free. The charac-  
3 teristic flow of the vector field  $b^\varepsilon \cdot \nabla_{x, \tilde{v}} = \varepsilon \tilde{v} \cdot \nabla_x + \omega_c \cdot \nabla_{\tilde{v}}$

$$\frac{d\mathcal{X}^\varepsilon}{ds} = \varepsilon \tilde{\mathcal{V}}^\varepsilon(s; x, \tilde{v}), \quad \frac{d\tilde{\mathcal{V}}^\varepsilon}{ds} = \omega_c \cdot \tilde{\mathcal{V}}^\varepsilon(s; x, \tilde{v}), \quad \mathcal{X}^\varepsilon(0; x, \tilde{v}) = x, \quad \tilde{\mathcal{V}}^\varepsilon(0; x, \tilde{v}) = \tilde{v}$$

4 is given by

$$\mathcal{X}^\varepsilon(s; x, \tilde{v}) = x + \varepsilon [I_2 - \mathcal{R}(-\omega_c s)] \frac{^\perp \tilde{v}}{\omega_c}, \quad \tilde{\mathcal{V}}^\varepsilon(s; x, \tilde{v}) = \mathcal{R}(-\omega_c s) \tilde{v}, \quad \omega_c = \frac{qB}{m}. \quad (8)$$

5 It is periodic, and has the same period as the characteristic flow ( $\mathcal{X}(s; x, \tilde{v}) = x, \tilde{\mathcal{V}}(s; x, \tilde{v}) =$   
6  $\mathcal{R}(-s\omega_c) \tilde{v}$ ) of the vector field  $b(x, \tilde{v}) \cdot \nabla_{x, \tilde{v}} = \omega_c \cdot \nabla_{\tilde{v}}$

$$S^\varepsilon(x, \tilde{v}) = S(x, \tilde{v}) = S, \quad S := \frac{2\pi}{|\omega_c|}, \quad (x, \tilde{v}) \in \mathbb{R}^2 \times \mathbb{R}^2, \quad \varepsilon > 0.$$

7 The properties of these flows are summarized below. The proof details are left to the  
8 reader (see also Proposition 3.1 [9]).

9 **Proposition 2.1**

10 We denote by  $(\mathcal{X}^\varepsilon(s; x, \tilde{v}), \tilde{\mathcal{V}}^\varepsilon(s; x, \tilde{v}))$  the characteristic flow of the autonomous vector  
11 field  $b^\varepsilon(x, \tilde{v}) \cdot \nabla_{x, \tilde{v}}$

$$\frac{d\mathcal{X}^\varepsilon}{ds} = \varepsilon \tilde{\mathcal{V}}^\varepsilon(s; x, \tilde{v}), \quad \frac{d\tilde{\mathcal{V}}^\varepsilon}{ds} = \omega_c \cdot \tilde{\mathcal{V}}^\varepsilon(s; x, \tilde{v}), \quad \mathcal{X}^\varepsilon(0; x, \tilde{v}) = x, \quad \tilde{\mathcal{V}}^\varepsilon(0; x, \tilde{v}) = \tilde{v}$$

12 and by  $(\mathcal{X}(s; x, \tilde{v}), \tilde{\mathcal{V}}(s; x, \tilde{v}))$  the characteristic flow of the autonomous vector field  
13  $b(x, \tilde{v}) \cdot \nabla_{x, \tilde{v}} = \omega_c \cdot \nabla_{\tilde{v}}$

$$\frac{d\mathcal{X}}{ds} = 0, \quad \frac{d\tilde{\mathcal{V}}}{ds} = \omega_c \cdot \tilde{\mathcal{V}}(s; x, \tilde{v}), \quad \mathcal{X}(0; x, \tilde{v}) = x, \quad \tilde{\mathcal{V}}(0; x, \tilde{v}) = \tilde{v}.$$

14 1. For any  $(x, \tilde{v}) \in \mathbb{R}^2 \times \mathbb{R}^2$  and  $\varepsilon > 0$ , the characteristic  $s \rightarrow (\mathcal{X}^\varepsilon, \tilde{\mathcal{V}}^\varepsilon)(s; x, \tilde{v})$  is  
15  $S$ -periodic, with  $S = 2\pi/\omega_0$ .

1 2. For any  $(x, \tilde{v}) \in \mathbb{R}^2 \times \mathbb{R}^2$  and  $\varepsilon > 0$  we have

$$|\mathcal{X}^\varepsilon(s; x, \tilde{v}) - \mathcal{X}(s; x, \tilde{v})| = |\mathcal{X}^\varepsilon(s; x, \tilde{v}) - x| \leq \varepsilon \frac{2}{\omega_0} |\tilde{v}|, \quad s \in \mathbb{R}$$

2 and

$$\tilde{\mathcal{V}}^\varepsilon(s; x, \tilde{v}) = \tilde{\mathcal{V}}(s; x, \tilde{v}) = \mathcal{R}(-s\omega_c)\tilde{v}, \quad s \in \mathbb{R}.$$

3 3. For any continuous function  $u \in C(\mathbb{R}^2 \times \mathbb{R}^2)$  we define the averages along the  
4 flows of  $b \cdot \nabla_{x, \tilde{v}}$ ,  $b^\varepsilon \cdot \nabla_{x, \tilde{v}}$

$$\langle u \rangle(x, \tilde{v}) = \frac{1}{S} \int_0^S u(\mathcal{X}(s; x, \tilde{v}), \tilde{\mathcal{V}}(s; x, \tilde{v})) \, ds, \quad (x, \tilde{v}) \in \mathbb{R}^2 \times \mathbb{R}^2$$

5

$$\langle u \rangle_\varepsilon(x, \tilde{v}) = \frac{1}{S} \int_0^S u(\mathcal{X}^\varepsilon(s; x, \tilde{v}), \tilde{\mathcal{V}}^\varepsilon(s; x, \tilde{v})) \, ds, \quad (x, \tilde{v}) \in \mathbb{R}^2 \times \mathbb{R}^2.$$

6 For any  $R_x, R_{\tilde{v}} \in \mathbb{R}_+$  we have

$$\|\langle u \rangle\|_{L^\infty(B(R_x) \times B(R_{\tilde{v}}))} \leq \|u\|_{L^\infty(B(R_x) \times B(R_{\tilde{v}}))}$$

7

$$\|\langle u \rangle_\varepsilon\|_{L^\infty(B(R_x) \times B(R_{\tilde{v}}))} \leq \|u\|_{L^\infty(B(R_x^\varepsilon) \times B(R_{\tilde{v}}))}, \quad R_x^\varepsilon = R_x + 2\varepsilon R_{\tilde{v}}/\omega_0$$

8

where  $B(R)$  stands for the closed ball of radius  $R$  in  $\mathbb{R}^2$ .

4. If  $u$  is Lipschitz continuous, then for any  $(x, \tilde{v}) \in \mathbb{R}^2 \times \mathbb{R}^2$  and  $\varepsilon > 0$  we have

$$\frac{|\langle u \rangle_\varepsilon(x, \tilde{v}) - \langle u \rangle(x, \tilde{v})|}{\varepsilon} \leq \text{Lip}(u) \frac{2}{\omega_0} |\tilde{v}|.$$

9

5. For any function  $u \in C_c^1(\mathbb{R}^2 \times \mathbb{R}^2)$  we have the inequality

$$\|u - \langle u \rangle\|_{L^2(\mathbb{R}^2 \times \mathbb{R}^2)} \leq \frac{2\pi}{\omega_0} \|b \cdot \nabla_{x, \tilde{v}} u\|_{L^2(\mathbb{R}^2 \times \mathbb{R}^2)}.$$

10

6. For any function  $u \in C^1(\mathbb{R}^2 \times \mathbb{R}^2)$ , we have  $\langle u \rangle \in C^1(\mathbb{R}^2 \times \mathbb{R}^2)$  and

$$\langle \nabla_x u \rangle = \nabla_x \langle u \rangle, \quad \tilde{v} \cdot \nabla_{\tilde{v}} \langle u \rangle = \langle \tilde{v} \cdot \nabla_{\tilde{v}} u \rangle, \quad {}^\perp \tilde{v} \cdot \nabla_{\tilde{v}} \langle u \rangle = 0.$$

11 In order to filter out the fast oscillations corresponding to the vector field  $\frac{b^\varepsilon(x, \tilde{v})}{\varepsilon} \cdot \nabla_{x, \tilde{v}}$ ,  
12 we perform one more change of coordinates

$$\tilde{f}^\varepsilon(t, x, \tilde{v}) = \tilde{F}^\varepsilon(t, X, \tilde{V}), \quad (X, \tilde{V}) = (\mathcal{X}^\varepsilon, \tilde{\mathcal{V}}^\varepsilon)(-t/\varepsilon; x, \tilde{v}). \quad (9)$$

13 By applying the chain rule, we obtain

$$\partial_t \tilde{F}^\varepsilon + \varepsilon \partial_{x, \tilde{v}} (\mathcal{X}^\varepsilon, \tilde{\mathcal{V}}^\varepsilon)(-t/\varepsilon; (\mathcal{X}^\varepsilon, \tilde{\mathcal{V}}^\varepsilon)(t/\varepsilon; X, \tilde{V})) a^\varepsilon[\tilde{F}^\varepsilon(t)_{-t/\varepsilon}]_{t/\varepsilon} \cdot \nabla_{X, \tilde{V}} \tilde{F}^\varepsilon = 0 \quad (10)$$

14 where we have used the notations

$$\tilde{F}^\varepsilon(t)_{-t/\varepsilon} = \tilde{F}^\varepsilon(t) \circ (\mathcal{X}^\varepsilon, \tilde{\mathcal{V}}^\varepsilon)\left(-\frac{t}{\varepsilon}\right), \quad a^\varepsilon[\tilde{F}^\varepsilon(t)_{-t/\varepsilon}]_{t/\varepsilon} = a^\varepsilon[\tilde{F}^\varepsilon(t) \circ (\mathcal{X}^\varepsilon, \tilde{\mathcal{V}}^\varepsilon)\left(-\frac{t}{\varepsilon}\right)] \circ (\mathcal{X}^\varepsilon, \tilde{\mathcal{V}}^\varepsilon)\left(\frac{t}{\varepsilon}\right).$$

- 1 As the characteristic flow  $(\mathcal{X}^\varepsilon, \tilde{V}^\varepsilon)$  in (8) is linear, the jacobian matrix simply writes  
 2 for any  $(x, \tilde{v}) \in \mathbb{R}^2 \times \mathbb{R}^2$

$$\partial_{x, \tilde{v}}(\mathcal{X}^\varepsilon, \tilde{V}^\varepsilon)(-t/\varepsilon; x, \tilde{v}) = \begin{pmatrix} I_2 & \frac{\varepsilon}{\omega_c} \mathcal{R}(-\pi/2)[I_2 - \mathcal{R}(\omega_c t/\varepsilon)] \\ O_2 & \mathcal{R}(\omega_c t/\varepsilon) \end{pmatrix}$$

and therefore (10) becomes

$$\begin{aligned} \partial_t \tilde{F}^\varepsilon + \varepsilon \left\{ a_x^\varepsilon[\tilde{F}^\varepsilon(t)_{-t/\varepsilon}]_{t/\varepsilon} + \frac{\varepsilon}{\omega_c} \mathcal{R}(-\pi/2)[I_2 - \mathcal{R}(\omega_c t/\varepsilon)] a_{\tilde{v}}^\varepsilon[\tilde{F}^\varepsilon(t)_{-t/\varepsilon}]_{t/\varepsilon} \right\} \cdot \nabla_X \tilde{F}^\varepsilon \\ + \varepsilon \mathcal{R}(\omega_c t/\varepsilon) a_{\tilde{v}}^\varepsilon[\tilde{F}^\varepsilon(t)_{-t/\varepsilon}]_{t/\varepsilon} \cdot \nabla_{\tilde{V}} \tilde{F}^\varepsilon = 0, \quad (t, X, \tilde{V}) \in \mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2. \end{aligned}$$

- 3 We have obtained a two scale problem and we expect that the asymptotic behavior  
 4 when  $\varepsilon$  becomes small will follow by averaging with respect to the fast time variable  
 5  $s = t/\varepsilon$ . As we are looking for second order approximations, we only need to average,  
 6 with respect to  $s$ , when  $\varepsilon$  is small

$$a_x^\varepsilon[\tilde{F}^\varepsilon(t)_{-s}]_s \cdot \nabla_X \tilde{F}^\varepsilon \quad \text{and} \quad \mathcal{R}(\omega_c s) a_{\tilde{v}}^\varepsilon[\tilde{F}^\varepsilon(t)_{-s}]_s \cdot \nabla_{\tilde{V}} \tilde{F}^\varepsilon$$

up to terms of order  $\varepsilon$ . By the second statement in Proposition 2.1, observe that

$$\begin{aligned} E[\tilde{F}^\varepsilon(t)_{-s}]_s(X, \tilde{V}) &= \frac{q}{2\pi\varepsilon_0} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \tilde{F}^\varepsilon(t, (\mathcal{X}^\varepsilon, \tilde{V}^\varepsilon)(-s; x', \tilde{v}')) \frac{\mathcal{X}^\varepsilon(s; X, \tilde{V}) - x'}{|\mathcal{X}^\varepsilon(s; X, \tilde{V}) - x'|^2} d\tilde{v}' dx' \\ &= \frac{q}{2\pi\varepsilon_0} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \tilde{F}^\varepsilon(t, X', \tilde{V}') \frac{\mathcal{X}^\varepsilon(s; X, \tilde{V}) - \mathcal{X}^\varepsilon(s; X', \tilde{V}')}{|\mathcal{X}^\varepsilon(s; X, \tilde{V}) - \mathcal{X}^\varepsilon(s; X', \tilde{V}')|^2} d\tilde{V}' dX' \\ &= E[\tilde{F}^\varepsilon(t)](X) + \mathcal{O}(\varepsilon) \end{aligned} \quad (11)$$

- 7 and therefore we deduce that

$$\frac{\omega_c}{2\pi} \int_0^{2\pi/\omega_c} a_x^\varepsilon[\tilde{F}^\varepsilon(t)_{-s}]_s \cdot \nabla_X \tilde{F}^\varepsilon(t) ds = \frac{\perp E[\tilde{F}^\varepsilon(t)](X)}{B} \cdot \nabla_X \tilde{F}^\varepsilon + \mathcal{O}(\varepsilon). \quad (12)$$

We concentrate now on the average of  $\mathcal{R}(\omega_c s) a_{\tilde{v}}^\varepsilon[\tilde{F}^\varepsilon(t)_{-s}]_s \cdot \nabla_{\tilde{V}} \tilde{F}^\varepsilon$ . We only need to consider the contributions of order 1 in  $a_{\tilde{v}}^\varepsilon[\tilde{F}^\varepsilon(t)_{-s}]_s$ , that is, those of

$$\begin{aligned} \frac{q}{2\pi\varepsilon_0 B} \left( \operatorname{div}_x \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{\perp(\cdot - x')}{|\cdot - x'|^2} \otimes \tilde{v}' \tilde{F}^\varepsilon(t, (\mathcal{X}^\varepsilon, \tilde{V}^\varepsilon)(-s; x', \tilde{v}')) d\tilde{v}' dx' \right) (\mathcal{X}^\varepsilon(s; X, \tilde{V})) \\ = \frac{q}{2\pi\varepsilon_0 B} \operatorname{div}_X \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{\perp(\mathcal{X}^\varepsilon(s; X, \tilde{V}) - \mathcal{X}^\varepsilon(s; X', \tilde{V}'))}{|\mathcal{X}^\varepsilon(s; X, \tilde{V}) - \mathcal{X}^\varepsilon(s; X', \tilde{V}')|^2} \otimes \mathcal{R}(-\omega_c s) \tilde{V}' \tilde{F}^\varepsilon(t, X', \tilde{V}') d\tilde{V}' dX' \end{aligned}$$

- 8 and

$$-\left( \partial_x \left( \frac{\perp E[\tilde{F}^\varepsilon(t)_{-s}]}{B} \right) \tilde{v} \right)_s (X, \tilde{V}) = \frac{\mathcal{R}(\pi/2)}{B} \partial_X E[\tilde{F}^\varepsilon(t)_{-s}]_s \mathcal{R}(-\omega_c s) \tilde{V}.$$

Notice that, up to terms of order  $\varepsilon$ , the average of the first contribution writes

$$\begin{aligned}
& \frac{\omega_c}{2\pi} \int_0^{\frac{2\pi}{\omega_c}} \mathcal{R}(\omega_c s) \frac{q}{2\pi\varepsilon_0 B} \operatorname{div}_X \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{\perp(X - X')}{|X - X'|^2} \otimes \mathcal{R}(-\omega_c s) \tilde{V}' \tilde{F}^\varepsilon(t, X', \tilde{V}') d\tilde{V}' dX' ds \\
&= \frac{q\mathcal{R}(-\pi/2)}{2\pi\varepsilon_0 B} \operatorname{div}_X \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{\omega_c}{2\pi} \int_0^{\frac{2\pi}{\omega_c}} \frac{\mathcal{R}(\omega_c s)(X - X')}{|X - X'|^2} \otimes \mathcal{R}(-\omega_c s) \tilde{V}' ds \tilde{F}^\varepsilon(t, X', \tilde{V}') d\tilde{V}' dX' \\
&= \frac{q\mathcal{R}(-\pi/2)}{2\pi\varepsilon_0 B} \operatorname{div}_X \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{(X - X') \otimes \tilde{V}' - \perp(X - X') \otimes \perp\tilde{V}'}{2|X - X'|^2} \tilde{F}^\varepsilon(t, X', \tilde{V}') d\tilde{V}' dX' \\
&= \frac{q\mathcal{R}(-\pi/2)}{2\pi\varepsilon_0 B} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{(X - X') \otimes \tilde{V}' - \perp(X - X') \otimes \perp\tilde{V}'}{2|X - X'|^2} \nabla_{X'} \tilde{F}^\varepsilon(t, X', \tilde{V}') d\tilde{V}' dX'.
\end{aligned}$$

- 1 By using the properties of the fundamental solution  $z \rightarrow -\frac{1}{2\pi} \ln|z|$ , it is easily seen  
2 that

$$\operatorname{div}_{X'} \frac{(X - X') \otimes \tilde{V}' - \perp(X - X') \otimes \perp\tilde{V}'}{2|X - X'|^2} = 0, \quad X \neq X'$$

and thus the above computations lead to

$$\begin{aligned}
& \frac{q\mathcal{R}(-\pi/2)}{2\pi\varepsilon_0 B} \lim_{r \searrow 0} \int_{\mathbb{R}^2} \int_{|X - X'| > r} \frac{(X - X') \otimes \tilde{V}' - \perp(X - X') \otimes \perp\tilde{V}'}{2|X - X'|^2} \nabla_{X'} \tilde{F}^\varepsilon(t, X', \tilde{V}') dX' d\tilde{V}' \\
&= \frac{q\mathcal{R}(-\pi/2)}{2\pi\varepsilon_0 B} \lim_{r \searrow 0} \int_{\mathbb{R}^2} \int_{|X - X'| = r} \tilde{F}^\varepsilon \frac{(X - X') \otimes \tilde{V}' - \perp(X - X') \otimes \perp\tilde{V}'}{2|X - X'|^2} \frac{X - X'}{|X - X'|} d\sigma(X') d\tilde{V}' \\
&= \frac{q\mathcal{R}(-\pi/2)}{2\pi\varepsilon_0 B} \lim_{r \searrow 0} \frac{1}{2r} \int_{|X - X'| = r} \int_{\mathbb{R}^2} \tilde{F}^\varepsilon(t, X', \tilde{V}') \tilde{V}' d\tilde{V}' d\sigma(X') \\
&= \frac{\perp j[\tilde{F}^\varepsilon(t)](X)}{2\varepsilon_0 B}. \tag{13}
\end{aligned}$$

Similarly, up to terms of order  $\varepsilon$ , the second contribution is, thanks to (11)

$$\begin{aligned}
& \frac{\omega_c}{2\pi B} \int_0^{2\pi/\omega_c} \mathcal{R}(\pi/2) \mathcal{R}(\omega_c s) \partial_X E[\tilde{F}^\varepsilon(t)] \mathcal{R}(-\omega_c s) \tilde{V} ds = -\frac{\operatorname{trace}(\partial_X E[\tilde{F}^\varepsilon(t)]) \perp \tilde{V}}{2B} \\
&= -\frac{\operatorname{div}_X E[\tilde{F}^\varepsilon(t)] \perp \tilde{V}}{2B} = -\frac{\rho[\tilde{F}^\varepsilon(t)] \perp \tilde{V}}{2\varepsilon_0 B}. \tag{14}
\end{aligned}$$

Combining (13), (14) we obtain

$$\frac{\omega_c}{2\pi} \int_0^{2\pi/\omega_c} \mathcal{R}(\omega_c s) a_v^\varepsilon[\tilde{F}^\varepsilon(t) - s] \cdot \nabla_{\tilde{V}} \tilde{F}^\varepsilon ds = \frac{\perp \{j[\tilde{F}^\varepsilon(t)] - \rho[\tilde{F}^\varepsilon(t)] \tilde{V}\}}{2\varepsilon_0 B} \cdot \nabla_{\tilde{V}} \tilde{F}^\varepsilon + \mathcal{O}(\varepsilon). \tag{15}$$

- 3 Thanks to (12), (15) we are led to the model

$$\partial_t \tilde{F} + \varepsilon \frac{\perp E[\tilde{F}(t)]}{B} \cdot \nabla_X \tilde{F} + \varepsilon \frac{\perp \{j[\tilde{F}(t)] - \rho[\tilde{F}(t)] \tilde{V}\}}{2\varepsilon_0 B} \cdot \nabla_{\tilde{V}} \tilde{F} = 0, \quad (t, X, \tilde{V}) \in \mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2 \tag{16}$$

- 4 which is supplemented by the initial condition

$$\tilde{F}(0, X, \tilde{V}) = f_{\text{in}} \left( X - \varepsilon \frac{\perp \tilde{V}}{\omega_c}, \tilde{V} + \varepsilon \frac{\perp E[f_{\text{in}}](X)}{B} \right), \quad (X, \tilde{V}) \in \mathbb{R}^2 \times \mathbb{R}^2. \tag{17}$$

### 3 The effective Vlasov-Poisson equations

We expect that solving (16) together with the initial condition (17), will provide a second order approximation for (1), (2). Although the above solution depends on  $\varepsilon$ , we use the notation  $\tilde{F}$ , saying that it is an approximation, when  $\varepsilon$  becomes small. The well posedness of the limit model (16), (17) is a direct consequence of the well posedness of the vorticity formulation for the 2D incompressible Euler equations, see also Lemma 3.3 [25]. Indeed, integrating (16) with respect to the velocity leads to the Euler equations

$$\partial_t \rho[\tilde{F}] + \frac{\perp E}{B^\varepsilon} \cdot \nabla_X \rho[\tilde{F}] = 0, E(t, X) = \frac{1}{2\pi\varepsilon_0} \int_{\mathbb{R}^2} \rho[\tilde{F}(t)](X') \frac{X - X'}{|X - X'|^2} dX', (t, X) \in \mathbb{R}_+ \times \mathbb{R}^2$$

which allows us to determine  $\rho$  and  $E$ . Multiplying (16) by  $\tilde{V}$  and integrating with respect to the velocity give a transport equation for the current density as well

$$\partial_t j[\tilde{F}] + \left( \frac{\perp E}{B^\varepsilon} \cdot \nabla_X \right) j[\tilde{F}] = 0$$

and finally the particle density  $\tilde{F}$  comes by solving the linear transport equation (16), with smooth advection field

$$\frac{\perp E}{B^\varepsilon} \cdot \nabla_X + \frac{1}{2\varepsilon_0 B^\varepsilon} \perp (j - \rho \tilde{V}) \cdot \nabla_{\tilde{V}}.$$

The proof details are left to the reader.

**Theorem 3.1** *Consider a non negative, smooth, compactly supported initial particle density  $\tilde{F}_{\text{in}} \in C_c^1(\mathbb{R}^2 \times \mathbb{R}^2)$  and a uniform magnetic field  $B^\varepsilon = \frac{B}{\varepsilon} \neq 0$ . There is a unique particle density  $\tilde{F} \in C^1(\mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2)$  whose restriction on  $[0, T] \times \mathbb{R}^2 \times \mathbb{R}^2$  is compactly supported for any  $T \in \mathbb{R}_+$ , whose Poisson electric field is smooth  $E[\tilde{F}] \in C^1(\mathbb{R}_+ \times \mathbb{R}^2)$ , satisfying (16), (17). Moreover, if for some integer  $k \geq 2$  we have  $\tilde{F}_{\text{in}} \in C_c^k(\mathbb{R}^2 \times \mathbb{R}^2)$ , then  $\tilde{F} \in C^k(\mathbb{R}_+ \times \mathbb{R}^2 \times \mathbb{R}^2)$  and  $E[\tilde{F}] \in C^k(\mathbb{R}_+ \times \mathbb{R}^2)$ .*

The properties of the above limit model are summarized below : conservation of the total kinetic energy, conservation of the total electric energy, invariance of  $\frac{1}{2}|j[\tilde{F}] - \rho[\tilde{F}]\tilde{V}|^2$ , invariance under the rotations in the velocity space.

#### Remark 3.1

1. The total kinetic energy is conserved. Indeed, multiplying (16) by  $m|\tilde{V}|^2/2$  and integrating with respect to velocity yields

$$\begin{aligned} \partial_t \int_{\mathbb{R}^2} m \frac{|\tilde{V}|^2}{2} \tilde{F} d\tilde{V} + \frac{\perp E[\tilde{F}(t)]}{B^\varepsilon} \cdot \nabla_X \int_{\mathbb{R}^2} m \frac{|\tilde{V}|^2}{2} \tilde{F} d\tilde{V} \\ = \frac{m}{2\varepsilon_0 B^\varepsilon} \int_{\mathbb{R}^2} \perp (j[\tilde{F}(t)] - \rho[\tilde{F}(t)]\tilde{V}) \cdot \tilde{V} \tilde{F} d\tilde{V} = 0 \end{aligned}$$

and our conclusion follows by integrating also with respect to  $X$ .

2. The total electric energy is conserved. Using the fundamental solution of the Poisson equation in  $\mathbb{R}^2$ , we have

$$\begin{aligned}
& \frac{d}{dt} \frac{1}{2\varepsilon_0} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} e(X - X') \rho[\tilde{F}(t)](X) \rho[\tilde{F}(t)](X') \, dX' dX \\
&= \frac{1}{\varepsilon_0} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} e(X - X') \rho[\tilde{F}(t)](X') \partial_t \rho[\tilde{F}(t)](X) \, dX' dX \\
&= \int_{\mathbb{R}^2} \Phi[\tilde{F}(t)](X) \partial_t \rho[\tilde{F}(t)](X) \, dX - \int_{\mathbb{R}^2} \Phi[\tilde{F}(t)](X) \operatorname{div}_X \left( \rho[\tilde{F}(t)] \frac{\perp E[\tilde{F}(t)]}{B^\varepsilon} \right) \, dX \\
&= \int_{\mathbb{R}^2} \nabla_X \Phi[\tilde{F}(t)] \cdot \frac{\perp E[\tilde{F}(t)]}{B^\varepsilon} \rho[\tilde{F}(t)](X) \, dX = 0.
\end{aligned}$$

3. The function  $(X, \tilde{V}) \rightarrow \frac{1}{2} |j[\tilde{F}(t)](X) - \rho[\tilde{F}(t)](X) \tilde{V}|^2$  is an invariant of the transport operator in (16). Indeed, thanks to the mass and momentum balances, we deduce that

$$\begin{aligned}
& \left( \partial_t + \frac{\perp E[\tilde{F}(t)]}{B^\varepsilon} \cdot \nabla_X + \frac{\perp (j[\tilde{F}(t)] - \rho[\tilde{F}(t)] \tilde{V})}{2\varepsilon_0 B^\varepsilon} \cdot \nabla_{\tilde{V}} \right) \frac{|j[\tilde{F}(t)] - \rho[\tilde{F}(t)] \tilde{V}|^2}{2} \\
&= -\rho[\tilde{F}(t)] \frac{\perp (j[\tilde{F}(t)] - \rho[\tilde{F}(t)] \tilde{V})}{2\varepsilon_0 B^\varepsilon} \cdot (j[\tilde{F}(t)] - \rho[\tilde{F}(t)] \tilde{V}) = 0.
\end{aligned}$$

1 4. The model (16), (17) is invariant under rotation in the velocity space. More  
2 exactly, if  $\tilde{F}$  solves (16), (17), then  $\tilde{F}_\theta(t, X, \tilde{V}) = \tilde{F}(t, X, \mathcal{R}(\theta)\tilde{V})$  solves (16)  
3 together with the initial condition

$$\tilde{F}_\theta(0, X, \tilde{V}) = \tilde{F}_{\text{in}}(X, \mathcal{R}(\theta)\tilde{V}), \quad (X, \tilde{V}) \in \mathbb{R}^2 \times \mathbb{R}^2.$$

4 In particular, if the initial particle density satisfies  $b \cdot \nabla_{X, \tilde{V}} \tilde{F}_{\text{in}} = 0$ , then, thanks  
5 to the uniqueness, we have  $b \cdot \nabla_{X, \tilde{V}} \tilde{F}(t) = 0$  at any time  $t \in \mathbb{R}_+$ .

## 6 4 The error analysis

7 The solution of (16), (17) will allow us to describe the asymptotic behavior of the  
8 family  $(f^\varepsilon)_{\varepsilon>0}$  corresponding to the initial condition  $f_{\text{in}}$ , when  $\varepsilon$  becomes small. Let us  
9 introduce the particle density

$$\tilde{f}(t, s, x, \tilde{v}) = \tilde{F}(t, (\mathcal{X}, \tilde{\mathcal{V}})(-s; x, \tilde{v})), \quad (t, s, x, \tilde{v}) \in \mathbb{R}_+ \times \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2$$

10 where  $\tilde{F}$  solves (16), (17) and  $(\mathcal{X}, \tilde{\mathcal{V}})$  is the characteristic flow associated to the vector  
11 field  $b(x, \tilde{v}) \cdot \nabla_{x, \tilde{v}} = \omega_c \perp \tilde{v} \cdot \nabla_{\tilde{v}}$ , see Proposition 2.1. The idea is to compare  $\tilde{f}^\varepsilon(t, x, \tilde{v})$  with  
12 respect to  $\bar{f}(t, x + \varepsilon \perp \tilde{v} / \omega_c, \tilde{v})$ , where  $\bar{f}(t, x, \tilde{v}) = \tilde{f}(t, t/\varepsilon, x, \tilde{v})$ . By direct computation  
13 we check that

$$\partial_t \bar{f} + \varepsilon \left( \frac{\perp E[\bar{f}(t)]}{B} \cdot \nabla_x \bar{f} + \frac{\perp (j[\bar{f}(t)] - \rho[\bar{f}(t)] \tilde{v})}{2\varepsilon_0 B} \cdot \nabla_{\tilde{v}} \bar{f} \right) + \frac{b(x, \tilde{v})}{\varepsilon} \cdot \nabla_{x, \tilde{v}} \bar{f} = 0. \quad (18)$$

We introduce the application  $T^\varepsilon(x, \tilde{v}) = (x + \varepsilon \frac{\perp \tilde{v}}{\omega_c}, \tilde{v})$ . Notice that  $\bar{f}(0) \circ T^\varepsilon$  is a second order approximation of  $\tilde{f}^\varepsilon(0)$  in  $L^2(\mathbb{R}^2 \times \mathbb{R}^2)$

$$\begin{aligned}
& \left\{ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left[ \tilde{f}^\varepsilon(0, x, \tilde{v}) - \bar{f} \left( 0, x + \varepsilon \frac{\perp \tilde{v}}{\omega_c}, \tilde{v} \right) \right]^2 d\tilde{v} dx \right\}^{1/2} \\
&= \left\{ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left[ f_{\text{in}} \left( x, \tilde{v} + \varepsilon \frac{\perp E[f_{\text{in}}]}{B}(x) \right) - \tilde{F} \left( 0, x + \varepsilon \frac{\perp \tilde{v}}{\omega_c}, \tilde{v} \right) \right]^2 d\tilde{v} dx \right\}^{1/2} \\
&= \left\{ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left[ f_{\text{in}} \left( x, \tilde{v} + \varepsilon \frac{\perp E[f_{\text{in}}]}{B}(x) \right) - f_{\text{in}} \left( x, \tilde{v} + \varepsilon \frac{\perp E[f_{\text{in}}]}{B}(x + \varepsilon \frac{\perp \tilde{v}}{\omega_c}) \right) \right]^2 d\tilde{v} dx \right\}^{1/2} \\
&\leq C\varepsilon^2.
\end{aligned} \tag{19}$$

1 By introducing a corrector term, we will prove that  $\bar{f} \circ T^\varepsilon$  is a second order approxi-  
2 mation of  $\tilde{f}^\varepsilon$  in  $L_{\text{loc}}^\infty(\mathbb{R}_+; L^2(\mathbb{R}^2 \times \mathbb{R}^2))$ . We mention that the asymptotic behavior for a  
3 very similar problem has been investigated in Theorem 1.2 [25], but without indicating  
4 the convergence rate. Our goal is to complete the asymptotic analysis by justifying  
5 the second order approximation. The arguments developed for well prepared initial  
6 particle densities in [9] apply for general initial particle densities as well, justifying the  
7 robustness of our method. For any smooth particle density  $\tilde{f} \in C_c^1(\mathbb{R}^2 \times \mathbb{R}^2)$  we use  
8 the notations

$$a[\tilde{f}] \cdot \nabla_{x, \tilde{v}} = \frac{\perp E[\tilde{f}]}{B} \cdot \nabla_x + \left[ \frac{\text{div}_x}{2\pi\varepsilon_0 B} \int_{\mathbb{R}^2} \frac{\perp(x-x')}{|x-x'|^2} \otimes j[\tilde{f]}(x') dx' - \partial_x \left( \frac{\perp E[\tilde{f}]}{B} \right) \tilde{v} \right] \cdot \nabla_{\tilde{v}}$$

9 and

$$\langle a \rangle [\tilde{f}] \cdot \nabla_{x, \tilde{v}} = \frac{\perp E[\tilde{f}]}{B} \cdot \nabla_x + \frac{\perp(j[\tilde{f}] - \rho[\tilde{f}]\tilde{v})}{2\varepsilon_0 B} \cdot \nabla_{\tilde{v}}.$$

10 For constructing the corrector, we need essentially to invert the transport operator  $\partial_s +$   
11  $b \cdot \nabla_{x, \tilde{v}}$  on the subspace of functions with zero average with respect to the characteristic  
12 flow of the vector field  $\partial_s + b \cdot \nabla_{x, \tilde{v}}$ , see [7, 8]. The expression of the corrector is explicit  
13 and follows by direct computations. Its smoothness and uniform boundedness with  
14 respect to the fast time variable will be crucial when establishing the error estimate.  
15 We consider particle densities depending also on the fast time variable and therefore  
16 we work in the phase space  $(s, x, \tilde{v}) \in \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2$ .

17 **Proposition 4.1** *Assume that  $\tilde{f} = \tilde{f}(s, x, \tilde{v}) \in C^1(\mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2)$  is  $S = \frac{2\pi}{|\omega_c|}$  periodic*  
18 *and uniformly compactly supported in  $(x, \tilde{v})$  with respect to  $s \in \mathbb{R}$ , such that*

$$\partial_s \tilde{f} + b(x, \tilde{v}) \cdot \nabla_{x, \tilde{v}} \tilde{f} = 0, \quad (s, x, \tilde{v}) \in \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2.$$

19 *Then we have the equality*

$$a[\tilde{f}(s)] \cdot \nabla_{x, \tilde{v}} \tilde{f} - \langle a \rangle [\tilde{f}(s)] \cdot \nabla_{x, \tilde{v}} \tilde{f} + (\partial_s + b(x, \tilde{v}) \cdot \nabla_{x, \tilde{v}}) \tilde{f}^2 = 0$$

where

$$\begin{aligned}
\tilde{f}^2 &= \frac{\cos(2\omega_c s)}{8\omega_c \pi \varepsilon_0 B} \operatorname{div}_x \int_{\mathbb{R}^2} \left[ \frac{\perp(x-x')}{|x-x'|^2} \otimes \perp j[\tilde{F}] + \frac{x-x'}{|x-x'|^2} \otimes j[\tilde{F}] \right] dx' \cdot \nabla_{\tilde{V}} \tilde{F}(x, \mathcal{R}(\omega_c s) \tilde{v}) \\
&+ \frac{\sin(2\omega_c s)}{8\omega_c \pi \varepsilon_0 B} \operatorname{div}_x \int_{\mathbb{R}^2} \left[ \frac{(x-x')}{|x-x'|^2} \otimes \perp j[\tilde{F}] - \frac{\perp(x-x')}{|x-x'|^2} \otimes j[\tilde{F}] \right] dx' \cdot \nabla_{\tilde{V}} \tilde{F}(x, \mathcal{R}(\omega_c s) \tilde{v}) \\
&+ \frac{\cos(2\omega_c s)}{4\omega_c} \left[ \mathcal{R}(\pi/2) \partial_x \left( \frac{E[\tilde{F}]}{B} \right) + \partial_x \left( \frac{E[\tilde{F}]}{B} \right) \mathcal{R}(-\pi/2) \right] : \perp \nabla_{\tilde{V}} \tilde{F} \otimes \mathcal{R}(\omega_c s) \tilde{v} \\
&- \frac{\sin(2\omega_c s)}{4\omega_c} \left[ \partial_x \left( \frac{E[\tilde{F}]}{B} \right) - \mathcal{R}(\pi/2) \partial_x \left( \frac{E[\tilde{F}]}{B} \right) \mathcal{R}(-\pi/2) \right] : \perp \nabla_{\tilde{V}} \tilde{F} \otimes \mathcal{R}(\omega_c s) \tilde{v}.
\end{aligned}$$

- 1 **Proof.** The particle density  $\tilde{f}$  satisfies the constraint  $(\partial_s + b \cdot \nabla_{x,\tilde{v}}) \tilde{f} = 0$  and therefore  
2 we have

$$\tilde{f}(s, x, \tilde{v}) = \tilde{f}(0, (\mathcal{X}, \tilde{\mathcal{V}})(-s; x, \tilde{v})), \quad (s, x, \tilde{v}) \in \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2.$$

- 3 Therefore there is a function  $\tilde{F} \in C_c^1(\mathbb{R}^2 \times \mathbb{R}^2)$  such that

$$\tilde{f}(s, x, \tilde{v}) = \tilde{F}(x, \mathcal{R}(\omega_c s) \tilde{v}), \quad (s, x, \tilde{v}) \in \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2.$$

- 4 Observe that  $\rho[\tilde{f}(s)] = \rho[\tilde{F}]$ ,  $j[\tilde{f}(s)] = \mathcal{R}(-\omega_c s) j[\tilde{F}]$  and

$$\nabla_{\tilde{v}} \tilde{f}(s, x, \tilde{v}) = \mathcal{R}(-\omega_c s) \nabla_{\tilde{V}} \tilde{F}(x, \mathcal{R}(\omega_c s) \tilde{v}), \quad (s, x, \tilde{v}) \in \mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2.$$

Notice that

$$\begin{aligned}
&\frac{1}{2\pi \varepsilon_0 B} \operatorname{div}_x \int_{\mathbb{R}^2} \frac{\perp(x-x')}{|x-x'|^2} \otimes j[\tilde{f}(s)](x') dx' \cdot \nabla_{\tilde{v}} \tilde{f}(s) - \frac{\perp j[\tilde{f}(s)]}{2\varepsilon_0 B} \cdot \nabla_{\tilde{v}} \tilde{f}(s) \quad (20) \\
&= \frac{\operatorname{div}_x}{2\pi \varepsilon_0 B} \int_{\mathbb{R}^2} \mathcal{R}(\omega_c s) \frac{\perp(x-x')}{|x-x'|^2} \otimes \mathcal{R}(-\omega_c s) j[\tilde{F}](x') dx' \cdot (\nabla_{\tilde{V}} \tilde{F})(x, \mathcal{R}(\omega_c s) \tilde{v}) \\
&- \frac{\perp j[\tilde{F}](x)}{2\varepsilon_0 B} \cdot (\nabla_{\tilde{V}} \tilde{F})(x, \mathcal{R}(\omega_c s) \tilde{v}).
\end{aligned}$$

- 5 As already observed in (13), the average with respect to  $s$  of

$$\frac{1}{2\pi \varepsilon_0 B} \operatorname{div}_x \int_{\mathbb{R}^2} \mathcal{R}(\omega_c s) \frac{\perp(x-x')}{|x-x'|^2} \otimes \mathcal{R}(-\omega_c s) j[\tilde{F}](x') dx'$$

coincides with  $\perp j[\tilde{F}]/(2\varepsilon_0 B)$ , and therefore we have

$$\begin{aligned}
&\frac{\operatorname{div}_x}{2\pi \varepsilon_0 B} \int_{\mathbb{R}^2} \mathcal{R}(\omega_c s) \frac{\perp(x-x')}{|x-x'|^2} \otimes \mathcal{R}(-\omega_c s) j[\tilde{F}](x') dx' - \frac{\perp j[\tilde{F}](x)}{2\varepsilon_0 B} \\
&= \frac{\operatorname{div}_x}{2\pi \varepsilon_0 B} \int_{\mathbb{R}^2} \left[ \cos^2(\omega_c s) - \frac{1}{2} \right] \frac{\perp(x-x')}{|x-x'|^2} \otimes j[\tilde{F}] + \left[ \sin^2(\omega_c s) - \frac{1}{2} \right] \frac{x-x'}{|x-x'|^2} \otimes \perp j[\tilde{F}] dx' \\
&+ \frac{\operatorname{div}_x}{2\pi \varepsilon_0 B} \int_{\mathbb{R}^2} \cos(\omega_c s) \sin(\omega_c s) \left[ \frac{\perp(x-x')}{|x-x'|^2} \otimes \perp j[\tilde{F}] + \frac{(x-x')}{|x-x'|^2} \otimes j[\tilde{F}] \right] dx'
\end{aligned}$$

whose zero average primitive with respect to  $s$  is

$$\begin{aligned} & \frac{\sin(2\omega_c s)}{8\omega_c \pi \varepsilon_0 B} \operatorname{div}_x \int_{\mathbb{R}^2} \left[ \frac{\perp(x-x')}{|x-x'|^2} \otimes j[\tilde{F}] - \frac{x-x'}{|x-x'|^2} \otimes \perp j[\tilde{F}] \right] dx' \\ & - \frac{\cos(2\omega_c s)}{8\omega_c \pi \varepsilon_0 B} \operatorname{div}_x \int_{\mathbb{R}^2} \left[ \frac{\perp(x-x')}{|x-x'|^2} \otimes \perp j[\tilde{F}] + \frac{x-x'}{|x-x'|^2} \otimes j[\tilde{F}] \right] dx'. \end{aligned}$$

Coming back to (20), and taking into account that  $\nabla_{\tilde{v}} \tilde{F}(x, \mathcal{R}(\omega_c s) \tilde{v})$  belongs to the kernel of  $\partial_s + b \cdot \nabla_{x, \tilde{v}}$ , since it depends only on the invariants  $x, \mathcal{R}(\omega_c s) \tilde{v}$  of  $\partial_s + b \cdot \nabla_{x, \tilde{v}}$ , it is easily seen that

$$\frac{\operatorname{div}_x}{2\pi \varepsilon_0 B} \int_{\mathbb{R}^2} \frac{\perp(x-x')}{|x-x'|^2} \otimes j[\tilde{f}(s)] dx' \cdot \nabla_{\tilde{v}} \tilde{f}(s) - \frac{\perp j[\tilde{f}(s)]}{2\varepsilon_0 B} \cdot \nabla_{\tilde{v}} \tilde{f}(s) + (\partial_s + b \cdot \nabla_{x, \tilde{v}}) \tilde{f}_I^2 = 0$$

where

$$\begin{aligned} \tilde{f}_I^2 &= \frac{\cos(2\omega_c s)}{8\omega_c \pi \varepsilon_0 B} \operatorname{div}_x \int_{\mathbb{R}^2} \left[ \frac{\perp(x-x')}{|x-x'|^2} \otimes \perp j[\tilde{F}] + \frac{x-x'}{|x-x'|^2} \otimes j[\tilde{F}] \right] dx' \cdot \nabla_{\tilde{v}} \tilde{F}(x, \mathcal{R}(\omega_c s) \tilde{v}) \\ &+ \frac{\sin(2\omega_c s)}{8\omega_c \pi \varepsilon_0 B} \operatorname{div}_x \int_{\mathbb{R}^2} \left[ \frac{(x-x')}{|x-x'|^2} \otimes \perp j[\tilde{F}] - \frac{\perp(x-x')}{|x-x'|^2} \otimes j[\tilde{F}] \right] dx' \cdot \nabla_{\tilde{v}} \tilde{F}(x, \mathcal{R}(\omega_c s) \tilde{v}). \end{aligned} \quad (21)$$

1 Similarly we obtain

$$-\partial_x \left( \frac{\perp E[\tilde{f}(s)]}{B} \right) \tilde{v} \cdot \nabla_{\tilde{v}} \tilde{f} + \frac{\rho[\tilde{f}(s)]}{2\varepsilon_0 B} \perp \tilde{v} \cdot \nabla_{\tilde{v}} \tilde{f} + (\partial_s + b \cdot \nabla_{x, \tilde{v}}) \tilde{f}_{II}^2 = 0$$

where

$$\begin{aligned} \tilde{f}_{II}^2 &= \frac{\cos(2\omega_c s)}{4\omega_c} \left[ \mathcal{R}(\pi/2) \partial_x \left( \frac{E[\tilde{F}]}{B} \right) + \partial_x \left( \frac{E[\tilde{F}]}{B} \right) \mathcal{R}(-\pi/2) \right] : \perp \nabla_{\tilde{v}} \tilde{F} \otimes \mathcal{R}(\omega_c s) \tilde{v} \\ &- \frac{\sin(2\omega_c s)}{4\omega_c} \left[ \partial_x \left( \frac{E[\tilde{F}]}{B} \right) - \mathcal{R}(\pi/2) \partial_x \left( \frac{E[\tilde{F}]}{B} \right) \mathcal{R}(-\pi/2) \right] : \perp \nabla_{\tilde{v}} \tilde{F} \otimes \mathcal{R}(\omega_c s) \tilde{v}. \end{aligned} \quad (22)$$

2 Our conclusion follows by combining (21), (22).  $\square$

3 **Remark 4.1** Notice that  $\tilde{f}^2 = \tilde{f}^2(s, x, \tilde{v})$  is continuous,  $S = \frac{2\pi}{|\omega_c|}$  periodic and uni-  
4 formly compactly supported in  $(x, \tilde{v})$  with respect to  $s$ . Moreover, if  $\tilde{f} \in C^k(\mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2)$   
5 for some integer  $k \geq 2$ , then  $\tilde{f}^2 \in C^{k-1}(\mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2)$ .

6 We appeal to the application  $T^\varepsilon(x, \tilde{v}) = (x + \varepsilon \perp \tilde{v} / \omega_c, \tilde{v})$ . We have

$$\partial T^\varepsilon b^\varepsilon = b \circ T^\varepsilon, \quad b \cdot \nabla_{x, \tilde{v}} = \omega_c \perp \tilde{v} \cdot \nabla_{\tilde{v}}, \quad b^\varepsilon \cdot \nabla_{x, \tilde{v}} = \varepsilon \tilde{v} \cdot \nabla_x + \omega_c \perp \tilde{v} \cdot \nabla_{\tilde{v}}. \quad (23)$$

7 Finally we are ready to prove that  $\bar{f}(t) \circ T^\varepsilon$ , where  $\bar{f}(t, x, \tilde{v}) = \tilde{F}(t, x, \mathcal{R}(\omega_c t / \varepsilon) \tilde{v})$ , is a  
8 second order approximation of  $\tilde{f}^\varepsilon(t, x, \tilde{v})$ .

1 **Proof.** (of Theorem 1.1)

2 Estimating the error between  $\tilde{f}^\varepsilon(t), \bar{f}(t) \circ T^\varepsilon$  is not of all obvious. A direct comparison  
3 between the above densities does not lead to the desired error estimate. We have to  
4 use a corrector term. More exactly, for any  $t \in \mathbb{R}_+$ , the particle density  $(s, x, \tilde{v}) \rightarrow$   
5  $\tilde{f}(t, s, x, \tilde{v}) = \tilde{F}(t, (\mathcal{X}, \tilde{\mathcal{V}})(-s; x, \tilde{v}))$  belongs to  $C^2(\mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2)$ , is  $S = \frac{2\pi}{|\omega_c|}$  periodic  
6 with respect to  $s$  and uniformly compactly supported in  $(x, \tilde{v})$ , with respect to  $s \in \mathbb{R}$   
7 and  $t \in [0, T], T \in \mathbb{R}_+$ . By definition, for any  $t \in \mathbb{R}_+$ , the particle density  $\tilde{f}(t)$  satisfies  
8 the constraint  $(\partial_s + b \cdot \nabla_{x, \tilde{v}})\tilde{f}(t) = 0$ . Thanks to Proposition 4.1, there is  $\tilde{f}^2(t, s, x, \tilde{v})$   
9 such that

$$a[\tilde{f}(t, s)] \cdot \nabla_{x, \tilde{v}} \tilde{f} - \langle a \rangle [\tilde{f}(t, s)] \cdot \nabla_{x, \tilde{v}} \tilde{f} + (\partial_s + b \cdot \nabla_{x, \tilde{v}})\tilde{f}^2 = 0. \quad (24)$$

It is easily seen, by the explicit formula of  $\tilde{f}^2$ , that the corrector  $\tilde{f}^2$  belongs to  $C^1(\mathbb{R}_+ \times$   
 $\mathbb{R} \times \mathbb{R}^2 \times \mathbb{R}^2)$ , is  $S = \frac{2\pi}{|\omega_c|}$  periodic with respect to  $s$ , and uniformly compactly supported  
in  $(x, \tilde{v})$  with respect to  $s \in \mathbb{R}$  and  $t \in [0, T], T \in \mathbb{R}_+$ . Taking  $s = t/\varepsilon$  in (24),  
multiplying by  $\varepsilon$  and combining with (18) yield

$$\begin{aligned} \frac{d}{dt} \left\{ \bar{f}(t) + \varepsilon^2 \tilde{f}^2(t, t/\varepsilon) \right\} + \varepsilon a[\bar{f}(t)] \cdot \nabla_{x, \tilde{v}} \bar{f}(t) + \frac{b}{\varepsilon} \cdot \nabla_{x, \tilde{v}} \left\{ \bar{f}(t) + \varepsilon^2 \tilde{f}^2(t, t/\varepsilon) \right\} \\ = \varepsilon^2 \partial_t \tilde{f}^2(t, t/\varepsilon, x, \tilde{v}). \end{aligned}$$

Using the corrector  $\tilde{f}^2$  led to a model similar to (7), nevertheless the transport operators  
appearing in the terms  $\frac{b}{\varepsilon} \cdot \nabla_{x, \tilde{v}} \left\{ \bar{f}(t) + \varepsilon^2 \tilde{f}^2(t, t/\varepsilon) \right\}, \frac{b^\varepsilon}{\varepsilon} \cdot \nabla_{x, \tilde{v}} \tilde{f}^\varepsilon$  are different, and the  
key point is to take advantage of the map  $T^\varepsilon$ . After composition with  $T^\varepsilon$ , the above  
equation becomes, thanks to (23)

$$\begin{aligned} \frac{d}{dt} \left\{ \bar{f}(t) \circ T^\varepsilon + \varepsilon^2 \tilde{f}^2(t, t/\varepsilon) \circ T^\varepsilon \right\} + \varepsilon a[\bar{f}(t)] \circ T^\varepsilon \cdot (\nabla \bar{f}(t)) \circ T^\varepsilon \\ + \frac{b^\varepsilon}{\varepsilon} \cdot \nabla \left\{ \bar{f}(t) \circ T^\varepsilon + \varepsilon^2 \tilde{f}^2(t, t/\varepsilon) \circ T^\varepsilon \right\} = \varepsilon^2 \partial_t \{ \tilde{f}^2(t, t/\varepsilon) \circ T^\varepsilon \}. \end{aligned}$$

Combining with (7) and using the notation  $r^\varepsilon(t) = \tilde{f}^\varepsilon(t) - \bar{f}(t) \circ T^\varepsilon - \varepsilon^2 \tilde{f}^2(t, t/\varepsilon) \circ T^\varepsilon$ ,  
lead to

$$\partial_t r^\varepsilon + \varepsilon a^\varepsilon[\tilde{f}^\varepsilon(t)] \cdot \nabla_{x, \tilde{v}} \tilde{f}^\varepsilon - \varepsilon(a[\bar{f}(t)] \cdot \nabla \bar{f}(t)) \circ T^\varepsilon + \frac{b^\varepsilon}{\varepsilon} \cdot \nabla_{x, \tilde{v}} r^\varepsilon = -\varepsilon^2 \partial_t \tilde{f}^2(t, t/\varepsilon) \circ T^\varepsilon. \quad (25)$$

As the magnetic field is uniform, the vector fields  $a^\varepsilon[\tilde{f}^\varepsilon], b^\varepsilon$  are divergence free and  
therefore

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \varepsilon a^\varepsilon[\tilde{f}^\varepsilon(t)] \cdot \nabla_{x, \tilde{v}} \tilde{f}^\varepsilon r^\varepsilon \, d\tilde{v} dx = \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \varepsilon a^\varepsilon[\tilde{f}^\varepsilon(t)] \cdot \nabla (\bar{f}(t) \circ T^\varepsilon + \varepsilon^2 \tilde{f}^2(t, t/\varepsilon) \circ T^\varepsilon) r^\varepsilon \, d\tilde{v} dx$$

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{b^\varepsilon}{\varepsilon} \cdot \nabla_{x, \tilde{v}} r^\varepsilon r^\varepsilon \, d\tilde{v} dx = 0.$$

Multiplying (25) by  $r^\varepsilon(t, x, \tilde{v})$  and integrating with respect to  $(x, \tilde{v})$  imply

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|r^\varepsilon(t)\|_{L^2}^2 + \varepsilon \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} a^\varepsilon[\tilde{f}^\varepsilon(t)] \cdot \nabla(\bar{f}(t) \circ T^\varepsilon) r^\varepsilon \, d\tilde{v} dx \\ & - \varepsilon \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} (a[\bar{f}(t)] \cdot \nabla \bar{f}(t)) \circ T^\varepsilon r^\varepsilon \, d\tilde{v} dx \\ & = -\varepsilon^2 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \partial_t \tilde{f}^2(t, t/\varepsilon) \circ T^\varepsilon r^\varepsilon \, d\tilde{v} dx - \varepsilon^3 \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} a^\varepsilon[\tilde{f}^\varepsilon(t)] \cdot \nabla(\tilde{f}^2(t, t/\varepsilon) \circ T^\varepsilon) r^\varepsilon \, d\tilde{v} dx \end{aligned}$$

and by Bellman lemma one gets

$$\begin{aligned} \|r^\varepsilon(t)\|_{L^2} & \leq \|r^\varepsilon(0)\|_{L^2} + \varepsilon \int_0^t \|a^\varepsilon[\tilde{f}^\varepsilon(t')] \cdot \nabla(\bar{f}(t') \circ T^\varepsilon) - (a[\bar{f}(t')] \cdot \nabla \bar{f}(t')) \circ T^\varepsilon\|_{L^2} \, dt' \\ & + \varepsilon^2 \int_0^t \|\partial_t \tilde{f}^2(t', t'/\varepsilon) \circ T^\varepsilon\|_{L^2} \, dt' + \varepsilon^3 \int_0^t \|a^\varepsilon[\tilde{f}^\varepsilon(t')] \cdot \nabla(\tilde{f}^2(t', t'/\varepsilon) \circ T^\varepsilon)\|_{L^2} \, dt'. \end{aligned} \quad (26)$$

- 1 We are working for  $t \in [0, T]$ ,  $T \in \mathbb{R}_+$ , and we denote by  $C$  any constant depending on  
2  $m, \varepsilon_0, q, T, B$  and the initial particle density  $f_{\text{in}}$ , but not on  $\varepsilon$ . Thanks to (19) we have

$$\|r^\varepsilon(0)\|_{L^2} \leq \|\tilde{f}^\varepsilon(0) - \bar{f}(0) \circ T^\varepsilon\|_{L^2} + \varepsilon^2 \|\tilde{f}^2(0, 0) \circ T^\varepsilon\|_{L^2}$$

- 3 and thus clearly  $\|r^\varepsilon(0)\|_{L^2} \leq C\varepsilon^2$ . Using the  $C_c^1$  regularity of  $\tilde{f}^2$  which comes from the  
4  $C_c^2$  regularity of  $\tilde{F}$ , it is straightforward that

$$\varepsilon^2 \int_0^T \|\partial_t \tilde{f}^2(t, t/\varepsilon) \circ T^\varepsilon\|_{L^2} \, dt + \varepsilon^3 \int_0^T \|a^\varepsilon[\tilde{f}^\varepsilon(t)] \cdot \nabla(\tilde{f}^2(t, t/\varepsilon) \circ T^\varepsilon)\|_{L^2} \, dt \leq C\varepsilon^2.$$

We claim that

$$\begin{aligned} & \varepsilon \int_0^t \|a^\varepsilon[\tilde{f}^\varepsilon(t')] \cdot \nabla(\bar{f}(t') \circ T^\varepsilon) - (a[\bar{f}(t')] \cdot \nabla \bar{f}(t')) \circ T^\varepsilon\|_{L^2} \, dt' \leq C\varepsilon^2 \quad (27) \\ & + C\varepsilon \int_0^t \|\tilde{f}^\varepsilon(t') - \bar{f}(t') \circ T^\varepsilon\|_{L^2}, \quad t \in [0, T]. \end{aligned}$$

- 5 This part of the proof relies on the smoothness and the uniform estimates of the  
6 densities  $(\tilde{f}^\varepsilon)_{\varepsilon>0}$ , together with the elliptic regularity results. Thanks to the uniform  
7 bounds

$$\sup_{\varepsilon>0, t \in [0, T]} \{\|\tilde{f}^\varepsilon(t)\|_{C^1(\mathbb{R}^2 \times \mathbb{R}^2)} + \|E[\tilde{f}^\varepsilon(t)]\|_{C^1(\mathbb{R}^2)}\} < +\infty$$

- 8 it is easily seen that

$$\|(a^\varepsilon[\tilde{f}^\varepsilon(t)] - a[\tilde{f}^\varepsilon(t)]) \cdot \nabla(\bar{f}(t) \circ T^\varepsilon)\|_{L^2} \leq C\varepsilon, \quad t \in [0, T], \quad \varepsilon > 0.$$

- 9 Thanks to elliptic regularity results, the quantity

$$\|(a[\tilde{f}^\varepsilon(t)] - a[\bar{f}(t) \circ T^\varepsilon]) \cdot \nabla(\bar{f}(t) \circ T^\varepsilon)\|_{L^2}$$

- 10 is bounded by the  $L^2$  norms of the charge and current densities

$$\|\rho[\tilde{f}^\varepsilon(t)] - \rho[\bar{f}(t) \circ T^\varepsilon]\|_{L^2(\mathbb{R}^2)} + \|j[\tilde{f}^\varepsilon(t)] - j[\bar{f}(t) \circ T^\varepsilon]\|_{L^2(\mathbb{R}^2)}$$

1 and thus by the  $L^2$  norm of the particle densities  $\|\tilde{f}^\varepsilon(t) - \bar{f}(t) \circ T^\varepsilon\|_{L^2(\mathbb{R}^2 \times \mathbb{R}^2)}$ . We obtain  
 2 the inequality

$$\|(a[\tilde{f}^\varepsilon(t)] - a[\bar{f}(t) \circ T^\varepsilon]) \cdot \nabla(\bar{f}(t) \circ T^\varepsilon)\|_{L^2} \leq C \|\tilde{f}^\varepsilon(t) - \bar{f}(t) \circ T^\varepsilon\|_{L^2(\mathbb{R}^2 \times \mathbb{R}^2)}, \quad t \in [0, T], \quad \varepsilon > 0.$$

3 The inequality (27) follows immediately, noticing that

$$\|a[\bar{f}(t) \circ T^\varepsilon] \cdot \nabla(\bar{f}(t) \circ T^\varepsilon) - (a[\bar{f}(t)] \cdot \nabla \bar{f}(t)) \circ T^\varepsilon\|_{L^2(\mathbb{R}^2 \times \mathbb{R}^2)} \leq C\varepsilon, \quad t \in [0, T], \quad \varepsilon > 0.$$

4 Finally (26) writes

$$\|r^\varepsilon(t)\|_{L^2} \leq C\varepsilon^2 + C\varepsilon \int_0^t \|\tilde{f}^\varepsilon(t') - \bar{f}(t') \circ T^\varepsilon\|_{L^2} dt'$$

implying that

$$\begin{aligned} \|\tilde{f}^\varepsilon(t) - \bar{f}(t) \circ T^\varepsilon\|_{L^2} &\leq \|r^\varepsilon(t)\|_{L^2} + \varepsilon^2 \|\tilde{f}^2(t, t/\varepsilon) \circ T^\varepsilon\|_{L^2} \\ &\leq C\varepsilon^2 + C\varepsilon \int_0^t \|\tilde{f}^\varepsilon(t') - \bar{f}(t') \circ T^\varepsilon\|_{L^2} dt', \quad t \in [0, T], \quad \varepsilon > 0. \end{aligned}$$

5 By Gronwall lemma we deduce

$$\|\tilde{f}^\varepsilon(t) - \bar{f}(t) \circ T^\varepsilon\|_{L^2} \leq C\varepsilon^2 \exp(C\varepsilon t), \quad t \in [0, T], \quad \varepsilon > 0.$$

Clearly we have

$$\begin{aligned} \|E[f^\varepsilon(t)] - E[\tilde{F}(t)]\|_{L^2} &= \|E[\tilde{f}^\varepsilon(t)] - E[\bar{f}(t)]\|_{L^2} \leq \|E[\tilde{f}^\varepsilon(t)] - E[\bar{f}(t) \circ T^\varepsilon]\|_{L^2} \\ &\quad + \|E[\bar{f}(t) \circ T^\varepsilon] - E[\bar{f}(t)]\|_{L^2} \leq C\varepsilon \end{aligned}$$

and therefore

$$\begin{aligned} &\left\{ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left[ \tilde{F} \left( t, x + \varepsilon \frac{\perp v}{\omega_c}, \mathcal{R}(\omega_c t/\varepsilon) \left( v - \varepsilon \frac{\perp E[\tilde{F}(t)]}{B} \right) \right) - f^\varepsilon(t, x, v) \right]^2 dv dx \right\}^{1/2} \\ &= \left\{ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left[ \tilde{F} \left( t, x + \frac{\varepsilon}{\omega_c} \left( \perp \tilde{v} - \varepsilon \frac{E[f^\varepsilon(t)]}{B} \right), \mathcal{R}(\omega_c t/\varepsilon) \left( \tilde{v} + \varepsilon \frac{\perp E[f^\varepsilon(t)] - \perp E[\tilde{F}(t)]}{B} \right) \right) \right. \right. \\ &\quad \left. \left. - \tilde{f}^\varepsilon(t, x, \tilde{v}) \right]^2 d\tilde{v} dx \right\}^{1/2} \\ &\leq \|\tilde{f}^\varepsilon(t) - \bar{f}(t) \circ T^\varepsilon\|_{L^2} + \left\{ \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \left[ \tilde{F} \left( t, x + \varepsilon \frac{\perp \tilde{v}}{\omega_c}, \mathcal{R}(\omega_c t/\varepsilon) \tilde{v} \right) - \right. \right. \\ &\quad \left. \left. \tilde{F} \left( t, x + \varepsilon \frac{\perp \tilde{v}}{\omega_c} - \frac{\varepsilon^2}{\omega_c} \frac{E[f^\varepsilon(t)]}{B}, \mathcal{R}(\omega_c t/\varepsilon) \tilde{v} + \varepsilon \mathcal{R}(\omega_c t/\varepsilon) \frac{\perp E[f^\varepsilon(t)] - \perp E[\tilde{F}(t)]}{B} \right) \right]^2 d\tilde{v} dx \right\}^{1/2} \\ &\leq C\varepsilon^2 + C\varepsilon \|E[f^\varepsilon(t)] - E[\tilde{F}(t)]\|_{L^2} \leq C\varepsilon^2, \quad t \in [0, T], \quad \varepsilon > 0. \end{aligned}$$

6

□

## 5 Conclusions

We presented regular reformulations for the Vlasov-Poisson equations with uniform magnetic fields and general initial conditions. The effective model comes by averaging over one cyclotronic period, once we have determined a periodic fast dynamics. Certainly, in the framework of the magnetic confinement, a much more interesting case is that of curved magnetic fields.

The above results extend to the three dimensional Vlasov-Poisson system, with strong external curved magnetic field. The same arguments lead to regular reformulations and second order estimates for both well prepared and general initial particle densities. In the three dimensional setting, we emphasize a fast periodic dynamics leaving invariant not only the guiding center and the modulus of the perpendicular velocity, but also the parallel velocity. Averaging over one period allows us to obtain completely explicit effective models. Nevertheless the analysis is much more elaborated due to the combination between the parallel and perpendicular dynamics and to the curvature effects. These studies will be the topic of future works [10].

Another interesting issue will be to handle models with self-consistent magnetic field, that is the Vlasov-Maxwell equations, perturbed by a strong external magnetic field.

## Acknowledgement

This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

This work has been carried out thanks to the support of the A\*MIDEX project (no ANR-11-IDEX-0001-02) funded by the Investissements d’Avenir French Government program, managed by the French National Research Agency (ANR).

## References

- [1] A.A. Arsen’ev, Global existence of weak solution of Vlasov’s system of equations, *Z. Vychisl. Mat. Fiz.* 15(1975) 136-147.
- [2] J. Batt, Global symmetric solutions of the initial value problem in stellar dynamics, *J. Differential Equations* 25(1977) 342-364.
- [3] M. Bostan, Transport equations with disparate advection fields. Application to the gyrokinetic models in plasma physics, *J. Differential Equations* 249(2010) 1620-1663.
- [4] M. Bostan, Gyro-kinetic Vlasov equation in three dimensional setting. Second order approximation, *SIAM J. Multiscale Model. Simul.* 8(2010) 1923-1957.
- [5] M. Bostan, A. Finot, M. Hauray, The effective Vlasov-Poisson system for strongly magnetized plasmas, *C. R. Acad. Sci. Paris, Ser. I* 354(2016) 771-777.

- 1 [6] M. Bostan, A. Finot, The effective Vlasov-Poisson system for the finite Larmor  
2 radius regime, *SIAM J. Multiscale Model. Simul.* 14(2016) 1238-1275.
- 3 [7] M. Bostan, Transport of charged particles under fast oscillating magnetic fields,  
4 *SIAM J. Math. Anal.* 44(2012) 1415-1447.
- 5 [8] M. Bostan, Multi-scale analysis for linear first order PDEs. The finite Larmor  
6 radius regime, *SIAM J. Math. Anal.* 48(2016) 2133-2188.
- 7 [9] M. Bostan, Asymptotic behavior for the Vlasov-Poisson equations with strong  
8 external magnetic field. Straight magnetic field lines, to appear in *SIAM J. Math.*  
9 *Anal.* (2019).
- 10 [10] M. Bostan, Asymptotic behavior for the Vlasov-Poisson equations with strong  
11 external curved magnetic field, in preparation.
- 12 [11] Y. Brenier, Convergence of the Vlasov-Poisson system to the incompressible Euler  
13 equations, *Comm. Partial Differential Equations* 25(2000) 737-754.
- 14 [12] N. Crouseilles, M. Lemou, F. Méhats, Asymptotic preserving schemes for highly  
15 oscillatory Vlasov-Poisson equations, *J. Comput. Phys.* 248(2013) 287-308.
- 16 [13] P. Degond, F. Filbet, On the asymptotic limit of the three dimensional Vlasov-  
17 Poisson system for large magnetic field: formal derivation, *J. Stat. Phys.* 165(2016)  
18 765-784.
- 19 [14] F. Filbet, L. Rodrigues, Asymptotically preserving particle-in-cell methods for  
20 inhomogeneous strongly magnetized plasmas, *SIAM J. Numer. Anal.* 55(2017)  
21 2416-2443.
- 22 [15] F. Filbet, S. Jin, A class of asymptotic-preserving schemes for kinetic equations  
23 and related problems with stiff sources, *J. Comput. Phys.* 229(2010) 7625-7648.
- 24 [16] E. Frénod, E. Sonnendrücker, Homogenization of the Vlasov equation and of  
25 the Vlasov-Poisson system with strong external magnetic field, *Asymptotic Anal.*  
26 18(1998) 193-213.
- 27 [17] E. Frénod, E. Sonnendrücker, Long time behavior of the two-dimensional Vlasov-  
28 equation with strong external magnetic field, *Math. Models Methods Appl. Sci.*  
29 10(2000) 539-553.
- 30 [18] E. Frénod, E. Sonnendrücker, The finite Larmor radius approximation, *SIAM J.*  
31 *Math. Anal.* 32(2001) 1227-1247.
- 32 [19] F. Golse, L. Saint-Raymond, The Vlasov-Poisson system with strong magnetic  
33 field, *J. Math. Pures Appl.* 78(1999) 791-817.
- 34 [20] D. Han-Kwan, Effect of the polarization drift in a strongly magnetized plasma,  
35 *Math. Model. Numer. Anal.* 46(2012) 929-947.
- 36 [21] P.-L. Lions, B. Perthame, Propagation of moments and regularity for the 3-  
37 dimensional Vlasov-Poisson system, *Invent. Math.* 105(1991) 415-430.

- 1 [22] E. Miot, On the gyrokinetic limit for the two-dimensional Vlasov-Poisson system,  
2 arXiv:1603.04502.
- 3 [23] K. Pfaffelmoser, Global classical solutions of the Vlasov-Poisson system in 3 di-  
4 mensions for general initial data, J. Differential Equations 95(1992) 281-303.
- 5 [24] L. Saint-Raymond, The gyro-kinetic approximation for the Vlasov-Poisson system,  
6 Math. Models Methods Appl. Sci. 10(2000) 1305-1332.
- 7 [25] L. Saint-Raymond, Control of large velocities in the two-dimensional gyro-kinetic  
8 approximation, J. Math. Pures Appl. 81(2002) 379-399.
- 9 [26] T. Ukai, S. Okabe, On the classical solution in the large time of the two dimensional  
10 Vlasov equations, Osaka J. Math. 5(1978) 245-261.