



Truly sedentary? The multi-range tactic as a response to resource heterogeneity and unpredictability in a large herbivore

Ophélie Couriot, A.J. Mark Hewison, Sonia Said, Francesca Cagnacci, Simon Chamaillé-Jammes, John Linnell, Atle Mysterud, Wibke Peters, Ferdinando Urbano, Marco Heurich, et al.

► To cite this version:

Ophélie Couriot, A.J. Mark Hewison, Sonia Said, Francesca Cagnacci, Simon Chamaillé-Jammes, et al.. Truly sedentary? The multi-range tactic as a response to resource heterogeneity and unpredictability in a large herbivore. *Oecologia*, 2018, 187 (1), pp.47-60. 10.1007/s00442-018-4131-5 . hal-02311434

HAL Id: hal-02311434

<https://hal.science/hal-02311434>

Submitted on 13 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Truly sedentary? The multi-range tactic as a response to resource
heterogeneity and unpredictability in a large herbivore**

Ophélie Couriot, A. J. Mark Hewison, Sonia Saïd, Francesca Cagnacci, Simon Chamaillé-Jammes, John D. C. Linnell, Atle Mysterud, Wibke Peters, Ferdinando Urbano, Marco Heurich, Petter Kjellander, Sandro Nicoloso, Anne Berger, Pavel Sustr, Max Kroeschel, Leif Soennichsen, Robin Sandfort, Benedikt Gehr and Nicolas Morellet

Ophélie Couriot, CEFS, Université de Toulouse, INRA, Castanet Tolosan, France, email: o.couriot@gmail.com, tel. +33 (0) 561 28 51 25, corresponding author

A. J. Mark Hewison, CEFS, Université de Toulouse, INRA, Castanet Tolosan, France, email: mark.hewison@inra.fr

Sonia Saïd, Office National de la Chasse et de la Faune Sauvage, Direction Recherche et Expertise, Unité Ongulés Sauvages, « Monfort » 01330 Birieux, France, email: sonia.said@oncfs.gouv.fr

Francesca Cagnacci, Biodiversity and Molecular Ecology Department, Research and Innovation Centre, Fondazione Edmund Mach, Via Mach 1, 38010 San Michele all'Adige, TN, Italy, email: cagnacci.fem@gmail.com

Simon Chamaillé-Jammes, Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175, CNRS - Université de Montpellier - Université Paul-Valéry Montpellier - EPHE, 34293 Montpellier Cedex 5, France, email: simon.chamaille@cefe.cnrs.fr

John D. C. Linnell, Norwegian Institute for Nature Research, PO Box 5685 Torgard, NO-7485 Trondheim, Norway, email: john.linnell@nina.no

Atle Mysterud, Centre for Ecological and Evolutionary Synthesis (CEES), University of Oslo, Blindern, P.O. Box 1066, 0316 Oslo, Norway, email: atle.mysterud@ibv.uio.no

26 Wibke Peters, Bavarian State Institute of Forestry (LWF), Hans-Carl-von-Carlowitz-Platz 1,
27 85354 Freising, Germany, email: wibkepeters@gmail.com

28 Ferdinando Urbano, Freelance Eurodeer Database Manager, Milan, Italy, email:
29 ferdi.urbano@gmail.com

30 Marco Heurich, Bavarian Forest National Park, Department of Conservation and Research,
31 Freyunger Str. 2, 94481 Grafenau, Germany & Chair of Wildlife Ecology and Management,
32 University of Freiburg, Tennenbacher Straße 4, 79106 Freiburg, Germany, email:
33 marco.heurich@npv-bw.bayern.de

34 Petter Kjellander, Grimsö Wildlife Research Station, Department of Ecology, Swedish
35 University of Agricultural Sciences, Riddarhyttan, Sweden, email: petter.kjellander@slu.se

36 Sandro Nicoloso, Research, Ecology and Environment Dimension (D.R.E.Am. Italia), Via
37 Enrico Bindi 14, 51100 Pistoia, PT, Italy, email: nicoloso@dream-italia.it

38 Anne Berger, Leibniz-Institute for Zoo and Wildlife Research, Alfred-Kowalke-Str. 17,
39 10315 Berlin, Germany, email: berger@izw-berlin.de

40 Pavel Sustr, Department of Biodiversity Research, Global Change Research Centre, Academy
41 of Sciences of the Czech Republic, Brno, Czech Republic, Šumava National Park, Vimperk,
42 Czech Republic, email: krakonosnasumave@gmail.com

43 Max Kroeschel, Chair of Wildlife Ecology and Management, University of Freiburg,
44 Tennenbacher Straße 4, 79106 Freiburg, Germany & Forest Research Institute of Baden-
45 Wuerttemberg, Wildlife Ecology, Wonnhaldestraße 4, 79100 Freiburg, Germany, email:
46 max.kroeschel@wildlife.uni-freiburg.de

47 Leif Soennichsen, Mammal Research Institute, Polish Academy of Sciences, Białowieża,
48 Poland, email: l_soennichsen@ibs.bialowieza.pl

49 Robin Sandfort, Institute of Wildlife Biology and Game Management, Department of
50 Integrative Biology and Biodiversity Research, University of Natural Resources and Life
51 Sciences, Vienna, email: robin.sandfort@boku.ac.at
52 Benedikt Gehr, Department of Evolutionary Biology and Environmental Studies, University
53 of Zurich, Winterthurerstrasse 190, 8057 Zurich, email: benedikt.gehr@ieu.uzh.ch
54 Nicolas Morellet, CEFS, Université de Toulouse, INRA, Castanet Tolosan, France, email:
55 nicolas.morellet@inra.fr

Author contributions: OC, NM and AJMH formulated the idea. AJMH, FC, JDCL, AM, MH, PK, SN, AB, PS, MK, LS, RS and BG provided data. FU created and updated the database. OC, NM, and AJMH developed the methodology, OC and NM performed the statistical analyses and wrote the manuscript with assistance from AJMH. SS, FC, SCJ, JDCL, AM, WP, FU, MH, PK, SN, AB, PS, MK, LS, RS and BG commented on and assisted in revising the manuscript.

Abstract

Much research on large herbivore movement has focused on the annual scale to distinguish between resident and migratory tactics, commonly assuming that individuals are sedentary at the within-season scale. However, apparently sedentary animals may occupy a number of sub-seasonal functional home ranges (sfHR), particularly when the environment is spatially heterogeneous and/or temporally unpredictable. The roe deer (*Capreolus capreolus*) experiences sharply contrasting environmental conditions due to its widespread distribution, but appears markedly sedentary over much of its range. Using GPS monitoring from 15 populations across Europe, we evaluated the propensity of this large herbivore to be truly sedentary at the seasonal scale in relation to variation in environmental conditions. We studied movement using net square displacement to identify the possible use of sfHR. We expected that roe deer should be less sedentary within seasons in heterogeneous and unpredictable environments, while migratory individuals should be seasonally more sedentary than residents. Our analyses revealed that, across the 15 populations, all individuals adopted a multi-range tactic, occupying between two and nine sfHR during a given season. In addition, we showed that (i) the number of sfHR was only marginally influenced by variation in resource distribution, but decreased with increasing sfHR size; and (ii) the distance between sfHR increased with increasing heterogeneity and predictability in resource distribution, as well as with increasing sfHR size. We suggest that the multi-range tactic is likely widespread among large herbivores, allowing animals to track spatio-temporal variation in resource distribution and, thereby, to cope with changes in their local environment.

Key words: migration, residency, sub-seasonal functional home range, plasticity, roe deer

Introduction

Movement is a fundamental characteristic of life which influences the survival and reproduction of organisms and, more generally, individual fitness and population dynamics (Turchin 1991; Revilla and Wiegand 2008). Following the Marginal Value Theorem (Charnov 1976), while foraging, an individual moves within a patch (*sensu* Wiens 1976), searching intensively for food, before leaving to search more widely for a new patch when the energetic benefit of the first patch has decreased below the average value of the alternative patches (Charnov 1976; Van Moorter et al. 2009). In slowly regenerating environments, animals should engage in thorough exploration to locate suitable patches. However, in rapidly regenerating environments, individuals are assumed to memorise the value and location of a given patch (Bracis et al. 2015; Riotte-Lambert et al. 2015), returning to previously visited patches periodically. This process leads to the emergence of a stable home range to which individuals restrict their movements in order to maximise resource acquisition (Brown and Orians 1970; Riotte-Lambert et al. 2015). Therefore, individuals of many species appear to be sedentary at this spatio-temporal scale, occupying a stable home range over a long time span (season and year). Site fidelity is widespread in the animal kingdom and has fundamental consequences for ecological processes (Börger et al. 2008).

An individual's lifetime track is an aggregation of successive elementary units with potentially different functionality (Baguette et al. 2014). Indeed, as stated by Van Moorter et al. (2016) "animals do not move for the sake of changing their geographic location, but rather for changing environmental conditions associated with changes in location". In general, an individual decides to move in order to satisfy its requirements in terms of refuge and resources (Nathan et al. 2008) which encompass changes in environmental space (Van Moorter et al. 2013). For example, since favourable sites for feeding or taking refuge do not necessarily occur at the same location, and since conditions vary in space and time, animals

have to move to cope with spatio-temporal heterogeneity in their environment (Pyke 1984; Mueller and Fagan 2008; Chapman et al. 2014). As a result of variation in conditions over time, an animal must shift from one suitable patch to another in order to fulfil its requirements, in which case it cannot be considered truly sedentary, even at the seasonal scale (Chapin et al. 1980; Barraquand and Benhamou 2008). Indeed, animals may use several spatio-temporally distinct suitable units, particularly in spatially heterogeneous and/or temporally unpredictable environments. While there has been considerable focus in recent years on movements associated with seasonal migration between seasonally distinct home ranges (Cagnacci et al. 2011; Cagnacci et al. 2015; Peters et al. 2017), there has been relatively little work on finer scale movements at the within season scale. The use of sub-seasonal functional home ranges (sfHR) has previously been described in two African herbivores, the sable antelope (*Hippotragus niger*) and the African savanna buffalo (*Syncerus caffer brachyceros*) (Owen-Smith et al. 2010; Cornélis et al. 2011; Benhamou 2014). However, no study has yet attempted to link the propensity of individuals to adopt this multi-range tactic with spatial and temporal variation in the prevailing environmental conditions.

Here, we used the EURODEER database (<http://www.eurodeer.org>, see methods) to analyse space use of roe deer (*Capreolus capreolus*) across widely contrasting environments, from the southern part of their geographic range in Italy to the northern part of their range, in Scandinavia. We focused on the roe deer as it is Europe's most widespread large wild herbivore and is considered highly sedentary over the majority of its range (Hewison et al. 1998). However, this species also exhibits a considerable degree of behavioural plasticity (Jepsen and Topping 2004), and is described as partially migratory in more extreme environments (Cagnacci et al. 2011). Hence, we first analysed whether roe deer are truly sedentary within a given season, or whether they adopt a movement tactic based on the use of a series of sfHR. Secondly, we hypothesised that the propensity of an animal to adopt this

130 multi-range tactic should depend on spatio-temporal variations in environmental conditions.
131 More specifically, we predicted that individuals should be less sedentary in spatially
132 heterogeneous compared to homogeneous environments (Mueller and Fagan 2008; Mueller et
133 al. 2011). Analogously to the nomadic movement tactic (Mueller and Fagan 2008), we also
134 expected individuals to be less sedentary in temporally unpredictable environments, or at least
135 in environments where resources vary more markedly over time at the within-season scale,
136 compared to more predictable environments. Finally, we expected that, within a given
137 seasonal range, migratory animals would be more sedentary than residents since they migrate
138 during spring and fall so that they are able to adjust their habitat use to seasonal variations in
139 food resource abundance and/or quality at this scale (Fryxell and Sinclair 1988).

Material and methods

Study areas and GPS data

This study was based on the database assembled by the EURODEER consortium, a data sharing project to investigate the movement ecology of European deer along environmental gradients (<http://eurodeer.org>, accessed on April 2016). We analysed data on 251 adult roe deer (286 individual-years) from 15 study sites (see Table 1) encompassing widely contrasting environmental conditions (latitude varied from 38.2°N to 60.7°N; longitude varied from 0.9 °E to 23.5 °E; Fig. 1). Roe deer were captured from 2003 to 2014 using drive nets, net traps or box traps depending on study site. All capture and marking procedures were done in accordance with local and European animal welfare laws. Deer were equipped with GPS collars programmed to obtain a GPS fix with intervals ranging from 10 minutes to 12 hours. In order to standardise the data for inter-population comparisons, for each individual, we restricted monitoring to the period from the 15th of February to the 15th of November, and retained the two locations per day that were closest to noon and midnight.

Discrimination of individual movement tactics

Of the recently published emerging tools for determining movement tactic (Gurarie et al. 2017; Spitz et al. 2017), we first used a common method proposed by Börger and Fryxell (2012), which has been proven useful for large mammals (Bunnefeld et al. 2011; Bischof et al. 2012), based on the net squared displacement (NSD), *i.e.* the Euclidian distance between the starting location and all subsequent locations of an individual over time (Turchin 1998), to determine each individual's annual movement tactic: migration, residency or dispersal. We considered two models of range residency, one with a constant NSD (the mean), and one with a linear increase of NSD before reaching an asymptote; we considered one model of migration including approximate dates of departure and return between seasonal ranges, and a

165 model of dispersal with an approximate date of departure (see Bunnefeld et al. 2011 and
166 Börger and Fryxell 2012 for more details on these models). In order to identify which of these
167 models best described the movement behaviour of a given individual, we used the system of
168 non-linear mixed models proposed by Börger and Fryxell (2012) which links theoretical
169 expectations to movement data. For model selection, as recommended by Börger and Fryxell
170 (2012), we retained the model with the largest concordance correlation (CC), expressing the
171 goodness of fit for each model (Huang et al. 2009). Because the assigned movement tactic
172 using this method did not always closely fit the data, we also visually examined the NSD
173 trajectories to determine each individual's annual movement tactic by eye (Bischof et al.
174 2012). We based our visual classification on the patterns of NSD typically observed for
175 migratory individuals, residents and dispersers, following Börger and Fryxell (2012). That is,
176 we assumed that the individual was resident when the NSD was relatively constant, or
177 initially increased linearly before rapidly reaching an asymptote. We assumed that the
178 individual was migratory when the NSD was constant before increasing rapidly during spring
179 to reach a plateau during summer, then decreased during fall, returning to its initial value.
180 Finally, we assumed that the individual had dispersed when the NSD was constant before
181 increasing rapidly to reach a plateau with no further increase or decrease. We then verified
182 that individuals which were classified as dispersers did not return to their point of departure
183 during subsequent monitoring, after the 15th of November. If they did (53 of 65 animals
184 originally classified as dispersers), these individuals were considered as migratory. We
185 excluded the remaining dispersers (N = 12) from subsequent analyses as movement patterns
186 during dispersal are governed by different ultimate causes than those involved in range
187 residency or migration (Bowler and Benton 2005; Chapman et al. 2014). After visual
188 reclassification, our data set included 193 residents and 93 migratory individuals. Note that
189 subsequent analyses based on this visual classification of individual movement tactics

generated results that were similar to those based on the classification using Borger and Fryxell's (2012) method (not shown).

Subsequently, for each migratory individual, we segmented the NSD using Lavielle's method (Lavielle 1999), which detects change points in a time series, to identify the dates of departure and return from, and to, the winter range (if any) and to define individual-based seasonal ranges. Dates of departure from the winter range ranged from the 6th of March to the 3rd of August (median = 4th of May, sd = 32 days), while return ranged from the 7th of May to the 28th of October (median = 10th of September, sd = 42 days). We then used the median departure and return dates across all migratory individuals to establish equivalent seasonal phases for resident individuals. As a result, we subsequently analysed movement behaviour of all deer during the winter period (prior to departure) and the summer period (after departure and prior to return), excluding the three days prior to and following departure and return to avoid the transience phase. We did not analyse data from the post-return period, (the second winter), since monitoring periods were too short to characterise individual movement during this period.

Detecting sub-seasonal functional home ranges

We then tested the assumption that roe deer were truly sedentary within the above defined seasonal periods or, whether their seasonal ranges were composed of several sub-seasonal functional home ranges (sfHR, Benhamou 2014). To do so, we segmented each individual's movement path (*i.e.* the temporal sequence of locations) for each seasonal period using Lavielle's (1999) method on the mean NSD to identify fine scale stationary states. Following several studies which have examined variation of weekly home ranges in response to variation in resource distribution (Rivrud et al. 2010; Van Beest et al. 2011; Morellet et al. 2013), we considered 14 locations (*i.e.* 7 days) as the minimum number required to describe a stationary

state. Because this choice of a minimum time span is somewhat arbitrary, we repeated the segmentation approach twice using minimum values of 5 and of 9 days (*i.e.* 7 ± 2 days) to assess the sensitivity of our results to the length of the temporal window. Finally, we retained the most parsimonious number of segments comprising each seasonal range for each individual (Figs. 2 & 3). We again repeated this step when considering 5 and 9 days as the minimum temporal windows.

We generated a non-linear mixed model describing the use of more than one sfHR within a given season. This model was based on the mean NSD per stationary state (*i.e.* the number of segments defined by Lavielle's method) for each seasonal period (Supplementary materials, Appendix 1, Eq. A1). To determine whether individuals were truly sedentary, we compared this model with the two models of range residency developed by Börger and Fryxell (2012) used above, adapted for each seasonal period (Supplementary materials, Appendix 1, Eq. A2 & A3), and retained the model with the largest CC (See Supplementary materials, Appendix 1, Table A1).

Describing site-specific environmental conditions

In order to explore the influence of environmental conditions on the propensity of roe deer to use a number of sfHR within a given season, we described forage resources at each site in terms of spatial heterogeneity and temporal predictability for both the winter and summer periods. To identify the spatial limits of each site, we used the 100% minimum convex polygon of the locations of all GPS monitored individuals at that site to which we added a buffer of 1 kilometre.

To quantify broad scale site-specific heterogeneity in resource distribution, we used the normalized difference vegetation index (NDVI). NDVI is widely used in ecological research and management as a proxy for biomass (Kerr and Ostrovsky 2003; Pettorelli et al.

2011) and forage quality available for ungulates (Hamel et al. 2009; Borowik et al. 2013; Lendrum et al. 2014; Garrouette et al. 2016) and particularly for roe deer (Morellet et al. 2013; Peters et al. 2017). NDVI was used because of its simplicity, consistent coverage of all study areas, high temporal resolution and demonstrated utility in a variety of ecosystems. Given the linear relationship between NDVI and biomass, a given variation in NDVI should index a similar relative reduction in forage biomass across study sites. Hence, it provides a useful and integrative measure of variation in primary productivity which we assume to be suitable for our purpose of comparing roe deer populations inhabiting widely contrasting environmental conditions.

We obtained weekly values for the NDVI (pixel size = ca. 250m x 250m) from the University of Natural Resources and Life Sciences, Vienna (BOKU) using near real-time filtered products (Whittaker smoother, (Vuolo et al. 2012; Klisch and Atzberger 2016)) of the NDVI time series generated by the NASA Moderate Resolution Imaging Spectroradiometer (MODIS). We used the REFMID values, supplied by BOKU, which are the most stable extrapolated values of NDVI (<http://ivfl-info.boku.ac.at/index.php/eo-data-processing/real-time-modis-data-eu-only> for more details).

To describe spatial heterogeneity in resource distribution, we measured spatial heterogeneity of the NDVI at the study site level using the standard deviation of weekly values of the NDVI over all pixels of each study site (Coops et al. 1998; Coops and Culvenor 2000). Then, we averaged these values across weeks for each period to obtain a proxy of spatial heterogeneity per seasonal period, defined above, and per year (from 2003 to 2014) for each study site. Spatial heterogeneity in NDVI ranged from 0.03 (low heterogeneity) to 0.23 (high heterogeneity).

To index temporal predictability in resource distribution, we calculated temporal constancy of the NDVI values across years (*i.e.* from 2003 to 2014) for each period at a given

site following the approach proposed by Colwell (1974). Constancy is a component of temporal predictability (Colwell 1974) that ranges from 0 (no temporal predictability or high temporal variability) to 1 (perfect temporal predictability or high temporal stability). We calculated a value of temporal constancy for each period at each study site across weeks and years (using weekly average values of NDVI across all pixels of a given study site). For this, we built a matrix with NDVI values sorted into 10 equal 0.1 interval classes between 0 and 1 in rows, as indicated by English et al. (2012), and the number of the weeks in columns (irrespective of the year). We counted the number of years for which we observed each class of NDVI for each week, and then we calculated constancy as follows:

$$Constancy = 1 + \frac{\sum \frac{\sum NDVI_{ij}}{\sum \sum NDVI_{ij}} \log(\frac{\sum NDVI_{ij}}{\sum \sum NDVI_{ij}})}{\log(s)}$$

where i represents the NDVI class, j is the week of the NDVI measure, irrespective of the year, and s is the number of NDVI classes.

As resource heterogeneity was negatively correlated with resource predictability during both seasonal periods ($r = -0.50$, $p < 0.001$ during winter, $r = -0.88$, $p < 0.001$ during summer), we used the residuals of the linear regression between resource predictability and resource heterogeneity to index temporal predictability in resource distribution. Hence, positive values indicate that temporal predictability is higher than expected for a given level of resource heterogeneity.

Statistical analyses

To analyse the link between space use behaviour of roe deer with spatial heterogeneity and temporal predictability in resource distribution, we exploited the extremely marked variation among study sites in environmental conditions, performing the analyses at the population level. We were unable to analyse within-population (*i.e.* individual level) variation in these

relationships due to the difficulty in measuring the distribution of resources within each study site at a sufficiently fine-scale. To describe space use behaviour, we used two individual-based metrics. First, we calculated the number of sfHR used by each individual during each period to describe the degree to which an individual was truly sedentary within a given seasonal range. Second, we calculated the median distance between the centres of all pairs of sfHR for each individual and each period to index the degree of spatial separation among functional ranges (sfHR distance). We log-transformed this quantity to achieve normality.

How many sub-seasonal functional home ranges?

We used Generalised Linear Mixed-Effects Models (glmm), with a Poisson distribution and a log-link function, to analyse variation in the number of sfHR within an individual's seasonal range (wintering or breeding period) in relation to annual movement tactic (resident vs. migratory), sex (male vs. female, because males are seasonally territorial in this species, Vanpé *et al.*, 2009) and two continuous descriptors of site-specific resource distribution: spatial heterogeneity and temporal predictability. We included the individual's period-specific log-transformed median sfHR size as an additive factor to control for variation in absolute resource availability between individuals (Morellet *et al.* 2013). Thus, the most complex model contained two three-way interactions among movement tactic, sex and resource heterogeneity and among movement tactic, sex and resource predictability, plus the logarithm of the median sfHR size as a fixed effect, and the study site as a random effect to control for repeated measures (individuals) per population. As the number of sfHR occupied by an individual ranged from 2 to 9, we subtracted 2 from the total number of sfHR per individual per season in order to conform to a Poisson distribution (Haight 1967). The number of sfHR then ranged from 0 to 7. We estimated sfHR size using fixed-kernel methods considering (i) an ad hoc approach to select the optimal smoothing parameter h for each sfHR estimate and

(ii) a constant h set to the overall median of the smoothing parameters estimated by the ad hoc approach for each individual sfHR estimate. Finally, sfHR size was estimated at both 90% and 50% isopleths.

Spatial separation among sub-seasonal functional home ranges

We used Linear Mixed-Effects models (lme) to investigate variation in log-transformed inter-sfHR distance, including period, movement tactic, sex and resource heterogeneity and resource predictability for each study site. We built the equivalent set of models described above for analysing variation in inter-sfHR distance.

For model selection, we used Akaike's Information Criterion (AIC, Burnham and Anderson 2002) and the number of parameters to select the most parsimonious model that best described the data. All analyses were performed in R version 3.1.2 (R Development Core Team 2014) using *glmer* and *lmer* functions in the *lme4* library.

Results

Our overall results were robust with respect to the choice of method to estimate sfHR size, the isopleth used (see Supplementary materials, Appendix 1, Fig. A7 to A15), and the minimum number of days required to describe a sfHR (see Supplementary materials, Appendix 1, Fig. A1 to A6). Thus, in the following, we only present results using the 90% fixed-kernel method based on a constant h and 7 days as the minimum temporal window for a sfHR.

Roe deer use sub-seasonal functional home ranges: the multi-range tactic

All individuals occupied more than one sfHR during a given season. Indeed, the model based on the occupation of more than one sfHR received more support than the two residency models (the CC value was highest for the multi-range model in all 572 cases, corresponding

to 286 individuals for the two periods, Supplementary materials, Appendix 1, Table A1). Individuals occupied a sfHR for at least 7 days (by definition, see Methods), and up to 186 days during the wintering period (median = 10 days), and for at least 7 days, and up to 223 days, during the breeding period (median = 24 days).

How many sub-seasonal functional home ranges?

The most parsimonious model describing variation in the number of sfHR during the winter period included the two-way interaction between resource predictability and movement tactic and the log-transformed sfHR size ($\Delta AIC = 0.0$, AIC weight = 0.319, $df = 6$) (Supplementary materials, Appendix 1, Table A2 & A4). During the winter period, the results were partially in accordance with our expectations. Firstly, in accordance with our second prediction, resident individuals occupied 1.5 fewer sfHR in the most predictable study site compared to the most unpredictable one. But contrary to this prediction, migrants tended to occupy one more sfHR in the most predictable study site compared to the most unpredictable one. However, in accordance with our third prediction, except in the most predictable study site where residents and migrants occupied the same number of sfHR, a resident individual used more sfHR on average than a migrant individual (Fig. 4a). Secondly, the number of sfHR that individuals used decreased as the median size of the sfHR they occupied increased. Indeed, an individual that occupied a large (sfHR size = 410 ha) sfHR had 4 fewer sfHR than an individual that occupied a small (sfHR size = 22 ha) sfHR (Fig. 4b).

The most parsimonious model describing variation in the number of sfHR during the summer period included sex and the log-transformed sfHR size ($\Delta AIC = 0.0$, AIC weight = 0.167, $df = 4$) (Supplementary materials, Appendix 1, Table A2 & A4). During the summer period, the difference in the number of sfHR occupied by males and females was low. Indeed, males used, on average, 0.4 sfHR more than females (Fig. 4c). Finally, as above, the number

of sfHR that individuals used decreased as the median size of the sfHR they occupied increased. Indeed, an individual that occupied a large (sfHR size = 450 ha) sfHR had 2 sfHR less than an individual that occupied a small (sfHR size = 20 ha) sfHR (Fig. 4c).

Spatial separation among sub-seasonal functional home ranges

The most parsimonious model describing variation in the log-transformed distance among sfHR during the wintering period included the two two-way interactions between resource heterogeneity and movement tactic and between resource heterogeneity and sex, with the additive effect of the log-transformed sfHR size ($\Delta AIC = 0.6$, AIC weight = 0.274, $df = 9$) (Supplementary materials, Appendix 1, Table A3 & A4). First, in accordance with our first prediction, spatial separation between pairs of sfHR during the wintering period increased with increasing resource heterogeneity, except among resident females (Fig. 5a and b). Indeed, when resource heterogeneity was high (resource heterogeneity = 0.26), the distance among sfHR of an individual was, on average, 700 meters (365 meters for resident males and 1050 meters for migratory males), while for an individual living in the least heterogeneous study site (resource heterogeneity = 0.08) it was, on average, 300 meters. However, for resident females, this distance was 350 meters lower when resource heterogeneity was high. Finally, spatial separation among sfHR during the winter period was 3980 meters for an individual which occupied the largest sfHR (median sfHR size = 410 ha) compared to 80 meters for an individual that occupied the smallest sfHR (median sfHR size = 22 ha) (Fig. 5c).

The best model describing variation in the log-transformed inter-sfHR distance during the summer period included resource predictability in addition to sex and the median log-transformed sfHR size ($\Delta AIC = 1.86$, AIC weight = 0.143, $df = 6$) (Supplementary materials, Appendix 1, Table A3 & A4). Firstly, contrary to our second prediction, the distance between sfHR was 384 meters in the most predictable study site compared to 200 meters in the most

unpredictable one (Fig. 6a). Secondly, the distance between sfHR was 308 meters for females and 177 meters for males (Fig. 6b). Finally, the spatial separation among sfHR during the summer period was 4450 meters for an individual inhabiting the largest sfHR (median sfHR size = 450 ha) compared to 76 meters for an individual inhabiting the smallest sfHR (median sfHR size = 20 ha) (Fig. 6c).

Discussion

Animals are considered sedentary when their routine movements are centred on revisited areas (Papi 1992), leading to the emergence of a stable home range which may be occupied for a season, or for several years (Börger et al. 2008). Sedentary behaviour is a defining feature of the resident movement tactic, however, migratory animals may also be seasonally sedentary within each of their distinct seasonal ranges (Börger et al. 2008; Mueller and Fagan 2008; Van Moorter et al. 2009). Here, we focused on movements at the within-season scale, when many large herbivores (Börger et al. 2008) and, in particular, roe deer (Hewison et al. 1998), are presumed to be sedentary. Based on a comprehensive analysis of movement behaviour of 15 populations across Europe, we demonstrated that roe deer are never truly sedentary at the seasonal scale. Instead, within any given season, both migratory individuals and residents occupied at least two (and up to nine) spatially distinct sfHR. We suggest that this constitutes an overlooked movement mode, the multi-range tactic, which allows large herbivores to track fine-scaled spatio-temporal variation in the distribution of available resources and, thereby, to cope with changing environmental conditions. Indeed, we found that this space-use behaviour varied in relation to variation in environmental conditions across the European continent.

At the annual scale, we were able to assign a given individual to either a residency or a migratory tactic. However, when we analysed space use behaviour at the finer temporal scale

within seasons, we found that all roe deer in this study adopted the multi-range tactic during a given season. A similar pattern of space use behaviour has been documented in two African large herbivores, the sable antelope and the African savannah buffalo (Owen-Smith et al. 2010; Cornélis et al. 2011; Benhamou 2014). These authors showed that the ranges of these animals were composed of several distinct areas which were exploited for several days or weeks. Thus, the use of multiple sfHR to track available food resources seems to be potentially widespread among large herbivores. Here, to understand the proximal drivers of this seasonal space use behaviour, we explored how spatial heterogeneity and temporal predictability of resource distribution influenced variation in the number and the spatial distribution of the sub-seasonal ranges that an individual exploits.

Mueller et al. (2011), focusing on movements at the annual scale, documented longer seasonal migrations among species inhabiting areas where large-scale primary productivity was spatially heterogeneous, but shorter migrations for species inhabiting environments with relatively low spatial heterogeneity. Similarly, but at the seasonal scale, we expected individuals to be less sedentary in heterogeneous environments than in homogeneous environments, occupying a higher number of more spatially distant sfHR. Our analyses only partially supported this prediction as, in heterogeneous study areas, the distance between sfHR was indeed higher compared to homogeneous sites, but only during winter. Furthermore, the number of sfHR that deer occupied did not vary in relation to resource heterogeneity. One explanation for this discrepancy could be linked to the relatively coarse spatial resolution of the NDVI metric that limits our ability to quantify small-scale variations in resource distribution at a level that is informative for individual animals. Indeed, Van Moorter et al. (2013) have shown that movements at a particular scale are driven by changes in the net profitability of trophic resources at the corresponding scale. As a result, a finer-scale measure of spatial heterogeneity could help us to better understand why roe deer use a number

of spatially distinct sub-seasonal functional home ranges, even in apparently homogeneous habitats.

Because nomadism is considered to be a response to unpredictability (Mueller and Fagan 2008), we expected roe deer to be less sedentary in unpredictable environments than in predictable environments. We found little support for this prediction. Our results indicated that, while controlling for resource heterogeneity, resource predictability had only a weak effect on the number of sfHR and the distance among sfHR. During summer, roe deer moved slightly further when in predictable environments than in unpredictable ones. During winter, resident roe deer adopted a space use behaviour that was somewhat similar to a nomadic tactic when in more unpredictable environments, occupying slightly more sfHR than in predictable environments. Migratory individuals, on the contrary, tended to occupy slightly less sfHR in predictable environments than in unpredictable ones. However, these relationships were weak, with the main difference being that resident individuals occupied more sfHR than migrants during winter. Seasonal migration is a tactic designed to cope with seasonal changes in the spatial distribution of resources (Fryxell and Sinclair 1988; Mueller and Fagan 2008). As a result, it seems that migratory individuals do not have to shift their ranges at the finer within-season scale, as they are able to adjust their movements to spatial variation in the distribution of available forage at the larger between-season scale. Given the lack of support for our predictions, we suggest that other factors such as fine-scale variation in forage plant distribution, predation risk, or human disturbance could drive roe deer to switch periodically from one locality to another. Theory also predicts that prey should perform frequent random movements to minimise the probability of encountering predators (Mitchell and Lima 2002). Indeed, previous studies have shown that herbivores modify their habitat selection or switch location after an encounter with a predator (Latombe et al. 2014; Lone et al. 2016).

From the point of view of landscape complementation (Dunning et al. 1992), an animal's home range must necessarily contain a combination of all non-substitutable resources required for survival and reproduction. Indeed, variation in home range size has previously been shown to reflect variation in resource availability, integrating interactions among local weather, climate and environmental seasonality (Morellet et al. 2013). Hence, all things being equal, animals will occupy larger home ranges when resources are more sparsely distributed in space (Saïd et al. 2005; Van Beest et al. 2011). Our analyses demonstrated strong relationships between sub-seasonal range size with both the number of sfRH and the distance between them. These results indicate that when resources are abundant, deer sequentially exploit a number of short-term functional ranges that are within close proximity. In contrast, when resources are limiting, deer tend to relocate less frequently, possibly due to the costs of doing so in terms of predation risk, mortality or energy (Hein et al. 2012; Johansson et al. 2014), but move greater distances when they do so to locate a new functional sub-seasonal range. This individual variation in space use is further demonstration of the extensive behavioural plasticity that roe deer express in relation to prevailing environmental conditions (Cagnacci et al. 2011; Morellet et al. 2013; Lone et al. 2016) which has driven the undoubted recent success story of this species across Europe (Linnell et al. 1998).

In conclusion, we suggest that the multi-range tactic is an individual-level behavioural response to cope with spatio-temporal variation in the distribution of resources at the within seasonal scale. This tactic appears to be ubiquitous in the roe deer, occurring across its entire European distribution, and encompassing a wide gradient of environmental conditions. We suggest that large herbivores may adopt this finer-scale tactic when environmental conditions fluctuate spatially and temporally, independently of seasonal variations. In the present context of climate change, predictions of more frequent and intense climatic events (IPCC 2014) may mean that an increasing number of large herbivore populations adopt the multi-range tactic,

combining more frequent spasmodic movements interspersed with short periods of
sedentarism, to track available resources.

Acknowledgments

This study was conducted within the EURODEER collaborative project. The co-authors are
grateful to all members for their support for the initiative. The EURODEER data base is
hosted by Fondazione Edmund Mach. For France, we would like to thank the local hunting
associations and the Fédération Départementale des Chasseurs de la Haute Garonne for
allowing us to work in the Comminges, as well as numerous co-workers and volunteers for
their assistance. The Norwegian data collection was funded by the Norwegian Environment
Agency and the county administration of Buskerud county. J Linnell was also funded by the
Research Council of Norway (grants 212919 and 251112). Financial support for GPS data
collection in the Bavarian Forest was provided by the EU-programme INTERREG IV (EFRE
Ziel 3) and the Bavarian Forest National Park Administration. We also thank two anonymous
reviewers for their constructive comments on an earlier version of this manuscript.

Ethical approval: All applicable institutional and/or national guidelines for the care and
use of animals were followed.

Data accessibility

Data used for this study are accessible on EURODEER website (<http://eurodeer.org>) on
demand.

References

Baguette M, Stevens VM, Clobert J (2014) The pros and cons of applying the movement

ecology paradigm for studying animal dispersal. *Mov Ecol* 2:2:13.

Barraquand F, Benhamou S (2008) Animal movements in heterogeneous landscapes: identifying profitable places and homogeneous movement bouts. *Ecology* 89:3336–3348.

Benhamou S (2014) Of scales and stationarity in animal movements. *Ecol Lett* 17:261–272.

Bischof R, Loe LE, Meisingset EL, et al (2012) A migratory northern ungulate in the pursuit of spring: jumping or surfing the green wave? *Am Nat* 180:407–424.

Börger L, Dalziel BD, Fryxell JM (2008) Are there general mechanisms of animal home range behaviour? A review and prospects for future research. *Ecol Lett* 11:637–650.

Börger L, Fryxell JM (2012) Quantifying individual differences in dispersal using net squared displacement. In: Clobert J, Baguette M, Benton TJ, Bullock J (eds) *Dispersal Ecology and Evolution*, Oxford Uni. pp 222–230

Borowik T, Pettorelli N, Sönichsen L, Jedrzejewska B (2013) Normalized difference vegetation index (NDVI) as a predictor of forage availability for ungulates in forest and field habitats. *Eur J Wildl Res* 59:675–682.

Bowler DE, Benton TG (2005) Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. *Biol Rev* 80:205–225.

Bracis C, Gurarie E, Van Moorter B, Goodwin RA (2015) Memory effects on movement behavior in animal foraging. *PLoS One* 10:1–21.

Brown JL, Orians GH (1970) Spacing patterns in mobile animals. *Annu Rev Ecol Syst* 1:239–262.

Bunnefeld N, Börger L, van Moorter B, et al (2011) A model-driven approach to quantify migration patterns: individual, regional and yearly differences. *J Anim Ecol* 80:466–476.

Burnham KP, Anderson DR (2002) *Model selection and multimodel inference: A practical information-theoretic approach* (2nd ed). Springer

Cagnacci F, Focardi S, Ghisla A, et al (2015) How many routes lead to migration?

538 Comparison of methods to assess and characterise migratory movements. *J Anim Ecol*
539 85:54–68.

540 Cagnacci F, Focardi S, Heurich M, et al (2011) Partial migration in roe deer: migratory and
541 resident tactics are end points of a behavioural gradient determined by ecological factors.
542 *Oikos* 120:1790–1802.

543 Chapin FS, Johnson DA, McKendrick JD (1980) Seasonal movement of nutrients in plants of
544 differing growth form in an Alaskan tundra ecosystem: implications for herbivory. *J Ecol*
545 68:189–209.

546 Chapman BB, Hulthén K, Wellenreuther M, et al (2014) Patterns of animal migration. In:
547 Hansson L-A, Åkesson S (eds) *Animal movement across scales*. Oxford University
548 Press, pp 11–35

549 Charnov EL (1976) Optimal foraging, the marginal value theorem. *Theor. Popul. Biol.* 9:129–
550 136.

551 Colwell RK (1974) Predictability, constancy, and contingency of periodic phenomena.
552 *Ecology* 55:1153.

553 Coops N, Culvenor D (2000) Utilizing local variance of simulated high spatial resolution
554 imagery to predict spatial pattern of forest stands. *Remote Sens Environ* 71:248–260.

555 Coops N, Culvenor D, Preston R, Catling P (1998) Procedures for predicting habitat and
556 structural attributes in eucalypt forests using high spatial resolution remotely sensed
557 imagery. *Aust For* 61:244–252.

558 Cornélis D, Benhamou S, Janeau G, et al (2011) Spatiotemporal dynamics of forage and water
559 resources shape space use of West African savanna buffaloes. *J Mammal* 92:1287–1297.

560 Dunning JB, Danielson BJ, Pulliam HR (1992) Ecological processes that affect populations in
561 complex landscapes. *Oikos* 65:169–175.

562 English AK, Chauvenet ALM, Safi K, Pettorelli N (2012) Reassessing the determinants of

563 breeding synchrony in Ungulates. PLoS One 7:e41444.

564 Fryxell JM, Sinclair ARE (1988) Causes and consequences of migration by large herbivores.

565 Trends Ecol Evol 3:237–241.

566 Garrouste EL, Hansen AJ, Lawrence RL (2016) Using NDVI and EVI to map spatiotemporal

567 variation in the biomass and quality of forage for migratory elk in the Greater

568 Yellowstone Ecosystem. Remote Sens. doi: 10.3390/rs8050404

569 Gurarie E, Cagnacci F, Peters W, et al (2017) A framework for modeling range shifts and

570 migrations: Asking whether, whither, when, and will it return. J Anim 943–959. doi:

571 10.1111/ijlh.12426

572 Haight FA (1967) Handbook of the Poisson distribution. Wiley

573 Hamel S, Garel M, Festa-Bianchet M, et al (2009) Spring Normalized Difference Vegetation

574 Index (NDVI) predicts annual variation in timing of peak faecal crude protein in

575 mountain ungulates. J Appl Ecol 46:582–589. doi: 10.1111/j.1365-2664.2009.01643.x

576 Hein AM, Hou C, Gillooly JF (2012) Energetic and biomechanical constraints on animal

577 migration distance. Ecol Lett 15:104–110. doi: 10.1111/j.1461-0248.2011.01714.x

578 Hewison AJM, Vincent J-P, Reby D (1998) Social organisation of European roe deer. In:

579 Andersen R, Duncan P, Linnell JDC (eds) The European roe deer: The Biology of

580 Success. Scandinavian University Press, Oslo, pp 189–219

581 Huang S, Meng SX, Yang Y (2009) Assessing the goodness of fit of forest models estimated

582 by nonlinear mixed-model methods. Can J For Res 39:2418–2436.

583 IPCC (2014) Climate change 2014: Synthesis Report.

584 Jepsen JU, Topping CJ (2004) Modelling roe deer (*Capreolus capreolus*) in a gradient of

585 forest fragmentation: behavioural plasticity and choice of cover. Can J Zool 82:1528–

586 1541. doi: Doi 10.1139/Z04-131

587 Johansson CL, Muijres FT, Hedenström A (2014) The physics of animal locomotion. In:

588 Hansson L-A, Åkesson S (eds) Animal movement across scales. Oxford University
589 Press, pp 232–258

590 Kerr JT, Ostrovsky M (2003) From space to species: Ecological applications for remote
591 sensing. *Trends Ecol Evol* 18:299–305.

592 Klisch A, Atzberger C (2016) Operational drought monitoring in Kenya using MODIS NDVI
593 time series. *Remote Sens* 8:267.

594 Latombe G, Fortin D, Parrott L (2014) Spatio-temporal dynamics in the response of woodland
595 caribou and moose to the passage of grey wolf. *J Anim Ecol* 83:185–198.

596 Lavielle M (1999) Detection of multiple changes in a sequence of dependent variables. *Stoch*
597 *Process their Appl* 83:79–102.

598 Lendrum PE, Anderson CR, Monteith KL, et al (2014) Relating the movement of a rapidly
599 migrating ungulate to spatiotemporal patterns of forage quality. *Mamm Biol* 79:369–375.

600 Linnell JDC, Duncan P, Andersen R (1998) The European roe deer: A portrait of a successful
601 species. In: Andersen R, Duncan P, Linnell JDC (eds) *The European roe deer: The*
602 *Biology of Success*. Scandinavian University Press, Oslo, pp 11–22

603 Lone K, Mysterud A, Gobakken T, et al (2016) Temporal variation in habitat selection breaks
604 the catch-22 of spatially contrasting predation risk from multiple predators. *Oikos*
605 126:624–632.

606 Mitchell WA, Lima SL (2002) Predator-prey shell games: large-scale movement and its
607 implications for decision-making by prey. *Oikos* 99:249–259.

608 Morellet N, Bonenfant C, Börger L, et al (2013) Seasonality, weather and climate affect home
609 range size in roe deer across a wide latitudinal gradient within Europe. *J Anim Ecol*
610 82:1326–1339.

611 Mueller T, Fagan WF (2008) Search and navigation in dynamic environments - from
612 individual behaviors to population distributions. *Oikos* 117:654–664.

613 Mueller T, Olson KA, Dressler G, et al (2011) How landscape dynamics link individual- to
 614 population-level movement patterns: a multispecies comparison of ungulate relocation
 615 data. *Glob Ecol Biogeogr* 20:683–694.

616 Nathan R, Getz WM, Revilla E, et al (2008) A movement ecology paradigm for unifying
 617 organismal movement research. *Proc Natl Acad Sci* 105:19052–19059.

618 Owen-Smith N, Fryxell JM, Merrill EH (2010) Foraging theory upscaled: the behavioural
 619 ecology of herbivore movement. *Philos Trans R Soc London* 365:2267–2278.

620 Papi F (1992) *Animal homing*. Chapman & Hall

621 Peters W, Hebblewhite M, Musterud A, et al (2017) Migration in geographic and ecological
 622 space by a large herbivore. *Ecol Monogr* 87:297–320.

623 Pettorelli N, Ryan S, Mueller T, et al (2011) The Normalized Difference Vegetation Index
 624 (NDVI): Unforeseen successes in animal ecology. *Clim Res* 46:15–27. doi:
 625 10.3354/cr00936

626 Pyke GH (1984) Optimal foraging theory: a critical review. *Annu Rev Ecol Syst* 15:523–575.

627 R Development Core Team (2014) *R: a language and environment for statistical computing*.

628 Revilla E, Wiegand T (2008) Individual movement behavior, matrix heterogeneity, and the
 629 dynamics of spatially structured populations. *Proc Natl Acad Sci USA* 105:19120–
 630 19125.

631 Riotte-Lambert L, Benhamou S, Chamaillé-Jammes S (2015) How Memory-Based Movement
 632 Leads to Nonterritorial Spatial Segregation. *Am Nat* 185:E103-16.

633 Rivrud IM, Loe LE, Myrsetrud A (2010) How does local weather predict red deer home range
 634 size at different temporal scales? *J Anim Ecol* 79:1280–1295.

635 Saïd S, Gaillard J-M, Duncan P, et al (2005) Ecological correlates of home-range size in
 636 spring–summer for female roe deer (*Capreolus capreolus*) in a deciduous woodland. *J*
 637 *Zool* 267:301–308.

638 Spitz DB, Hebblewhite M, Stephenson TR (2017) “MigrateR”: extending model-driven
639 methods for classifying and quantifying animal movement behavior. *Ecography* (Cop)
640 40:788–799.

641 Turchin P (1991) Translating foraging movements in heterogeneous environments into the
642 spatial distribution of foragers. *Ecology* 72:1253–1266.

643 Turchin P (1998) Quantitative analysis of movement: measuring and modeling population
644 redistribution in animals and plants. Sinauer Associates, Sunderland, Massachusetts,
645 USA

646 Van Beest FM, Rivrud IM, Loe LE, et al (2011) What determines variation in home range
647 size across spatiotemporal scales in a large browsing herbivore? *J Anim Ecol* 80:771–
648 785.

649 Van Moorter B, Bunnefeld N, Panzacchi M, et al (2013) Understanding scales of movement:
650 animals ride waves and ripples of environmental change. *J Anim Ecol* 82:770–780.

651 Van Moorter B, Rolandsen CM, Basille M, Gaillard J-M (2016) Movement is the glue
652 connecting home ranges and habitat selection. *J Anim Ecol* 85:21–31.

653 Van Moorter B, Visscher D, Benhamou S, et al (2009) Memory keeps you at home: a
654 mechanistic model for home range emergence. *Oikos* 118:641–652.

655 Vanpé C, Gaillard J, Morellet N, et al (2009) Age-Specific Variation in Male Breeding
656 Success of a Territorial Ungulate Species, the European Roe Deer. *J Mammal* 90:661–
657 665.

658 Vuolo F, Mattiuzzi M, Klisch A, Atzberger C (2012) Data service platform for MODIS
659 Vegetation Indices time series processing at BOKU Vienna: current status and future
660 perspectives. *SPIE Proc* 8538:85380A.

661 Wiens JA (1976) Population responses to patchy environments. *Annu Rev Ecol Syst* 7:81–
662 120.

663

Tables

Table 1: Number of males, females, migratory individuals, and residents per study site.

Study site id	Study site name	Migrants	Residents	Females	Males
a	Aspromonte - Italy	4	2	4	2
b	Aurignac - France	20	86	66	40
c	Baden-Wurttemberg - Germany	5	19	22	2
d	Bavarian Forest National Park - Germany	28	35	36	27
e	Šumava National Park – Czech republic	0	2	1	1
f	Bernese Oberland - Switzerland	9	15	19	5
g	Bialowieza National Park - Poland	2	5	7	0
h	Brandenburg - Germany	2	0	2	0
i	Grimso - Sweden	1	0	0	1
j	Koberg - Sweden	5	9	10	4
k	Leoben - Austria	4	4	5	3
l	Matese - Italy	0	3	3	0
m	Monte Bondone - Italy	4	4	4	4
n	Rendena Giudicarie – Italy	6	3	6	3
o	Østafjells west - Norway	3	6	5	4
Total		93	193	190	96

Figure legends

Fig. 1 Location of study sites where roe deer were monitored using GPS collars. Letters correspond to the 'study site id' in Table 1, with the corresponding number of individuals in italics

Fig. 2 Example of the segmentation approach using Lavielle's Method (1999) on the movement trajectory of a migratory individual (individual 1367 monitored in 2008) during the winter and the summer seasons (top) and the corresponding sub-seasonal functional home ranges (sfHR) (bottom). The X-axis indicates the chronological order of successive locations and the Y-axis shows the Net Squared Displacement (NSD). Vertical lines correspond to cut-off points defining the sequential sfHR

Fig. 3 Example of the segmentation approach using Lavielle's Method (1999) on the movement trajectory of a resident individual (individual 2296 monitored in 2013), during the winter and the summer seasons (top) and the corresponding sub-seasonal functional home ranges (sfHR) (bottom). The X-axis indicates the chronological order of successive locations and the Y-axis shows the Net Squared Displacement (NSD). Vertical lines correspond to cut-off points defining the sequential sfHR

Fig. 4 Graphical representation of the best models describing variation in the number of sub-seasonal home ranges (sfHR) during the winter season, as a function of the two-way interaction between resource predictability and movement tactic (a) and the log-transformed sfHR size (b) ; and during the summer season as a function of sex and the log-transformed sfHR size (c). In order to better visualize the raw data, we plotted the mean number of sfHR for each study site. Dotted lines represent the 95% confidence intervals resulting from the model predictions, error bars represent the standard deviation in number of sfHR, and the letters correspond to the 'study site id' in Table 1. We fixed movement tactic as resident,

resource predictability and the log-transformed sfHR size at their mean when not represented in the figures

Fig. 5 Graphical representation of the best model describing variation in the log-transformed distance among sub-seasonal home ranges (sfHR) during the winter, as a function of the two-way interaction between resource heterogeneity and movement tactic and between resource heterogeneity and sex, represented as a three-way interaction (a: females, b: males), and the log-transformed median sfHR size (c). In order to better visualize the raw data, we plotted the mean log-transformed sfHR distance for each study site. Dotted lines represent the 95% confidence intervals resulting from the model predictions, error bars represent the standard deviation of log-transformed inter-sfHR distance, and the letters correspond to the ‘study site id’ in Table 1. We fixed resource heterogeneity and the log-transformed sfHR size at their mean, movement tactic as resident and sex as female when not represented in the figures.

Fig. 6 Graphical representation of the best model describing variation in the log-transformed distance among sub-seasonal home ranges (sfHR) during the summer season, as a function of resource predictability (a), sex (b) and the log-transformed median sfHR size (c). In order to better visualize the raw data, we plotted the mean log-transformed sfHR distance for each study site. Dotted lines represent the 95% confidence intervals resulting from the model predictions, error bars represent the standard deviation of log-transformed inter-sfHR distance, and the letters correspond to the ‘study site id’ in Table 1. We fixed resource predictability and the log-transformed sfHR size at their mean and sex as female when not represented in the figures

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5








Fig. 6







2296:2013 summer



2296:2013 winter

