



## Chemical probes of turbulence in the diffuse medium: the TDR model (Corrigendum)

B. Godard, E. Falgarone, G. Pineau Des Forêts

### ► To cite this version:

B. Godard, E. Falgarone, G. Pineau Des Forêts. Chemical probes of turbulence in the diffuse medium: the TDR model (Corrigendum). *Astronomy & Astrophysics - A&A*, 2016, 593, pp.C1. 10.1051/0004-6361/201423526e . hal-02318125

HAL Id: hal-02318125

<https://hal.science/hal-02318125v1>

Submitted on 17 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Chemical probes of turbulence in the diffuse medium: the TDR model (*Corrigendum*)

B. Godard<sup>1</sup>, E. Falgarone<sup>1</sup>, and G. Pineau des Forêts<sup>2,1</sup>

<sup>1</sup> LERMA, CNRS UMR 8112, École Normale Supérieure & Observatoire de Paris, 75005 Paris, France  
e-mail: benjamin.godard@obspm.fr

<sup>2</sup> Institut d’Astrophysique Spatiale, CNRS UMR 8617, Université Paris-Sud, 91405 Orsay, France

A&A 570, A27 (2014), DOI: [10.1051/0004-6361/201423526](https://doi.org/10.1051/0004-6361/201423526)

**Key words.** astrochemistry – turbulence – ISM: molecules – ISM: kinematics and dynamics – ISM: structure – errata, addenda

We have noticed two typos in our original paper. In Sect. 3.1, Eq. (2), which gives the expression of the effective temperature of an ion-neutral reaction, should read (Pineau des Forêts et al. 1986)

$$T_{\text{eff}} = \frac{m_1 T_2 + m_2 T_1}{m_1 + m_2} + \frac{1}{3k} \frac{m_1 m_2}{m_1 + m_2} u_{\text{D}}^2. \quad (2)$$

In Sect. 5.2, the destruction timescale of the  $\text{CH}^+$  by reaction with  $\text{H}_2$  should read  $\sim 1$  yr  $f_{\text{H}_2}^{-1}$  ( $50 \text{ cm}^{-3}/n_{\text{H}}$ ). These correct

expressions are those that have been used in all the calculations performed in our original paper. These typos have therefore no bearing on the paper results/conclusions.

## References

- Pineau des Forêts, G., Flower, D. R., Hartquist, T. W., & Dalgarno, A. 1986,  
[MNRAS, 220, 801](https://doi.org/10.1080/002612386182101)