



HAL
open science

Pharmacodynamic modelling of β -lactam/ β -lactamase inhibitor checkerboard data: illustration with aztreonam–avibactam

Alexia Chauzy, Julien Buyck, B.L.M. de Jonge, Sandrine Marchand, N. Grégoire, W. Couet

► To cite this version:

Alexia Chauzy, Julien Buyck, B.L.M. de Jonge, Sandrine Marchand, N. Grégoire, et al.. Pharmacodynamic modelling of β -lactam/ β -lactamase inhibitor checkerboard data: illustration with aztreonam–avibactam. *Clinical Microbiology and Infection*, 2019, 25 (4), pp.515.e1-515.e4. 10.1016/j.cmi.2018.11.025 . hal-02321069

HAL Id: hal-02321069

<https://hal.science/hal-02321069>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

**Pharmacodynamic modeling of β -lactam / β -lactamase inhibitor checkerboard data:
illustration with aztreonam-avibactam**

Alexia Chauzy¹, Julien Buyck¹, Boudewijn L.M. de Jonge², Sandrine Marchand^{1,3}, Nicolas Grégoire¹,
William Couet^{1,3#}

INSERM, U1070, UFR de Médecine Pharmacie, Université de Poitiers, 1 rue Georges Bonnet, TSA
51106, 86073 Poitiers Cedex 9, France¹; Pfizer Essential Health, New York, USA²; Laboratoire de
Toxicologie-Pharmacocinétique, CHU de Poitiers, 2 rue de la Milétrie, 86000 Poitiers, France³

Intended category: Research note

Corresponding author:

William Couet

Université de Poitiers

INSERM U1070 « Pharmacologie des Anti-Infectieux »

Pôle Biologie Santé

1 rue Georges Bonnet – BP 633

86022 Poitiers cedex, France

william.couet@univ-poitiers.fr

+33 (0)5 49 45 43 79 (telephone)

+33 (0) 5 49 45 43 78 (fax)

Abstract

Objectives: Checkerboard experiments followed by Fractional Inhibitory Concentration (FIC) index determinations are commonly used to assess *in vitro* pharmacodynamic interactions between combined antibiotics, but FIC index cannot be determined in case of antibiotic/non-active compound combinations. The aim of this study was to use a simple modeling approach to quantify the *in vitro* activity of aztreonam-avibactam, a new β -lactam- β -lactamase inhibitor combination.

Methods: MIC checkerboard experiments were performed with 12 *Enterobacteriaceae* with diverse β -lactamases profiles. Aztreonam MICs in the absence and presence of avibactam at different concentrations (ranging from 0.0625 to 4 mg/L) were determined. Aztreonam MIC versus avibactam concentrations were fitted by an inhibitory E_{\max} model with a baseline effect parameter.

Results: A concentration-dependent relationship was observed with a steep initial reduction of aztreonam MIC at low avibactam concentrations and reaching a maximum at higher avibactam concentrations, that was adequately fitted by the model. Maximum avibactam effect was characterized by the ratio of aztreonam MICs in the absence of avibactam (MIC_0) and when avibactam concentration tends toward infinity (MIC_{∞}), and this ratio ranged between 90 and 10068 depending on the strain. Avibactam potency was characterized by avibactam concentrations corresponding to 50% of the maximum effect (IC_{50} values between 0.00022 and 0.053 mg/L).

Conclusions: An inhibitory E_{\max} model with a baseline effect could quantify maximum avibactam effect and potency among various strains. This simple modeling approach can be used to compare the activity of other combinations of antibiotics with non-antibiotic drugs when FIC index is inappropriate.

Keywords: Checkerboard, Combinations, β -lactam antibiotic, β -lactamase inhibitor

Introduction

Infections caused by carbapenem-resistant *Enterobacteriaceae* (CRE) represent a major threat to human health worldwide (1). These pathogens are resistant to most prescribed antibiotics due to the production of various types of β -lactamases and are extremely hard to eradicate. Combining a β -lactam antibiotic with a β -lactamase inhibitor to inactivate β -lactamase activity constitutes an interesting option for treating CRE infections (2). Aztreonam-avibactam is one of the new β -lactam- β -lactamase inhibitor combinations under development. Aztreonam is a monobactam antibiotic which is not hydrolyzed by metallo- β -lactamases (MBLs) that inactivate most β -lactam antibiotics (3). Yet it is hydrolyzed by some serine β -lactamases, such as extended-spectrum β -lactamases (ESBLs) and serine carbapenemases including KPC (3). However, these serine β -lactamases are efficiently inhibited by the β -lactamase inhibitor avibactam (4). Therefore, combining aztreonam with avibactam is expected to restore the clinical utility of aztreonam against CRE.

Pharmacodynamic interaction between antibiotics used in combination is usually investigated by performing checkerboard experiments followed by Fractional Inhibitory Concentration (FIC) index determinations (5). This is based on the comparison of the MIC of each antibiotic alone (MIC_A and MIC_B for antibiotics A and B, respectively) with the combination-derived MICs ($MIC_{A/B}$ and $MIC_{B/A}$) (Equation 1).

$$FIC\ index = \frac{MIC_{A/B}}{MIC_A} + \frac{MIC_{B/A}}{MIC_B}$$

(Equation 1)

It is a simple approach that can be used for rapid screening of antibiotic combinations in the presence of various strains. However, if an antibiotic is combined with a non-antibiotic compound, such as a β -lactamase inhibitor, the FIC index cannot be determined since the non-antibiotic drug has *a priori* no antimicrobial activity and cannot be characterized by an MIC value. Antimicrobial activity has been reported for non-antibiotic compounds, but with MIC values much higher than concentrations clinically achievable (6, 7). Accordingly, antibiotic/non-antibiotic combinations have been defined as synergistic (27th European Congress of Clinical Microbiology and Infectious Diseases, abstract P0655) but for non-antibiotic drug concentrations corresponding to toxic concentrations (8).

The objective of the present study was to use a simple but meaningful modeling approach for the analysis of checkerboard-type data obtained with β -lactam- β -lactamase inhibitor combinations, or any combination of an active antibiotic with a non-active compound, based on the E_{max} model previously used in that context (9) or to describe the pro-convulsant effect of biphenyl acetic acid on fluoroquinolones (10).

Methods

Chemicals. Aztreonam (Sigma-Aldrich, St Quentin Fallavier, France) and avibactam (provided as a dry powder by AstraZeneca, Macclesfield, United Kingdom) were used to prepare stock solutions of 50 mg/mL of aztreonam in methanol/dimethyl sulfoxide (50/50, v/v) and 1 mg/mL of avibactam in sterile water. Aztreonam and avibactam stock solutions were further diluted in cation-adjusted Mueller-Hinton broth (CAMHB).

Bacterial strains. Twelve *Enterobacteriaceae* strains, resistant to aztreonam, including 4 *Klebsiella pneumoniae*, 2 *Escherichia coli*, 2 *Citrobacter freundii*, 3 *Enterobacter cloacae* and 1 *Enterobacter aerogenes* were studied (Table 1). The β -lactamase content of each strain was characterized by IHMA laboratory (International Health Management Associates, Inc., Schaumburg, IL, USA) through PCR.

In vitro susceptibility testing. For each bacterial strain, MICs of aztreonam and avibactam alone were determined according to the EUCAST guidelines for broth microdilutions (11). Aztreonam MICs in the presence of avibactam were also determined using checkerboard method. Checkerboards were set up with 2-fold dilutions of aztreonam (0.0078 to 512 mg/L) and avibactam (0.0625 to 4 mg/L) in such a way that different combinations of aztreonam and avibactam concentrations were obtained in each well. After drug dilution, 96-well plates were inoculated with each organism to yield the appropriate density (0.5×10^6 CFU/mL) and incubated for 16 to 20 h at 37°C. The MIC was recorded as the lowest concentration of antimicrobial agent that completely inhibited visible growth of the organism in microdilution wells. Negative growth controls were performed in wells containing only

CAMHB. All experiments were performed in duplicate, second replicate was used to validate the first one. Only the first replicate was used for data analysis.

Data analysis. The antibacterial effect of aztreonam in combination with avibactam was investigated by modeling the interaction between both compounds, using WinNonlin software (version 6.2, Certara, Princeton, NJ, USA). Aztreonam MICs versus avibactam concentrations were fitted according to an inhibitory E_{max} model with a baseline effect parameter (Equation 2). A Gauss-Newton minimization method with the Levenberg and Hartley modification was used for data fitting. Data were not log-transformed before analysis and we used the weighted least squares method for minimization based on observations (Y_{obs_i}) i.e. the weight applied to each observation (w_i) was: $w_i = 1/(Y_{obs_i})^2$.

$$MIC_{C_{AVI}} = MIC_0 - \frac{(MIC_0 - MIC_{\infty}) \times C_{AVI}}{(C_{AVI} + IC_{50})}$$

(Equation 2)

Where, MIC_0 is the MIC of aztreonam in the absence of avibactam and MIC_{∞} the asymptotic value of MIC when avibactam concentration (C_{AVI}) tends toward infinity. MIC_0/MIC_{∞} ratio characterizes avibactam maximum effect, traditionally referred as efficacy in the E_{max} model and expressed here as a percentage, while IC_{50} , the avibactam concentration corresponding to 50% of the maximum inhibitory effect, characterizes the potency of avibactam. Then, avibactam IC_{90} , corresponding to the avibactam concentration which produces 90% of the maximum effect, was derived from the model for the different strains studied.

Results

The inhibitory E_{max} model with a baseline effect parameter adequately described the steep initial decay of aztreonam MIC in the presence of very low avibactam concentrations before reaching a plateau at higher avibactam concentrations, as shown in the individual plots (Figure 1). Weighted residuals were evenly distributed around 0. However, for *E. coli* 1266865, a slightly different profile was observed, with a much shallower decrease of aztreonam MIC in the presence of increasing avibactam concentrations and no plateau within this range of avibactam concentrations. For the other strains,

estimated MIC_0 were close to measured values, maximum avibactam effect (MIC_0/MIC_{∞}) ranged between roughly 100 and 10 000, and potency (IC_{50}) between 0.00022 and 0.053 mg/L (Table 1).

Discussion

An inhibitory E_{max} model with a baseline effect parameter successfully described the effect of avibactam on aztreonam MIC in all but one strain. The presence of avibactam dramatically reduced aztreonam MIC, up to roughly 100 to 10 000-fold and starting at low concentrations (most IC_{50} values \ll 0.1 mg/L). Because avibactam possesses some intrinsic antibacterial activity at relatively high levels ($MIC \geq 8$ mg/L) (12), the maximum concentration of avibactam used in these checkerboard experiments was 4 mg/L in order to restrict this investigation to the effect of avibactam on aztreonam MIC, presumably due to β -lactamases inhibition. Maximum effect and potency of avibactam were highly variable between strains, reflecting the wide range of β -lactamases produced by the different strains and the difference of affinity of β -lactamases for avibactam (13). However, model-derived aztreonam MIC in the absence of avibactam was higher for *E.coliaceae* 1318536 ($MIC = 510$ mg/L) than for *E.coliaceae* 1285905 ($MIC = 64$ mg/L) although both strains express exactly the same β -lactamases, suggesting differences in efflux or outer membrane permeability as well as in expression levels of enzymes between the two strains.

Compared with FIC index that only concludes to synergism, additivity or antagonism, the E_{max} model can quantify efficacy and potency among various strains or β -lactam- β -lactamase inhibitor combinations and is therefore more informative. This simple E_{max} model was previously used to characterize the activity of an experimental BLI in combination with imipenem for selecting optimal dosing strategies of the combination (9). However, this objective may fall beyond the limits of these checkerboard experiments. Alternative approaches such as time-kill experiments combined with semi-mechanistic pharmacokinetic/pharmacodynamics (PK/PD) modeling to characterize aztreonam-avibactam combination are better suited for that (14).

Yet simple checkerboard experiments analyzed with an E_{max} model seem appropriate for comparing β -lactam- β -lactamase inhibitor combinations efficacy and potency.

Funding

This research project was funded by the Innovative Medicine Initiative Joint undertaking under COMBACTE-CARE grant agreement 115620 resources of which are composed of financial contribution from the European Union Seventh Framework Programme (FP7/2007-2013) and European Federation of Pharmaceutical Industries and Associations (EFPIA) companies in kind contribution. Avibactam dry powder was provided by AstraZeneca.

Acknowledgments

This study was presented in part at the annual meeting of the International Society of Anti-Infective Pharmacology, Vienna, Austria, 2017. We thank Emma Marquizeau for her technical assistance with the checkerboard experiments.

Conflict of interest

Boudewijn de Jonge is currently an employee of Pfizer.

References

- [1] Canton R, Akova M, Carmeli Y, Giske CG, Glupczynski Y, Gniadkowski M, et al. Rapid evolution and spread of carbapenemases among Enterobacteriaceae in Europe. *Clin Microbiol Infect.* 2012;18(5):413-31.
- [2] Perez F, El Chakhtoura NG, Papp-Wallace KM, Wilson BM, Bonomo RA. Treatment options for infections caused by carbapenem-resistant Enterobacteriaceae: can we apply "precision medicine" to antimicrobial chemotherapy? *Expert Opin Pharmacother.* 2016;17(6):761-81.
- [3] Biedenbach DJ, Kazmierczak K, Bouchillon SK, Sahm DF, Bradford PA. In vitro activity of aztreonam-avibactam against a global collection of Gram-negative pathogens from 2012 and 2013. *Antimicrob Agents Chemother.* 2015;59(7):4239-48.
- [4] Livermore DM, Mushtaq S, Warner M, Zhang J, Maharjan S, Doumith M, et al. Activities of NXL104 combinations with ceftazidime and aztreonam against carbapenemase-Producing Enterobacteriaceae. *Antimicrob Agents Chemother.* 2011;55(1):390-4.
- [5] White RL, Burgess DS, Manduru M, Bosso JA. Comparison of three different in vitro methods of detecting synergy: time-kill, checkerboard, and E test. *Antimicrob Agents Chemother.* 1996;40(8):1914-8.
- [6] Martins M, Dastidar SG, Fanning S, Kristiansen JE, Molnar J, Pages JM, et al. Potential role of non-antibiotics (helper compounds) in the treatment of multidrug-resistant Gram-negative infections: mechanisms for their direct and indirect activities. *Int J Antimicrob Agents.* 2008;31(3):198-208.
- [7] Nigam A, Gupta D, Sharma A. Treatment of infectious disease: beyond antibiotics. *Microbiol Res.* 2014;169(9-10):643-51.
- [8] Schulz M, Iwersen-Bergmann S, Andresen H, Schmoldt A. Therapeutic and toxic blood concentrations of nearly 1,000 drugs and other xenobiotics. *Crit Care.* 2012;16(4):R136.

- [9] Bhagunde P, Chang KT, Hirsch EB, Ledesma KR, Nikolaou M, Tam VH. Novel modeling framework to guide design of optimal dosing strategies for beta-lactamase inhibitors. *Antimicrob Agents Chemother.* 2012;56(5):2237-40.
- [10] Marchand S, Pariat C, Boulanger A, Bouquet S, Couet W. A pharmacokinetic/pharmacodynamic approach to show that not all fluoroquinolones exhibit similar sensitivity toward the proconvulsant effect of biphenyl acetic acid in rats. *J Antimicrob Chemother.* 2001;48(6):813-20.
- [11] European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European Society for Clinical Microbiology and Infectious Diseases (ESCMID). 2003. EUCAST Discussion Document E. Dis 5.1: determination of minimum inhibitory concentrations (MICs) of antibacterial agents by broth dilution. *Clin Microbiol Infect.*9:1-7.
- [12] Berkhout J, Melchers MJ, van Mil AC, Nichols WW, Mouton JW. In vitro activity of ceftazidime-avibactam combination in in vitro checkerboard assays. *Antimicrob Agents Chemother.* 2015;59(2):1138-44.
- [13] Ehmann DE, Jahic H, Ross PL, Gu RF, Hu J, Kern G, et al. Avibactam is a covalent, reversible, non-beta-lactam beta-lactamase inhibitor. *Proc Natl Acad Sci U S A.* 2012;109(29):11663-8.
- [14] Sy S, Zhuang L, Xia H, Beaudoin ME, Schuck VJ, Derendorf H. Prediction of in vivo and in vitro infection model results using a semimechanistic model of avibactam and aztreonam combination against multidrug resistant organisms. *CPT Pharmacometrics Syst Pharmacol.* 2017;6(3):197-207.

Figure 1. Aztreonam MICs versus avibactam concentrations (mg/L) for 12 *Enterobacteriaceae* isolates. The circles represent the aztreonam MICs determined during one checkerboard experiment and the solid lines the individual aztreonam MICs predicted by the E_{max} model with baseline effect.

Table 1. Parameter estimates of the inhibitory E_{max} model with baseline effect for 12 resistant *Enterobacteriaceae* isolates

Organism	IHMA No.	β -lactamases	ATM MIC (mg/L)	ATM-AVI MIC ^a (mg/L)	Parameter estimates (RSE%)				
					MIC ₀ (mg/L)	MIC _∞ (mg/L)	MIC ₀ /MIC _∞	AVI IC ₅₀ (mg/L)	AVI IC ₉₀ (mg/L)
<i>Escherichia coli</i>	1266865	TEM-OSBL(b), CMY-42, NDM-5	32	4	ND ^b	ND ^b	ND ^b	ND ^b	ND ^b
	1275629	CTX-M-15, VIM-23	128	0.0625	128 (22)	0.056 (22)	2262	0.00040 (27)	0.0036
<i>Klebsiella pneumoniae</i>	1277372	SHV-12(e), TEM-OSBL(b), CTX-M-15, NDM-7	128	0.0625	128 (20)	0.044 (22)	2894	0.00047 (24)	0.0043
	1289268	SHV-OSBL(b), TEM-OSBL(b), CTX-M-15, VIM-4	128	0.0625	127 (32)	0.033 (50)	3909	0.00064 (38)	0.0058
	1251604	SHV-5(e), VIM-26	256	0.125	256 (33)	0.12 (32)	2178	0.00053 (39)	0.0048
	947566	SHV-12(2be), VIM-42	64	0.125	64 (29)	0.064 (41)	995	0.0018 (36)	0.016
<i>Citrobacter cfreundii</i>	974673	SHV-12(2be), TEM-OSBL(2b), CTX-M-3, CMY-34, NDM-1	512	0.125	511 (32)	0.051 (83)	10068	0.00053 (38)	0.0048
	1080008	VIM-23	128	1	133 (38)	0.23 (166)	580	0.015 (49)	0.14
<i>Enterobacter aerogenes</i>	1286221	TEM-OSBL(b), CTX-M-15, NDM-1	128	0.03125	127 (46)	0.019 (64)	6808	0.00026 (58)	0.0023
<i>Enterobacter cloacae</i>	1285905	CTX-M-15, NDM-1	64	0.25	64 (30)	0.21 (32)	299	0.0037 (38)	0.034
	1318536	CTX-M-15, NDM-1	512	0.125	510 (35)	0.083 (41)	6182	0.00022 (43)	0.0020
	1251704	TEM-OSBL(b), VIM-1	32	1	33 (30)	0.37 (84)	90	0.053 (47)	0.55

^a aztreonam MIC determined in the presence of 4 mg/L avibactam^b The model could not be fitted to the data obtained from this isolateATM, aztreonam; AVI, avibactam; RSE, relative standard error; ND, not determined; OSBL, Original-spectrum β -lactamase; TEM, Temoneira; CMY, cephamycins; NDM, Nex Delhi metallo- β -lactamase; CTX-M, cefotaxime-hydrolyzing β -lactamase; VIM, Verona integrin-encoded metallo- β -lactamase; SHV, sulfhydryl variable

ATM MIC (mg/L)

