

MCSS Update biomolecular fragments

solvent models, scoring functions, ... and application examples to oligonucleotides

Fabrice Leclerc

Institute for Integrative Biology of the Cell (I2BC) CNRS-CEA-Univ. Paris Sud, Univ. Paris Saclay <u>fabrice.leclerc@u-psud.fr</u>

Fragment-Based Ligand Design



Fragment-Based Ligand Design



Fragment-Based Ligand Design



Success stories in FBLD



"SAR by NMR"

Shuker et al., 1996

Metalloproteinases (MMP-3) (arthritis and tumor metastasis)



BcI-2 (pro- and anti-apoptotic)

Hadjuk, Mol. Interv., 2006



MCSS Workflow







• CHARMM-GUI & CHARMMING integrations



- CHARMM-GUI & CHARMMING integrations
 - preparation of targets for MCSS calculations



- CHARMM-GUI & CHARMMING integrations
 - preparation of targets for MCSS calculations
 - HOOK, oligo(peptides), DLD, ..., Molpy (biomolecular residues)



- CHARMM-GUI & CHARMMING integrations
 - preparation of targets for MCSS calculations
 - HOOK, oligo(peptides), DLD, ..., Molpy (biomolecular residues)
- CHARMM force field parameters



- CHARMM-GUI & CHARMMING integrations
 - preparation of targets for MCSS calculations
 - HOOK, oligo(peptides), DLD, ..., Molpy (biomolecular residues)
- CHARMM force field parameters
 - from CHARMM27 to CHARMM36



- CHARMM-GUI & CHARMMING integrations
 - preparation of targets for MCSS calculations
 - HOOK, oligo(peptides), DLD, ..., Molpy (biomolecular residues)
- CHARMM force field parameters
 - from CHARMM27 to CHARMM36
- CHARMM solvent models



- CHARMM-GUI & CHARMMING integrations
 - preparation of targets for MCSS calculations
 - HOOK, oligo(peptides), DLD, ..., Molpy (biomolecular residues)
- CHARMM force field parameters
 - from CHARMM27 to CHARMM36
- CHARMM solvent models
 - ACE, FACTS, ..., explicit solvent



- CHARMM-GUI & CHARMMING integrations
 - preparation of targets for MCSS calculations
 - HOOK, oligo(peptides), DLD, ..., Molpy (biomolecular residues)
- CHARMM force field parameters
 - from CHARMM27 to CHARMM36
- CHARMM solvent models
 - ACE, FACTS, ..., explicit solvent
- automatic generation of fragments

Chemistry at HARvard Macromolecular Mechanics MCSS+Molpy Workflow





MCSS in Practice (1)





MCSS in Practice (2)





MCSS in Practice (3)



MCSS+Molpy in Practice



MCSS+Molpy in Practice





Fragments & Partition Schemes











Nucleic fragments library





Conformational & Connection library

Structure of Nucleotides																					
Nucleobases, Ex :	Ō	Α	Е																		
Mono-nucleotide, Ex :	D	A	Х	Ν	1	1	0														
Di-nucleotide, Ex :	R	Т	Y	S	1	0	1	R	Т	Y	S	0	1	1							
Tri-nucleotide, Ex :	R	С	X	Ν	1	0	2	R	С	X	Ν	0	0	2	R	С	Х	Ν	0	1	2

	Туре		Patches		Orientation			Pucker
0	None	Ε	Ethyl	X	Anti-Trans]	Ν	North
D	Deoxyribonucleic	Η	Hydrogen	Y	Syn-Cis		S	South
R	Ribonucleic	M	Methyl					

	Base		5' Patch		3' Patch	
0	None	0	None	0	None	0
Α	ADES	1	5Ter	1	3Ter	1
С	CYTS	2	5Pho	2	3Po3	2
G	GUAS	3	5Pom	3	3Pho	
Ι	INOS			4	3Pom	
Р	PSUS			5	2Ter	
Т	THYS			6	2Po3	
U	URAS			7	2Pho	
				8	2Pom	

	Connection
0	None
1	3 - 5
2	2 - 5









Clustering of initial poses

352 chains

10 12 14 16 18

12

8 RMSD (Å)



RRM (top 60 U&C)



KH (top 60 A&G)



Zf-CCCH (top 60 U&A)

Energy (kcal/mol) -30 chain search sequence: CU -20

chain search

-20







Best scored chain (1,3 Å)





16



Scoring functions: nucleotideprotein interactions





Other biomolecules

- amino acids
- sugars
- lipids
- etc



Other biomolecules

- amino acids
- sugars
- lipids
- etc

You're welcome to join and contribute ...



Acknowledgments

- Martin Karplus
- Coralie Rohmer, MSc.
- Manuel Simoes
- Nicolas Chevrollier, Ph.D. student
- Pierre-Yves Vits, Jean-François Kneib

