

Soil and substrate morphology as witnesses of present and former agricultural landscape management

Caroline Chartin, Sébastien Salvador-Blanes, Florent Hinschberger, Hocine Bourennane, Jean-Jacques Macaire

▶ To cite this version:

Caroline Chartin, Sébastien Salvador-Blanes, Florent Hinschberger, Hocine Bourennane, Jean-Jacques Macaire. Soil and substrate morphology as witnesses of present and former agricultural landscape management. EGU General Assembly, Apr 2009, Vienne, Austria. pp.19 - 24, 2009, Geophysical Research Abstracts, Vol. 11, EGU2009-11596. hal-02325495

HAL Id: hal-02325495 https://hal.science/hal-02325495

Submitted on 22 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Soil and substrate morphology as witnesses of present and former agricultural landscape management contact: caroline.chartin@etu.univ-tours.fr C. Chartin (1), S. Salvador-Blanes (1), F. Hinschberger (1), H. Bourennane (2), and J.-J. Macaire (1)

(1) Université François Rabelais – Tours, CNRS/INSU, Université d'Orléans, Institut des Sciences de la Terre d'Orléans – UMR 6113, Faculté des Sciences et Techniques, Parc Grandmont, 37200 TOURS, France (2) Institut National de la Recherche Agronomique – Unité de Recherche Science du Sol, 2163 avenue de la Pomme de Pin, CS 40001 ARDON, 45075 ORLEANS Cedex 2, France

Combination of water / tillage erosion and BACKGROUND successive regroupings of the lands, have significantly modified the morphology of cultivated hillslopes. Nowadays, we can still observe over cultivated landscapes various anthropogenic structures (e.g. lynchets) that correspond to former and/or present field limits.

To characterise the geometry of these various structures, and assess their relation with topography, surficial resistivity, or any other easily available indicator. \rightarrow existence of a strong mathematical relation between these indicators and soil thickness would allow us to extrapolate the quantification to larger areas.

STUDY SITE:

near Tours in the Parisian Basin (France)

- 17 ha hillslope in a chalky watershed
- elevation: from 35 to 90 m (SE facing slope)
- regrouping of the lands in the 1960s

1 – STRUCTURES CHARACTERISATION

Two types (Figs. 1&2)

- Lynchets: at lower limits of present fields mostly concave
- Secondary structures: at some former field limits (1950) mostly convex







Figure 1: Profile curvature and present and former (1950) field limits









interrogate about substrate morphology (Fig. 4).

→ Substrate morphology modified downslope present field → Could explain weak correlations between soil thickness

- Soil accumulations due to past and present field limits cannot be estimated by topographic attributes exclusively. - Irregularity in the substrate morphology due to tillage - Surficial resistivity has to be further treated from background noise due to substrate heterogeneity to find a better link with soil accumulations.



.OPE	-						
	-						
	-						
	-						
25							
55,5 55 -	UPS	UPSLOPE			DOWNSLOPE		
(m) 54,5 - 54 - 54 - 53,5 - 53 -			-				
0,01,01,01,01,01,01,01,01,01,01,01,01,01				<u> </u>			
	chalk						
52,5 -						<u> </u>	
52				<u>, , , , , , , , , , , , , , , , , , , </u>			
b) -30	0 -20	-20 -10 0 10 20 distance from former field limit (m)					