



**HAL**  
open science

# Uniform blow-up estimates for nonlinear heat equations and applications

Frank Merle, Hatem Zaag

► **To cite this version:**

Frank Merle, Hatem Zaag. Uniform blow-up estimates for nonlinear heat equations and applications. *Methods and Applications of Analysis*, 2001, 8, pp.551 - 556. hal-02326191

**HAL Id: hal-02326191**

**<https://hal.science/hal-02326191v1>**

Submitted on 24 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Uniform blow-up estimates for nonlinear heat equations and applications

Frank Merle

*IUF and Université de Cergy-Pontoise*

Hatem Zaag

*CNRS École Normale Supérieure*

We consider the following nonlinear heat equation

$$\begin{cases} u_t &= \Delta u + |u|^{p-1}u \\ u(0) &= u_0, \end{cases} \quad (1)$$

where  $u : (x, t) \in \mathbb{R}^N \times [0, T) \rightarrow \mathbb{R}$ ,

$$1 < p, (N - 2)p < N + 2 \text{ and either } u_0 \geq 0 \text{ or } (3N - 4)p < 3N + 8. \quad (2)$$

More general vector-valued heat equations can be considered with similar results (see [MZ99] for more details) :

$$\begin{cases} u_t &= \Delta u + F(|u|)u \\ u(0) &= u_0, \end{cases} \quad (3)$$

where  $u : \Omega \times [0, T) \rightarrow \mathbb{R}^M$ ,  $p$  satisfies (2),  $\Omega = \mathbb{R}^N$  or  $\Omega$  is a smooth bounded convex domain of  $\mathbb{R}^N$ ,  $F(|u|) \sim |u|^{p-1}$  as  $|u| \rightarrow +\infty$ , and  $M \in \mathbb{N}$ .

We are interested in the blow-up phenomenon for (1). Many authors has been interested in this topic. Let us mention for instance Friedman [Fri65], Fujita [Fuj66], Ball [Bal77], Bricmont and Kupiainen [BKL94], Chen and Matano [CM89], Galaktionov and Vázquez [GV95], Giga and Kohn [GK89], [GK87], [GK85], Herrero and Velázquez [HV93], [HV92].

In the following, we consider  $u(t)$  a blow-up solution of (1) and denote its blow-up time by  $T$ . We aim at finding sharp uniform estimates at blow-up and specifying the blow-up behavior of  $u(t)$ . Such a study is done considering equation (1) in its self-similar form : for all  $a \in \mathbb{R}^N$ , we define

$$y = \frac{x - a}{\sqrt{T - t}}, \quad s = -\log(T - t), \quad w_{a,T}(y, s) = (T - t)^{\frac{1}{p-1}} u(x, t). \quad (4)$$

Therefore,  $w_{a,T} = w$  satisfies  $\forall s \geq -\log T, \forall y \in \mathbb{R}^N$  :

$$\frac{\partial w}{\partial s} = \Delta w - \frac{1}{2}y \cdot \nabla w - \frac{w}{p-1} + |w|^{p-1}w. \quad (5)$$

Let us introduce the following Lyapunov functional associated with (5) :

$$E(w) = \int_{\mathbb{R}^N} \left( \frac{1}{2}|\nabla w|^2 + \frac{1}{2(p-1)}|w|^2 - \frac{1}{p+1}|w|^{p+1} \right) \rho(y) dy$$

where  $\rho(y) = \frac{e^{-\frac{|y|^2}{4}}}{(4\pi)^{N/2}}$ .

The question is to estimate  $w_{a,T}(s)$  as  $s \rightarrow +\infty$ , uniformly with respect to  $a$ , whether  $a$  is a blow-up point or not ( $a$  is called a blow-up point if there exists  $(a_n, t_n) \rightarrow (a, T)$  such that  $|u(a_n, t_n)| \rightarrow +\infty$ ).

Giga and Kohn showed that self-similar variables are convenient for describing the blow-up rate in the following sense : there exists  $\epsilon_0(u_0) > 0$  such that  $\forall s \geq s_0^*(u_0)$ ,

$$\epsilon_0 \leq |w(s)|_{L^\infty} \leq \frac{1}{\epsilon_0}. \quad (6)$$

We first aim at sharpening this result in order to obtain compactness properties in our problem.

## 1 A Liouville Theorem for equation (5)

We are interested in classifying all global and bounded solutions of (5), for all subcritical  $p$  :

$$p > 1 \text{ and } (N-2)p < N+2. \quad (7)$$

We claim the following :

**Theorem 1 (A Liouville Theorem for equation (5))** *Assume (7) and consider  $w$  a solution of (5) defined for all  $(y, s) \in \mathbb{R}^N \times \mathbb{R}$  such that  $\forall (y, s) \in \mathbb{R}^N \times \mathbb{R}, |w(y, s)| \leq C$ . Then, either  $w \equiv 0$ , or  $w \equiv \kappa$  or  $w(y, s) = \pm \varphi(s - s_0)$  where  $\kappa = (p-1)^{-\frac{1}{p-1}}$ ,  $s_0 \in \mathbb{R}$  and  $\varphi(s) = \kappa(1 + e^s)^{-\frac{1}{p-1}}$ .*

**Remark :**  $\varphi$  is in fact an  $L^\infty$  connection between two critical points of (5):  $\kappa$  and 0. Indeed,

$$\dot{\varphi} = -\frac{\varphi}{p-1} + \varphi^p, \quad \varphi(-\infty) = \kappa, \quad \varphi(+\infty) = 0.$$

**Remark :** A similar classification result can be obtained with a solution  $w$  defined only on  $(-\infty, s^*)$  (see [MZ98]).

Theorem 1 has the following corollary :

**Corollary 1 (A Liouville Theorem for equation (1))** *Assume (7) and consider  $u$  a solution of (1) defined for all  $(x, t) \in \mathbb{R}^N \times (-\infty, 0)$  such that  $\forall (x, t) \in \mathbb{R}^N \times (-\infty, 0)$ ,  $|u(x, t)| \leq C(-t)^{-\frac{1}{p-1}}$ . Then, either  $u \equiv 0$ , or  $u(x, t) = \pm \kappa(T^* - t)^{-\frac{1}{p-1}}$  for some  $T^* \geq 0$ .*

The proofs can be found in [MZ99] and [MZ98]. The key tools in the proof are the following :

- i) A classification of all possible linear behaviors of  $w(s)$  as  $s \rightarrow -\infty$  in  $L^2_\rho(\mathbb{R}^N)$  ( $L^\infty_{loc}(\mathbb{R}^N)$ ),
- ii) The following geometric transformations which keeps (5) invariant :

$$w(y, s) \rightarrow w_{a,b}(y, s) = w(y + ae^{\frac{s}{2}}, s + b),$$

where  $a \in \mathbb{R}^N$  and  $b \in \mathbb{R}$ ,

- iii) A blow-up criterion for (5) used for solutions close to the constant point  $\kappa$  (This criterion is also a blow-up criterion for (1) via the transformation (4)) :

*If for some  $s_0 \in \mathbb{R}$ ,  $I(w(s_0)) > 0$  where*

$$I(w) = -2E(w) + \frac{p-1}{p+1} \left( \int_{\mathbb{R}^N} |w(y)|^2 \rho(y) dy \right)^{\frac{p+1}{2}},$$

*then  $w(s)$  blows up in finite time.*

**Remark :** This criterion is sharp for solutions near constants. Indeed, if  $w(s_0) \equiv C_0$ , then

$$w \text{ blows-up in finite time} \Leftrightarrow |C_0| > \kappa \Leftrightarrow I(C_0) > 0.$$

**Remark :** The proof of the Liouville Theorem strongly relies on the existence of a Lyapunov functional for equation (5) and can not be extended to other systems where the nonlinearity is not a gradient. In [Zaa], we go beyond this restriction and introduce new tools to prove a Liouville Theorem of the same type for the following system

$$u_t = \Delta u + v^p, \quad v_t = \Delta v + u^q.$$

## 2 Localization at blow-up

We assume again (2). The estimate (6) of Giga and Kohn gives compactness in the problem. Using a compactness procedure in the singular zone of  $\mathbb{R}^N$  (which is, say  $\{y \mid |w(y, s)| \geq \frac{\kappa}{2}\}$ ), we find a solution satisfying the hypotheses of Theorem 1. Therefore,  $\Delta w$  is small with respect to  $|w|^p$  in this singular zone (or equivalently,  $\Delta u$  is small with respect to  $|u|^p$ ). A

subcritical localization procedure introduced by Zaag [Zaa98] (under the level of the constant  $\kappa$ ) allows us to propagate this estimate towards the intermediate zone between the singular and the regular one. We claim the following :

**Theorem 2 (Comparison with the associated ordinary differential equation)** *Assume (2) and consider  $T \leq T_0$  and  $\|u_0\|_{C^2(\mathbb{R}^N)} \leq C_0$ . Then,  $\forall \epsilon > 0$ , there is  $C(\epsilon, C_0, T_0)$  such that  $\forall (x, t) \in \mathbb{R}^N \times [0, T)$ ,*

$$|u_t - |u|^{p-1}u| \leq \epsilon |u|^p + C.$$

**Remark :** This way, we prove that the solution of the PDE (1) can be uniformly and globally in space-time compared to a solution of an ODE (localized by definition). Note that the condition  $u(0) \in C^2(\mathbb{R}^N)$  is not restrictive because of the regularizing effect of the Laplacian.

**Remark :** Many striking corollaries can be derived from this theorem. It implies in particular that no oscillation is possible near a blow-up point  $a$ , and that  $|u(x, t)| \rightarrow +\infty$  as  $(x, t) \rightarrow (a, T)$ . Moreover,  $\forall \epsilon_0 > 0$ , there exists  $t_0(\epsilon_0) < T$  such that for all  $b \in \mathbb{R}^N$ , if  $|u(b, t)| \leq (1 - \epsilon_0)\kappa(T - t)^{-\frac{1}{p-1}}$  for some  $t \in [t_0, T)$ , then  $b$  is not a blow-up point (this specifies more precisely a former result by Giga and Kohn where  $t_0 = t_0(\epsilon_0, a)$ ).

### 3 Optimal $L^\infty$ estimates at blow-up

We still assume (2). Using estimate (6) and the Liouville Theorem, we make a compactness argument to get the following sharp estimates :

**Theorem 3 ( $L^\infty$  refined estimates for  $w(s)$ )** *Assume that (2) holds. Then, there exist positive constants  $C_i$  for  $i = 1, 2, 3$  such that if  $u$  is a solution of (1) which blows-up at time  $T$  and satisfies  $u(0) \in C^3(\mathbb{R}^N)$ , then  $\forall \epsilon > 0$ , there exists  $s_1(\epsilon) \geq -\log T$  such that  $\forall s \geq s_1, \forall a \in \mathbb{R}^N$ ,*

$$\begin{aligned} \|w_{a,T}(s)\|_{L^\infty} &\leq \kappa + \left(\frac{N\kappa}{2p} + \epsilon\right)\frac{1}{s}, & \|\nabla w_{a,T}(s)\|_{L^\infty} &\leq \frac{C_1}{\sqrt{s}}, \\ \|\nabla^2 w_{a,T}(s)\|_{L^\infty} &\leq \frac{C_2}{s}, & \|\nabla^3 w_{a,T}(s)\|_{L^\infty} &\leq \frac{C_3}{s^{3/2}}. \end{aligned}$$

**Remark :** In the case  $N = 1$ , Herrero and Velázquez (Filippas and Kohn also) proved some related estimates, using a Sturm property introduced in particular by Chen et Matano (the number of space oscillations is a decreasing function of time).

**Remark :** The constant  $\frac{N\kappa}{2p}$  is optimal (see Herrero and Velázquez, Brimont and Kupiainen, Merle and Zaag).

## 4 Different notions of blow-up profiles and the stability question

We assume (2). We consider  $a \in \mathbb{R}^N$ , a blow-up point of  $u(t)$ , solution of (1). From translation invariance, we can assume  $a = 0$ . We would like to know whether  $u(t)$  (or  $w_{0,T}(s)$  defined in (4)) has a universal behavior or not, as  $t \rightarrow T$  (or  $s \rightarrow +\infty$ ).

Filippas, Kohn, Liu, Herrero et Velázquez prove that  $w$  behaves in two distinct ways :

- either  $\forall R > 0$ ,  $\sup_{|y| \leq R} \left| w(Qy, s) - \left[ \kappa + \frac{\kappa}{2ps} \left( l - \frac{1}{2} \sum_{k=1}^l y_k^2 \right) \right] \right| = O\left(\frac{1}{s^{1+\delta}}\right)$  as  $s \rightarrow +\infty$ , for some  $\delta > 0$  where  $l \in \{1, \dots, N\}$ ,  $Q$  is a  $N \times N$  orthogonal matrix and  $I_l$  is the  $l \times l$  identity matrix.
- or  $\forall R > 0$ ,  $\sup_{|y| \leq R} |w(y, s) - \kappa| \leq C(R)e^{-\lambda_0 s}$  for some  $\lambda_0 > 0$ .

From a physical point of view, these results do not tell us much about the transition between the singular zone ( $w \geq \alpha$  where  $\alpha > 0$ ) and the regular one ( $w \simeq 0$ ). In [MZ99], we specify this transition by proving the existence of a profile in the variable  $z = \frac{y}{\sqrt{s}}$ .

### Theorem 4 (Existence of a blow-up profile for equation (1))

*Assume (2) holds. There exists  $l \in \{0, 1, \dots, N\}$  and a  $N \times N$  orthogonal matrix  $Q$  such that  $w(Q(z)\sqrt{s}, s) \rightarrow f_l(z)$  uniformly on compact sets  $|z| \leq C$ , where  $f_l(z) = (p-1 + \frac{(p-1)^2}{4p} \sum_{i=1}^l |z_i|^2)^{-\frac{1}{p-1}}$  if  $l \geq 1$  and  $f_0(z) = \kappa = (p-1)^{-\frac{1}{p-1}}$ .*

This result has been proved by Velázquez in [Vel92]. However, the convergence speed depends on the considered blow-up point in [Vel92], whereas they are uniform in [MZ99]. This uniformity allows us to derive the stability of the profile  $f_N$  in [FKMZ].

Using renormalization theory, Bricmont and Kupiainen prove in [BK94] the existence of a solution of (5) such that

$$\forall s \geq s_0, \forall y \in \mathbb{R}^N, |w(y, s) - f_N\left(\frac{y}{\sqrt{s}}\right)| \leq \frac{C}{\sqrt{s}}.$$

Merle and Zaag prove the same result in [MZ97], thanks to a technique of finite-dimension reduction. They also prove the stability of such a behavior with respect to initial data, in a neighborhood of the constructed solution.

In [Zaa98] and [Vel92], it is proved that in this case,  $u(x, t) \rightarrow u^*(x)$  as  $t \rightarrow T$  uniformly on  $\mathbb{R}^N \setminus \{0\}$  and that  $u^*(x) \sim \left[ \frac{8p|\log|x||}{(p-1)^2|x|^2} \right]^{\frac{1}{p-1}}$  as  $x \rightarrow 0$ .

One interesting problem is to relate all known blow-up profiles' notions : profiles for  $|y|$  bounded,  $\frac{|y|}{\sqrt{s}}$  bounded or  $x \simeq 0$ . We prove in the following that all these descriptions are equivalent, in the case of single point blow-up with a non degenerate profile (generic case). This answers many questions which arose in former works.

**Theorem 5 (Equivalence of blow-up behaviors at a blow-up point)**

Assume (2) and consider  $a$  be an isolated blow-up point of  $u(t)$  solution of (1). The following behaviors of  $u(t)$  and  $w_{a,T}(s)$  (defined in (4)) are equivalent :

- i)  $\forall R > 0$ ,  $\sup_{|y| \leq R} \left| w(y, s) - \left[ \kappa + \frac{\kappa}{2ps} \left( N - \frac{1}{2}|y|^2 \right) \right] \right| = o\left(\frac{1}{s}\right)$  as  $s \rightarrow +\infty$ ,
- ii)  $\forall R > 0$ ,  $\sup_{|z| \leq R} |w(z\sqrt{s}, s) - f_0(z)| \rightarrow 0$  as  $s \rightarrow +\infty$  with  $f_0(z) = (p-1 + \frac{(p-1)^2}{4p}|z|^2)^{-\frac{1}{p-1}}$ ,
- iii)  $\exists \epsilon_0 > 0$  such that for all  $|x-a| \leq \epsilon_0$ ,  $u(x, t) \rightarrow u^*(x)$  as  $t \rightarrow T$  and  $u^*(x) \sim \left[ \frac{8p|\log|x-a||}{(p-1)^2|x-a|^2} \right]^{\frac{1}{p-1}}$  as  $x \rightarrow a$ .

A further application of the Liouville Theorem is the stability of the behavior described in Theorem 5, with respect to perturbations in initial data. Using a dynamical system approach, we prove in [FKMZ], with Fermanian the following :

**Theorem 6 (Stability of the blow-up profile)** Assume (7) and consider  $\tilde{u}(t)$  a blow-up solution of (1) with initial data  $\tilde{u}_0$  which blows-up at  $t = \tilde{T}$  at only one point  $\tilde{a} = 0$  and satisfies (6). Assume that

$$\text{for all } |x| \geq R \text{ and } t \in [0, \tilde{T}), \quad |\tilde{u}(x, t)| \leq M.$$

and that the function  $\tilde{w}_{0, \tilde{T}}(y, s)$  defined in (4) satisfies uniformly on compact sets of  $\mathbb{R}^N$

$$\tilde{w}_{0, \tilde{T}}(y, s) - \kappa \underset{s \rightarrow +\infty}{\sim} \frac{\kappa}{2ps} \left( N - \frac{|y|^2}{2} \right). \quad (8)$$

Then, there is a neighborhood  $\mathcal{V}$  in  $L^\infty$  of  $\tilde{u}_0$  such that for all  $u_0 \in \mathcal{V}$  the solution of (1) with initial data  $u_0$  blows-up at time  $T = T(u_0)$  at a unique point  $a = a(u_0)$  and the function  $w_{a,T}(y, s)$  defined in (4) satisfies uniformly on compact sets of  $\mathbb{R}^N$

$$w_{a,T}(y, s) - \kappa \underset{s \rightarrow +\infty}{\sim} \frac{\kappa}{2ps} \left( N - \frac{|y|^2}{2} \right).$$

Moreover,  $(a(u_0), T(u_0))$  goes to  $(0, \tilde{T})$  as  $u_0$  goes to  $\tilde{u}_0$ .

**Remark :** This results generalizes the stability result of [MZ97]. Note that unlike most applications of the Liouville Theorem, this result is valid for all subcritical  $p$ . In [FKZ], the same result is proved (only under the condition (2)), by a completely different approach based on the Liouville Theorem and on [MZ97].

**Remark :** In [FKMZ], we prove the stability with respect to initial data of the blow-up behavior with the minimal speed

$$\|u(t)\|_{L^\infty} \leq C(T - t)^{-\frac{1}{p-1}} \quad (9)$$

for all subcritical  $p$  (that is under the condition (7)). Note that this result is obvious under the weaker assumption (2), for Giga and Kohn proved in [GK87] that all blow-up solutions satisfy (9). No blow-up rate estimate is known if

$$u_0 \text{ has no sign and } (3N - 4)p \geq 3N - 8.$$

Therefore, our result is meaningful in this last case.

## References

- [Bal77] J. M. Ball. Remarks on blow-up and nonexistence theorems for nonlinear evolution equations. *Quart. J. Math. Oxford Ser. (2)*, 28(112):473–486, 1977.
- [BK94] J. Bricmont and A. Kupiainen. Universality in blow-up for nonlinear heat equations. *Nonlinearity*, 7(2):539–575, 1994.
- [BKL94] J. Bricmont, A. Kupiainen, and G. Lin. Renormalization group and asymptotics of solutions of nonlinear parabolic equations. *Comm. Pure Appl. Math.*, 47(6):893–922, 1994.
- [CM89] X. Y. Chen and H. Matano. Convergence, asymptotic periodicity, and finite-point blow-up in one-dimensional semilinear heat equations. *J. Differential Equations*, 78(1):160–190, 1989.
- [FKMZ] C. Fermanian Kammerer, F. Merle, and H. Zaag. Stability of the blow-up profile of non-linear heat equations from the dynamical system point of view. preprint.
- [FKZ] C. Fermanian Kammerer and H. Zaag. Boundedness till blow-up of the difference between two solutions to the semilinear heat equation. preprint.



- [Fri65] A. Friedman. Remarks on nonlinear parabolic equations. In *Proc. Sympos. Appl. Math., Vol. XVII*, pages 3–23. Amer. Math. Soc., Providence, R.I., 1965.
- [Fuj66] H. Fujita. On the blowing up of solutions of the Cauchy problem for  $u_t = \Delta u + u^{1+\alpha}$ . *J. Fac. Sci. Univ. Tokyo Sect. I*, 13:109–124, 1966.
- [GK85] Y. Giga and R. V. Kohn. Asymptotically self-similar blow-up of semilinear heat equations. *Comm. Pure Appl. Math.*, 38(3):297–319, 1985.
- [GK87] Y. Giga and R. V. Kohn. Characterizing blowup using similarity variables. *Indiana Univ. Math. J.*, 36(1):1–40, 1987.
- [GK89] Y. Giga and R. V. Kohn. Nondegeneracy of blowup for semilinear heat equations. *Comm. Pure Appl. Math.*, 42(6):845–884, 1989.
- [GV95] V. A. Galaktionov and J. L. Vázquez. Geometrical properties of the solutions of one-dimensional nonlinear parabolic equations. *Math. Ann.*, 303:741–769, 1995.
- [HV92] M. A. Herrero and J. J. L. Velázquez. Flat blow-up in one-dimensional semilinear heat equations. *Differential Integral Equations*, 5(5):973–997, 1992.
- [HV93] M. A. Herrero and J. J. L. Velázquez. Blow-up behaviour of one-dimensional semilinear parabolic equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 10(2):131–189, 1993.
- [MZ97] F. Merle and H. Zaag. Stability of the blow-up profile for equations of the type  $u_t = \Delta u + |u|^{p-1}u$ . *Duke Math. J.*, 86(1):143–195, 1997.
- [MZ98] F. Merle and H. Zaag. Optimal estimates for blowup rate and behavior for nonlinear heat equations. *Comm. Pure Appl. Math.*, 51(2):139–196, 1998.
- [MZ99] F. Merle and H. Zaag. A Liouville theorem for vector-valued nonlinear heat equations and applications. *Math. Annalen*, 1999. to appear.
- [Vel92] J. J. L. Velázquez. Higher-dimensional blow up for semilinear parabolic equations. *Comm. Partial Differential Equations*, 17(9-10):1567–1596, 1992.

- [Zaa] H. Zaag. A liouville theorem and blow-up behavior for a vector-valued nonlinear heat equation with no gradient structure. preprint.
- [Zaa98] H. Zaag. Blow-up results for vector-valued nonlinear heat equations with no gradient structure. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 15(5):581–622, 1998.

**Address :**

Département de mathématiques, Université de Cergy-Pontoise, 2 avenue Adolphe Chauvin, B.P. 222, Pontoise, 95 302 Cergy-Pontoise cedex, France.  
Département de mathématiques et applications, École Normale Supérieure, 45 rue d'Ulm, 75 230 Paris cedex 05, France.

e-mail : `merle@math.pst.u-cergy.fr`, `Hatem.Zaag@ens.fr`