



Study of the X-ray activity of Sagittarius A* during the 2011 XMM-Newton campaign (Corrigendum)

Enmanuelle Mossoux,, Nicolas Grosso,, H. Vincent, Frédéric, Delphine
Porquet,

► To cite this version:

Enmanuelle Mossoux,, Nicolas Grosso,, H. Vincent, Frédéric, Delphine Porquet,. Study of the X-ray activity of Sagittarius A* during the 2011 XMM-Newton campaign (Corrigendum). *Astronomy and Astrophysics - A&A*, 2015, 580, pp.C2. 10.1051/0004-6361/201424682e . hal-02327493

HAL Id: hal-02327493

<https://hal.science/hal-02327493v1>

Submitted on 15 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Study of the X-ray activity of Sagittarius A* during the 2011 *XMM-Newton* campaign (Corrigendum)

Enmanuelle Mossoux¹, Nicolas Grosso¹, Frédéric H. Vincent², and Delphine Porquet¹

¹ Observatoire Astronomique de Strasbourg, Université de Strasbourg, CNRS, UMR 7550, 11 rue de l’Université,
67000 Strasbourg, France

e-mail: enmanuelle.mossoux@astro.unistra.fr

² Nicolaus Copernicus Astronomical Center, ul. Bartycka 18, 00-716 Warszawa, Poland

A&A 573, A46 (2015), DOI: [10.1051/0004-6361/201424682](https://doi.org/10.1051/0004-6361/201424682)

Key words. Galaxy: center – X-rays: individuals: Sgr A* – radiation mechanisms: general – errata, addenda

The weight equation in Sect. 3.1 of [Mossoux et al. \(2015\)](#) unfortunately features a typo. The correct equation should read:

$$w = \frac{CR_{\text{src+bkg}}}{CR_{\text{src+bkg}} - CR_{\text{bkg}}} \quad (1)$$

with $CR_{\text{src+bkg}}$ the count rate of the Bayesian block of the source+background event list and CR_{bkg} the count rate of the corresponding Bayesian block of the background that is surface-corrected. As introduced in the first preprint version of the proceeding of [Scargle et al. \(2013\)](#), this weight equation adjusts the Voronoi time-interval in order to subtract the average background. The results presented in [Mossoux et al. \(2015\)](#) remain unchanged since they were all obtained following the recipe of [Scargle et al. \(2013\)](#) and using the correct equation for the Voronoi time-interval weighting.

Therefore, the “alternative” photon-weighting recently tested by [Worpel & Schwöpe \(2015\)](#) in their Sect. 4.6 ($w \equiv 1 - CR_{\text{bkg}}/CR_{\text{src+bkg}}$) is actually equivalent to the recipe of [Scargle et al. \(2013\)](#) that we used in [Mossoux et al. \(2015\)](#) since the photon-weighting is equal to the inverse of the Voronoi time-interval weighting. Indeed, the “alternative” photon-weighting of [Worpel & Schwöpe \(2015\)](#) is identical to those used in Eq. (1) of [Mossoux et al. \(2015\)](#) to compute smoothed light-curve.

References

- Mossoux, E., Grosso, N., Vincent, F. H., & Porquet, D. 2015, [A&A, 573, A46](#)
Scargle, J. D., Norris, J. P., Jackson, B., & Chiang, J. 2013, 2012 Fermi Symp. Proc. – eConf C121028 [[arXiv:1304.2818v1](https://arxiv.org/abs/1304.2818v1)]
Worpel, H., & Schwöpe, A. D. 2015, [A&A, 578, A80](#)