



## Substitutive normal sets

Christian Mauduit

### ► To cite this version:

Christiane Mauduit. Substitutive normal sets. Colloquia Mathematica Societatis Janos Bolyai, 1990.  
hal-02333726

HAL Id: hal-02333726

<https://hal.science/hal-02333726>

Submitted on 25 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## SUBSTITUTIVE NORMAL SETS

CHRISTIAN MAUDUIT

Let  $u = (u_n)_{n \in \mathbb{N}}$  be an infinite sequence of integers. Define  $B(u)$  as the set of those real  $\alpha$ 's such that the sequence  $(u_n \alpha)_{n \in \mathbb{N}}$  is uniformly distributed modulo 1. A set  $B$  of real numbers is said to be normal if there exists a sequence  $u$  such that  $B = B(u)$ . G. Rauzy has given a complete characterization of normal sets (cf. [4]), namely, a set  $B$  is normal if and only if (i)  $0 \notin B$ ,  $B+1=B$ ; (ii)  $\forall q \in \mathbb{Z}^*$ ,  $qB \subset B$ ; (iii) there exists a sequence  $(f_n)_{n \in \mathbb{N}}$  of continuous real functions such that  $\lim_{n \rightarrow \infty} f_n(x) = 0$  exactly for those  $x$ 's which are elements of  $B$ .

We will say that a set  $B$  of real numbers is  $\Sigma$ -normal (or a substitutive normal set) if there exists an increasing

sequence  $u$  generated by an irreducible substitution such that  $B = B(u)$ . We give the following characterization of  $\Sigma$ -normal sets (cf. [2]):

**THEOREM:**  $B$  is a  $\Sigma$ -normal set if and only if  $R \setminus B$  is an algebraic extension of finite degree of  $Q$ .

### I. SUBSTITUTION OVER A FINITE ALPHABET

Let  $g \geq 2$ ,  $A = (a_1, \dots, a_g)$  an alphabet and  $\Sigma$  a substitution over  $A$  (cf. [6]).

$\Sigma$  is given by the set  $\Sigma(a_i) = a_{\sigma_i(1)}, \dots, a_{\sigma_i(\ell_i(1))}$  ( $i \in \{1, \dots, g\}$ ), where  $\ell_i(n)$  is the length of the word  $\Sigma^n(a_i)$  ( $n \in \mathbb{N}$ ) and  $\sigma_i$  a map from  $\{1, \dots, \ell_i(1)\}$  to  $\{1, \dots, g\}$ .

We will suppose that  $\Sigma$  is irreducible, that is to say that for any  $(i, j) \in \{1, \dots, g\}^2$ , there exists an integer  $n$  such that the word  $\Sigma^n(a_i)$  contains the letter  $a_j$ .  
We remark that  $L(n) = M^n T(1, \dots, 1)$ , where  
 $L(n) = T(\ell_1(n), \dots, \ell_g(n))$  and

$$M = (M_{ij})_{(i,j) \in \{1, \dots, g\}^2} \text{ with } M_{ij} = \text{card}(\sigma_i^{-1}(\{j\})).$$

**Definition:** Let  $m \in A^\mathbb{N}$  and  $t \in \{1, \dots, g\}$ . We call support of  $m$  with respect to  $a_t$  and we denote by  $[m]_t$ , the increasing sequence of those integers  $n$  such that  $a_t$  is

the  $(n+1)$ -th letter of the word  $m$ .

We say that a sequence  $u$  of integers is generated by a substitution  $\Sigma$  if there exists  $m \in A^N$  such that  $\Sigma(m) = m$  and  $t \in \{1, \dots, g\}$  such that  $u$  is the support of  $m$  with respect to  $a_t$ .

### III. $\lambda$ FUNCTIONS ASSOCIATED TO $\Sigma$

Statistical properties of sequences generated by  $\Sigma$  are determined by the behaviour of the following functions.

If  $\sigma_i^{-1}(\{j\})$  is non empty, let us denote by  $k_1, \dots, k_{M_{ij}}$  the increasing sequence of its elements.

For any  $v \in \{1, \dots, M_{ij}\}$  we put  $\lambda_{ij}^{(v)} = \sum_{k' < k_v} \ell_{\sigma_i(k')}$

$$\Lambda_{ij} = \lambda_{ij}^{(v)}, v \in \{1, \dots, M_{ij}\} \setminus \{0\} \text{ and } \Lambda = \bigcup_{(i,j) \in \{1, \dots, g\}^2} \Lambda_{ij}.$$

Now we define the set  $E$ :

$$E = \{\alpha \in \mathbb{R}, \exists k \in \mathbb{Z}^*, \forall \lambda \in \Lambda \lim_{n \rightarrow \infty} e(k\lambda(n)\alpha) = 1\}$$

(where  $e(x) = \exp(2i\pi x)$ ) and we show that

$$E = \{\alpha \in \mathbb{R}, \exists k \in \mathbb{Z}^*, \forall i \in \{1, \dots, g\} \lim_{n \rightarrow \infty} e(k\lambda_i(n)\alpha) = 1\}$$

is an algebraic extension of finite degree of  $\mathbb{Q}$ .

Example 1. If  $\Sigma$  is a substitution of constant length (i.e.  $\ell_i(1) = r$  for any  $i \in \{1, \dots, g\}$ ), that is to say if  $u$  is generated by a finite automaton (cf. [1]), we have  $\Lambda = \{\lambda\}$  with  $\lambda(n) = r^n$ , and so  $E = Q$ .

Example 2. If  $\Sigma$  is defined by  $\Sigma(a_1) = a_1a_2a_2$  and  $\Sigma(a_2) = a_2a_1$ , we have  $\Lambda_{11} = \Lambda_{22} = \emptyset$ ,  $\Lambda_{12} = \{\ell_1, \ell_1 + \ell_2\}$  and  $\Lambda_{21} = \{\ell_2\}$  with  $\ell_1(0) = \ell_2(0) = 1$ ,  $\ell_1(1) = 3$ ,  $\ell_2(1) = 2$  and for any  $i \in \{1, 2\}$   $\ell_i(n+1) = 2\ell_i(n) + \ell_i(n-1)$ . We deduce from this that  $E = Q(\sqrt{2})$ .

### III. NORMAL SET WITH RESPECT TO A SUBSTITUTIVE SEQUENCE

First, we have to show that if  $u$  is generated by  $\Sigma$ , then  $B(u) = R \setminus E$ . We recall that, according to Weyl's criterium (cf. [5]), the sequence  $(u_n \cdot \alpha)_{n \in \mathbb{N}}$  is uniformly distributed modulo 1 if and only if, for any  $k \in \mathbb{Z}^*$ ,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n < N} e(k u_n \alpha) = 0.$$

Let us consider the Weyl sums  $s_i(\alpha, n) = \sum_{p \in [\Sigma^n(a_i)]} e(p\alpha)$

and the vector  $w(\alpha, n) = (w_i(\alpha, n))_{i \in \{1, \dots, g\}}$ ,

with  $w_i(\alpha, n) = \frac{s_i(\alpha, n)}{s_i(0, n)}$ .

Then  $w(\alpha, n+1) = A(\alpha, n) w(\alpha, n)$  where  $A(\alpha, n)$  is a  $g \times g$ -matrix. We remark that every coefficient of  $A(\alpha, n)$  depending on  $\alpha$  is a real linear combination of  $e(\lambda(n)\alpha)$  with  $\lambda \in \Lambda$ .

So, if  $\alpha \in \mathbb{R} \setminus E$  the arguments of these complex numbers are "well distributed" modulo  $2\pi$ , and we show that

$$\lim_{n \rightarrow \infty} B(\alpha, n) = 0 \text{ (we put } B(\alpha, n) = A(\alpha, n) A(\alpha, n-1) \dots \\ \dots A(\alpha, 0)).$$

We deduce from this that  $w(\alpha, n) \in o(1)$  and then

$$\sum_{p \in [m]} e(p\alpha) \in o(N), \text{ i.e. } (u_n \cdot \alpha)_{n \in \mathbb{N}} \text{ is uniformly}$$

distributed modulo 1.

Conversely, let  $\alpha \in E$ .

As  $(A(\alpha, n) - A(0, n))$  converges to 0 with an exponential speed,  $(B(\alpha, n) - B(0, n))$  is smaller than some constant related to the spectrum of  $M$ . By iterating  $\Sigma$ , this constant can be made arbitrarily small.

$$As w(\alpha, n+1) = B(\alpha, n) w(\alpha, n)$$

$$= T(1, \dots, 1) + (B(\alpha, n) - B(0, n))^T (0, \dots, 0, 1, 0, \dots, 0),$$

it follows that  $\lim |w_1(\alpha, n)| > 0$  and so the sequence  $(u_n \cdot \alpha)_{n \in \mathbb{N}}$  is not uniformly distributed modulo 1.

#### IV. CONVERSE

Let  $F$  be an algebraic extension of finite degree of  $\mathbb{Q}$ . We know that there exists a Pisot number  $\theta$  such that  $F = \mathbb{Q}(\theta)$  (cf. [3]).

On the other hand, we can show that for any Pisot number  $\theta$  of degree  $g$ , there exists an integer  $v$  and an irreducible non negative integral  $g \times g$  matrix  $M = (M_{ij})_{(i,j) \in \{1, \dots, g\}^2}$  such that the spectrum contains only  $\theta^v$  and its conjugates.

Now consider the substitution  $\Sigma$  given by

$$\forall i \in \{1, \dots, g\}, \Sigma(a_i) = \underbrace{a_1 \dots a_1}_{M_{11} \text{ times}} \dots \underbrace{a_j \dots a_j}_{M_{1j} \text{ times}} \dots \underbrace{a_g \dots a_g}_{M_{1g} \text{ times}}$$

By iterating  $\Sigma$ , we can suppose  $M_{ij} > 0$ . Then, let  $m$  be the fixed point of  $\Sigma$  beginning with  $a_1$  and  $u = [m]_1$  be the sequence generated by  $m$  and  $a_1$ .

Using the same method as in § III, we show that

$$E = \mathbb{Q}(\theta^v).$$

As  $\theta$  is a Pisot number,  $\mathbb{Q}(\theta^v) = \mathbb{Q}(\theta)$  and

$$B(u) = R \setminus E = R \setminus \mathbb{Q}(\theta^v) = R \setminus \mathbb{Q}(\theta) = R \setminus F.$$

So,  $R \setminus F$  is a  $\Sigma$ -normal set.

## REFERENCES

- [1] C. MAUDUIT: Automates finis et ensembles normaux,  
*Ann. Inst. Fourier*, t. 36, fasc. 2, pp. 1-25.
- [2] C. MAUDUIT: Caractérisation des ensembles normaux substitutifs, *Invent. Math.* (to appear).
- [3] C. PISOT: La répartition modulo 1 et les nombres algébriques, *Ann. Scuola Normale Sup. Pisa*, vol. 7, 1938, pp. 205-245.
- [4] G. RAUZY: Caractérisation des ensembles normaux, *Bull. Soc. Math. France*, vol. 98, 1970, pp. 401-414.
- [5] G. RAUZY: Propriétés statistiques de suites arithmétiques, *F.U.F.*, 1976.
- [6] G. ROZENBERG and A. SALOMAA: The mathematical theory of L-systems, *Academic Press*, 1980.

Christian Mauduit  
C.N.R.S. L.A. 225  
Université d'Aix-Marseille II  
Département Mathématique-Informatique  
70, route Léon-Lachamp  
13288 Marseille Cedex 9  
France