



HAL
open science

A non-invasive method to appraise time-dependent effects of venom toxins on the mouse neuromuscular excitability in vivo

Tânia C. Gonçalves, Michel Partiseti, Denis Servent, Evelyne Benoit

► To cite this version:

Tânia C. Gonçalves, Michel Partiseti, Denis Servent, Evelyne Benoit. A non-invasive method to appraise time-dependent effects of venom toxins on the mouse neuromuscular excitability in vivo. 23th Meeting on Toxinology “ Toxins: Immunity, Inflammation and Pain ” organized by the French Society of Toxinology (SFET), Toxicon, 149, pp.104, 2018, 10.1016/j.toxicon.2018.02.030 . hal-02334359

HAL Id: hal-02334359

<https://hal.science/hal-02334359>

Submitted on 15 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A non-invasive method to appraise time-dependent effects of venom toxins on the mouse neuromuscular excitability in vivo

Tânia C. GONÇALVES^{1,2}, Michel PARTISETI², Denis SERVENT¹, Evelyne BENOIT^{1,3,*}

¹CEA, Institut des Sciences du Vivant Frédéric Joliot, SIMOPRO, Université Paris-Saclay, 91191 Gif-sur-Yvette, France; ²Sanofi R&D, Integrated Drug Discovery Unit, In-Vitro Biology & Pharmacology A, 94403 Vitry-sur-Seine, France; ³Institut des Neurosciences Paris-Saclay, UMR CNRS/Université Paris-Sud 9197, 91198 Gif-sur-Yvette, France.

* Corresponding author; E-mail: evelyne.benoit@cea.fr

The conventional methods, usually performed *in vivo*, provide limited information regarding sensory and neuromuscular excitability properties. In the present work, we review the new minimally invasive methods that have been improved to supplement the conventional electrophysiological methods. In particular, the automated sequence of multiple excitability tests developed by Professor Bostock (Qtrac©, Institute of Neurology, London, UK) to assess sensory and neuromuscular excitability in clinical neurophysiology, has been adapted to animal models *in vivo*, initially in rats and, more recently, in mice. This method is particularly appropriate for venom toxin injections and repeated recordings from the same animal cohort, making it a promising tool to appraise the progression of toxin effects, and that of an eventual given treatment, on the sensory and neuromuscular excitability of animal models *in vivo*. The *in vivo* efficiency in mice and long-term follow-up of the effects of sub-lethal doses of spider and *conus* venom toxins will be emphasised. It is indubitable that this method brings specific information regarding changes in ion transfer and membrane properties induced by a given venom toxin which, thereby, will help to a better characterization of toxin action on the sensory and neuromuscular excitability properties, *in vivo*.

Keywords: *In vivo* electrophysiology, sensory and neuromuscular excitability, toxins