



HAL
open science

Hopf-Galois structures on ambiskew polynomial rings

Julien Bichon, Agustín García Iglesias

► **To cite this version:**

Julien Bichon, Agustín García Iglesias. Hopf-Galois structures on ambiskew polynomial rings. *Journal of Noncommutative Geometry*, 2021, 15 (4), pp.1409-1431. 10.4171/JNCG/441 . hal-02336997

HAL Id: hal-02336997

<https://hal.science/hal-02336997>

Submitted on 29 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HOPF-GALOIS STRUCTURES ON AMBISKEW POLYNOMIAL RINGS

JULIEN BICHON AND AGUSTÍN GARCÍA IGLESIAS

ABSTRACT. We provide necessary and sufficient conditions to extend the Hopf-Galois algebra structure on an algebra R to a generalized ambiskew ring based on R , in a way such that the added variables for the extension are skew-primitive in an appropriate sense. We show that the associated Hopf algebra is again a generalized ambiskew ring, based on a suitable Hopf algebra $\underline{H}(R)$. Several examples are examined, including the Hopf-Galois objects over $U_q(\mathfrak{sl}_2)$.

1. INTRODUCTION

Let R be an algebra over a field k and consider the following data:

- (1) A pair of commuting algebra automorphisms $\tau, \omega \in \text{Aut } R$; set $\sigma := \tau\omega = \omega\tau$.
- (2) A σ -central element c ; that is $c \in Z_\sigma(R) := \{x \in R : xr = \sigma(r)x, r \in R\}$.
- (3) A scalar $\xi \in k^*$.

The associated *generalized ambiskew polynomial algebra* $A = A(R, X, Y, \tau, \omega, c, \xi)$, see [9], is the quotient of the free product $R * k\langle X, Y \rangle$ by the relations

$$Xr = \tau(r)X, \quad Yr = \omega(r)Y, \quad XY - \xi YX = c,$$

for any $r \in R$.

When c is assumed to be central (so that $\sigma = \text{id}$ or $\omega = \tau^{-1}$) then $A = A(R, X, Y, \tau, \tau^{-1}, c, \xi)$ is called an *ambiskew polynomial algebra* [8]; we set $A(R, X, Y, \tau, c, \xi) := A(R, X, Y, \tau, \tau^{-1}c, \xi)$.

A common question is about the ring-theoretical properties such as simplicity, or the representation theory, of the resulting algebra A , see for example [7], [9], a useful summary of known results being given in [4, Section 6].

Another interesting question is to find necessary and sufficient conditions on these data so that A preserves some extra structure on R . When R is a Hopf algebra, Brown-Macaulay [4] discuss the case in which the Hopf structure on R extends to a Hopf structure on the ambiskew polynomial Hopf algebra $A_R = A(R, X, Y, \tau, c, \xi)$ via the formula

$$(1.1) \quad \Delta(X) = X \otimes 1 + g \otimes X, \quad \Delta(Y) = Y \otimes 1 + h \otimes Y$$

for some group-like elements $g, h \in G(R)$; here, c is central¹. This generalizes previous results from [6] in the case when R is commutative. As an example, $U_q(\mathfrak{sl}_2)$ can be constructed from $k\mathbb{Z}$ by this process.

In this article, we generalize these results in two directions:

- (A) We discuss the extension of the Hopf algebra structure on R to a generalized ambiskew polynomial algebra A .
- (B) We further extend this to the case when R is a Hopf-Galois object over some Hopf algebra.

Hopf-Galois objects are natural generalizations of Hopf algebras with a Galois-theoretic flavour, introduced in [12], and have proved to be fundamental tools in analysing tensor categories of comodules [16, 14] as well as in recent classification questions for pointed Hopf algebras [1, 2].

While our first goal was to extend the results of [4] to Hopf-Galois objects in the case of ambiskew polynomial rings, our analysis evidenced that the proper context for doing that was that of generalized ambiskew polynomial rings, leading to the results in item (A).

¹In loc.cit. the authors choose the notation (X_+, X_-) and (g_+, g_-) instead of (X, Y) and (g, h) , respectively; and use h instead of c , σ instead of τ .

The punchline for item (B) is that while having to deal simultaneously with R and the Hopf algebra coacting on it can look difficult, it turns out that, fortunately, results of Grunspan [10] and Schauenburg [15] ensure that there exists a definition of Hopf-Galois objects that does not use any Hopf algebra, in terms of an algebra map

$$\mu: R \rightarrow R \otimes R^{\text{op}} \otimes R,$$

subject to certain axioms, see Section 2. We start with a Hopf-Galois algebra R , and discuss the ingredients with necessary and sufficient conditions to extend the Hopf-Galois structure of R to $A(R, X, Y, \sigma, c, \xi)$ via the formula

$$(1.2) \quad \begin{aligned} \mu(X) &= X \otimes 1 \otimes 1 - g \otimes g^{-1} X \otimes 1 + g \otimes g^{-1} \otimes X, \\ \mu(Y) &= Y \otimes 1 \otimes 1 - h \otimes h^{-1} Y \otimes 1 + h \otimes h^{-1} \otimes Y, \end{aligned}$$

for some quasi-central group-like elements $g, h \in G_0(R)$, see Definition 2.4.

Given a Hopf-Galois algebra (R, μ) as above, there exists a canonical Hopf algebra $\underline{H}(R)$ such that R is an $\underline{H}(R)$ -Galois object [10, 15]. As for the first item (A), the need of a generalization of the results in [4] becomes evident when one considers the Hopf algebra associated to the Hopf-Galois algebra $A_R = A(R, X, Y, \tau, c, \xi)$ extending a Hopf-Galois algebra R : this Hopf algebra is an extension of $\underline{H}(R)$ not in terms of an ambiskew polynomial algebra but of a generalized ambiskew polynomial algebra A . See Corollary 5.5 and Subsection 6.5 for an example.

The paper is organized as follows. In Section 2 we recall the definition of Hopf-Galois algebras (R, μ) and their realization as Hopf-Galois objects over a Hopf algebra $\underline{H}(R)$; we discuss the concepts of group-like and skew-primitive elements in this setting. Section 3 is devoted to the answer of item (A) above: in Theorem 3.1 we find necessary and sufficient conditions so that a generalized ambiskew polynomial algebra A_R extends the structure of a Hopf algebra R in a way such that the added variables are skew-primitive; we recover some of the results in [4] as a corollary. Next, in Section 4 we find necessary and sufficient conditions so that a generalized ambiskew polynomial algebra extends the structure of a Hopf-Galois algebra R as in (1.2) and we provide an answer for item (B) in Theorem 4.1. In Section 5 we show in Theorem 5.3 that the Hopf algebra associated to a Hopf-Galois ambiskew polynomial algebra A_R is a Hopf ambiskew polynomial algebra, associated to $\underline{H}(R)$. We conclude the article with a series of examples in Section 6.

Notation and conventions. We work over a field k . All algebras, vector spaces and unadorned tensor products \otimes are assumed to be defined over k . We assume that the reader has familiarity with Hopf algebra theory, for which the book [13] is a convenient reference. We use standard notations, in particular Δ , ε and S stand respectively for the comultiplication, counit and antipode of a Hopf algebra, and Sweedler's notation $h \mapsto h_{(1)} \otimes h_{(2)}$, resp. $m \mapsto m_{(0)} \otimes m_{(1)}$ for the comultiplication, resp. the coaction on a (right) comodule M , of a Hopf algebra H ; $h \in H, m \in M$

Acknowledgements. The authors were supported by CONICET, FONCyT PICT-2015-2845, Secyt (UNC), the MathAmSud project GR2HOPF, and the Emergence Program of the I-SITE project CAP 20-25 of Clermont Auvergne University.

2. HOPF-GALOIS ALGEBRAS

2.1. Basic definitions. The definition below was introduced by Grunspan [10] with more axioms, and simplified later by Schauenburg [15], under the name *quantum torsor*. In view of Theorem 2.8 given in the next subsection, it seems natural to call such a structure a Hopf-Galois algebra.

Definition 2.1. *A Hopf-Galois algebra is a non-zero algebra R together with an algebra map*

$$\mu: R \longrightarrow R \otimes R^{\text{op}} \otimes R$$

such that

$$(2.1) \quad (\mu \otimes \text{id}_R \otimes \text{id}_R) \circ \mu = (\text{id}_R \otimes \text{id}_R \otimes \mu) \circ \mu,$$

$$(2.2) \quad (m \otimes \text{id}_R) \circ \mu = \eta \otimes \text{id}_R, \quad (\text{id}_R \otimes m) \circ \mu = \text{id}_R \otimes \eta,$$

where $m : R \otimes R \rightarrow R$ and $\eta : k \rightarrow R$ denote the respective multiplication and unit of R .

We shall write

$$\mu(r) = r_{(1)} \otimes r_{(2)} \otimes r_{(3)}, \quad r \in R.$$

Hence, for any $r \in R$,

$$r \otimes 1 = r_{(1)} \otimes r_{(2)}r_{(3)}, \quad 1 \otimes r = r_{(1)}r_{(2)} \otimes r_{(3)};$$

and we may set

$$r_{(1)} \otimes r_{(2)} \otimes r_{(3)} \otimes r_{(4)} \otimes r_{(5)} := \mu(r_{(1)}) \otimes r_{(2)} \otimes r_{(3)} = r_{(1)} \otimes r_{(2)} \otimes \mu(r_{(3)}).$$

Example 2.2. The basic example is a Hopf algebra H with

$$\mu(x) = x_{(1)} \otimes S(x_{(2)}) \otimes x_{(3)}, \quad x \in R.$$

We denote by $\underline{R}(H)$ the resulting Hopf-Galois algebra. This defines a functor from the category of Hopf algebras to the category of Hopf-Galois algebras (a morphism of Hopf-Galois algebras being an algebra map commuting with μ in the obvious way).

A Hopf-Galois algebra arises from a Hopf algebra as above if and only if there exists an algebra map $\alpha : R \rightarrow k$. Indeed, starting with R and such an α , the Hopf algebra H whose counit is α and with the other structure maps defined by

$$\Delta(x) = \alpha(x_{(2)})x_{(1)} \otimes x_{(3)}, \quad S(x) = \alpha(x_{(1)}x_{(3)})x_{(2)}$$

is such that $R = \underline{R}(H)$.

Example 2.3. A more significant example is the Weyl algebra

$$A_1(k) = k\langle x, y \mid xy - yx = 1 \rangle$$

with Hopf-Galois structure defined by

$$\mu(x) = x \otimes 1 \otimes 1 - 1 \otimes x \otimes 1 + 1 \otimes 1 \otimes x, \quad \mu(y) = y \otimes 1 \otimes 1 - 1 \otimes y \otimes 1 + 1 \otimes 1 \otimes y.$$

More examples will be discussed in the final section. We now introduce group-like elements and skew-primitives in Hopf-Galois algebras.

Definition 2.4. A group-like element in a Hopf-Galois algebra R is an invertible element $g \in R$ such that

$$(2.3) \quad \mu(g) = g \otimes g^{-1} \otimes g.$$

We denote by $G_0(R)$ the set of group-like elements in R ; $g \in G_0(R)$ is quasi-central if there exists a character α on $G_0(R)$ such that

$$xg = \alpha(x)gx, \quad \text{for any } x \in G_0(R).$$

If $g \in G_0(R)$, then we write $g \cdot r := grg^{-1}$.

Remark 2.5. It should be noticed that, contrary to the Hopf algebra case, if $g \in R$ is group-like, then λg is grouplike for any nonzero scalar $\lambda \in k$. If we consider $G_0(R)$ acting on R by conjugation, then g and λg define the same operator.

Definition 2.6. Let R be a Hopf-Galois algebra and let $g, h \in G_0(R)$. The set $\mathcal{P}_{g,h}^0(R)$ of (g, h) -skew primitive elements is the collection of those $x \in R$ such that

$$\mu(x) = x \otimes h^{-1} \otimes h - g \otimes g^{-1}xh^{-1} \otimes h + g \otimes g^{-1} \otimes x.$$

Remark 2.7. If $x \in \mathcal{P}_{g,h}^0(R)$, then $h^{-1}x \in \mathcal{P}_{h^{-1}g,1}^0(R)$ and $xg^{-1} \in \mathcal{P}_{1,hg^{-1}}^0(R)$. Notice also that $g, h \in \mathcal{P}_{g,h}^0(R)$. Indeed,

$$\mu(h) = h \otimes h^{-1} \otimes h = h \otimes h^{-1} \otimes h - g \otimes g^{-1}hh^{-1} \otimes h + g \otimes g^{-1} \otimes h.$$

and similarly for g .

2.2. From Hopf-Galois algebras to Hopf algebras. We recall that a (right) Hopf-Galois object R over a Hopf algebra H is a (right) comodule algebra R over H with trivial coinvariants $R^{\text{co}H} = k$ and such that the *Galois map*

$$\text{can}: R \otimes R \rightarrow R \otimes H, \quad \text{can}(r \otimes s) = rs_{(0)} \otimes s_{(1)}, \quad r, s \in R$$

is bijective.

The following result relates what we called Hopf-Galois algebras and classical Hopf-Galois objects.

Theorem 2.8 (Grunspan-Schauenburg, [10, 15]). *Let R be an algebra. The following assertions are equivalent.*

- (1) *There exists an algebra map $\mu: R \rightarrow R \otimes R^{\text{op}} \otimes R$ making R into a Hopf-Galois algebra.*
- (2) *R is a right Hopf-Galois object over some Hopf algebra $\underline{H}(R)$.*

Therefore the notion of Hopf-Galois algebra provides a definition of Hopf-Galois object which is Hopf algebra free.

The construction of $\underline{H}(R)$ in the theorem is as follows. Let $R = (R, \mu)$ be a Hopf-Galois algebra. Then R is a right Hopf-Galois object for the Hopf algebra $\underline{H}(R)$ defined as the subalgebra

$$(2.4) \quad \underline{H}(R) = \{x \otimes y : xy_{(1)} \otimes y_{(2)} \otimes y_{(3)} = 1 \otimes x \otimes y\} \subseteq R^{\text{op}} \otimes R$$

with coalgebra structure

$$(2.5) \quad \Delta(x \otimes y) = x \otimes y_{(1)} \otimes y_{(2)} \otimes y_{(3)}, \quad \varepsilon(x \otimes y) = xy.$$

The coaction is given by μ , as $\mu: R \rightarrow R \otimes \underline{H}(R)$.

Remark 2.9. Here, we follow Schauenburg in [15, §3] and we write a generic element $\sum x_i \otimes y_i \in R^{\text{op}} \otimes R$ as a single tensor $x \otimes y$, “in the spirit of Sweedler’s notation”. Definition (2.4) and formula (2.5) should be interpreted in this sense.

Conversely, let H be a Hopf algebra and let R be a right Hopf-Galois object, with coaction $r \mapsto \rho(r) = r_{(0)} \otimes r_{(1)}$, $r \in R$. Then R is a Hopf-Galois algebra with

$$\mu(r) = r_{(0)} \otimes \text{can}^{-1}(1 \otimes r_{(1)}).$$

We now use the construction $\underline{H}(R)$ to relate the group-likes and skew-primitives of the previous subsection with ordinary group-likes and skew-primitives in a Hopf algebra.

Lemma 2.10. *Let R be a Hopf-Galois algebra.*

- (1) *There is a short exact sequence of groups*

$$1 \rightarrow k^* \rightarrow G_0(R) \xrightarrow{\varphi} G(\underline{H}(R)) \rightarrow 1,$$

where $\varphi(g) = g^{-1} \otimes g$.

- (2) *For any $g \in G_0(R)$, there is a linear short exact sequence*

$$0 \rightarrow k \rightarrow \mathcal{P}_{g,1}^0(R) \xrightarrow{\psi} \mathcal{P}_{\varphi(g),1}(\underline{H}(R)) \rightarrow 0,$$

where $\psi(x) = g^{-1} \otimes x - g^{-1}x \otimes 1$.

Proof. (1) On the one hand, we see that $k^* \subset G_0(R)$, as in Remark 2.5. If $g \in G_0(R)$, then it is clear that $g^{-1} \otimes g \in G(\underline{H}(R))$. Moreover, $\varphi(\lambda) = 1 \otimes 1$ for any $\lambda \in k^*$.

Now, if $u = x \otimes y = \sum_i x_i \otimes y_i \in G(\underline{H}(R))$, then $\varepsilon(x \otimes y) = xy = \sum_i x_i y_i = 1$ and

$$\Delta(x \otimes y) = x \otimes y \otimes x \otimes y = \sum_{i,j} x_i \otimes y_i \otimes x_j \otimes y_j.$$

On the other hand, by definition

$$\Delta(x \otimes y) = x \otimes y_{(1)} \otimes y_{(2)} \otimes y_{(3)}.$$

Hence, by multiplying the factors in the middle, we get

$$x \otimes 1 \otimes y = \sum_i x_i \otimes 1 \otimes y_i = \sum_{i,j} x_i \otimes y_i x_j \otimes y_j.$$

If we assume that $\{x_i\}_i$ is linearly independent, we obtain, for any i , that $1 \otimes y_i = \sum_j y_i x_j \otimes y_j$ and hence

$$1 \otimes \mu(y_i) = \sum_j y_i x_j \otimes \mu(y_j).$$

Multiplying the left handed tensors, this gives

$$\mu(y_i) = \sum_j y_i x_j y_{j(1)} \otimes y_{j(2)} \otimes y_{j(3)} = \sum_j y_i \otimes x_j \otimes y_j = y_i \otimes u.$$

Applying the same reasoning to $u^{-1} = \sum_k x'_k \otimes y'_k$ yields

$$\mu(y'_k) = y'_k \otimes u^{-1}$$

for any k . Hence for any i, k , we have $\mu(y_i y'_k) = y_i y'_k \otimes 1 \otimes 1$ and $\mu(y'_k y_i) = y'_k y_i \otimes 1 \otimes 1$, so that there exists scalars $\lambda_{i,k}, \mu_{k,i}$ such that $y_i y'_k = \lambda_{i,k}$ and $y'_k y_i = \mu_{k,i}$. There exists j, l such that $y_j y'_l \neq 0$, so we see from the previous identities that $\lambda_{j,l} = \mu_{l,j}$ and that y'_l is invertible in R . Hence for any i we have $y_i = \lambda_{i,l} (y'_l)^{-1}$. We get $u = (\sum_i \lambda_{i,l} x_i) \otimes (y'_l)^{-1} = g^{-1} \otimes g$ for $g = (y'_l)^{-1}$, and we then see that $g \in G_0(R)$.

(2) We know that $k \subset \mathcal{P}_{g,1}^0(R)$, see Remark 2.7. Also, it is easy to check that $g^{-1} \otimes x - g^{-1} x \otimes 1 \in P_{\varphi(g),1}(\underline{H}(R))$ if $x \in \mathcal{P}_{1,g}^0(R)$, and that $\text{Ker}(\psi) = k$.

Now, let $u = x \otimes y = \sum x_i \otimes y_i \in \mathcal{P}_{\varphi(g),1}(\underline{H}(R))$. We assume that $\{y_i\}$ is linearly independent. We have $x y_{(1)} \otimes y_{(2)} \otimes y_{(3)} = 1 \otimes x \otimes y$ and

$$(2.6) \quad x \otimes y_{(1)} \otimes y_{(2)} \otimes y_{(3)} = x \otimes y \otimes 1 \otimes 1 + g^{-1} \otimes g \otimes x \otimes y$$

so that multiplying the factors in the middle as before:

$$\begin{aligned} x \otimes 1 \otimes y &= x \otimes y \otimes 1 + g^{-1} \otimes g x \otimes y, \text{ i.e.} \\ \sum_i x_i \otimes 1 \otimes y_i &= \sum_i x_i \otimes y_i \otimes 1 + \sum_i g^{-1} \otimes g x_i \otimes y_i. \end{aligned}$$

If $1 \notin \text{Span}\{y_i\}$, we get that $u = \sum_i x_i \otimes y_i = 0$. Otherwise, there exists for any i , a scalar λ_i such that

$$x_i \otimes 1 = g^{-1} \otimes g x_i + \lambda_i u$$

with $\lambda_k \neq 0$ for some k . This gives

$$u = g^{-1} \otimes z - g^{-1} z \otimes 1$$

for $z = -\lambda_k^{-1} g x_k$. Going back to the expression (2.6), we get

$$g^{-1} \otimes \mu(z) - g^{-1} z \otimes 1 \otimes 1 \otimes 1 = g^{-1} \otimes z \otimes 1 \otimes 1 - g^{-1} z \otimes 1 \otimes 1 \otimes 1 + g^{-1} \otimes g \otimes g^{-1} \otimes z - g^{-1} \otimes g \otimes g^{-1} z \otimes 1.$$

This shows that $z \in \mathcal{P}_{g,1}^0(R)$, and we have $u = \psi(z)$. \square

3. AMBISKEW HOPF ALGEBRAS

In this section we assume that R is a Hopf algebra and consider the generalized ambiskew algebra $A = A(R, E, F, \tau, \omega, c, \xi)$ as in the introduction. We look for necessary and sufficient conditions to extend the comultiplication in R to A via the formulas

$$(3.1) \quad \Delta(E) = E \otimes 1 + g \otimes E, \quad \Delta(F) = F \otimes 1 + h \otimes F,$$

for some (necessarily group-like) elements $g, h \in G(R)$.

When the element c is assumed to be central, or equivalently when $\omega = \tau^{-1}$ (so $\sigma = \text{id}$), this question has been addressed in [4]. Our result reads as follows.

Theorem 3.1. *Let H be a Hopf algebra, let $\tau, \omega \in \text{Aut } H$ be such that $\tau\omega = \omega\tau$ and define $\sigma = \tau\omega$. Let $c \in Z_\sigma(H)$ be σ -central and $\xi \in k^*$. Then the generalized ambiskew polynomial algebra $A(H, E, F, \tau, \omega, c, \xi)$ has a Hopf algebra structure extending that of H and such that (3.1) holds for some $g, h \in G(H)$ if and only if*

- there are characters α, β of H , with $\alpha * \beta = \beta * \alpha$ and

$$(3.2) \quad \tau(r) = \alpha(r_{(1)})r_{(2)}, \quad \omega(r) = \beta(r_{(1)})r_{(2)}.$$

$$(3.3) \quad \alpha(r_{(1)})r_{(2)} = g \cdot r_{(1)}\alpha(r_{(2)}), \quad \beta(r_{(1)})r_{(2)} = h \cdot r_{(1)}\beta(r_{(2)}) \quad r \in H,$$

- the group-like elements g, h are central in $G(H)$ and $\alpha(h) = \beta(g)^{-1} = \xi$,
- the σ -central element c is $(gh, 1)$ -skew primitive.

Proof. Assume $A = A(H, E, F, \tau, \omega, c, \xi)$ is a Hopf algebra as in the statement. Then, by comparing $\Delta(Xr)$ and $\Delta(\tau(r)X)$ we obtain that

$$\tau(r_{(1)}) \otimes r_{(2)} = \tau(r)_{(1)} \otimes \tau(r)_{(2)} = g \cdot r_{(1)} \otimes \tau(r_{(2)}).$$

Hence, if $\alpha = \varepsilon \circ \tau : H \rightarrow k$ we get that $\alpha(r_{(1)})r_{(2)} = g \cdot r_{(1)}\alpha(r_{(2)})$ and $\tau(r) = \alpha(r_{(1)})r_{(2)}$. Similarly, by comparing $\Delta(Yr)$ and $\Delta(\omega(r)Y)$ we obtain that $\omega(r) = \beta(r_{(1)})r_{(2)}$ for $\beta = \varepsilon \circ \omega$ and both identities in (3.3) hold. Also, since $\tau\omega = \omega\tau$, it follows that $\alpha * \beta = \beta * \alpha$.

Now, observe that (3.3) applied to $r = h$ gives $g \cdot h = h$, that is $gh = hg$. We use this to see that

$$\begin{aligned} \Delta(EF - \xi FE) &= (EF - \xi FE) \otimes 1 + gh \otimes (EF - \xi FE) \\ &\quad + (\alpha(h) - \xi)hE \otimes F + (1 - \xi\beta(g))gF \otimes E \\ &= \Delta(c) \in H \otimes H \end{aligned}$$

and thus, since A is a free R -module with basis $\{X^a Y^b : a, b \geq 0\}$, we have $\alpha(h) = \beta(g)^{-1} = \xi$ and $\Delta(c) = c \otimes 1 + gh \otimes c$, so $c = EF - \xi FE$ is a σ -central $(gh, 1)$ -skew primitive element. The previous computations allow as well to prove converse statement. \square

Remark 3.2. Notice that $g \cdot c = g\sigma(g)^{-1}c = \alpha(g)^{-1}\beta(g)^{-1}c = \xi\alpha(g)^{-1}c$. We can thus compute $\tau(c)$ in two ways and get

$$\xi(\alpha(g)^{-1} - \alpha(g))c = \alpha(c)(1 - gh).$$

So either $\alpha(g) = \pm 1$ (and $\alpha(c) = 0$ or $1 = gh$) or $c = \frac{\alpha(ghc)}{1 - \alpha(g)^2}(1 - gh)$.

Similarly, if we compute $\omega(c)$, we get that

$$\xi^{-1}(\beta(h)^{-1} - \beta(h))c = \beta(c)(1 - gh),$$

that is $\beta(h) = \pm 1$ (and $\beta(c) = 0$ or $1 = gh$) or $c = \frac{\beta(ghc)}{1 - \beta(h)^2}(1 - gh)$.

Theorem 3.1 extends the result in [4] as mentioned; we recall this here as a corollary.

Corollary 3.3. [4] *Assume c is central (i.e. $\omega = \tau^{-1}$ so $\sigma = \text{id}$). Then the ambiskew polynomial algebra $A(R, E, F, \tau, c, \xi) = A(R, E, F, \tau, \tau^{-1}, c, \xi)$ has a Hopf algebra structure extending that of H and such that (3.1) holds for some $g, h \in G(H)$ if and only if there are*

- a character χ of R ,
- central group-like elements $g, h \in Z(G(R))$ such that $gh \in Z(R)$ and

$$\chi(g) = \chi(h) = \xi,$$

- a central $(gh, 1)$ -skew primitive element c ,

such that the following² holds:

$$\chi(r_{(1)})r_{(2)} = g \cdot r_{(1)}\chi(r_{(2)}), \quad r \in R.$$

²In the notation from [4], this condition reads $\tau_\chi^l = \text{ad}_l(y_+) \circ \tau_\chi^r$ and then $\tau = \tau_\chi^l$.

In this setting,

$$\tau(r) = \chi(r_{(1)})r_{(2)} = g \cdot r_{(1)}\chi(r_{(2)}).$$

Moreover,

$$(3.4) \quad \begin{aligned} & \text{either } \bullet \xi = \pm 1; \text{ (and } gh = 1 \text{ or } \chi(c) = 0), \\ & \text{or else } \bullet \xi \neq \pm 1 \text{ and } c = \lambda(1 - gh), \text{ for } \lambda = \frac{\chi(c)}{1 - \xi^2} \in k. \end{aligned}$$

□

Remark 3.4. Conditions (3.4) are stated in [4, §3.1], in this setting they follow by Remark 3.2.

4. AMBISKEW HOPF-GALOIS ALGEBRAS

Let (R, μ) be a Hopf-Galois algebra and let us fix $A = A(R, X, Y, \tau, \omega, c, \xi)$ an associated generalized ambis skew polynomial algebra. We investigate when A is again a Hopf-Galois algebra, extending R .

Theorem 4.1. *Let $R = (R, \mu)$ be a Hopf-Galois algebra. Assume given a sextuple $(\tau, \omega, g, h, c, \xi)$ where*

$$(1) \quad \tau, \omega \in \text{Aut}(R) \text{ are commuting algebra automorphisms satisfying, for any } r \in R, \\ (4.1) \quad \tau(r)_{(1)} \otimes \tau(r)_{(2)} \otimes \tau(r)_{(3)} = \tau(r_{(1)}) \otimes r_{(2)} \otimes r_{(3)} = \tau(r_{(1)}) \otimes \tau(r_{(2)}) \otimes \tau(r_{(3)}), \\ (4.2) \quad \omega(r)_{(1)} \otimes \omega(r)_{(2)} \otimes \omega(r)_{(3)} = \omega(r_{(1)}) \otimes r_{(2)} \otimes r_{(3)} = \omega(r_{(1)}) \otimes \omega(r_{(2)}) \otimes \omega(r_{(3)}).$$

$$(2) \quad g, h \in G_0(R) \text{ are quasi-central group-like elements such that, for any } r \in R, \\ (4.3) \quad g \cdot r_{(1)} \otimes g \cdot r_{(2)} \otimes r_{(3)} = \tau(r_{(1)}) \otimes \tau(r_{(2)}) \otimes r_{(3)}, \\ (4.4) \quad h \cdot r_{(1)} \otimes h \cdot r_{(2)} \otimes r_{(3)} = \omega(r_{(1)}) \otimes \omega(r_{(2)}) \otimes r_{(3)}.$$

$$(3) \quad \xi \in k^* \text{ is a scalar such that } \tau(h) = \xi h, \omega(g) = \xi^{-1} g.$$

$$(4) \quad c \in Z_\sigma(R) \cap \mathcal{P}_{gh,1}(R), \text{ for } \sigma := \tau\omega; \text{ that is } c \text{ is a } \sigma\text{-central skew-primitive element:}$$

$$\mu(c) = c \otimes 1 \otimes 1 - gh \otimes (gh)^{-1}c \otimes 1 + gh \otimes (gh)^{-1} \otimes c$$

and $cr = \sigma(r)rc$, for all $r \in R$.

Then there is a unique Hopf-Galois algebra structure on $A = A(R, X, Y, \tau, \omega, c, \xi)$ extending μ and such that

$$(4.5) \quad \begin{aligned} \mu(X) &= X \otimes 1 \otimes 1 - g \otimes g^{-1}X \otimes 1 + g \otimes g^{-1} \otimes X, \\ \mu(Y) &= Y \otimes 1 \otimes 1 - h \otimes h^{-1}Y \otimes 1 + h \otimes h^{-1} \otimes Y. \end{aligned}$$

Conversely, given commuting $\tau, \omega \in \text{Aut}(R)$, $c \in Z_\sigma(R)$ for $\sigma = \tau\omega$ and $\xi \in k^*$, if (4.5) defines a Hopf-Galois algebra structure on $A(R, X, Y, \tau, \omega, c, \xi)$ extending μ , for some $g, h \in G_0(R)$, then conditions (1) to (4) are satisfied.

Proof. We assume there is an extension $\mu : A \rightarrow A \otimes A^{\text{op}} \otimes A$ and elements $g, h \in G(R)$ such that (4.5) holds. Recall the notation $x \cdot r := xrx^{-1}$, for $x \in G(R)$, $r \in R$.

◇ *The commutation rule.* On the one hand, we have, in $A \otimes A^{\text{op}} \otimes A$,

$$\mu(Xr) = \mu(\tau(r)X) = (\tau(r)_{(1)} \otimes \tau(r)_{(2)} \otimes \tau(r)_{(3)})\mu(X).$$

On the other,

$$\begin{aligned} \mu(Xr) &= Xr_{(1)} \otimes r_{(2)} \otimes r_{(3)} - gr_{(1)} \otimes r_{(2)}g^{-1}X \otimes r_{(3)} + gr_{(1)} \otimes r_{(2)}g^{-1} \otimes Xr_{(3)} \\ &= \tau(r_{(1)})X \otimes r_{(2)} \otimes r_{(3)} - (g \cdot r_{(1)})g \otimes g^{-1}X\tau^{-1}(g \cdot r_{(2)}) \otimes r_{(3)} \\ &\quad + (g \cdot r_{(1)})g \otimes g^{-1}(g \cdot r_{(2)}) \otimes \tau(r_{(3)})X \\ &= (\tau(r_{(1)}) \otimes r_{(2)} \otimes r_{(3)})(X \otimes 1 \otimes 1) - (g \cdot r_{(1)} \otimes \tau^{-1}(g \cdot r_{(2)}) \otimes r_{(3)})(g \otimes g^{-1}X \otimes 1) \\ &\quad + (g \cdot r_{(1)} \otimes g \cdot r_{(2)} \otimes \tau(r_{(3)}))(g \otimes g^{-1} \otimes X). \end{aligned}$$

Since A is a free R -module with basis $\{X^a Y^b : a, b \in \mathbb{N}\}$, we get the following identities:

$$\begin{aligned}\tau(r)_{(1)} \otimes \tau(r)_{(2)} \otimes \tau(r)_{(3)} &= \tau(r_{(1)}) \otimes r_{(2)} \otimes r_{(3)} = g \cdot r_{(1)} \otimes g \cdot r_{(2)} \otimes \tau(r_{(3)}) \\ g \cdot r_{(1)} \otimes g \cdot r_{(2)} \otimes r_{(3)} &= \tau(r_{(1)}) \otimes \tau(r_{(2)}) \otimes r_{(3)}\end{aligned}$$

We may combine these identities to show that $\mu \circ \tau = \tau^{\otimes 3} \circ \mu$:

$$\tau(r_{(1)}) \otimes r_{(2)} \otimes r_{(3)} = g \cdot r_{(1)} \otimes g \cdot r_{(2)} \otimes \tau(r_{(3)}) = \tau(r_{(1)}) \otimes \tau(r_{(2)}) \otimes \tau(r_{(3)})$$

and thus we see that our identities are then equivalent to (4.1) and (4.3). Similarly, from the identity $Yr = \omega(r)Y$ we obtain (4.2) and (4.4).

◇ *The group characters.* We make the following

Claim. There are characters $\alpha, \beta, \gamma, \delta \in \widehat{G_0(R)}$ such that for any $x \in G_0(R)$,

$$(4.6) \quad \tau(x) = \alpha(x)x \quad \omega(x) = \beta(x)x,$$

$$(4.7) \quad \tau(x) = \gamma(x)g \cdot x, \quad \omega(x) = \delta(x)h \cdot x.$$

Hence, $\alpha(h)\beta(g) = \gamma(h)\delta(g)$ and both g, h are quasi-central group-like elements in $G_0(R)$ with

$$(4.8) \quad gh(hg)^{-1} \in k^* \subset Z(R), \quad (gh) \otimes (gh)^{-1} = (hg) \otimes (hg)^{-1}.$$

As for the proof, notice that combining (2.2) and (4.1) we get

$$\tau(x)x^{-1} \otimes x = 1 \otimes \tau(x), \quad x \in G_0(R).$$

Hence $\tau(x)$ is a scalar multiple of x , and we get a character $\alpha \in \widehat{G_0(R)}$. Similarly for ω ; we call the corresponding character β and (4.6) follows.

Now, we see from (4.3) that there is a character $\chi \in \widehat{G_0(R)}$ such that $g \cdot x = \chi(x)x$; which shows that g is a quasi-central group-like element. Hence $\tau(x) = \gamma(x)g \cdot x$, for $\gamma(x) = \alpha(x)\chi(x)^{-1}$. Analogously, $h \in G(R)$ is quasi-central and $\omega(x) = \delta(x)h \cdot x$, for some $\delta \in \widehat{G_0}$; (4.7) follows.

Finally, we see that

$$(4.9) \quad gh = \alpha(h)\gamma(h)^{-1}hg,$$

whence $gh(hg)^{-1} \in k^\times$ and (4.8) follows.

◇ *The bracket rule.* Recall that, for any $t \in G_0(R)$, $Xt = \alpha(t)tX$ and $Yt = \beta(t)tY$. Also, observe that both XY and YX are σ -central, $\sigma = \tau\omega$, in A_R and hence $XY - \xi YX \in Z_\sigma(R)$. Now,

$$\begin{aligned}\mu(XY) &= XY \otimes 1 \otimes 1 - Xh \otimes h^{-1}Y \otimes 1 + Xh \otimes h^{-1} \otimes Y \\ &\quad - gY \otimes g^{-1}X \otimes 1 + gh \otimes h^{-1}Yg^{-1}X \otimes 1 - gh \otimes h^{-1}g^{-1}X \otimes Y \\ &\quad + gY \otimes g^{-1} \otimes X - gh \otimes h^{-1}Yg^{-1} \otimes X + gh \otimes h^{-1}g^{-1} \otimes XY \\ &= XY \otimes 1 \otimes 1 - \alpha(h)hX \otimes h^{-1}Y \otimes 1 + \alpha(h)hX \otimes h^{-1} \otimes Y \\ &\quad - gY \otimes g^{-1}X \otimes 1 + \beta(g^{-1})gh \otimes (gh)^{-1}YX \otimes 1 - gh \otimes (gh)^{-1}X \otimes Y \\ &\quad + gY \otimes g^{-1} \otimes X - \beta(g^{-1})gh \otimes (gh)^{-1}Y \otimes X + gh \otimes (gh)^{-1} \otimes XY.\end{aligned}$$

On the other hand, using (4.8),

$$\begin{aligned}\mu(YX) &= YX \otimes 1 \otimes 1 - \beta(g)gY \otimes g^{-1}X \otimes 1 + \beta(g)gY \otimes g^{-1} \otimes X \\ &\quad - hX \otimes h^{-1}Y \otimes 1 + \alpha(h^{-1})gh \otimes (gh)^{-1}XY \otimes 1 - gh \otimes (gh)^{-1}Y \otimes X \\ &\quad + hX \otimes h^{-1} \otimes Y - \alpha(h^{-1})gh \otimes (gh)^{-1}X \otimes Y + gh \otimes (gh)^{-1} \otimes YX.\end{aligned}$$

As we request $XY - \xi YX \in R$, we see that

$$(4.10) \quad \alpha(h) = \xi, \quad \beta(g) = \xi^{-1},$$

and there is a σ -central skew-primitive element $c \in Z_\sigma(R) \cap \mathcal{P}_{gh,1}(R)$, so that $XY - \xi YX = c$. As a result, $\tau(h) = \xi h$, $\omega(g) = \xi^{-1}g$.

◇ *On the converse.* The previous computations show that given a Hopf-Galois algebra (R, μ) , if there is a quintuple $(\tau, \omega, g, h, c, \xi)$ satisfying conditions (1) to (4), then (4.5) defines a Hopf-Galois structure on A that extends μ .

This ends the proof of the theorem. \square

We write down the case $\omega = \tau^{-1}$ for completeness. We start with a remark that will be of importance further on.

Remark 4.2. We point out that even in the case $\omega = \tau^{-1}$ we do not have $gh = hg$ as in the Hopf algebra case, cf. Theorem 3.1 and Corollary 3.3. Instead, we have that g, h commute up to a scalar, see (4.8). In particular,

$$(4.11) \quad gh \cdot r = \alpha(h)\gamma(h)^{-1} r, \quad r \in R.$$

Corollary 4.3. *Let $R = (R, \mu)$ be a Hopf-Galois algebra. Assume given a quintuple (τ, g, h, c, ξ) where*

(1) $\tau \in \text{Aut}(R)$ is an algebra automorphism satisfying, for any $r \in R$,

$$(4.12) \quad \tau(r)_{(1)} \otimes \tau(r)_{(2)} \otimes \tau(r)_{(3)} = \tau(r_{(1)}) \otimes r_{(2)} \otimes r_{(3)} = \tau(r_{(1)}) \otimes \tau(r_{(2)}) \otimes \tau(r_{(3)}).$$

(2) $g, h \in G_0(R)$ are quasi-central group-like elements such that, for any $r \in R$,

$$(4.13) \quad g \cdot r_{(1)} \otimes g \cdot r_{(2)} \otimes r_{(3)} = \tau(r_{(1)}) \otimes \tau(r_{(2)}) \otimes r_{(3)},$$

$$(4.14) \quad h \cdot r_{(1)} \otimes h \cdot r_{(2)} \otimes r_{(3)} = \tau^{-1}(r_{(1)}) \otimes \tau^{-1}(r_{(2)}) \otimes r_{(3)}.$$

(3) $\xi \in k^*$ is a scalar such that $\tau(g) = \xi g$ and $\tau(h) = \xi h$.

(4) $c \in Z(R) \cap \mathcal{P}_{gh,1}(R)$ is a central skew-primitive element, i.e.

$$\mu(c) = c \otimes 1 \otimes 1 - gh \otimes (gh)^{-1} c \otimes 1 + gh \otimes (gh)^{-1} \otimes c.$$

Then there is a unique Hopf-Galois algebra structure on $A = A(R, X, Y, \tau, c, \xi)$ extending μ and such that (4.5) holds.

Conversely, given $\tau \in \text{Aut}(R)$, $c \in Z(R)$ and $\xi \in k^*$, if (4.5) defines a Hopf-Galois algebra structure on $A(R, X, Y, \tau, c, \xi)$ extending μ , for some $g, h \in G_0(R)$, then conditions (1) to (4) are satisfied.

Proof. This is Theorem 4.1 for $\omega = \tau^{-1}$. In particular, notice that the four conditions (4.1)–(4.4) become just three, as we obtain (4.12) and (4.13), together with the additional (4.14). For the remaining identity in (4.2), namely

$$\tau^{-1}(r_{(1)}) \otimes r_{(2)} \otimes r_{(3)} = \tau^{-1}(r_{(1)}) \otimes \tau^{-1}(r_{(2)}) \otimes \tau^{-1}(r_{(3)}),$$

we remark that it follows by applying $\tau^{-2} \otimes \tau^{-1} \otimes \tau^{-1}$ to the corresponding identity in (4.1), as $\mu \circ \tau^{-1} = (\tau^{-1})^{\otimes 3} \circ \mu$. \square

5. THE HOPF ALGEBRA ASSOCIATED TO AN AMBISKEW HOPF-GALOIS ALGEBRA

Let (R, μ) be a Hopf-Galois algebra. Recall that this is equivalent to a structure of Hopf-Galois object over certain Hopf algebra $\underline{H}(R)$, by Theorem 2.8. In this section, we study the corresponding Hopf algebra $\underline{H}(A_R)$, where $A_R = A(R, X, Y, \tau, \omega, c, \xi)$ is the generalized ambiskew polynomial algebra that extends (R, μ) as in Theorem 4.1. Recall that $\sigma = \tau\omega = \omega\tau$.

We start with a technical lemma. We shall stick to the notation $r \otimes s := \sum_i r_i \otimes s_i \in \underline{H}(R)$ for a generic element in $\underline{H}(R)$, see Remark 2.9.

Lemma 5.1. *Let (R, μ_R) and A_R be as above. Then*

(1) $\underline{H}(R) \subset \underline{H}(A_R)$.

(2) $(g \otimes g^{-1})(h \otimes h^{-1}) = (h \otimes h^{-1})(g \otimes g^{-1})$.

(3) If $r \otimes s \in \underline{H}(R)$, then

$$\tau(r) \otimes \tau(s) = r \otimes s,$$

$$\omega(r) \otimes \omega(s) = r \otimes s.$$

(4) If $r \otimes s \in \underline{H}(R)$, then

$$g \cdot r \otimes \tau(s) \in \underline{H}(R), \quad h \cdot r \otimes \omega(s) \in \underline{H}(R).$$

In particular, $(gh \cdot r) \otimes \sigma(s) \in \underline{H}(R)$.

Proof. (1) This is clear, as $\mu_{A|R} = \mu_R$. (2) is (4.8).

(3) Recall that $r \otimes s \in \underline{H}(R)$ if and only if

$$rs_{(1)} \otimes s_{(2)} \otimes s_{(3)} = 1 \otimes r \otimes s.$$

If we apply $\tau^{\otimes 3}$ to this equality, as $\tau(1) = 1$, then we get

$$1 \otimes \tau(r) \otimes \tau(s) = \tau(rs_{(1)}) \otimes \tau(s_{(2)}) \otimes \tau(s_{(3)}) \stackrel{(4.1)}{=} \tau(rs_{(1)}) \otimes s_{(2)} \otimes s_{(3)} = 1 \otimes r \otimes s.$$

The result for ω is analogous.

(4) By (4.1) and (4.3):

$$(g \cdot r)\tau(s)_{(1)} \otimes \tau(s)_{(2)} \otimes \tau(s)_{(3)} = (g \cdot r)(g \cdot s_{(1)}) \otimes g \cdot s_{(2)} \otimes \tau(s_{(3)}) = g \cdot 1 \otimes g \cdot r \otimes \tau(s).$$

Thus $g \cdot r \otimes \tau(s) \in \underline{H}(R)$. The other case is similar. \square

We use Lemma 5.1 to define algebra automorphisms $\tau', \omega' \in \text{Aut } \underline{H}(R)$ via the formulas

$$(5.1) \quad \tau'(r \otimes s) = g \cdot r \otimes \tau(s), \quad \omega'(r \otimes s) = h \cdot r \otimes \omega(s), \quad r \otimes s \in \underline{H}(R).$$

Notice that $\tau'\omega' = \omega'\tau'$ as g, h are quasi-central and thus $gh \cdot r = hg \cdot r$, $r \in R$.

Remark 5.2. If $\tau\omega = \text{id}$, that is if $gh \cdot r \otimes s = r \otimes s$ for all $r \otimes s \in \underline{H}(R)$, then $gh \in Z(G_0(R))$ and $gh = hg$.

Proof. Indeed, let $t \in G_0(R)$ and fix $r \otimes s = t \otimes t^{-1} \in \underline{H}(R)$. Then $(gh \cdot t) \otimes t^{-1} = t \otimes t^{-1}$ which implies $gh \cdot t = t$, i.e. $gh \in Z(G_0(R))$. If $t = h$, then it follows that $gh = hg$. \square

Theorem 5.3. Let (R, μ) be a Hopf-Galois algebra, let $(\tau, \omega, g, h, c, \xi)$ be a sextuple satisfying the conditions of Theorem 4.1 and let $A_R = A(R, X, Y, \tau, \omega, c, \xi)$ be the associated Hopf-Galois algebra.

Consider $\tau', \omega' \in \text{Aut } \underline{H}(R)$ as in (5.1), the group-like elements $g^{-1} \otimes g, h^{-1} \otimes h \in G(\underline{H}(R))$, the scalar $\lambda \in k^*$ such that $gh = \lambda hg$, and the elements $c' \in \underline{H}(R)$ and $\xi' \in k^*$ defined by

$$c' = (gh)^{-1} \otimes c - (gh)^{-1}c \otimes 1, \quad \xi' = \xi\lambda^{-1}.$$

Then $A(\underline{H}(R), E, F, \tau', \omega', c', \xi')$ has a Hopf algebra structure as in Theorem 3.1, and we have a Hopf algebra isomorphism

$$\underline{H}(A_R) \simeq A(\underline{H}(R), E, F, \tau', \omega', c', \xi').$$

Proof. Step 1. We first have to show that the conditions in Theorem 3.1 are satisfied. Define $\alpha, \beta : \underline{H}(R) \rightarrow k$ by $\alpha(r \otimes s) = (g \cdot r)\tau(s)$ and $\beta(r \otimes s) = (h \cdot r)\omega(s)$. It is clear that α and β are characters on $\underline{H}(R)$. We have

$$\begin{aligned} \alpha((r \otimes s)_{(1)})(r \otimes s)_{(2)} &= \alpha(r \otimes s_{(1)})s_{(2)} \otimes s_{(3)} = (g \cdot r)\tau(s_{(1)})s_{(2)} \otimes s_{(3)} \\ &\stackrel{(4.1)}{=} (g \cdot r)\tau(s_{(1)})\tau(s_{(2)}) \otimes \tau(s_{(3)}) = g \cdot r \otimes \tau(s) = \tau'(r \otimes s) \end{aligned}$$

so that (3.2) holds for τ' , and for ω' . Similarly

$$\begin{aligned} g^{-1} \otimes g \cdot (r \otimes s)_{(1)}\alpha((r \otimes s)_{(2)}) &= g^{-1} \otimes g \cdot (r \otimes s_{(1)})\alpha(s_{(2)} \otimes s_{(3)}) = (g \cdot r \otimes g \cdot s_{(1)})g \cdot s_{(2)}\tau(s_{(3)}) \\ &\stackrel{(4.3)}{=} (g \cdot r \otimes \tau(s_{(1)}))\tau(s_{(2)})\tau(s_{(3)}) = g \cdot r \otimes \tau(s) = \tau'(r \otimes s) \end{aligned}$$

and (3.3) holds for τ' , and for ω' as well.

It is clear that τ' and ω' commute since τ and ω commute and g and h commute up to a scalar, so the characters α and β convolution commute.

We have

$$\alpha(h^{-1} \otimes h) = g \cdot h^{-1}\tau(h) = \lambda^{-1}h^{-1}\xi h = \xi'$$

and similarly $\beta(g^{-1} \otimes g) = \xi'^{-1}$. It is an immediate verification that $c' \in \underline{H}(R)$ and that c' is $((gh)^{-1} \otimes gh, 1)$ -primitive, with $(gh)^{-1} \otimes gh = (g^{-1} \otimes g)(h^{-1} \otimes h)$. We have finally, for $\sigma' = \tau'\omega'$,

$$\begin{aligned} c'(r \otimes s) &= r(gh)^{-1} \otimes cs - r(gh)^{-1}c \otimes s = (gh)^{-1}(gh) \cdot r \otimes \sigma(s)c - (gh)^{-1}(gh \cdot r)c \otimes s \\ &= (gh)^{-1}(gh) \cdot r \otimes \sigma(s)c - (gh)^{-1}c\sigma^{-1}(gh \cdot r) \otimes s \\ &= (gh)^{-1}(gh) \cdot r \otimes \sigma(s)c - (gh)^{-1}c(gh \cdot r) \otimes \sigma(s) \text{ (by Lemma 5.1)} \\ &= ((gh) \cdot r \otimes \sigma(s))c' = \sigma'(r \otimes s)c' \end{aligned}$$

and hence c' is σ' -central.

Step 2. We now check that $A_R = A(R, X, Y, \tau, \omega, c, \xi)$ has a natural $A(\underline{H}(R), E, F, \tau', \omega', c', \xi')$ -comodule algebra structure. We start with preliminary computations. We have

$$\begin{aligned} (g \otimes E + X \otimes 1 \otimes 1)r_{(1)} \otimes r_{(2)} \otimes r_{(3)} &= gr_{(1)} \otimes (E(r_{(2)} \otimes r_{(3)})) + Xr_{(1)} \otimes r_{(2)} \otimes r_{(3)} \\ &= gr_{(1)} \otimes (g \cdot r_{(2)} \otimes \tau(r_{(3)}E) + \tau(r_{(1)})X \otimes r_{(2)} \otimes r_{(3)}) \\ &\stackrel{(4.1)}{=} (g \cdot r_{(1)})g \otimes (g \cdot r_{(2)} \otimes \tau(r_{(3)}E)) + \tau(r_{(1)})X \otimes \tau(r_{(2)}) \otimes \tau(r_{(3)}) \\ &\stackrel{(4.3)}{=} \tau(r_{(1)})g \otimes (\tau(r_{(2)}) \otimes \tau(r_{(3)}E) + \tau(r_{(1)})X \otimes \tau(r_{(2)}) \otimes \tau(r_{(3)})) \\ &\stackrel{(4.1)}{=} \tau(r_{(1)})g \otimes (\tau(r_{(2)}) \otimes \tau(r_{(3)}E) + \tau(r_{(1)})X \otimes \tau(r_{(2)}) \otimes \tau(r_{(3)})) \\ &= \tau(r_{(1)}) \otimes \tau(r_{(2)}) \otimes \tau(r_{(3)}) (g \otimes E + X \otimes 1 \otimes 1) \end{aligned}$$

and similarly

$$(h \otimes F + Y \otimes 1 \otimes 1)r_{(1)} \otimes r_{(2)} \otimes r_{(3)} = \omega(r)_{(1)} \otimes \omega(r)_{(2)} \otimes \omega(r)_{(3)} (h \otimes F + Y \otimes 1 \otimes 1).$$

We also have

$$\begin{aligned} &(g \otimes E + X \otimes 1 \otimes 1)(h \otimes F + Y \otimes 1 \otimes 1) - \xi(h \otimes F + Y \otimes 1 \otimes 1)(g \otimes E + X \otimes 1 \otimes 1) \\ &= gh \otimes (EF - \xi'FE) + (XY - \xi YX) \otimes 1 \otimes 1 + (gY - \xi Yg) \otimes E + (Xh - \xi hX) \otimes F \\ &= gh \otimes c' + c \otimes 1 \otimes 1 = gh \otimes (gh)^{-1} \otimes c - gh \otimes (gh)^{-1}c \otimes 1 + c \otimes 1 \otimes 1 \\ &= c_{(1)} \otimes c_{(2)} \otimes c_{(3)} \end{aligned}$$

These computations show that we have an algebra map

$$A_R \longrightarrow A_R \otimes A(\underline{H}(R), E, F, \tau', \omega', c', \xi'),$$

defined by

$$R \ni r \longmapsto r_{(1)} \otimes r_{(2)} \otimes r_{(3)}, \quad X \longmapsto g \otimes E + X \otimes 1 \otimes 1, \quad Y \longmapsto h \otimes F + Y \otimes 1 \otimes 1$$

which is easily seen to furnish the expected comodule algebra structure.

Step 3. We now prove that the previous comodule algebra structure makes A_R a Hopf-Galois object over $A(\underline{H}(R)) = A(\underline{H}(R), E, F, \tau', \omega', c', \xi')$. For this, we claim that there exists an algebra map

$$\begin{aligned} \theta : A(\underline{H}(R)) &\longrightarrow A_R^{\text{op}} \otimes A_R \\ r \otimes s, E, F &\longmapsto r \otimes s, E' := g^{-1} \otimes X - g^{-1}X \otimes 1, F' := h^{-1} \otimes Y - h^{-1}Y \otimes 1 \end{aligned}$$

such that $r \otimes s \longmapsto r \otimes s$ and

$$E \longmapsto E' := g^{-1} \otimes X - g^{-1}X \otimes 1, \quad F \longmapsto F' := h^{-1} \otimes Y - h^{-1}Y \otimes 1.$$

To prove the claim, we have to check a number of identities. We have, for $r \otimes s \in \underline{H}(R)$,

$$\begin{aligned} E'(r \otimes s) &= rg^{-1} \otimes Xs - rg^{-1}X \otimes s = rg^{-1} \otimes \tau(s)X - rg^{-1}X \otimes s \\ &= (g \cdot r \otimes \tau(s))(g^{-1} \otimes X) - (\tau^{-1}(g \cdot r) \otimes s)(g^{-1}X \otimes 1) \\ &= (g \cdot r \otimes \tau(s))(g^{-1} \otimes X - g^{-1}X \otimes 1) \\ &= \tau'(r \otimes s)E' \end{aligned}$$

where the third equality holds since $g \cdot r \otimes \tau(s) = \tau^{-1}(g \cdot r) \otimes s$, by Lemma 5.1. Similarly $F'(r \otimes s) = \omega'(r \otimes s)F'$. Finally we see that:

$$\begin{aligned} E'F' &= h^{-1}g^{-1} \otimes XY - h^{-1}Yg^{-1} \otimes X - h^{-1}g^{-1}X \otimes Y + h^{-1}Yg^{-1}X \otimes 1 \\ &= (gh)^{-1} \otimes XY - \xi(gh)^{-1}Y \otimes X - (gh)^{-1}X \otimes Y + \xi(gh)^{-1}YX \otimes 1. \\ F'E' &= \lambda^{-1}(gh)^{-1} \otimes YX - \lambda^{-1}\xi^{-1}(gh)^{-1}X \otimes Y - \lambda^{-1}(gh)^{-1}Y \otimes X + \lambda^{-1}\xi^{-1}(gh)^{-1}XY \otimes 1. \end{aligned}$$

Hence we have

$$\begin{aligned} E'F' - \xi'F'E' &= (gh)^{-1} \otimes (XY - \xi YX) - (gh)^{-1}(XY - \xi YX) \otimes 1 \\ &= (gh)^{-1} \otimes c - (gh)^{-1}c \otimes 1 = c'. \end{aligned}$$

We have thus proved that the announced algebra map is well-defined. It is now an immediate verification that the composite map

$$A_R \otimes A(\underline{H}(R)) \xrightarrow{\text{id} \otimes \theta} A_R \otimes A_R \otimes A_R \xrightarrow{\text{mult} \otimes \text{id}} A_R \otimes A_R$$

is inverse to the Galois map in Section 2, so we conclude that A_R is a Hopf-Galois object over $A(\underline{H}(R))$.

Step 4. To finish the proof, first notice that the elements $E' = g^{-1} \otimes X - g^{-1}X \otimes 1$ and $F' = h^{-1} \otimes Y - h^{-1}Y \otimes 1$ from the previous step belong to $\underline{H}(A_R)$, and that the computations done in this previous step ensure the existence of a Hopf algebra map

$$f : A(\underline{H}(R)) \longrightarrow \underline{H}(R),$$

given by

$$r \otimes s \longmapsto r \otimes s, \quad E \longmapsto E', \quad F \longmapsto F'.$$

It is obvious that f commutes with the respective coactions on A_R , so using the bijective canonical maps of our two Hopf-Galois objects, we conclude that f is an isomorphism. \square

While the group-like and skew-primitive elements in a Hopf-Galois algebra R can be described from those of $\underline{H}(R)$ thanks to Lemma 2.10, the proof of the previous theorem also provides the description of the automorphisms in Theorem 4.1 in terms of characters of $\underline{H}(R)$:

Corollary 5.4. *Let (R, μ) be Hopf-Galois algebra, let $(\tau, \omega, g, h, c, \xi)$ be a sextuple satisfying the conditions of Theorem 4.1. Then there are convolution commuting characters $\alpha, \beta : \underline{H}(R) \rightarrow k$ such that for $r \in R$,*

$$\tau(r) = \alpha(r_{(2)} \otimes r_{(3)})g \cdot r_{(1)}, \quad \omega(r) = \beta(r_{(2)} \otimes r_{(3)})h \cdot r_{(1)}$$

and for $r \otimes s \in \underline{H}(R)$,

$$\begin{aligned} \alpha((r \otimes s)_{(1)})(r \otimes s)_{(2)} &= g^{-1} \otimes g \cdot (r \otimes s)_{(1)}\alpha((r \otimes s)_{(2)}), \\ \beta((r \otimes s)_{(1)})(r \otimes s)_{(2)} &= h^{-1} \otimes h \cdot (r \otimes s)_{(1)}\beta((r \otimes s)_{(2)}). \end{aligned}$$

Proof. We have, for any $r \in R$, by the construction of τ' ,

$$r_{(1)} \otimes \tau'(r_{(2)} \otimes r_{(3)}) = r_{(1)} \otimes g \cdot r_{(2)} \otimes \tau(r_{(3)})$$

hence

$$g \cdot r_{(1)} \otimes \tau'(r_{(2)} \otimes r_{(3)}) = g \cdot r_{(1)} \otimes g \cdot r_{(2)} \otimes \tau(r_{(3)}).$$

Multiplying everything, we get, by the definition of α in the previous proof the announced identity for τ and α , and similarly for ω and β . The final identities have been proved in the previous proof as well. \square

Notice that when $\omega = \tau^{-1}$, that is when $A_R = A(R, X, Y, \tau, c, \xi)$ is an ambiskew polynomial algebra, for some $\tau \in \text{Aut } R$, then we do not necessarily have $\omega' = \tau'^{-1}$, as we get

$$\tau'\omega'(g^{-1} \otimes g) = gh \cdot g^{-1} \otimes g = \lambda g^{-1} \otimes g.$$

Thus it becomes evident that we need to work with generalized ambiskew polynomial algebras rather than plain ambiskew polynomial algebras. We resume this situation in the next corollary, see §6.5 for a concrete example.

Corollary 5.5. *Let (R, μ) be a Hopf-Galois algebra and assume that $A_R = A(R, X, Y, \tau, c, \xi)$ is a Hopf-Galois algebra. Then the corresponding Hopf algebra $\underline{H}(A_R)$ is a generalized ambiskew polynomial algebra $A(\underline{H}(R), \tau', \omega', c', \xi')$, where $\tau', \omega' \in \text{Aut } \underline{H}(R)$ are commuting automorphisms not necessarily inverse to each other.*

6. EXAMPLES

6.1. Baby example: $R = k$. It is immediate that Theorem 4.1 produces only two examples: the Hopf algebra $k[X, Y]$ with primitive X, Y , and the Weyl algebra $A_1(k)$ as in Example 2.3.

6.2. Almost general example: cleft Galois objects. Before going into other specific examples, we examine the case of cleft Galois objects, i.e. those obtained by deforming the multiplication of a Hopf algebra by a Hopf 2-cocycle. This is the most studied type of Hopf-Galois objects, but there exist some that are not of this type, see [3].

Let H be a Hopf algebra and let $\pi : H \otimes H \rightarrow k$ be a 2-cocycle on H , i.e. π is convolution invertible, $\pi(x, 1) = \varepsilon(x) = \pi(1, x)$ for any $x \in H$, and we have, for any $x, y, z \in H$,

$$\pi(x_{(1)}, y_{(1)})\pi(x_{(2)}y_{(2)}, z) = \pi(y_{(1)}, z_{(1)})\pi(x, y_{(2)}z_{(2)}).$$

The algebra ${}_{\pi}H$ has H as underlying vector space, and product defined by

$$x \cdot y = \pi(x_{(1)}, y_{(1)})x_{(2)}y_{(2)}.$$

It follows from standard cocycle identities [5] that ${}_{\pi}H$ is a Hopf-Galois algebra with

$$\begin{aligned} \mu : {}_{\pi}H &\longrightarrow {}_{\pi}H \otimes {}_{\pi}H^{\text{op}} \otimes {}_{\pi}H \\ x &\longmapsto \pi^{-1}(S(x_{(3)}), x_{(4)})x_{(1)} \otimes S(x_{(2)}) \otimes x_{(5)} \end{aligned}$$

and that the map

$$\begin{aligned} H &\longrightarrow \underline{H}({}_{\pi}H) \subset {}_{\pi}H^{\text{op}} \otimes {}_{\pi}H \\ x &\longmapsto \pi^{-1}(S(x_{(2)}), x_{(3)})S(x_{(1)}) \otimes x_{(4)} \end{aligned}$$

is a Hopf algebra isomorphism. We thus see from Lemma 2.10 that

$$G_0({}_{\pi}H) = \{\lambda g, \lambda \in k^*, g \in G(H)\}$$

and for $g, h \in G(H)$,

$$\mathcal{P}_{g,h}^0({}_{\pi}H) = \{\lambda 1 + x, \lambda \in k, x \in \mathcal{P}_{g,h}(H)\}.$$

Using this description and Corollary 5.4, we then see that the sextuples $(\tau, \omega, g, h, c, \xi)$ as in Theorem 4.1 can be chosen as follows:

- (1) g, h are central group-like elements in H ;
- (2) the automorphisms $\tau, \omega \in \text{Aut } {}_{\pi}H$ are of the form

$$\tau(x) = \alpha(x_{(2)})g \cdot x_{(1)}, \quad \omega(x) = \beta(x_{(2)})h \cdot x_{(1)}$$

for some convolution commuting algebra maps $\alpha, \beta : H \rightarrow k$ that also satisfy

$$\alpha(x_{(2)})x_{(1)} = \alpha(x_{(1)})g \cdot x_{(2)}, \quad \beta(x_{(2)})x_{(1)} = \beta(x_{(1)})h \cdot x_{(2)}$$

(these last identities are in H) and $\alpha(h) = \beta(g)^{-1}$, $\xi = \alpha(h)\lambda$, where λ is such that $gh = \lambda hg$ in ${}_{\pi}H$.

- (3) c is σ -central for $\sigma = \tau\omega$, and $c = \gamma 1 + x$, for $\gamma \in k$ and $x \in \mathcal{P}_{gh,1}(H)$;

6.3. The example $R = k[T]$. We now examine the case when $R = \underline{R}(k[T])$. The analysis of the previous subsection applies: we have $G_0(R) = k^*$ and $\mathcal{P}_{1,1}^0(R) = k + kT$ and for the sextuples $(\tau, \omega, g, h, c, \xi)$ of Theorem 4.1, the automorphisms τ, ω are of the form $\tau(T) = T + \alpha 1$, $\omega(T) = T + \beta 1$ for some $\alpha, \beta \in k$. Moreover we must have $\beta = -\alpha$ if $c \neq 0$ because of the σ -centrality of c .

At the end, after suitable changes in the generators, we find that a Hopf-Galois algebra $A = A(R, \tau, \omega, g, h, c, \xi)$ as in Theorem 4.1 belongs to one of the following classes.

- (1) A is the enveloping Hopf algebra of the Lie algebra \mathfrak{sl}_2 :

$$A = U(\mathfrak{sl}_2) = k\langle T, X, Y \mid [X, T] = X, [Y, T] = -Y, [X, Y] = T \rangle.$$

- (2) A is the enveloping Hopf algebra of the 3-dimensional Heisenberg Lie algebra \mathfrak{h} :

$$A = U(\mathfrak{h}) = k\langle T, X, Y \mid [X, T] = 0 = [Y, T], [X, Y] = T \rangle.$$

- (3) $A = U_\beta$, $\beta \in k$, is the enveloping algebra of a solvable 3-dimensional Lie algebra:

$$A = U_\beta = k\langle T, X, Y \mid [X, T] = X, [Y, T] = \beta Y, [X, Y] = 0 \rangle.$$

- (4) $A = k[T, X, Y]$ is abelian.

- (5) A is a Hopf-Galois algebra and not a Hopf algebra, with

$$A = U_{\alpha, \lambda} = k\langle T, X, Y \mid [X, T] = \alpha X, [Y, T] = -\alpha Y, [X, Y] = 1 + \lambda T \rangle, \quad \alpha \in \{0, 1\}, \quad \lambda \in k.$$

Moreover, the Hopf algebra $\underline{H}(U_{\alpha, \lambda})$ is

- (a) $U(\mathfrak{sl}_2)$ if $\alpha = 1$ and $\lambda \neq 0$;
- (b) U_{-1} if $\alpha = 1$ and $\lambda = 0$;
- (c) $U(\mathfrak{h})$ if $\alpha = 0$ and $\lambda \neq 0$;
- (d) $k[T, X, Y]$ if $\alpha = 0$ and $\lambda = 0$.

6.4. The example $R = k[T, T^{-1}] = k\mathbb{Z}$. In this part, we set $R = \underline{R}(H)$, for $H = k\mathbb{Z} = k\langle K^{\pm 1} \rangle$. As above, we choose two (central) group-like elements $g = K^n$, $h = K^m$, $m, n \in \mathbb{Z}$. We fix two characters $\alpha, \beta: H \rightarrow k$, which are determined by nonzero scalars $\alpha := \alpha(K)$, $\beta := \beta(K)$, satisfying $\xi = \alpha^m = \beta^{-n}$ (as $\lambda = 1$). If $c \neq 0$, then there is $\gamma, \mu \in k$ such that $c = \gamma + \mu(1 - gh)$. As c is σ -central, it follows that in this case $\beta = \alpha^{-1}$ (hence $\alpha^n = \alpha^m = \xi$).

Thus, a Hopf-Galois algebra $A = A(R, \tau, \omega, g, h, c, \xi)$ as in Theorem 4.1, with $g = K^n$ and $h = K^m$, belongs to one of the following classes.

- (1) For each α, β with $\xi = \alpha^m = \beta^{-n}$:

$$A = k\langle K, X, Y \mid XK = \alpha KX, YK = \beta KY, XY - \xi YX = 0 \rangle.$$

- (2) For each α with $\xi = \alpha^m = \alpha^n$ and $\gamma, \mu \in k$:

$$A = k\langle K, X, Y \mid XK = \alpha KX, YK = \alpha^{-1}KY, XY - \xi YX = \gamma + \mu(1 - K^{n+m}) \rangle.$$

In the last class, we recognize the Hopf algebra $U_q(\mathfrak{sl}_2)$ when $n = m = -1$, $\gamma = 0$, $\mu = \frac{1}{q-q^{-1}}$ and $\xi = q^2 = \alpha^{-1}$. On the other hand, when $\gamma \neq 0$, we obtain the $U_q(\mathfrak{sl}_2)$ -cleft objects $A_{(a)}$ as computed in [11, Lemma 16], for $a = \gamma + \mu$ and $n = m = -1$, $\mu = \frac{1}{q-q^{-1}}$ as before.

6.5. The example $R = k\langle T_1^{\pm 1}, T_2^{\pm 1} \mid T_1 T_2 = q T_2 T_1 \rangle$. Here $q \in k^*$ and $R = \pi H$ is a twisted group algebra of the group algebra $H = k\mathbb{Z}^2 = k\langle K_1^{\pm 1}, K_2^{\pm 1} \mid K_1 K_2 = K_2 K_1 \rangle$ and π such that $\frac{\pi(K_1, K_2)}{\pi(K_2, K_1)} = q$. Subsection 6.2 applies to give the complete description of all the generalized ambiskew polynomial algebras that are Hopf-Galois as in Theorem 4.1. We focus on giving an example to illustrate Corollary 5.5, see §6.5.1 below. In particular, we assume that $q \neq 1$.

We choose $g = K_1, h = K_2 \in G(H)$, so $gh = qhg$ in R , that is, $\lambda = q$. Now, we get that

$$\tau(T_1) = \alpha(K_1)T_1, \quad \tau(T_2) = q\alpha(K_2)T_2, \quad \omega(T_1) = q^{-1}\beta(K_1)T_1, \quad \omega(T_2) = \beta(K_2)T_2,$$

for some algebra maps $\alpha, \beta: H \rightarrow k$ such that $\alpha(K_2) = \xi q^{-1} = \beta(K_1)^{-1}$.

In particular, $\sigma(T_1) = \xi^{-1}\alpha(K_1)T_1$ and $\sigma(T_2) = \xi\beta(K_2)T_2$.

Now $c = \gamma + \mu(1 - T_1 T_2)$. If $c \neq 0$, then being σ -central amounts to

- (1) If $\mu = 0$, then $\xi = \alpha(K_1) = \beta(K_2)^{-1}$. Thus, $\omega = \tau^{-1}$ and hence $\sigma = \text{id}$.
- (2) If $\mu \neq 0$ and $\gamma = -\mu$, then $\alpha(K_1) = q^{-1}\xi$ and $\beta(K_2) = q\xi^{-1}$. Hence $\sigma \neq \text{id}$.
- (3) Otherwise, $q = 1$ and this is a contradiction.

Next we analyze cases (1) and (2).

6.5.1. *Case (1)*. Fix $\mu = 0$, $\gamma = 1$, so $\tau\omega = \text{id}$ and $c = 1$. This will provide an example for Corollary 5.5. We have

$$\tau(T_1) = \xi T_1, \quad \tau(T_2) = \xi T_2, \quad \omega(T_1) = \xi^{-1}T_1, \quad \omega(T_2) = \xi^{-1}T_2.$$

Let $A_R = A(R, X, Y, \tau, c, \xi)$ be the corresponding Hopf-Galois algebra. This is generated by $T_1^{\pm 1}, T_2^{\pm 1}, X$ and Y with relations

$$T_1 T_2 = q T_2 T_1, \quad X T_1 = \xi T_1 X, \quad X T_2 = \xi T_2 X, \quad Y T_1 = \xi^{-1} T_1 Y, \quad Y T_2 = \xi^{-1} T_2 Y, \quad X Y - \xi Y X = 1.$$

Now, by Theorem 5.3, the Hopf algebra $\underline{H}(A_R)$ is a generalized ambiskew polynomial algebra $A(\underline{H}(R), \tau', \omega', c', \xi')$.

In this case, $\underline{H}(R) \simeq H$ via the assignment $T_1^{-1} \otimes T_1 \rightsquigarrow K_1, T_2^{-1} \otimes T_2 \rightsquigarrow K_2$. Recall that

$$\xi' = \xi \lambda^{-1} = q^{-1} \xi, \quad c' = (gh)^{-1} \otimes c - (gh)^{-1} c \otimes 1 = 0.$$

Finally, $\tau', \omega' \in \text{Aut } \underline{H}(R)$ are the automorphisms given by

$$\begin{aligned} \tau'(T_1^{-1} \otimes T_1) &= T_1 \cdot T_1^{-1} \otimes \tau(T_1) = \xi T_1^{-1} \otimes T_1, \\ \omega'(T_1^{-1} \otimes T_1) &= T_2 \cdot T_1^{-1} \otimes \tau^{-1}(T_1) = q \xi^{-1} T_1^{-1} \otimes T_1, \\ \tau'(T_2^{-1} \otimes T_2) &= T_1 \cdot T_2^{-1} \otimes \tau(T_2) = \xi q^{-1} T_2^{-1} \otimes T_2, \\ \omega'(T_2^{-1} \otimes T_2) &= T_2 \cdot T_2^{-1} \otimes \tau^{-1}(T_2) = \xi^{-1} T_2^{-1} \otimes T_2. \end{aligned}$$

In other words, under the identification above, we have that

$$\tau'(K_1) = \xi K_1, \quad \tau'(K_2) = \xi q^{-1} K_1, \quad \omega'(K_1) = \xi^{-1} q K_1, \quad \omega'(K_2) = \xi^{-1} K_2.$$

In particular, we see that τ' and ω' are not inverse to each other and the Hopf algebra $\underline{H}(A_R)$ is generated by $K_1^{\pm 1}, K_2^{\pm 1}, E$ and F with relations

$$\begin{aligned} K_1 K_2 &= K_2 K_1, \quad E K_1 = \xi K_1 E, \quad E K_2 = \xi q^{-1} K_2 E, \quad F K_1 = \xi^{-1} q K_1 F, \quad F K_2 = \xi^{-1} K_2 F, \\ E F - q^{-1} \xi F E &= 0. \end{aligned}$$

6.5.2. *Case (2)*. We have $c = \gamma T_1 T_2$ and

$$\tau(T_1) = q^{-1} \xi T_1, \quad \tau(T_2) = \xi T_2, \quad \omega(T_1) = \xi^{-1} T_1, \quad \omega(T_2) = q \xi^{-1} T_2.$$

Hence $A_R = A(R, X, Y, \tau, c, \xi)$ is generated by $T_1^{\pm 1}, T_2^{\pm 1}, X$ and Y with relations

$$\begin{aligned} T_1 T_2 &= q T_2 T_1, \quad X T_1 = q^{-1} \xi T_1 X, \quad X T_2 = \xi T_2 X, \quad Y T_1 = \xi^{-1} T_1 Y, \quad Y T_2 = q \xi^{-1} T_2 Y, \\ X Y - \xi Y X &= \gamma T_1 T_2. \end{aligned}$$

For the Hopf algebra $\underline{H}(A_R)$, we get $c' = \gamma(1 - K_1 K_2)$ and

$$\tau'(K_1) = q^{-1} \xi K_1, \quad \tau'(K_2) = \xi q^{-1} K_1, \quad \omega'(K_1) = \xi^{-1} q K_1, \quad \omega'(K_2) = q \xi^{-1} K_2.$$

In this case, $\omega' \tau' = \text{id}$. The Hopf algebra is now generated by $K_1^{\pm 1}, K_2^{\pm 1}, E$ and F with relations

$$K_1 K_2 = K_2 K_1, \quad E K_1 = q^{-1} \xi K_1 E, \quad E K_2 = \xi q^{-1} K_2 E, \quad F K_1 = \xi^{-1} q K_1 F, \quad F K_2 = q \xi^{-1} K_2 F,$$

$$E F - q^{-1} \xi F E = \gamma(1 - K_1 K_2).$$

6.6. The example $R = A_1(k)$. Here $R = k\langle U, V \mid UV - VU = 1 \rangle$ is a cleft object for the polynomial Hopf algebra $H = k[u, v]$, with $G(H) = \{1\}$ and $\mathcal{P}_{1,1}(H) = ku + kv$. As above, we obtain that the automorphisms τ, ω are determined by algebra maps $\alpha, \beta: H \rightarrow k$, as $\tau(U) = U + \alpha(u)$, $\omega(U) = U + \beta(u)$, similarly for $\tau(V), \omega(V)$. We have that $c = \gamma + c_1U + c_2V$, for some $\gamma, c_1, c_2 \in k$. The fact that c is $\sigma = \tau\omega$ -central determines that $c_1 = \alpha(v) + \beta(v)$ and $c_2 = -\alpha(u) - \beta(u)$. Therefore $A_R = A(R, \tau, \omega, g, h, c, \xi)$ with $g = h = 1$ is determined by scalars $\gamma, \alpha_u, \alpha_v, \beta_u, \beta_v \in k$ and relations

$$\begin{aligned} [U, V] &= 1, & [X, U] &= \alpha_u X, & [X, V] &= \alpha_v X, & [Y, U] &= \beta_u Y, & [Y, V] &= \beta_v Y \\ [X, Y] &= \gamma + (\alpha_v + \beta_v)U - (\alpha_u + \beta_u)V. \end{aligned}$$

If $c = \gamma + (\alpha_v + \beta_v)U - (\alpha_u + \beta_u)V = 0$, then this is a Hopf algebra. When $c \neq 0$, if we identify $\underline{H}(R)$ with $H = k[u, v]$ via $1 \otimes U - U \otimes 1 \rightsquigarrow u$, $1 \otimes V - V \otimes 1 \rightsquigarrow v$ then for the corresponding Hopf algebra $\underline{H}(A_R)$ it follows that $\xi' = \xi$, $\tau' = \tau$ and $\omega' = \omega$, in such a way that $\underline{H}(A_R) = A(H, \tau, \omega, c', \xi)$ for $c' = (\alpha_v + \beta_v)u - (\alpha_u + \beta_u)v$.

REFERENCES

- [1] N. Andruskiewitsch, I. Angiono, A. García Iglesias, A. Masuoka, C. Vay, Lifting via cocycle deformation, *J. Pure Appl. Alg.* **218** (4) (2014), 684-703.
- [2] I. Angiono, A. García Iglesias, Liftings of Nichols algebras of diagonal type II. All liftings are cocycle deformations, *Selecta Math.*, to appear (2019).
- [3] J. Bichon, The representation category of the quantum group of a non-degenerate bilinear form, *Comm. Algebra* **31** (2003), no. 10, 4831-4851.
- [4] K.A. Brown, M. Macauley, Ambiskew Hopf algebras, *J. Pure Appl. Algebra* **217** (2013), no. 1, 59-74.
- [5] Y. Doi, Braided bialgebras and quadratic bialgebras, *Comm. Algebra* **21** (1993), no. 5, 1731-1749.
- [6] J.T. Hartwig, Hopf structures on ambiskew polynomial rings, *J. Pure Appl. Algebra* **212** (2008), no. 4, 863-883.
- [7] D.A. Jordan, A simple localization of the quantized Weyl algebra, *J. Algebra* **174** (1995), no. 1, 267-281.
- [8] ———, Down-up algebras and ambiskew polynomial rings *J. Algebra* **228** (2000), no. 1, 311-346.
- [9] D.A. Jordan, I.E. Wells, Simple ambiskew polynomial rings, *J. Algebra* **382** (2013), 46-70.
- [10] C. Grunspan, Quantum torsors, *J. Pure Appl. Algebra* **184** (2003), no. 2-3, 229-255.
- [11] R. Günther, Crossed products for pointed Hopf algebras, *Comm. Algebra*, **27** (1999), 4389-4410.
- [12] H.F. Kreimer, M. Takeuchi, Hopf algebras and Galois extensions of an algebra, *Indiana Univ. Math. J.* **30** (1981), no. 5, 675-692.
- [13] S. Montgomery, Hopf algebras and their actions on rings, American Mathematical Society, 1993.
- [14] P. Schauenburg, Hopf bigalois extensions, *Comm. Algebra* **24**, No. 12 (1996), 3797-3825.
- [15] ———, Quantum torsors with fewer axioms, [arXiv:math/0302003](https://arxiv.org/abs/math/0302003) (2003).
- [16] K.H. Ulbrich, Fibre functors of finite-dimensional comodules, *Manuscripta Math.* **65** (1989), no. 1, 39-46.

JULIEN BICHON: LABORATOIRE DE MATHÉMATIQUES BLAISE PASCAL, UNIVERSITÉ CLERMONT AUVERGNE & CNRS, COMPLEXE UNIVERSITAIRE DES CÉZEAUX, 63178 AUBIÈRE CEDEX, FRANCE
E-mail address: julien.bichon@uca.fr

AGUSTÍN GARCÍA IGLESIAS: FACULTAD DE MATEMÁTICA, ASTRONOMÍA, FÍSICA Y COMPUTACIÓN (FAMAF), MEDINA ALLENDE S/N, UNIVERSIDAD NACIONAL DE CÓRDOBA, CIUDAD UNIVERSITARIA, CÓRDOBA (X5000HUA), ARGENTINA.
E-mail address: aigarcia@famaf.unc.edu.ar