



**HAL**  
open science

# Density properties for fractional Sobolev spaces with variable exponents

Azeddine Baalal, Mohamed Berghout

► **To cite this version:**

Azeddine Baalal, Mohamed Berghout. Density properties for fractional Sobolev spaces with variable exponents. *Annals of Functional Analysis*, 2019, 10 (3), pp.308-324. 10.1215/20088752-2018-0031 . hal-02338668

**HAL Id: hal-02338668**

**<https://hal.science/hal-02338668>**

Submitted on 30 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Ann. Funct. Anal.  $\diamond$

<https://doi.org/10.1215/20088752-2018-0031>

ISSN: 2008-8752 (electronic)

<http://projecteuclid.org/afa>

## DENSITY PROPERTIES FOR FRACTIONAL SOBOLEV SPACES WITH VARIABLE EXPONENTS

AZEDDINE BAALAL<sup>1</sup> and MOHAMED BERGHOUT<sup>2\*</sup>

Communicated by S. Barza

ABSTRACT. In this article we show some density properties of smooth and compactly supported functions in fractional Sobolev spaces with variable exponents. The additional difficulty in this nonlocal setting is caused by the fact that the variable exponent Lebesgue spaces are not translation-invariant.

### 1. Introduction

Function spaces with variable exponents have been the subject of intensive investigation in recent years. Examples of such spaces include Lebesgue and Sobolev spaces with variable exponents. Introduced by Orlicz [20] in 1931, their properties were further developed by Nakano [19] as special cases of the theory of modular spaces. In the ensuing decades they were primarily considered as important examples of modular spaces or the class of Musielak–Orlicz spaces. Initially of theoretical interest, by the end of the twentieth century these function spaces moved beyond theory and into the area of variational problems and studies of the  $p(x)$ -Laplacian operator, which in turn further fueled the development of this theory. Also stimulating the wide-ranging investigation of these spaces were their application to various problems in applied mathematics, for example, in the areas of nonlinear elasticity theory, fluid mechanics, and mathematical modeling of physical phenomena. (For more details on these spaces, see the monographs [6], [17], [18]; see also [22].)

---

Copyright 2019 by the Tusi Mathematical Research Group.

Received Aug. 12, 2018; Accepted Nov. 5, 2018.

$\diamond$ Advance publication—final volume, issue, and page numbers to be assigned.

\*Corresponding author.

2010 *Mathematics Subject Classification*. Primary 46E35; Secondary 35A15.

*Keywords*. fractional Sobolev spaces with variable exponents, density properties.

Let  $\Omega$  be a bounded domain of  $\mathbb{R}^n$ . We fix  $s \in (0, 1)$ , and we consider two variable exponents—that is,  $q : \Omega \rightarrow [1, +\infty)$  and  $p : \Omega \times \Omega \rightarrow [1, +\infty)$ —to be two continuous functions. In the following, we use the symbol  $:=$  to make the left-hand side equal by definition to the right-hand side.

The variable exponent Lebesgue space  $L^{q(\cdot)}(\Omega)$  is defined by

$$L^{q(\cdot)}(\Omega) := \left\{ u : \Omega \rightarrow \mathbb{R} \text{ measurable, } \int_{\Omega} |u(x)|^{q(x)} dx < \infty \right\}.$$

We define a norm, the so-called *Luxembourg norm*, in this space by

$$\|u\|_{L^{q(\cdot)}(\Omega)} := \inf \left\{ \lambda > 0 : \int_{\Omega} \left| \frac{u(x)}{\lambda} \right|^{q(x)} dx \leq 1 \right\}.$$

We define the fractional Sobolev space with variable exponents via the Gagliardo approach as follows:

$$W^{s,q(\cdot),p(\cdot,\cdot)}(\Omega) := \left\{ u \in L^{q(\cdot)}(\Omega), \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)}}{\lambda^{p(x,y)} |x - y|^{n+sp(x,y)}} dx dy < \infty, \text{ for some } \lambda > 0 \right\}.$$

Let

$$[u]^{s,p(\cdot,\cdot)}(\Omega) := \inf \left\{ \lambda > 0 : \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)}}{\lambda^{p(x,y)} |x - y|^{n+sp(x,y)}} dx dy \leq 1 \right\}$$

be the corresponding variable exponent Gagliardo seminorm. It is easy to see that  $W^{s,q(\cdot),p(\cdot,\cdot)}(\Omega)$  is a Banach space with the norm

$$\|u\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\Omega)} := \|u\|_{L^{q(\cdot)}(\Omega)} + [u]^{s,p(\cdot,\cdot)}(\Omega).$$

The definition of the spaces  $L^{q(\cdot)}(\mathbb{R}^n)$  and  $W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$  is analogous to that of  $L^{q(\cdot)}(\Omega)$  and  $W^{s,q(\cdot),p(\cdot,\cdot)}(\Omega)$ ; one just changes every occurrence of  $\Omega$  by  $\mathbb{R}^n$  (for more detail, see [1], [2], [4], and [16]).

Fractional Sobolev spaces with variable exponents have major applications in variational problems related to a well-known fractional version of the  $p(x)$ -Laplace operator, given by  $\operatorname{div}(|\nabla u|^{p(x)-2} \nabla u)$ , that is associated with the variable exponent Sobolev space  $W^{1,p(\cdot)}(\Omega)$ . We refer, for instance, to [6] and [13].

The denseness of regular functions such as  $C^\infty(\Omega)$ , functions which are continuously differentiable arbitrarily many times (smooth functions), or  $C_0^\infty(\Omega)$ , the subset of  $C^\infty(\Omega)$  of functions which have compact support, was one of the central tools in the function spaces setting—it might have some additional ramifications in the variable exponent Sobolev spaces. Moreover, the density of regular functions was one of the questions considered early (from 1986) in the context of minimizers of variational integrals. It is well known that the regularity of the Dirichlet energy integral minimizer is related to the density of smooth functions in the corresponding function space (see [14], [23]–[26]).

The denseness of smooth functions in the variable exponent Sobolev spaces  $W^{k,p(\cdot)}(\Omega)$  has proved to be a more difficult problem. Nowadays, denseness is a topic of intense focus and study by many researchers (we refer the reader to [6], [8]–[10], [15], [21], [27] and the references therein).

We recall that the continuity of the shift operator  $T_h : L^{q(\cdot)} \rightarrow L^{q(\cdot)}$ ,  $T_h f(x) := f(x+h)$  may fail (see [5]). In connection with the density problem for smooth function, we introduce the most important condition on the exponent in the study of variable exponent spaces, the well-known log-Hölder continuity condition: a function  $q : \Omega \rightarrow \mathbb{R}$  is called *log-Hölder continuous* on  $\Omega$  if there exists  $C > 0$  such that

$$|q(x) - q(y)| \leq \frac{C}{-\log|x-y|}, \quad |x-y| \leq \frac{1}{2}. \quad (1.1)$$

We say that a function  $p : \Omega \times \Omega \rightarrow \mathbb{R}$  satisfies condition (B-B) on  $\Omega \times \Omega$  if there exists  $C > 0$  such that

$$|p(x, y) - p(x', y')| \leq \frac{C}{-\log(|x-x'| + |y-y'|)}, \quad (1.2)$$

for all  $(x, y)$  and  $(x', y') \in \Omega \times \Omega$  such that  $|x-x'| + |y-y'| \leq \frac{1}{2}$ .

Notice that conditions (1.1) and (1.2) ensure that the typical mollifier function approximate function in Lebesgue Spaces with variable exponents (we refer the reader to [5], [21], [2] and [7]).

We define the following class of variable exponents

$$\mathcal{P}^{\log}(\Omega) := \{q : \Omega \rightarrow \mathbb{R} : q \text{ is measurable and log-Hölder continuous}\}$$

and

$$\begin{aligned} &\mathcal{P}^{\log}(\Omega \times \Omega) \\ &:= \{p : \Omega \times \Omega \rightarrow \mathbb{R} : p \text{ is measurable and satisfies condition (B-B)}\}. \end{aligned}$$

We set  $p^- := \operatorname{ess\,inf}_{(x,y) \in \Omega \times \Omega} p(x, y)$ ,  $p^+ := \operatorname{ess\,sup}_{(x,y) \in \Omega \times \Omega} p(x, y)$ ,  $q^- := \operatorname{ess\,inf}_{x \in \Omega} q(x)$ , and  $q^+ := \operatorname{ess\,sup}_{x \in \Omega} q(x)$ .

We assume that

$$1 < p^- \leq p(x, y) \leq p^+ < \infty, \quad (1.3)$$

$$1 < q^- \leq q(x) \leq q^+ < \infty, \quad (1.4)$$

$$p((x, y) - (z, z)) = p(x, y), \quad \forall (x, y), (z, z) \in \Omega \times \Omega. \quad (1.5)$$

The aim of this paper is to prove a number of density properties of smooth and compactly supported functions in fractional Sobolev spaces with variable exponent. The first result is an approximation with a continuous and compactly supported function.

**Theorem 1.1.** *Let  $u \in W^{s, q(\cdot), p(\cdot, \cdot)}(\mathbb{R}^n)$ . Then for any fixed  $\delta > 0$ , there exists a continuous and compactly supported function  $u_\delta$  such that*

$$\|u - u_\delta\|_{W^{s, q(\cdot), p(\cdot, \cdot)}(\mathbb{R}^n)} \longrightarrow 0 \quad \text{as } \delta \longrightarrow 0.$$

The second result is an approximation with smooth and compactly supported functions. More precisely, we have the following.

**Theorem 1.2.** *Let  $q \in \mathcal{P}^{\log}(\Omega)$  and  $p \in \mathcal{P}^{\log}(\Omega \times \Omega)$ . Then the space  $C_0^\infty(\mathbb{R}^n)$  is dense in  $W^{s, q(\cdot), p(\cdot, \cdot)}(\mathbb{R}^n)$ .*

We consider  $\Omega \subset \mathbb{R}^n$  to be a  $W^{s,q(\cdot),p(\cdot,\cdot)}$ -extension domain if there exists a continuous linear extension operator

$$\mathcal{E} : W^{s,q(\cdot),p(\cdot,\cdot)}(\Omega) \longrightarrow W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$$

such that  $\mathcal{E}u|_{\Omega} = u$  for each  $u \in W^{s,q(\cdot),p(\cdot,\cdot)}(\Omega)$ . From Theorem 1.2, we derive the density of  $C^\infty(\overline{\Omega})$  also in  $W^{s,q(\cdot),p(\cdot,\cdot)}$ -extension domains, which include in particular domains with Lipschitz boundary (see [1]).

**Theorem 1.3.** *Let  $q \in \mathcal{P}^{\log}(\Omega)$ ,  $p \in \mathcal{P}^{\log}(\Omega \times \Omega)$  and suppose that  $\Omega$  is a  $W^{s,q(\cdot),p(\cdot,\cdot)}$ -extension domain. Then the space  $C^\infty(\overline{\Omega})$  is dense in  $W^{s,q(\cdot),p(\cdot,\cdot)}(\Omega)$ .*

This paper is organized as follows. In Section 2, we describe some properties of fractional Sobolev spaces with variable exponents that will be useful in our exposition. In Section 3, we prove the density results for fractional Sobolev spaces with variable exponents.

## 2. Notation and preliminary results

In this section we give some notation and prove several useful properties of fractional Sobolev spaces with variable exponents.

We use the following notation throughout this article.  $\mathbb{R}^n$  is the  $n$ -dimensional Euclidean space, and  $n \in \mathbb{N}$  always stands for the dimension of the space. A domain  $\Omega \subset \mathbb{R}^n$  is a connected open set equipped with the  $n$ -dimensional Lebesgue measure. For constants, we use the letter  $C$  whose value may change even within a string of estimates. A ball with radius  $R$  and center 0 will be denoted by  $B_R$ . The closure of a set  $A$  is denoted by  $\overline{A}$ . We use the usual convention of identifying two  $\mu$ -measurable functions on  $A$  (almost everywhere (a.e.) in  $A$ , for short) if they agree almost everywhere—that is, if they agree up to a set of  $\mu$ -measure zero. The Lebesgue integral of a Lebesgue measurable function  $f : \Omega \longrightarrow \mathbb{R}$  is defined in the standard way and denoted by  $\int_{\Omega} f(x) dx$ . By  $\omega_{n-1}$ , we denote the  $(n-1)$ -dimensional measure of the unit sphere  $S^{n-1}$ . By  $\text{supp } f$ , we denote the support of  $f$ , which is the complement of the biggest open set on which it vanishes; in other words,  $\text{supp } f$  is the closure of the set  $\{x; f(x) \neq 0\}$ . We denote by  $C(\overline{\Omega})$  the space of uniformly continuous functions equipped with the supremum norm  $\|f\|_{\infty} = \sup_{x \in \overline{\Omega}} |f(x)|$ . By  $C^k(\overline{\Omega})$ ,  $k \in \mathbb{N}$ , we denote the space of all functions  $f$  such that  $\partial_{\alpha} f := \frac{\partial^{|\alpha|} f}{\partial^{\alpha_1} x_1 \cdots \partial^{\alpha_n} x_n} \in C(\overline{\Omega})$  for all multi-indices  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$ ,  $|\alpha| := \alpha_1 + \alpha_2 + \cdots + \alpha_n \leq k$ . The space is equipped with the norm  $\sup_{|\alpha| \leq k} \|\partial_{\alpha} f\|_{\infty}$ ,  $C^\infty(\overline{\Omega}) = \bigcap_k C^k(\overline{\Omega})$ . The set of smooth functions in  $\Omega$  is denoted by  $C^\infty(\Omega)$ —it consists of functions in  $\Omega$  which are continuously differentiable arbitrarily many times. The set  $C_0^\infty(\Omega)$  is the subset of  $C^\infty(\Omega)$  of functions which have compact support.

The important role of manipulating Lebesgue–Sobolev spaces with variable exponents is played by the modular of the  $L^{q(\cdot)}(\Omega)$  space, which is the mapping  $\rho : L^{q(\cdot)}(\Omega) \longrightarrow \mathbb{R}^+$  defined by

$$\rho(u) = \int_{\Omega} |u(x)|^{q(x)} dx.$$

From [10, Theorems 1.4, 1.8], we obtain the following proposition.

**Proposition 2.1.** *Let  $\Omega$  be an open subset of  $\mathbb{R}^n$ , and let  $u_n, u \in L^{q(\cdot)}(\Omega)$  be such that  $n \in \mathbb{N}$ . Then*

- (1)  $C_0^\infty(\Omega)$  is dense in the space  $(L^{q(\cdot)}(\Omega), \|\cdot\|_{L^{q(\cdot)}(\Omega)})$ ,
- (2)  $\lim_{n \rightarrow +\infty} \rho(u_n - u) \rightarrow 0 \iff \lim_{n \rightarrow +\infty} \|u_n - u\|_{L^{q(\cdot)}(\Omega)} = 0$ .

Next, we prove several properties of  $W^{s, q(\cdot), p(\cdot, \cdot)}(\mathbb{R}^n)$  that are needed to obtain our main results.

**Lemma 2.2.** *Let  $\varphi \in C_0^\infty(\mathbb{R}^n)$ . Then there exists a positive constant  $C$  such that*

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|\varphi(x) - \varphi(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy \leq C.$$

*Proof.* Let  $\varphi \in C_0^\infty(\mathbb{R}^n)$  with  $\text{supp } \varphi \subseteq B_R$ . Therefore, if  $x, y \in \mathbb{R}^n \setminus B_R$ , then  $\varphi(x) = \varphi(y) = 0$ . Thus

$$\begin{aligned} I &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|\varphi(x) - \varphi(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy \\ &= \int_{B_R} \int_{\mathbb{R}^n} \frac{|\varphi(x) - \varphi(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy \\ &= I_1 + I_2, \end{aligned}$$

where

$$I_1 := \int_{B_R} \int_{B_{2R}} \frac{|\varphi(x) - \varphi(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy$$

and

$$I_2 := \int_{B_R} \int_{\mathbb{R}^n \setminus B_{2R}} \frac{|\varphi(x) - \varphi(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy.$$

We first estimate  $I_1$ . We have

$$\begin{aligned} I_1 &= \int_{B_R} \int_{B_{2R}} \frac{|\varphi(x) - \varphi(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy \\ &= \int_{B_R} \int_{B_{2R}} \frac{|\varphi(x) - \varphi(y)|^{p(x,y)}}{|x - y|^{p(x,y)}} \times \frac{|x - y|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy. \end{aligned}$$

Hence

$$I_1 \leq C \int_{B_{2R}} \int_{B_{2R}} |x - y|^{-n+(1-s)p(x,y)} dx dy,$$

where  $C$  depends on the  $C^1$ -norm of  $\varphi$ ,  $p^+$ , and  $p^-$ .

We set

$$d := \sup\{|x - y| : (x, y) \in B_{2R} \times B_{2R}\}.$$

Observe that

$$|x - y|^{-n+(1-s)p(x,y)} \leq \max\{^{-n+(1-s)p^+}, d^{-n+(1-s)p^-}\}.$$

Therefore,

$$I_1 \leq C \int_{B_{2R}} \int_{B_{2R}} dx dy \leq C.$$

Now we estimate  $I_2$ . For this, we observe that if  $x \in B_R$  and  $y \in \mathbb{R}^n \setminus B_{2R}$ , then

$$|x - y| \geq |y| - |x| \geq \frac{|y|}{2}.$$

Thus

$$I_2 \leq C(n, s, p^+, p^-, \|\varphi\|_{L^\infty(\mathbb{R}^n)}) \int_{B_R} \left( \int_{\mathbb{R}^n \setminus B_{2R}} \frac{dy}{|y|^{n+sp(x,y)}} \right) dx.$$

Now, if  $|y| > 1$ , then

$$\int_{\mathbb{R}^n \setminus B_{2R}} \frac{dy}{|y|^{n+sp(x,y)}} \leq \int_{\mathbb{R}^n \setminus B_{2R}} \frac{dy}{|y|^{n+sp^-}} = \omega_{n-1} C(R, s, p^-) \leq C,$$

where  $\omega_{n-1}$  denotes the  $(n-1)$ -dimensional measure of the unit sphere  $S^{n-1}$ .

Now, we assume that  $|y| < 1$ . We claim that

$$\int_{\mathbb{R}^n \setminus B_{2R}} \frac{dy}{|y|^{n+sp(x,y)}} \leq \int_{\mathbb{R}^n \setminus B_{2R}} \frac{dy}{|y|^{n+sp^+}} = \omega_{n-1} C(R, s, p^+) \leq C.$$

Hence,  $I$  is bounded.  $\square$

As an obvious consequence of Lemma 2.2 and Proposition 2.1, we have  $C_0^\infty(\mathbb{R}^n) \subseteq W^{s,q(\cdot);p(\cdot)}(\mathbb{R}^n)$ . Now we give two approximation results.

**Lemma 2.3.** *Let  $u \in L^{q(\cdot)}(\mathbb{R}^n)$ . Then there exists a sequence of functions  $u_m \in L^{q(\cdot)}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  such that*

$$\|u - u_m\|_{L^{q(\cdot)}(\mathbb{R}^n)} \longrightarrow 0 \quad \text{as } m \longrightarrow +\infty.$$

*Proof.* We set

$$u_m(x) := \begin{cases} m & \text{if } u(x) \geq m, \\ u(x) & \text{if } u(x) \in (-m, m), \\ -m & \text{if } u(x) \leq -m. \end{cases}$$

We have

$$u_m \longrightarrow u \quad \text{a.e. in } \mathbb{R}^n$$

and

$$|u_m(x)|^{q(x)} \leq |u(x)|^{q(x)} \in L^1(\mathbb{R}^n).$$

By the dominated convergence theorem and Proposition 2.1, the claim follows.  $\square$

**Lemma 2.4.** *Let  $u \in L^{p(\cdot, \cdot)}(\mathbb{R}^n \times \mathbb{R}^n)$ . Then there exists a sequence of functions  $u_M \in L^{p(\cdot, \cdot)}(\mathbb{R}^n \times \mathbb{R}^n) \cap L^\infty(\mathbb{R}^n \times \mathbb{R}^n)$  such that*

$$\|u - u_M\|_{L^{p(\cdot, \cdot)}(\mathbb{R}^n \times \mathbb{R}^n)} \longrightarrow 0 \quad \text{as } M \longrightarrow +\infty.$$

*Proof.* We set

$$u_M(x, y) := \begin{cases} M & \text{if } u(x, y) \geq M, \\ u(x, y) & \text{if } u(x, y) \in (-M, M), \\ -M & \text{if } u(x, y) \leq -M. \end{cases}$$

We have

$$u_M \longrightarrow u \quad \text{a.e. in } \mathbb{R}^n \times \mathbb{R}^n$$

and

$$|u_M(x, y)|^{p(x, y)} \leq |u(x, y)|^{p(x, y)} \in L^1(\mathbb{R}^n \times \mathbb{R}^n).$$

The claim follows from the dominated convergence theorem and Proposition 2.1.  $\square$

The proofs of density properties of smooth and compactly supported functions are mainly based on a basic technique of convolution (which makes functions  $C^\infty$ ), joined with a cutoff (which makes their support compact). In the remainder of this article, we describe properties of these operations with respect to the norm in fractional Sobolev spaces with variable exponents.

Let  $\eta \in C_0^\infty(\mathbb{R}^n)$  be such that  $\eta \geq 0$  in  $\mathbb{R}^n$  and  $\text{supp } \eta \subseteq B_1$ . We also assume that

$$\int_{B_1} \eta(x) dx = 1.$$

Let  $\varepsilon > 0$ , and let  $\eta_\varepsilon$  be the mollifier defined as

$$\eta_\varepsilon(x) = \frac{1}{\varepsilon^n} \eta\left(\frac{x}{\varepsilon}\right), \quad x \in \mathbb{R}^n.$$

For any  $u \in W^{s, q(\cdot), p(\cdot, \cdot)}(\mathbb{R}^n)$ , we will denote by  $u_\varepsilon$  the function defined as the convolution between  $u$  and  $\eta_\varepsilon$ ; that is,

$$u_\varepsilon(x) = (u * \eta_\varepsilon)(x) = \int_{\mathbb{R}^n} u(x - z) \eta_\varepsilon(z) dz, \quad x \in \mathbb{R}^n.$$

Note that, by construction,  $u_\varepsilon \in C^\infty(\mathbb{R}^n)$ . For small  $\varepsilon$ , the convolution does not change the norm too much, according to the following result (see, e.g., [2, Lemma 3.2]), which will be useful in the rest of this article.

**Lemma 2.5.** *Let  $u \in W^{s, q(\cdot), p(\cdot, \cdot)}(\mathbb{R}^n)$ ,  $q \in \mathcal{P}^{\log}(\Omega)$  and  $p \in \mathcal{P}^{\log}(\Omega \times \Omega)$ . We assume that (1.3), (1.4) and (1.5) hold. Then  $\|u - u_\varepsilon\|_{W^{s, q(\cdot), p(\cdot, \cdot)}(\mathbb{R}^n)} \rightarrow 0$  as  $\varepsilon \rightarrow 0$ .*

*Proof.* Let  $u \in W^{s, q(\cdot), p(\cdot, \cdot)}(\mathbb{R}^n)$ . Since  $u \in L^{q(\cdot)}(\Omega)$  and  $q \in \mathcal{P}^{\log}(\Omega)$ , by [21, Corollary], we know that

$$\|u - u_\varepsilon\|_{L^{q(\cdot)}(\mathbb{R}^n)} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0.$$

Hence, from Proposition 2.1, it suffices to prove that



$$\int_{\Omega \times \Omega} |u_\varepsilon(x) - u(x) - u_\varepsilon(y) + u(y)|^{p(x,y)} K(x, y) dx dy \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0,$$

where  $K(x, y) = |x - y|^{-n-sp(x,y)}$ . By the same way in [2, Lemma 3.2], we can prove that

$$\begin{aligned} & \int_{\Omega \times \Omega} |u_\varepsilon(x) - u(x) - u_\varepsilon(y) + u(y)|^{p(x,y)} K(x, y) dx dy \\ & \leq |B_1|^{p^- + p^+ - 1} \int_{\Omega \times \Omega \times B_1} |u(x - \varepsilon z) - u(y - \varepsilon z) - u(x) + u(y)|^{p(x,y)} \\ & \quad \times K(x, y) (\eta(z)^{p^+} + \eta(z)^{p^-}) dx dy dz. \end{aligned}$$

Fix  $z \in B_1$  and put  $w = (z, z) \in \Omega \times \Omega$ . We define the function  $v : \Omega \times \Omega \rightarrow \mathbb{R}$  by

$$v(x, y) := (u(x) - u(y)) (K(x, y))^{\frac{1}{p(x,y)}}, \quad \forall (x, y) \in \Omega \times \Omega.$$

Then  $v \in L^{p(\cdot, \cdot)}(\Omega \times \Omega)$ . Let  $\varepsilon' > 0$ . Since  $p \in \mathcal{P}^{\log}(\Omega \times \Omega)$ , by [21, Corollary], there exists  $g \in C_0^\infty(\Omega \times \Omega)$  with  $\|v - g\|_{L^{p(\cdot, \cdot)}(\Omega \times \Omega)} \leq \frac{\varepsilon'}{3}$ , hence

$$\begin{aligned} & \|v(\cdot - \varepsilon w) - v\|_{L^{p(\cdot, \cdot)}(\Omega \times \Omega)} \\ & \leq \|v(\cdot - \varepsilon w) - g(\cdot - \varepsilon w)\|_{L^{p(\cdot, \cdot)}(\Omega \times \Omega)} \\ & \quad + \|g(\cdot - \varepsilon w) - g\|_{L^{p(\cdot, \cdot)}(\Omega \times \Omega)} + \|v - g\|_{L^{p(\cdot, \cdot)}(\Omega \times \Omega)} \\ & \leq \frac{\varepsilon'}{3} + \frac{\varepsilon'}{3} + \frac{\varepsilon'}{3} = \varepsilon', \end{aligned}$$

with  $\varepsilon$  is sufficiently small. Hence

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega \times \Omega} |u(x - \varepsilon z) - u(y - \varepsilon z) - u(x) + u(y)|^{p(x,y)} \times K(x, y) dx dy = 0.$$

Moreover, for a.e.  $z \in B_1$ , there exists a positive constant  $C$  such that

$$\begin{aligned} & (\eta(z)^{p^+} + \eta(z)^{p^-}) \int_{\Omega \times \Omega} |u(x - \varepsilon z) - u(y - \varepsilon z) - u(x) + u(y)|^{p(x,y)} \\ & \quad \times K(x, y) dx dy \\ & \leq 2C(\eta(z)^{p^+} + \eta(z)^{p^-}) \int_{\Omega \times \Omega} |u(x) - u(y)|^{p(x,y)} \times K(x, y) dx dy \in L^\infty(B_1), \end{aligned}$$

for any  $\varepsilon > 0$ . Hence, using the dominated convergence theorem, we deduce that

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \int_{B_1} \eta(z)^{p(x,y)} \int_{\Omega \times \Omega} |u(x - \varepsilon z) - u(y - \varepsilon z) - u(x) + u(y)|^{p(x,y)} \\ & \quad \times K(x, y) dx dy dz = 0. \end{aligned}$$

Consequently

$$\int_{\Omega \times \Omega} |u_\varepsilon(x) - u(x) - u_\varepsilon(y) + u(y)|^{p(x,y)} K(x, y) dx dy \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0,$$

which concludes the proof.  $\square$

Now, we will discuss the cutoff technique needed for the density argument. For any  $j \in \mathbb{N}$ , let  $\tau_j \in C^\infty(\mathbb{R}^n)$  be such that

$$\begin{aligned} 0 &\leq \tau_j(x) \leq 1, \quad \forall x \in \mathbb{R}^n, \\ \tau_j(x) &= \begin{cases} 1 & \text{if } x \in B_j, \\ 0 & \text{if } x \in \mathbb{R}^n \setminus B_{j+1}, \end{cases} \end{aligned} \quad (2.1)$$

where  $B_j$  denotes the ball centered at zero with radius  $j$ . We have the following result.

**Lemma 2.6.** *Let  $u \in W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$ . Then  $\tau_j u \in W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$ .*

*Proof.* Let  $u \in W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$ . It is clear that  $\tau_j u \in L^{q(\cdot)}(\mathbb{R}^n)$  since  $|\tau_j| \leq 1$ . Furthermore,

$$\begin{aligned} I &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|\tau_j(x)u(x) - \tau_j(y)u(y)|^{p(x,y)}}{\lambda^{p(x,y)}|x-y|^{n+sp(x,y)}} dx dy \\ &\leq 2^{p^+-1} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|\tau_j(x)(u(x) - u(y))|^{p(x,y)}}{\lambda^{p(x,y)}|x-y|^{n+sp(x,y)}} dx dy \\ &\quad + 2^{p^+-1} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(y)(\tau_j(x) - \tau_j(y))|^{p(x,y)}}{\lambda^{p(x,y)}|x-y|^{n+sp(x,y)}} dx dy \\ &\leq 2^{p^+-1} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^{p(x,y)}}{\lambda^{p(x,y)}|x-y|^{n+sp(x,y)}} dx dy \\ &\quad + 2^{p^+-1} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(y)(\tau_j(x) - \tau_j(y))|^{p(x,y)}}{\lambda^{p(x,y)}|x-y|^{n+sp(x,y)}} dx dy, \end{aligned}$$

where the integral

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^{p(x,y)}}{\lambda^{p(x,y)}|x-y|^{n+sp(x,y)}} dx dy$$

is finite since  $u \in W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$ .

According to Lemma 2.3, we can assume that  $u \in L^\infty(\mathbb{R}^n)$ . Hence,

$$|u(y)|^{p(x,y)} \leq \|u\|_{L^\infty(\mathbb{R}^n)}^{p^+} + \|u\|_{L^\infty(\mathbb{R}^n)}^{p^-}.$$

Therefore,

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(y)(\tau_j(x) - \tau_j(y))|^{p(x,y)}}{\lambda^{p(x,y)}|x-y|^{n+sp(x,y)}} dx dy \leq \frac{C}{\alpha} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|(\tau_j(x) - \tau_j(y))|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} dx dy,$$

where  $\alpha = \min\{\lambda^{p^+}, \lambda^{p^-}\}$  and the constant  $C$  depends on  $p^+$ ,  $p^-$ , and  $\|u\|_{L^\infty(\mathbb{R}^n)}$ . Finally, using Lemma 2.2, we get

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(y)(\tau_j(x) - \tau_j(y))|^{p(x,y)}}{\lambda^{p(x,y)}|x-y|^{n+sp(x,y)}} dx dy < \infty.$$

This concludes the proof.  $\square$

Now, we will prove that we can approximate a function in  $W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$  with another function with compact support, by keeping the error small. More precisely, we have the following result.

**Lemma 2.7.** *Let  $u \in W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$ . Then  $\text{supp}(\tau_j u) \subseteq \bar{B}_{j+1} \cap \text{supp } u$ , and*

$$\|\tau_j u - u\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)} \longrightarrow 0 \quad \text{as } j \longrightarrow +\infty.$$

*Proof.* By (2.1) and [11, Lemma 9], we get

$$\text{supp}(\tau_j u) \subseteq \bar{B}_{j+1} \cap \text{supp } u.$$

Now, let us show that

$$\|\tau_j u - u\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)} \longrightarrow 0 \quad \text{as } j \longrightarrow +\infty.$$

From Proposition 2.1, it suffices to prove that

$$\int_{\mathbb{R}^n} |\tau_j(x)u(x) - u(x)|^{q(x)} dx \longrightarrow 0 \quad \text{as } j \longrightarrow +\infty$$

and

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|\tau_j(x)u(x) - u(x) - \tau_j(y)u(y) + u(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy \longrightarrow 0 \quad \text{as } j \longrightarrow +\infty.$$

We observe that

$$\begin{aligned} |\tau_j(x)u(x) - u(x)|^{q(x)} &= |\tau_j(x) - 1|^{q(x)} |u(x)|^{q(x)} \\ &\leq 2^{q^+ - 1} (|\tau_j(x)|^{q(x)} + 1) |u(x)|^{q(x)} \\ &\leq 2^{q^+} |u(x)|^{q(x)} \in L^1(\mathbb{R}^n). \end{aligned}$$

Moreover, by (2.1)

$$|\tau_j(x)u(x) - u(x)|^{q(x)} \longrightarrow 0 \quad \text{as } j \longrightarrow +\infty \text{ a.e. in } \mathbb{R}^n.$$

Then, by the dominated convergence theorem, we get

$$\int_{\mathbb{R}^n} |\tau_j(x)u(x) - u(x)|^{q(x)} \longrightarrow 0 \quad \text{as } j \longrightarrow +\infty.$$

Now, let us show that

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|\tau_j(x)u(x) - u(x) - \tau_j(y)u(y) + u(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy \longrightarrow 0 \quad \text{as } j \longrightarrow +\infty.$$

We set  $\eta_j = 1 - \tau_j$ . Then  $\eta_j u = u - \tau_j u$ . Moreover,

$$\begin{aligned} &|\tau_j(x)u(x) - u(x) - \tau_j(y)u(y) + u(y)| \\ &= |\eta_j(x)(u(x) - u(y)) - (\tau_j(y) - \tau_j(x))u(y)|. \end{aligned}$$

Therefore,

$$\begin{aligned}
& \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|\tau_j(x)u(x) - u(x) - \tau_j(y)u(y) + u(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} dx dy \\
& \leq 2^{p^+-1} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|\tau_j(x) - \tau_j(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} |u(y)|^{p(x,y)} dx dy \\
& \quad + 2^{p^+-1} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} \eta_j^{p(x,y)}(x) dx dy.
\end{aligned}$$

Note that by Lemma 2.3, we can assume that  $u \in L^\infty(\mathbb{R}^n)$ . Hence,

$$\frac{|\tau_j(x) - \tau_j(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} |u(y)|^{p(x,y)} \leq C(\|u\|_{L^\infty(\mathbb{R}^n)}, p^+, p^-) \frac{|\tau_j(x) - \tau_j(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}}.$$

By Lemma 2.2, we deduce that

$$\frac{|\tau_j(x) - \tau_j(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} \in L^1(\mathbb{R}^n \times \mathbb{R}^n).$$

Furthermore,

$$\frac{|\tau_j(x) - \tau_j(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} |u(y)|^{p(x,y)} \longrightarrow 0 \quad \text{as } j \longrightarrow \infty \text{ a.e. in } \mathbb{R}^n \times \mathbb{R}^n.$$

Hence, by the dominated convergence theorem,

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|\tau_j(x) - \tau_j(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} |u(y)|^{p(x,y)} dx dy \longrightarrow 0 \quad \text{as } j \longrightarrow \infty.$$

Also,

$$\frac{|u(x) - u(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} (\eta_j(x))^{p(x,y)} \leq (\eta_j(x))^{p^+} + \eta_j(x)^{p^-} \frac{|u(x) - u(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}}$$

and

$$\frac{|u(x) - u(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} \in L^1(\mathbb{R}^n \times \mathbb{R}^n).$$

Again by (2.1),

$$\frac{|u(x) - u(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} (\eta_j(x))^{p(x,y)} \longrightarrow 0 \quad \text{as } j \longrightarrow \infty \text{ a.e. in } \mathbb{R}^n \times \mathbb{R}^n.$$

Hence, by the dominated convergence theorem,

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^{p(x,y)}}{|x-y|^{n+sp(x,y)}} (\eta_j(x))^{p(x,y)} dx dy \longrightarrow 0 \quad \text{as } j \longrightarrow \infty.$$

This concludes the proof.  $\square$

### 3. Approximation by smooth and compactly supported functions

In this section we give some density properties of smooth and compactly supported functions in fractional Sobolev spaces with variable exponents. The additional difficulty in this nonlocal setting is caused by the fact that the variable exponent Lebesgue spaces are not translation-invariant (see [6, Section 3.6]). To overcome this difficulty, we exploit the techniques developed in Section 2.

**Theorem 3.1.** *Let  $u \in W^{s,q(\cdot),p(\cdot)}(\mathbb{R}^n)$ . Then for any fixed  $\delta > 0$ , there exists a continuous and compactly supported function  $u_\delta$  such that*

$$\|u - u_\delta\|_{W^{s,q(\cdot),p(\cdot)}(\mathbb{R}^n)} \longrightarrow 0 \quad \text{as } \delta \longrightarrow 0.$$

*Proof.* Let  $u \in W^{s,q(\cdot),p(\cdot)}(\mathbb{R}^n)$ . According to Lemma 2.2, we can approximate  $u$  with a sequence of bounded functions. Consequently, we can also assume that  $u \in L^\infty(\mathbb{R}^n)$ .

Let  $\tau_j \in C^\infty(\mathbb{R}^n)$  be as in Section 2, with  $\tau_j(P) = 1$  if  $|P| \leq j$  and  $\tau_j(P) = 0$  if  $|P| \geq j + 1$ . Let  $u_j = \tau_j u$ . Then

$$u_j \longrightarrow u \quad \text{a.e. in } \mathbb{R}^n \text{ as } j \longrightarrow \infty,$$

and

$$|u(x) - u_j(x)|^{q(x)} \leq 2^{q^+} |u(x)|^{q(x)} \in L^1(\mathbb{R}^n).$$

As a consequence, by the dominated convergence theorem,

$$\int_{\mathbb{R}^n} |u(x) - u_j(x)|^{q(x)} dx \longrightarrow 0 \quad \text{as } j \longrightarrow \infty.$$

Hence, for any fixed  $\delta > 0$ , there exists  $j_\delta \in \mathbb{N}$  such that

$$\int_{\mathbb{R}^n} |u(x) - u_{j_\delta}(x)|^{q(x)} dx \leq \delta. \quad (3.1)$$

Note that  $u_{j_\delta}$  is supported in  $\overline{B}_{j_\delta+1}$  and that  $\mu(A) = \int_A dx$  is finite over compact sets. Hence, we can use Lusin's theorem (see [12, Theorem 7.10, p. 121] for the definition of the uniform norm). We obtain that there exist a closed set  $E_\delta \subset \mathbb{R}^n$  and a continuous and compactly supported function  $u_\delta : \mathbb{R}^n \longrightarrow \mathbb{R}$  such that

$$u_\delta = u_{j_\delta} \quad \text{in } \mathbb{R}^n \setminus E_\delta, \quad \mu(E_\delta) \leq \delta \quad \text{and} \quad \|u_{j_\delta}\|_{L^\infty(\mathbb{R}^n)} \leq \|u\|_{L^\infty(\mathbb{R}^n)}.$$

In particular, since  $0 \leq \tau_{j_\delta}(x) \leq 1$ , we have

$$\|u_\delta\|_{L^\infty(\mathbb{R}^n)} \leq \|u\|_{L^\infty(\mathbb{R}^n)} < \infty.$$

Therefore,

$$\begin{aligned} \int_{\mathbb{R}^n} |u_{j_\delta}(x) - u_\delta(x)|^{q(x)} dx &= \int_{E_\delta} |u_{j_\delta}(x) - u_\delta(x)|^{q(x)} dx \\ &\leq 2^{q^+ - 1} \left( \int_{E_\delta} |u_{j_\delta}(x)|^{q(x)} dx + \int_{E_\delta} |u_\delta(x)|^{q(x)} dx \right) \\ &\leq C(q^+, q^-, \|u\|_{L^\infty(\mathbb{R}^n)}) \mu(E_\delta). \end{aligned}$$

Hence

$$\int_{\mathbb{R}^n} |u_{j_\delta}(x) - u_\delta(x)|^{q(x)} dx \leq C\delta. \quad (3.2)$$

On the other hand,

$$\begin{aligned} & |u(x) - u_\delta(x)|^{q(x)} \\ & \leq 2^{q^+-1} (|u(x) - u_{j_\delta}(x)|^{q(x)} + |u_{j_\delta}(x) - u_\delta(x)|^{q(x)}). \end{aligned}$$

Hence

$$\begin{aligned} \int_{\mathbb{R}^n} |u(x) - u_\delta(x)|^{q(x)} dx & \leq 2^{q^+-1} \int_{\mathbb{R}^n} |u(x) - u_{j_\delta}(x)|^{q(x)} dx \\ & \quad + 2^{q^+-1} \int_{\mathbb{R}^n} |u_{j_\delta}(x) - u_\delta(x)|^{q(x)} dx. \end{aligned}$$

Now, from (3.1) and (3.2), we deduce that

$$\int_{\mathbb{R}^n} |u(x) - u_\delta(x)|^{q(x)} dx \longrightarrow 0 \quad \text{as } \delta \longrightarrow 0.$$

Therefore, by Proposition 2.1, we obtain

$$\|u - u_\delta\|_{L^{q(\cdot)}(\mathbb{R}^n)} \longrightarrow 0 \quad \text{as } \delta \longrightarrow 0.$$

Moreover,

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|(u - u_j)(x) - (u - u_j)(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy \longrightarrow 0 \quad \text{as } j \longrightarrow \infty.$$

Hence, for any fixed  $\delta > 0$ , there exists  $j_\delta \in \mathbb{N}$  such that

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|(u - u_{j_\delta})(x) - (u - u_{j_\delta})(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy \leq \delta.$$

Notice that

$$v_{j_\delta}(x, y) = \frac{|u_{j_\delta}(x) - u_{j_\delta}(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}}$$

is supported in  $\{P \in \mathbb{R}^n \times \mathbb{R}^n; |P| \leq j_\delta + 1\}$  and  $\mu(A) = \int_A \int_A dx dy$  is finite over compact sets. Hence, we can again use Lusin's theorem. We get that there exist a closed set  $E_\delta \subset \mathbb{R}^n \times \mathbb{R}^n$  and a continuous and compactly supported function  $u_\delta : \mathbb{R}^n \times \mathbb{R}^n \longrightarrow \mathbb{R}$  such that

$$u_\delta = v_{j_\delta} \quad \text{in } \mathbb{R}^n \times \mathbb{R}^n \setminus E_\delta, \quad \mu(E_\delta) \leq \delta \quad \text{and}$$

$$\|u_{j_\delta}\|_{L^\infty(\mathbb{R}^n \times \mathbb{R}^n)} \leq \|v_{j_\delta}\|_{L^\infty(\mathbb{R}^n \times \mathbb{R}^n)}.$$

In particular, since  $0 \leq \tau_{j_\delta}(x) \leq 1$ , we have

$$\|u_\delta\|_{L^\infty(\mathbb{R}^n \times \mathbb{R}^n)} \leq \left\| \frac{|u(x) - u(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} \right\|_{L^\infty(\mathbb{R}^n \times \mathbb{R}^n)}.$$

Now, by putting

$$v(x, y) := \frac{|u(x) - u(y)|^{p(x,y)}}{|x - y|^{\frac{n}{p(x,y)} + s}}$$

in Lemma 2.4, we can assume that

$$\frac{|u(x) - u(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} \in L^\infty(\mathbb{R}^n \times \mathbb{R}^n).$$

Therefore

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|(u - u_\delta)(x) - (u - u_\delta)(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)}} dx dy \longrightarrow 0 \quad \text{as } \delta \longrightarrow 0,$$

which concludes the proof.  $\square$

**Theorem 3.2.** *Let  $q \in \mathcal{P}^{\log}(\Omega)$  and  $p \in \mathcal{P}^{\log}(\Omega \times \Omega)$ . Then the space  $C_0^\infty(\mathbb{R}^n)$  is dense in  $W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$ .*

*Proof.* We show that for any  $u \in W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$ , there exists a sequence  $\rho_\varepsilon \in C_0^\infty(\mathbb{R}^n)$  such that

$$\|\rho_\varepsilon - u\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)} \longrightarrow 0 \quad \text{as } \varepsilon \longrightarrow 0.$$

Let  $u \in W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$ , and let us fix  $\delta > 0$ . Let  $\tau_j$  be as in Section 2. By Lemma 2.7, we know that

$$\|u - \tau_j u\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)} \leq \frac{\delta}{2},$$

for  $j$  large enough.

For any  $\varepsilon > 0$ , let us consider

$$\rho_\varepsilon := \tau_j u * \eta_\varepsilon,$$

where  $\eta_\varepsilon$  is the mollifier function defined in Section 2. By construction,  $\rho_\varepsilon \in C^\infty(\mathbb{R}^n)$ . Furthermore, by [3, Proposition IV.18],

$$\text{supp } \rho_\varepsilon \subseteq \text{supp}(\tau_j u) + \overline{B_\varepsilon}.$$

Also, by Lemma 2.7,

$$\text{supp}(\tau_j u) \subseteq \overline{B_{j+1}} \cap \text{supp } u.$$

Hence,

$$\text{supp } \rho_\varepsilon \subseteq (\overline{B_{j+1}} \cap \text{supp } u) + \overline{B_\varepsilon}.$$

As a consequence of this,

$$\rho_\varepsilon \in C_0^\infty(\mathbb{R}^n),$$

for  $\varepsilon$  small enough. Furthermore, Lemma 2.5 gives

$$\|\rho_\varepsilon - \tau_j u\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)} \leq \frac{\delta}{2}$$

for  $\varepsilon$  small enough. Therefore,

$$\begin{aligned} \|u - \rho_\varepsilon\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)} &\leq \|u - \tau_j u\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)} + \|\tau_j u - \rho_\varepsilon\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)} \\ &\leq \frac{\delta}{2} + \frac{\delta}{2} \\ &\leq \delta. \end{aligned}$$

Since  $\delta$  can be taken arbitrarily small, this concludes the proof.  $\square$

**Theorem 3.3.** *Let  $q \in \mathcal{P}^{\log}(\Omega)$ ,  $p \in \mathcal{P}^{\log}(\Omega \times \Omega)$  and suppose that  $\Omega$  is a  $W^{s,q(\cdot),p(\cdot,\cdot)}$ -extension domain. Then  $C^\infty(\overline{\Omega})$  is dense in  $W^{s,q(\cdot),p(\cdot,\cdot)}(\Omega)$ .*

*Proof.* Let  $u \in W^{s,q(\cdot),p(\cdot,\cdot)}(\Omega)$ . Since  $\Omega$  is a  $W^{s,q(\cdot),p(\cdot,\cdot)}$ -extension domain, we find that  $\tilde{u} \in W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)$  with  $\tilde{u}(x) = u(x)$  for all  $x \in \Omega$  and

$$\|\tilde{u}\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)} \leq C\|u\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\Omega)}.$$

Due to Theorem 3.2, we can choose  $\tilde{u}_\varepsilon \in C_0^\infty(\mathbb{R}^n)$  with

$$\tilde{u}_\varepsilon \longrightarrow u \quad \text{in } W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n).$$

We set  $u_\varepsilon := \tilde{u}_\varepsilon|_\Omega$ . Then

$$\|u - u_\varepsilon\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\Omega)} \leq \|\tilde{u} - \tilde{u}_\varepsilon\|_{W^{s,q(\cdot),p(\cdot,\cdot)}(\mathbb{R}^n)} \longrightarrow 0.$$

Hence  $u_\varepsilon \in C^\infty(\overline{\Omega})$  are the required approximating functions.  $\square$

**Acknowledgment.** The authors thank Lars Diening for insightful comments and for giving some very useful suggestions for the improvement of our paper.

## References

1. A. Baalal and M. Berghout, *Traces and fractional Sobolev extension domains with variable exponent*, Int. J. Math. Anal. (N.S.) **12** (2018), no. 2, 85–98, available online at <http://www.m-hikari.com/ijma/ijma-2018/ijma-1-4-2018/p/berghoutIJMA1-4-2018.pdf>. 2, 4
2. A. Bahrouni and V. D. Rădulescu, *On a new fractional Sobolev space and applications to nonlocal variational problems with variable exponent*, Discrete Contin. Dyn. Syst. Ser. S **11** (2018), no. 3, 379–389. Zbl 1374.35158. MR3732173. DOI 10.3934/dcdss.2018021. 2, 3, 7, 8
3. H. Brezis, *Analyse fonctionnelle: Théorie et applications*, Masson, Paris, 1983. Zbl 0511.46001. MR0697382. 14
4. L. Del Pezzo and J. D. Rossi, *Traces for fractional Sobolev spaces with variable exponents*, Adv. Oper. Theory **2** (2017), no. 4, 435–446. Zbl 1386.46035. MR3730039. DOI 10.22034/aot.1704-1152. 2
5. L. Diening, *Maximal function on generalized Lebesgue spaces  $L^{p(\cdot)}$* , Math. Inequal. Appl. **7** (2004), no. 2, 245–253. Zbl 1071.42014. DOI 10.7153/mia-07-27. 3
6. L. Diening, P. Harjulehto, P. Hästö, and M. Růžička, *Lebesgue and Sobolev spaces with variable exponents*, Lecture Notes in Math. **2017**, Springer, Heidelberg, 2011. Zbl 1222.46002. MR2790542. DOI 10.1007/978-3-642-18363-8. 1, 2, 12
7. S. Dipierro and E. Valdinoci, *A density property for fractional weighted Sobolev spaces*, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. **26** (2015), no. 4, 397–422. Zbl 1346.46024. MR3420496. DOI 10.4171/RLM/712. 3
8. D. E. Edmunds and J. Rákosník, *Density of smooth functions in  $W^{k,p(x)}(\Omega)$* , Proc. Roy. Soc. London Ser. A **437** (1992), no. 1899, 229–236. Zbl 0779.46027. MR1177754. DOI 10.1098/rspa.1992.0059. 2
9. X. Fan, S. Wang, and D. Zhao, *Density of  $C^\infty(\Omega)$  in  $W^{1,p(x)}(\Omega)$  with discontinuous exponent  $p(x)$* , Math. Nachr. **279** (2006), no. 1–2, 142–149. Zbl 1100.46020. MR2193613. DOI 10.1002/mana.200310351. 2
10. X. Fan and D. Zhao, *On the spaces  $L^{p(x)}(\Omega)$  and  $W^{m,p(x)}(\Omega)$* , J. Math. Anal. Appl. **263** (2001), no. 2, 424–446. Zbl 1028.46041. MR1866056. DOI 10.1006/jmaa.2000.7617. 2, 5
11. A. Fiscella, R. Servadei, and E. Valdinoci, *Density properties for fractional Sobolev spaces*, Ann. Acad. Sci. Fenn. Math. **40** (2015), no. 1, 235–253. Zbl 1346.46025. MR3310082. DOI 10.5186/aasfm.2015.4009. 10



12. G. B. Folland, *Real Analysis: Modern Techniques and Their Applications*, Pure Appl. Math., Wiley, New York, 1984. [Zbl 0549.28001](#). [MR0767633](#). [12](#)
13. P. Harjulehto, P. Hästö, Ú. V. Lê, and M. Nuortio, *Overview of differential equations with non-standard growth*, *Nonlinear Anal.* **72** (2010), no. 12, 4551–4574. [Zbl 1188.35072](#). [MR2639204](#). [DOI 10.1016/j.na.2010.02.033](#). [2](#)
14. P. A. Hästö, “Counter-examples of regularity in variable exponent Sobolev spaces” in *The  $p$ -harmonic Equation and Recent Advances in Analysis (Manhattan, 2003)*, *Contemp. Math.* **370**, Amer. Math. Soc., Providence, 2005, 133–143. [Zbl 1084.46025](#). [MR2126704](#). [DOI 10.1090/conm/370/06832](#). [2](#)
15. P. A. Hästö, *On the density of continuous functions in variable exponent Sobolev space*, *Rev. Mat. Iberoam.* **23** (2007), no. 1, 213–234. [Zbl 1144.46031](#). [MR2351132](#). [DOI 10.4171/RMI/492](#). [2](#)
16. U. Kaufmann, J. D. Rossi, and R. Vidal, *Fractional Sobolev spaces with variable exponents and fractional  $p(x)$ -Laplacians*, *Electron. J. Qual. Theory Differ. Equ.* **2017**, no. 76. [Zbl 06931307](#). [MR3725474](#). [DOI 10.14232/ejqtde.2017.1.76](#). [2](#)
17. V. Kokilashvili, A. Meskhi, H. Rafeiro, and S. Samko, *Integral Operators in Non-Standard Function Spaces, Vol. 1: Variable Exponent Lebesgue and Amalgam Spaces*, *Oper. Theory Adv. Appl.* **248**, Birkhäuser/Springer, Cham, 2016. [Zbl 1385.47001](#). [MR3559400](#). [1](#)
18. V. Kokilashvili, A. Meskhi, H. Rafeiro, and S. Samko, *Integral Operators in Non-Standard Function Spaces, Vol. 2: Variable Exponent Hölder, Morrey-Campanato and Grand Spaces*, *Oper. Theory Adv. Appl.* **249**, Birkhäuser/Springer, Cham, 2016. [Zbl 1367.47004](#). [MR3559401](#). [DOI 10.1007/978-3-319-21018-6\\_1](#). [1](#)
19. H. Nakano, *Modulated Semi-Ordered Linear Spaces*, Maruzen, Tokyo, 1950. [Zbl 0041.23401](#). [MR0038565](#). [1](#)
20. W. Orlicz, *Über konjugierte Exponentenfolgen*, *Studia Math.* **3** (1931), 200–211. [Zbl 0003.25203](#). [1](#)
21. S. Samko, *Density of  $C_0^\infty(\mathbb{R}^n)$  in the generalized Sobolev spaces  $W^{m,p(x)}(\mathbb{R}^n)$*  (in Russian), *Dokl. Ross. Akad. Nauk* **369** (1999), no. 4, 451–454; English translation in *Dokl. Math.* **60** (1999), no. 3, 382–385. [Zbl 1052.46028](#). [MR1748959](#). [2](#), [3](#), [7](#), [8](#)
22. A. Scapellato, *Homogeneous Herz spaces with variable exponents and regularity results*, *Electron. J. Qual. Theory Differ. Equ.* **2018**, no. 82. [Zbl 07011129](#). [MR3863876](#). [1](#)
23. V. V. Zhikov, *Averaging of functionals of the calculus of variations and elasticity theory* (in Russian), *Izv. Ross. Akad. Nauk Ser. Mat.* **50** (1986), no. 4, 675–710; English translation in *Izv. Math.* **29** (1987), no. 1, 33–66. [Zbl 0599.49031](#). [MR0864171](#). [2](#)
24. V. V. Zhikov, *On Lavrentiev’s phenomenon*, *Russ. J. Math. Phys.* **3** (1995), no. 2, 249–269. [Zbl 0910.49020](#). [MR1350506](#). [2](#)
25. V. V. Zhikov, *Meyer-type estimates for solving the nonlinear Stokes system* (in Russian), *Differ. Uravn.* **33** (1997), no. 1, 107–114; English translation in *Differ. Equ.* **33** (1997), no. 1, 108–115. [Zbl 0911.35089](#). [MR1607245](#). [2](#)
26. V. V. Zhikov, *On some variational problems*, *Russ. J. Math. Phys.* **5** (1997), no. 1, 105–116. [Zbl 0917.49006](#). [MR1486765](#). [2](#)
27. V. V. Zhikov, *On the density of smooth functions in Sobolev-Orlicz spaces* (in Russian), *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **310** (2004), 67–81; English translation in *J. Math. Sci. (N.Y.)* **132** (2006), no. 3, 285–294. [Zbl 1086.46026](#). [MR2120185](#). [DOI 10.1007/s10958-005-0497-0](#). [2](#)

<sup>1</sup>DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, HASSAN II UNIVERSITY OF CASABLANCA, FACULTY OF SCIENCES AÏN CHOCK, B.P. 5366 MAARIF, CASABLANCA, MOROCCO.

*E-mail address:* [abaalal@gmail.com](mailto:abaalal@gmail.com)

<sup>2</sup>DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, HASSAN II UNIVERSITY OF CASABLANCA, FACULTY OF SCIENCES AÏN CHOCK, B.P. 5366 MAARIF, CASABLANCA, MOROCCO.

*E-mail address:* [moh.berghout@gmail.com](mailto:moh.berghout@gmail.com)