



**HAL**  
open science

# Propagation of chaos for mean field rough differential equations

Ismaël Bailleul, Rémi Catellier, François Delarue

► **To cite this version:**

Ismaël Bailleul, Rémi Catellier, François Delarue. Propagation of chaos for mean field rough differential equations. *Annals of Probability*, 2021, 49 (2), pp.944-996. 10.1214/20-AOP1465 . hal-02339372

**HAL Id: hal-02339372**

**<https://hal.science/hal-02339372>**

Submitted on 30 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Propagation of chaos for mean field rough differential equations

Ismaël Bailleul \*

*Université Rennes 1  
IRMAR - UMR 6625,  
35000 Rennes, France*

*e-mail:* [ismael.bailleul@univ-rennes1.fr](mailto:ismael.bailleul@univ-rennes1.fr)

Rémi Catellier

*Université Côte d'Azur,  
Laboratoire J.A.Dieudonné,  
06108 Nice, France*

*e-mail:* [remi.catellier@unice.fr](mailto:remi.catellier@unice.fr)

François Delarue †

*Université Côte d'Azur,  
Laboratoire J.A.Dieudonné,  
06108 Nice, France*

*e-mail:* [francois.delarue@unice.fr](mailto:francois.delarue@unice.fr)

**Abstract:** We address propagation of chaos for large systems of rough differential equations associated with random rough differential equations of mean field type

$$dX_t = V(X_t, \mathcal{L}(X_t))dt + F(X_t, \mathcal{L}(X_t))dW_t,$$

where  $W$  is a random rough path and  $\mathcal{L}(X_t)$  is the law of  $X_t$ . We prove propagation of chaos, and provide also an explicit optimal convergence rate. The analysis is based upon the tools we developed in our companion paper [1] for solving mean field rough differential equations and in particular upon a corresponding version of the Itô-Lyons continuity theorem. The rate of convergence is obtained by a coupling argument developed first by Sznitman for particle systems with Brownian inputs.

**MSC 2010 subject classifications:** Primary 60H10, 60G99.

**Keywords and phrases:** random rough differential equations, particle system, mean field interaction, propagation of chaos, convergence rate.

## 1. Introduction

The study of mean field stochastic dynamics and interacting diffusions / Markov processes finds its roots in Kac's simplified approach to kinetic theory [25] and McKean's work [29] on nonlinear parabolic equations. It provides the description

---

\*I.Bailleul thanks the Centre Henri Lebesgue ANR-11-LABX-0020-01 for its stimulating mathematical research programs, and the U.B.O. for their hospitality, part of this work was written there. Partial support from the ANR-16-CE40-0020-01.

†F. Delarue thanks the Institut Universitaire de France.

of evolutions  $(\mu_t)_{t \geq 0}$  in the space of probability measures under the form of a pathspace random dynamics

$$\begin{aligned} dX_t(\omega) &= V(X_t(\omega), \mu_t)dt + F(X_t(\omega), \mu_t)dW_t(\omega), \\ \mu_t &:= \mathcal{L}(X_t), \end{aligned} \tag{1.1}$$

where  $\mathcal{L}(A)$  stands for the law of a random variable  $A$  over a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  containing  $\omega$  and relates it to the empirical behaviour of large systems of interacting dynamics. The main emphasis of subsequent works has been on proving propagation of chaos and other limit theorems, and giving stochastic representations of solutions to nonlinear parabolic equations under more and more general settings; see for instance [32, 33, 22, 15, 16, 30, 24, 5, 6]. Classical stochastic calculus makes sense of equation (1.1) only when the process  $W$  is a semi-martingale under  $\mathbb{P}$ , for some filtration, and the integrand is predictable. However, this setting happens to be too restrictive in a number of situations, especially when the diffusivity is random. This prompted several authors to address equation (1.1) by means of rough paths theory. Indeed, one may understand rough paths theory as a natural framework for providing probabilistic models of interacting populations, beyond the realm of Itô calculus. Cass and Lyons [12] did the first study of mean field random rough differential equations and proved the well-posed character of equation (1.1), and propagation of chaos for an associated system of interacting particles, under the crucial assumption that there is no mean field interaction in the diffusivity, *i.e.*  $F(x, \mu) = F(x)$ , and that the drift depends linearly on the mean field interaction. Bailleul extended partly these results in [3] by proving well-posedness of the mean field rough differential equation (1.1) in the case where the drift depends nonlinearly on the interaction term and the diffusivity is still independent of the interaction, and by proving an existence result when the diffusivity depends on the interaction. Another breakthrough came with our earlier arXiv deposit [2], in which we explained how to handle the case when  $F$  truly depends on the interaction term by making a systematic use of Lions' approach to differential calculus on Wasserstein space. To make the content more accessible, we eventually decided to split [2] into two parts: While the current work is mainly inspired from the second half of [2], our companion article [1] corresponds to the first half of [2]; Therein, we address the well-posedness of the mean field rough equation (1.1) for a genuinely nonlinear  $F$ .

In fact, as explained in [1], the general case may be easily reduced to the study of the simpler equation

$$dX_t(\omega) = F(X_t(\omega), \mathcal{L}(X_t))dW_t(\omega), \tag{1.2}$$

which is precisely the version we address in this paper. To make it clear, the purpose of the present article is to prove that, under suitable assumptions, the solution of (1.2) coincides with the limit (in a convenient sense), as  $n$  tends to  $\infty$ , of the  $n$ -particle system

$$X_t^i(\omega) = X_0^i(\omega) + \int_0^t F\left(X_s^i(\omega), \frac{1}{n} \sum_{j=1}^n \delta_{X_s^j(\omega)}\right) dW_s^i(\omega), \quad t \geq 0, \tag{1.3}$$

for  $1 \leq i \leq n$ , where  $(X_0^i(\cdot), W^i(\cdot))_{1 \leq i \leq n}$  is a collection of independent and identically distributed variables with the same distribution as  $(X_0(\cdot), W(\cdot))$ , the first component being regarded as a random variable with values in  $\mathbb{R}^d$  and the second one as a random variable with values in the space of continuous functions. Of course, equation (1.3) must be understood as a rough differential equation driven by the signal  $(W^1(\omega), \dots, W^n(\omega))$  with  $(X^1(\omega), \dots, X^n(\omega))$  as output. As it is well-known, this requires to lift  $(W^1(\omega), \dots, W^n(\omega))$  into an enhanced rough path  $\mathbf{W}^{(n)}(\omega)$  and henceforth to define the various iterated integrals. Asking the paths  $W(\omega)$ ,  $\omega \in \Omega$ , to have a finite  $p$ -variation for  $2 \leq p < 3$ , this prompts us to assume that, instead of  $((X_0^1(\cdot), W^1(\cdot)), \dots, (X_0^n(\cdot), W^n(\cdot)))$ , we have in fact  $n$  independent copies  $(X_0^i(\cdot), W^i(\cdot), \mathbb{W}^i(\cdot))_{1 \leq i \leq n}$  of the triple  $(X_0(\cdot), W(\cdot), \mathbb{W}(\cdot))$ , where  $\mathbb{W}(\omega)$  is the iterated integral of  $W(\omega)$  and  $\mathbb{W}^i(\omega)$  is the iterated integral of  $W^i(\omega)$ . Of course, it is also needed to define the iterated integrals of  $W^j(\omega)$  with respect to  $W^i(\omega)$ , for  $j \neq i$ . Not only we assume below that such iterated integrals do indeed exist, but we make the additional assumption that there is a measurable map  $\mathcal{I}$  giving  $\mathbb{W}^{i,j}(\omega)$  from  $W^i(\omega)$  and  $W^j(\omega)$ , that is

$$\mathbb{W}^{i,j}(\omega) = \mathcal{I}(W^i(\omega), W^j(\omega)), \quad i \neq j. \quad (1.4)$$

In words, (1.4) says that there exists a measurable way to construct the iterated integral of two independent copies of the signal in the limiting equation (1.2). Hence, (1.4) should be really regarded as an intrinsic property of (1.2) and not as a specific feature of the particle system (1.3).

More generally, it is in fact a key point in the subsequent analysis to draw a parallel between the underlying rough path used to give a meaning to (1.3) and the notion of *extended*<sup>1</sup> rough set-up used in [1] to address (1.2). We provide a reminder of the latter notion in Section 2. Basically, it says that, in order to solve (1.2), we must not only lift, for a given  $\omega \in \Omega$ , the trajectory  $W(\omega)$  into an enhanced rough path  $(W(\omega), \mathbb{W}(\omega))$ , but we must in fact lift the whole trajectory  $(W(\omega), W(\cdot))$ , the second component being seen as a path with values in some  $\mathbb{L}^q(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^m)$  space, where  $m$  is the dimension of the signal. Then, we call *extended* rough path set-up the enhancement of  $(W(\omega), W(\cdot))$ .

The striking fact of our analysis is then based upon an observation noticed first by Tanaka in his seminal work [34] on limit theorems for mean field type diffusions, and used crucially by Cass and Lyons in their seminal work [12]. We refer to it as Tanaka's trick. It says that, for a given  $\omega \in \Omega$ , the particle system (1.3) itself may be interpreted as a mean field equation, but with respect to the empirical measure of the driving noise. Transposed to the rough paths theory, it says that, for any fixed  $\omega \in \Omega$ , the path

$$\mathbf{W}^{(n)}(\omega) = \left( (W^i(\omega))_{1 \leq i \leq n}, (\mathbb{W}^{i,j}(\omega))_{1 \leq i, j \leq n} \right) =: \left( W^{(n)}(\omega), \mathbb{W}^{(n)}(\omega) \right),$$

which underpins the rough structure used to solve (1.3), may be seen as an extended rough set-up on its own – below, we just say a rough set-up) but

<sup>1</sup>In fact, the term *extended* does not appear in [1], but it is here of a convenient use to distinguish from the standard rough set-up used to solve the particle system (1.3).

defined on the finite probability space

$$\left( \{1, \dots, n\}, \mathcal{P}(\{1, \dots, n\}), \frac{1}{n} \sum_{i=1}^n \delta_i \right),$$

where  $\mathcal{P}(\{1, \dots, n\})$  denotes the collection of subsets of  $\{1, \dots, n\}$ , instead of the former probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . We call this set-up the **empirical rough set-up**, and we make its construction entirely clear in the sequel of the paper. For sure, given the iterated integrals of the signal  $(W^1(\omega), \dots, W^n(\omega))$ , the rough integral (1.3) should be interpreted in the usual sense, as given by standard Lyons' rough paths theory. In short, this requires to expand locally the integrand in (1.3), which in turns requires to have a convenient notion of derivative with respect to the measure argument. In this regard, a crucial fact in [1] is to use Lions' approach [27, 7, 9] to differential calculus on the space  $\mathcal{P}_2(\mathbb{R}^d)$  of probability measures on  $\mathbb{R}^d$  with a finite square moment, the so-called  $d$ -dimensional Wasserstein space,  $d$  denoting here and throughout the dimension of the output in (1.2). The core of our analysis in Section 4 is that, whenever Wasserstein derivatives on  $\mathcal{P}_2(\mathbb{R}^d)$  are projected, through empirical measures, into classical derivatives on  $(\mathbb{R}^d)^n$ , as it is needed to differentiate the integrand in (1.3), the resulting solution for (1.3), as given by standard rough paths theory, coincides with the solution obtained by interpreting (1.3) as a mean field rough equation driven by the aforementioned empirical rough set-up – see Section 3 for reminders on solvability results for mean field rough equations. In this way, the convergence of solutions of (1.3) to solutions of (1.2) as  $n$  tends to  $\infty$  is reduced to a form of continuity of the solutions to mean field rough differential equations with respect to the underlying rough set-up. We called the latter *continuity of the Itô-Lyons solution map*, see Theorem 5.4 of our companion work [1]. Our first main result, Theorem 4.3, shows that, for a sufficiently large class of input signals, propagation of chaos is in fact a consequence of the *continuity of the Itô-Lyons solution map* for mean field rough differential equations. At this stage, it is worth mentioning that it is precisely in the requirements of the *continuity of the Itô-Lyons map* that the structure condition (1.4) about the cross-iterated integrals comes in. In [1], a *rough set-up* that satisfies (1.4) is said to be *strong*.

While the proofs of both our first main result and the underlying *continuity property of the Itô-Lyons solution map* are mostly based on compactness arguments, our second main result is to elucidate, under slightly stronger assumptions the convergence rate in the propagation of chaos; see Theorem 5.1 in Section 5. The strategy is directly inspired from original Sznitman's coupling argument for mean field systems driven by Brownian signals, see [32]. Although the proof is much more involved than in the Brownian setting, we recover the same rate of convergence: It coincides with the rate of convergence (in Wasserstein metric) of the empirical measure of an  $n$ -sample of (sufficiently integrable) i.i.d. variables to their common distribution. In particular, the speed decays with the dimension.

As in [1], our analysis holds for continuous rough paths whose  $p$ -variation, for some  $p \in [2, 3)$ , is finite and has sub-exponential tails and for which the so-called *local accumulated variation* –that counts the increments of the signal of a given size over a bounded interval– has super-exponential tails, see [1, Theorem 1.1]. Among others, our results apply to continuous centred Gaussian signals defined over some time interval  $[0, T]$  that have independent components and whose covariance function has finite  $\rho$ -two dimensional variation, for some  $\rho \in [1, 3/2)$ .

The present work leaves wide open the question of refining the strong law of large numbers given by the propagation of chaos result stated in Theorem 4.3. A central limit theorem for the fluctuations of the empirical measure of the particle system is expected to hold under reasonable conditions on the common law of the rough drivers. Large and moderate deviation results would also be most welcome. In a different direction, it would be interesting to investigate the propagation of chaos phenomenon for systems of interacting rough dynamics subject to a common noise. Very interesting things happen in the Itô setting in relation with mean field games [8, 26]. Also, one would get a more realistic model of natural phenomena by working with systems of particles driven by non-independent noises. Individuals with close initial conditions could have drivers strongly correlated while individuals started far apart could have (almost-)independent drivers. Limit mean field dynamics are likely to be different from the results obtained here – see [14] for a result in this direction in the Itô setting. We invite the reader to make her/his own mind about these problems.

The paper is organized as follows. We recall in Section 2 the construction of a rough set-up, as introduced in [1]. We provide in Section 3 a sketchy presentation of related solvability results for equation (1.2), including a review of the main assumptions that we need on the diffusivity  $F$ . Convergence of the particle system (1.3) is established in Section 4. The convergence rate is addressed in Section 5, under additional regularity assumptions on  $F$  and integrability assumptions on the signal. Proofs of some technical results are given in Appendix A.1 and A.2.

**Notations.** We gather here a number of notations that will be used throughout the text.

- We set  $\mathcal{S}_2 := \{(s, t) \in [0, \infty)^2 : s \leq t\}$ , and  $\mathcal{S}_2^T := \{(s, t) \in [0, T]^2 : s \leq t\}$ .
- We denote by  $(\Omega, \mathcal{F}, \mathbb{P})$  an atomless Polish probability space,  $\mathcal{F}$  standing for the completion of the Borel  $\sigma$ -field under  $\mathbb{P}$ , and denote by  $\langle \cdot \rangle$  the expectation operator, by  $\langle \cdot \rangle_r$ , for  $r \in [1, +\infty]$ , the  $\mathbb{L}^r$ -norm on  $(\Omega, \mathcal{F}, \mathbb{P})$  and by  $\langle \cdot \rangle$  and  $\langle \cdot \rangle_r$  the expectation operator and the  $\mathbb{L}^r$ -norm on  $(\Omega^2, \mathcal{F}^{\otimes 2}, \mathbb{P}^{\otimes 2})$ . When  $r$  is finite,  $\mathbb{L}^r(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R})$  is separable as  $\Omega$  is Polish.
- As for processes  $X_\bullet = (X_t)_{t \in I}$ , defined on a time interval  $I$ , we often write  $X$  for  $X_\bullet$ .

## 2. From Probabilistic Rough Structures to Rough Integrals

### 2.1. Overview on Probabilistic Rough Structure

We here provide a brief reminder of the content of Section 2 in [1]. We refer the reader to the paper for a complete review. Throughout the section, we work on a finite time horizon  $[0, T]$ , for a given  $T > 0$ .

The first level of the rough path structure used to give a meaning to (1.2) is defined as an  $\omega$ -indexed pair of paths

$$(W_t(\omega), W_t(\cdot))_{0 \leq t \leq T}, \quad (2.1)$$

where  $(W_t(\cdot))_{0 \leq t \leq T}$  is a collection of  $q$ -integrable  $\mathbb{R}^m$ -valued random variables on  $(\Omega, \mathcal{F}, \mathbb{P})$ , which we regard as a deterministic  $\mathbb{L}^q(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^m)$ -valued path, for some exponent  $q \geq 8$ , and  $(W_t(\omega))_{0 \leq t \leq T}$  stands for the realizations of these random variables along the outcome  $\omega \in \Omega$ ; so the pair (2.1) takes values in  $\mathbb{R}^m \times \mathbb{L}^q(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^m)$ . The second level has the form of an  $\omega$ -dependent two-index path with values in  $(\mathbb{R}^m \times \mathbb{L}^q(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^m))^{\otimes 2}$  and is encoded in matrix form as

$$\begin{pmatrix} W_{s,t}(\omega) & W_{s,t}^{\perp}(\omega, \cdot) \\ W_{s,t}^{\perp}(\cdot, \omega) & W_{s,t}^{\perp}(\cdot, \cdot) \end{pmatrix}_{0 \leq s \leq t \leq T}, \quad (2.2)$$

where

- $W_{s,t}(\omega)$  is in  $(\mathbb{R}^m)^{\otimes 2} \simeq \mathbb{R}^{m \times m}$ ,
- $W_{s,t}^{\perp}(\omega, \cdot)$  is in  $\mathbb{R}^m \otimes \mathbb{L}^q(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^m) \simeq \mathbb{L}^q(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^{m \times m})$ ,
- $W_{s,t}^{\perp}(\cdot, \omega)$  is in  $\mathbb{L}^q(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^m) \otimes \mathbb{R}^m \simeq \mathbb{L}^q(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^{m \times m})$ ,
- $W_{s,t}^{\perp}(\cdot, \cdot)$  is in  $\mathbb{L}^q(\Omega^{\otimes 2}, \mathcal{F}^{\otimes 2}, \mathbb{P}^{\otimes 2}; \mathbb{R}^{m \times m})$ , the realizations of which read in the form  $\Omega^2 \ni (\omega, \omega') \mapsto W_{s,t}^{\perp}(\omega, \omega') \in \mathbb{R}^{m \times m}$  and the two sections of which are precisely given by  $W_{s,t}^{\perp}(\omega, \cdot) : \Omega \ni \omega' \mapsto W_{s,t}^{\perp}(\omega, \omega')$ , and  $W_{s,t}^{\perp}(\cdot, \omega) \ni \omega' \mapsto W_{s,t}^{\perp}(\omega', \omega)$ , for  $\omega \in \Omega$ .

A convenient form of Chen's relations is required, for any  $\omega \in \Omega$ ,

$$\begin{aligned} W_{r,t}(\omega) &= W_{r,s}(\omega) + W_{s,t}(\omega) + W_{r,s}(\omega) \otimes W_{s,t}(\omega), \\ W_{r,t}^{\perp}(\cdot, \omega) &= W_{r,s}^{\perp}(\cdot, \omega) + W_{s,t}^{\perp}(\cdot, \omega) + W_{r,s}(\cdot) \otimes W_{s,t}(\omega), \\ W_{r,t}^{\perp}(\omega, \cdot) &= W_{r,s}^{\perp}(\omega, \cdot) + W_{s,t}^{\perp}(\omega, \cdot) + W_{r,s}(\omega) \otimes W_{s,t}(\cdot), \\ W_{r,t}^{\perp}(\cdot, \cdot) &= W_{r,s}^{\perp}(\cdot, \cdot) + W_{s,t}^{\perp}(\cdot, \cdot) + W_{r,s}(\cdot) \otimes W_{s,t}(\cdot), \end{aligned} \quad (2.3)$$

for any  $0 \leq r \leq s \leq t \leq T$ , with notation  $f_{r,s} := f_s - f_r$ , for a function  $f$  from  $[0, \infty)$  into a vector space. In (2.3), we denoted by  $X(\cdot) \otimes Y(\cdot)$ , for any two  $X$  and  $Y$  in  $\mathbb{L}^q(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^m)$ , the random variable  $(\omega, \omega') \mapsto (X_i(\omega)Y_j(\omega'))_{1 \leq i, j \leq m}$  defined on  $\Omega^2$ . It is in  $\mathbb{L}^q(\Omega^2, \mathcal{F}^{\otimes 2}, \mathbb{P}^{\otimes 2}; \mathbb{R}^{m \times m})$ . The notation  $\perp$  in  $W^{\perp}$  is used to indicate that  $W_{s,t}^{\perp}(\cdot, \cdot)$  should be thought of as the random variable

$$(\omega, \omega') \mapsto \int_s^t (W_r(\omega) - W_s(\omega)) \otimes dW_r(\omega').$$

Since  $\Omega^2 \ni (\omega, \omega') \mapsto (W_t(\omega))_{t \geq 0}$  and  $\Omega^2 \ni (\omega, \omega') \mapsto (W_t(\omega'))_{t \geq 0}$  are independent under  $\mathbb{P}^{\otimes 2}$ , we then understand  $\mathbb{W}_{s,t}^\perp$  as an iterated integral for two independent copies of the noise. We refer to Examples 2.3 and 2.5 in [1]. In the end, we denote by  $\mathbf{W}(\omega)$  the so-called rough set-up specified by the  $\omega$ -dependent collection of maps given by (2.1) and (2.2).

## 2.2. Regularity of the Rough Set-Up

Following [1], we use the notion of  $p$ -variation to handle the regularity of the various trajectories in hand. Throughout, the exponent  $p$  is taken in the interval  $[2, 3)$ . For a continuous function  $\mathbb{G}$  from the simplex  $\mathcal{S}_2^T$  into some  $\mathbb{R}^\ell$ , we set, for any  $p' \geq 1$ ,

$$\|\mathbb{G}\|_{[0,T],p'-v}^{p'} := \sup_{0=t_0 < t_1 \dots < t_n=T} \sum_{i=1}^n |\mathbb{G}_{t_{i-1}, t_i}|^{p'},$$

and define for any function  $g$  from  $[0, T]$  into  $\mathbb{R}^\ell$ ,  $\|g\|_{[0,T],p-v}^p := \|\mathbb{G}\|_{[0,T],p-v}^p$  as the  $p$ -variation semi-norm of its associated two index function  $\mathbb{G}_{s,t} := g_t - g_s$ . Similarly, for a random variable  $\mathbb{G}(\cdot)$  on  $\Omega$  with values in  $\mathcal{C}(\mathcal{S}_2^T; \mathbb{R}^\ell)$ , and  $p' \geq 1$ , we define its  $p'$ -variation in  $\mathbb{L}^q$  as

$$\langle \mathbb{G}(\cdot) \rangle_{q;[0,T],p'-v}^{p'} := \sup_{0=t_0 < t_1 \dots < t_n=T} \sum_{i=1}^n \langle \mathbb{G}_{t_{i-1}, t_i}(\cdot) \rangle_q^{p'}, \quad (2.4)$$

and define for a random variable  $G(\cdot)$  on  $\Omega$ , with values in  $\mathcal{C}([0, T]; \mathbb{R}^\ell)$ ,

$$\langle G(\cdot) \rangle_{q;[0,T],p'-v}^{p'} := \langle \mathbb{G}(\cdot) \rangle_{q;[0,T],p'-v}^{p'},$$

as the  $p'$ -variation semi-norm in  $\mathbb{L}^q$  of its associated two-index function  $\mathcal{S}_2^T \ni (s, t) \mapsto \mathbb{G}_{s,t}(\cdot) = G_t(\cdot) - G_s(\cdot)$ . Lastly, for a random variable  $\mathbb{G}(\cdot, \cdot)$  from  $(\Omega^2, \mathcal{F}^{\otimes 2})$  into  $\mathcal{C}(\mathcal{S}_2^T; \mathbb{R}^\ell)$ , we set

$$\langle\langle \mathbb{G}(\cdot, \cdot) \rangle\rangle_{q;[0,T],p'/2-v}^{p'/2} := \sup_{0=t_0 < t_1 \dots < t_n=T} \sum_{i=1}^n \langle\langle \mathbb{G}_{t_{i-1}, t_i}(\cdot, \cdot) \rangle\rangle_q^{p'/2}. \quad (2.5)$$

Given these definitions, we require from the rough set-up  $\mathbf{W}$  that

- For any  $\omega \in \Omega$ , the path  $W(\omega)$  is in the space  $\mathcal{C}([0, T]; \mathbb{R}^m)$ , and the map  $W : \Omega \ni \omega \mapsto W(\omega) \in \mathcal{C}([0, T]; \mathbb{R}^m)$  is Borel-measurable and  $q$ -integrable.
- For any  $\omega \in \Omega$ , the two-index path  $\mathbb{W}(\omega)$  is in  $\mathcal{C}(\mathcal{S}_2^T; \mathbb{R}^{m \times m})$ , and the map  $\mathbb{W} : \Omega \ni \omega \mapsto \mathbb{W}(\omega) \in \mathcal{C}(\mathcal{S}_2^T; \mathbb{R}^{m \times m})$  is Borel-measurable and  $q$ -integrable.
- For any  $(\omega, \omega') \in \Omega^2$ , the two-index path  $\mathbb{W}^\perp(\omega, \omega')$  is an element of  $\mathcal{C}(\mathcal{S}_2^T; \mathbb{R}^{m \times m})$ , and the map  $\mathbb{W}^\perp : \Omega^2 \ni (\omega, \omega') \mapsto \mathbb{W}^\perp(\omega, \omega') \in \mathcal{C}(\mathcal{S}_2^T; \mathbb{R}^{m \times m})$  is Borel-measurable and  $q$ -integrable.

Moreover, we may set, for some fixed  $p \in [2, 3)$  and for all  $0 \leq s \leq t \leq T$  and  $\omega \in \Omega$ ,

$$\begin{aligned} v(s, t, \omega) := & \|W(\omega)\|_{[s,t],p-v}^p + \langle W(\cdot) \rangle_{q:[s,t],p-v}^p \\ & + \|\mathbb{W}(\omega)\|_{[s,t],p/2-v}^{p/2} + \langle \mathbb{W}^\perp(\omega, \cdot) \rangle_{q:[s,t],p/2-v}^{p/2} \\ & + \langle \mathbb{W}^\perp(\cdot, \omega) \rangle_{q:[s,t],p/2-v}^{p/2} + \langle\langle \mathbb{W}^\perp(\cdot, \cdot) \rangle\rangle_{q:[s,t],p/2-v}^{p/2}, \end{aligned} \quad (2.6)$$

and we assume that, for any positive finite time  $T$  and any  $\omega \in \Omega$ , the quantity  $v(0, T, \omega)$  is finite. Importantly,  $\omega \mapsto (v(s, t, \omega))_{(s,t) \in \mathcal{S}_2^T}$  is a random variable with values in  $\mathcal{C}(\mathcal{S}_2^T; \mathbb{R}_+)$  and is super-additive, namely, for any  $0 \leq r \leq s \leq t \leq T$ , and  $\omega \in \Omega$ ,

$$v(r, t, \omega) \geq v(r, s, \omega) + v(s, t, \omega).$$

We then assume  $\langle v(0, T, \cdot) \rangle_q < \infty$ , which implies, by Lebesgue's dominated convergence theorem, that the function  $\mathcal{S}_2^T \ni (s, t) \mapsto \langle v(s, t, \cdot) \rangle_q$  is continuous. We assume that it is of bounded variation on  $[0, T]$ , *i.e.*

$$\langle v(\cdot) \rangle_{q:[s,t],1-v} := \sup_{0 \leq t_1 < \dots < t_K \leq T} \sum_{i=1}^K \langle v(t_{i-1}, t_i, \cdot) \rangle_q < \infty. \quad (2.7)$$

We then call a control any family of random variables  $(\omega \mapsto w(s, t, \omega))_{(s,t) \in \mathcal{S}_2^T}$  that is jointly continuous in  $(s, t)$  and that satisfies,

$$w(s, t, \omega) \geq v(s, t, \omega) + \langle v(\cdot) \rangle_{q:[s,t],1-v}, \quad (2.8)$$

together with

$$\begin{aligned} \langle w(s, t, \cdot) \rangle_q & \leq 2 w(s, t, \omega), \\ w(r, t, \omega) & \geq w(r, s, \omega) + w(s, t, \omega), \quad r \leq s \leq t. \end{aligned} \quad (2.9)$$

A typical choice to get (2.10) and (2.9) is to choose

$$w(s, t, \omega) := v(s, t, \omega) + \langle v(\cdot) \rangle_{q:[s,t],1-v}. \quad (2.10)$$

### 2.3. Controlled Trajectories

With a rough set-up at hands on a given finite time interval  $[0, T]$ , we define an associated notion of controlled path and rough integral in the spirit of Gubinelli [23]. Again, we refer to [1] for details, see Definition 3.1 therein.

**Definition 2.1.** *An  $\omega$ -dependent continuous  $\mathbb{R}^d$ -valued path  $(X_t(\omega))_{0 \leq t \leq T}$  is called an  $\omega$ -controlled path on  $[0, T]$  if its increments can be decomposed as*

$$X_{s,t}(\omega) = \delta_x X_s(\omega) W_{s,t}(\omega) + \mathbb{E}[\delta_\mu X_s(\omega, \cdot) W_{s,t}(\cdot)] + R_{s,t}^X(\omega), \quad (2.11)$$

where  $(\delta_x X_t(\omega))_{0 \leq t \leq T}$  belongs to  $\mathcal{C}([0, T]; \mathbb{R}^{d \times m})$  and  $(\delta_\mu X_t(\omega, \cdot))_{0 \leq t \leq T}$  to  $\mathcal{C}([0, T]; \mathbb{L}^{4/3}(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^{d \times m}))$ ,  $(R_{s,t}^X(\omega))_{s,t \in \mathcal{S}_2^T}$  is in the space  $\mathcal{C}(\mathcal{S}_2^T; \mathbb{R}^d)$ , and

$$\begin{aligned} \|X(\omega)\|_{\star, [0, T], w, p} &:= |X_0(\omega)| + |\delta_x X_0(\omega)| + \langle \delta_\mu X_0(\omega, \cdot) \rangle_{4/3} \\ &+ \|X(\omega)\|_{[0, T], w, p} < \infty, \end{aligned}$$

where

$$\begin{aligned} \|X(\omega)\|_{[0, T], w, p} &:= \|X(\omega)\|_{[0, T], w, p} + \|\delta_x X(\omega)\|_{[0, T], w, p} \\ &+ \langle \delta_\mu X(\omega, \cdot) \rangle_{[0, T], w, p, 4/3} + \|R^X(\omega)\|_{[0, T], w, p/2}, \end{aligned}$$

with

$$\begin{aligned} \|X(\omega)\|_{[0, T], w, p} &:= \sup_{\emptyset \neq (s,t) \subset [0, T]} \frac{|X_{s,t}(\omega)|}{w(s, t, \omega)^{1/p}}, \quad \text{and similarly for } \partial_x X \\ \langle \delta_\mu X(\omega, \cdot) \rangle_{[0, T], w, p, 4/3} &:= \sup_{\emptyset \neq (s,t) \subset [0, T]} \frac{\langle \delta_\mu X_{s,t}(\omega, \cdot) \rangle_{4/3}}{w(s, t, \omega)^{1/p}}, \\ \|R^X(\omega)\|_{[0, T], w, p/2} &:= \sup_{\emptyset \neq (s,t) \subset [0, T]} \frac{|R_{s,t}^X(\omega)|}{w(s, t, \omega)^{2/p}}. \end{aligned}$$

We call  $\delta_x X(\omega)$  and  $\delta_\mu X(\omega, \cdot)$  in (2.11) the derivatives of  $X(\omega)$ .

We then define the notion of random controlled trajectory, which consists of a collection of  $\omega$ -controlled trajectories indexed by the elements of  $\Omega$ .

**Definition 2.2.** A family of  $\omega$ -controlled paths  $(X(\omega))_{\omega \in \Omega}$  such that  $X$ ,  $\delta_x X$ ,  $\delta_\mu X$  and  $R^X$  are measurable from  $\Omega$  into  $\mathcal{C}([0, T]; \mathbb{R}^d)$ ,  $\mathcal{C}([0, T]; \mathbb{R}^{d \times m})$ ,  $\mathcal{C}([0, T]; \mathbb{L}^{4/3}(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^{d \times m}))$  and  $\mathcal{C}(\mathcal{S}_2^T; \mathbb{R}^d)$ , and satisfy

$$\langle X_0(\cdot) \rangle_2 + \langle \|X(\cdot)\|_{[0, T], w, p} \rangle_8 < \infty \quad (2.12)$$

is called a random controlled path on  $[0, T]$ .

It is proven in [1, Lemma 3.3] that a random controlled trajectory induces a continuous path  $t \mapsto X_t(\cdot)$  from  $[0, T]$  to  $L^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^d)$ .

#### 2.4. Rough Integral

As for the construction, of the rough integral, we recall the following statement from [1, Theorem 3.4].

**Theorem 2.3.** There exists a universal constant  $c_0$  and, for any  $\omega \in \Omega$ , there exists a continuous linear map

$$(X_t(\omega))_{0 \leq t \leq T} \mapsto \left( \int_s^t X_{s,u}(\omega) \otimes d\mathbf{W}_u(\omega) \right)_{(s,t) \in \mathcal{S}_2^T}$$

from the space of  $\omega$ -controlled trajectories equipped with the norm  $\|\cdot\|_{\star,[0,T],p}$ , onto the space of continuous functions from  $\mathcal{S}_2^T$  into  $\mathbb{R}^d \otimes \mathbb{R}^m$  with finite norm  $\|\cdot\|_{[0,T],w,p/2}$ , with  $w$  in the latter norm being evaluated along the realization  $\omega$ , that satisfies for any  $0 \leq r \leq s \leq t \leq T$  the identity

$$\begin{aligned} & \int_r^t X_{r,u}(\omega) \otimes d\mathbf{W}_u(\omega) \\ &= \int_r^s X_{r,u}(\omega) \otimes d\mathbf{W}_u(\omega) + \int_s^t X_{s,u}(\omega) \otimes d\mathbf{W}_u(\omega) + X_{r,s}(\omega) \otimes W_{s,t}(\omega), \end{aligned}$$

together with the estimate

$$\begin{aligned} & \left| \int_s^t X_{s,u}(\omega) \otimes d\mathbf{W}_u(\omega) - \delta_x X_s(\omega) \mathbb{W}_{s,t}(\omega) - \mathbb{E}[\delta_\mu X_s(\omega, \cdot) \mathbb{W}_{s,t}^\perp(\cdot, \omega)] \right| \\ & \leq c_0 \|X(\omega)\|_{[0,T],w,p} w(s, t, \omega)^{3/p}. \end{aligned} \quad (2.13)$$

Above,  $\delta_x X_s(\omega) \mathbb{W}_{s,t}(\omega)$  is the product of a  $d \times m$  matrix and an  $m \times m$  matrix, so it gives back a  $d \times m$  matrix, with components  $(\delta_x X_s(\omega) \mathbb{W}_{s,t}(\omega))_{i,j} = \sum_{k=1}^m (\delta_x X_s^i(\omega))_k (\mathbb{W}_{s,t}(\omega))_{k,j}$ , for  $i \in \{1, \dots, d\}$  and  $j \in \{1, \dots, m\}$ , and similarly for  $\mathbb{E}[\delta_\mu X_s(\omega, \cdot) \mathbb{W}_{s,t}^\perp(\cdot, \omega)]$ . As usual, the above construction allows us to define an additive process setting

$$\int_s^t X_u(\omega) \otimes d\mathbf{W}_u(\omega) := \int_s^t X_{s,u}(\omega) \otimes d\mathbf{W}_u(\omega) + X_s(\omega) \otimes W_{s,t}(\omega),$$

for  $0 \leq t \leq T$ . We can thus consider the integral process  $(\int_0^t X_s(\omega) \otimes d\mathbf{W}_s(\omega))_{0 \leq t \leq T}$  as an  $\omega$ -controlled trajectory with values in  $\mathbb{R}^{d \times m}$ , with

$$\left( \delta_x \left[ \int_0^\cdot X_s(\omega) \otimes d\mathbf{W}_s(\omega) \right]_t \right)_{(i,j),k} = (X_t(\omega))_i \delta_{j,k},$$

for  $i \in \{1, \dots, d\}$  and  $j, k \in \{1, \dots, m\}$ , where  $\delta_{j,k}$  stands for the usual Kronecker symbol, and with null  $\mu$ -derivative.

When the trajectory  $X(\omega)$  takes in values in  $\mathbb{R}^d \otimes \mathbb{R}^m$  rather than  $\mathbb{R}^d$ , the integral  $\int_0^t X_s(\omega) \otimes d\mathbf{W}_s(\omega)$  belongs to  $\mathbb{R}^d \otimes \mathbb{R}^m \otimes \mathbb{R}^m$ . We then set for  $i \in \{1, \dots, d\}$

$$\left( \int_0^t X_s(\omega) d\mathbf{W}_s(\omega) \right)_i := \sum_{j=1}^m \left( \int_0^t X_s(\omega) \otimes d\mathbf{W}_s(\omega) \right)_{i,j,j},$$

and consider  $\int_0^t X_s(\omega) d\mathbf{W}_s(\omega)$  as an element of  $\mathbb{R}^d$ .

## 2.5. Stability of Controlled Paths under Nonlinear Maps

A key fact in [1] is to use regularity properties of functions defined on Wasserstein space through a lifting procedure to an  $L^2$  space standing above the probability space. We refer the reader to Lions' lectures [27], to the lecture notes [7] of Cardaliaguet or to Carmona and Delarue's monograph [9, Chapter 5] for basics on the subject.

- Recall  $(\Omega, \mathcal{F}, \mathbb{P})$  stands for an atomless probability space, with  $\Omega$  a Polish space and  $\mathcal{F}$  its Borel  $\sigma$ -algebra. Fix a finite dimensional space  $E = \mathbb{R}^k$  and denote, for  $r \geq 1$ , by  $L^r := \mathbb{L}^r(\Omega, \mathcal{F}, \mathbb{P}; E)$  the space of  $E$ -valued random variables on  $\Omega$  with finite  $r$  moment. We equip the space  $\mathcal{P}_r(E) := \{\mathcal{L}(Z); Z \in L^r\}$  with the  $r$ -Wasserstein distance

$$\mathbf{d}_r(\mu_1, \mu_2) := \inf \left\{ \|Z_1 - Z_2\|_r; \mathcal{L}(Z_1) = \mu_1, \mathcal{L}(Z_2) = \mu_2 \right\}. \quad (2.14)$$

When  $r = 2$ , an  $\mathbb{R}^k$ -valued function  $u$  defined on  $\mathcal{P}_2(E)$  is canonically extended into  $L^2$  by setting, for any  $Z \in L^2$ ,  $U(Z) := u(\mathcal{L}(Z))$ .

- The function  $u$  is then said to be differentiable at  $\mu \in \mathcal{P}_2(E)$  if its canonical lift is Fréchet differentiable at some point  $Z$  such that  $\mathcal{L}(Z) = \mu$ ; we denote by  $\nabla_Z U \in (L^2)^k$  the gradient of  $U$  at  $Z$ . The function  $U$  is then differentiable at any other point  $Z' \in L^2$  such that  $\mathcal{L}(Z') = \mu$ , and the laws of  $\nabla_Z U$  and  $\nabla_{Z'} U$  are equal, for any such  $Z'$ .

- The function  $u$  is said to be of class  $C^1$  if its canonical lift is of class  $C^1$ . If  $u$  is of class  $C^1$  on  $\mathcal{P}_2(E)$ , then  $\nabla_Z U$  is  $\sigma(Z)$ -measurable and given by an  $\mathcal{L}(Z)$ -dependent function  $Du$  from  $E$  to  $E^k$  such that

$$\nabla_Z U = (Du)(Z). \quad (2.15)$$

In order to emphasize the fact that  $Du$  depends upon  $\mathcal{L}(Z)$ , we shall write  $D_\mu u(\mathcal{L}(Z))(\cdot)$  instead of  $Du(\cdot)$ . Importantly, this representation is independent of the choice of the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  and can be easily transported from one probability space to another.

Throughout the paper, we regard the function  $F$  in (1.2) as a map from  $\mathbb{R}^d \times \mathbb{L}^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^d)$  into the space  $L(\mathbb{R}^m, \mathbb{R}^d) \cong \mathbb{R}^d \otimes \mathbb{R}^m$  of linear mappings from  $\mathbb{R}^m$  to  $\mathbb{R}^d$ . Intuitively, we identify the coefficient driving equation (1.2) with its lift  $\hat{F}$ . Following [1, Subsection 3.3], we require  $F$  to satisfy the following regularity assumptions.

**Regularity assumptions 1** – Assume that  $F$  is continuously differentiable in the joint variable  $(x, Z)$ , that  $\partial_x F$  is also continuously differentiable in  $(x, Z)$  and that there is some positive finite constant  $\Lambda$  such that  $|F(x, \mu)|$ ,  $|\partial_x F(x, \mu)|$ ,  $|\partial_x^2 F(x, \mu)|$ ,  $\|\nabla_Z F(x, Z)\|_2$  and  $\|\partial_x \nabla_Z F(x, Z)\|_2$  are bounded by  $\Lambda$ , for any  $x \in \mathbb{R}^d$ ,  $\mu \in \mathcal{P}_2(\mathbb{R}^d)$  and  $Z \in \mathbb{L}^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^d)$ . Assume moreover that, for any  $x \in \mathbb{R}^d$ , the mapping  $Z \mapsto \nabla_Z F(x, Z)$  is a  $\Lambda$ -Lipschitz function of  $Z \in \mathbb{L}^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^d)$ .

We recall below that, for an  $\omega$ -controlled path  $X(\omega)$  and for an  $\mathbb{R}^d$ -valued random controlled path  $Y(\cdot)$ ,  $F(X(\omega), Y(\cdot)) := (F(X_t(\omega), Y_t(\cdot)))_{0 \leq t \leq T}$  may be also expanded in the form of an  $\omega$ -controlled trajectory. As explained in [1, (3.8)], it suffices for our purpose to provide the form of the expansion when  $\delta_\mu X(\omega) \equiv 0$  and  $\delta_\mu Y(\cdot) \equiv 0$ .

**Proposition 2.4.** *Let  $X(\omega)$  be an  $\omega$ -controlled path and  $Y(\cdot)$  be an  $\mathbb{R}^d$ -valued random controlled path. Assume that  $\delta_\mu X(\omega) \equiv 0$  and  $\delta_\mu Y(\cdot) \equiv 0$  and that  $\sup_{0 \leq t \leq T} (|\delta_x X_t(\omega)| \vee \langle \delta_x Y_t(\cdot) \rangle_\infty) < \infty$ . Then,  $F(X(\omega), Y(\cdot))$  is an  $\omega$ -controlled path with*

$$\delta_x \left( F(X(\omega), Y(\cdot)) \right)_t = \partial_x F(X_t(\omega), Y_t(\cdot)) \delta_x X_t(\omega),$$

which is understood as  $(\sum_{\ell=1}^d \partial_{x_t} F^{i,j}(X_t(\omega), Y_t(\cdot)) (\delta_x X_t^\ell(\omega))_k)_{i,j,k}$ , with  $i, k \in \{1, \dots, d\}$  and  $j \in \{1, \dots, m\}$ , and (with a similar interpretation for the product)

$$\delta_\mu \left( F(X(\omega), Y(\cdot)) \right)_t = D_\mu F(X_t(\omega), \mathcal{L}(Y_t)) (X_t(\cdot)) \delta_x Y_t(\cdot).$$

## 2.6. Local Accumulation

In order to proceed with the analysis of (1.3), we make use of the notion of local accumulation. Following [1], we define it as follows. Given a nondecreasing<sup>2</sup> continuous positive valued function  $\varpi$  on  $\mathcal{S}_2$ , a non-negative parameter  $s$  and a positive threshold  $\alpha$ , we define inductively a sequence of times setting  $\tau_0(s, \alpha) := s$ , and

$$\tau_{n+1}^\varpi(s, \alpha) := \inf \left\{ u \geq \tau_n^\varpi(s, \alpha) : \varpi(\tau_n^\varpi(s, \alpha), u) \geq \alpha \right\}, \quad (2.16)$$

with the understanding that  $\inf \emptyset := +\infty$ . For  $t \geq s$ , set

$$N_\varpi([s, t], \alpha) := \sup \left\{ n \in \mathbb{N} : \tau_n^\varpi(s, \alpha) \leq t \right\}. \quad (2.17)$$

We call  $N_\varpi$  the local accumulation of  $\varpi$  (of size  $\alpha$  if we specify the value of the threshold):  $N_\varpi([s, t], \alpha)$  is the largest number of disjoint open sub-intervals  $(a, b)$  of  $[s, t]$  on which  $\varpi(a, b)$  is greater than or equal to  $\alpha$ . When  $\varpi(s, t) = w(s, t, \omega)^{1/p}$  with  $w$  a control satisfying (2.8) and (2.9) and when the framework makes it clear, we just write  $N([s, t], \omega, \alpha)$  for  $N_\varpi([s, t], \alpha)$ . Similarly, we also write  $\tau_n(s, \omega, \alpha)$  for  $\tau_n^\varpi(s, \alpha)$  when  $\varpi(s, t) = w(s, t, \omega)$ . We will also use the convenient notation

$$\tau_n^\varpi(s, t, \alpha) := \tau_n^\varpi(s, \alpha) \wedge t.$$

## 3. Analysis of the Mean Field Rough Differential Equation

### 3.1. Solving the Equation

The following notion of solution to (1.2) is taken from [1, Definition 4.1].

<sup>2</sup>In the sense that  $\varpi(a, b) \geq \varpi(a', b')$  if  $(a', b') \subset (a, b)$ .

**Definition 3.1.** Let  $W$  together with its enhancement  $\mathbf{W}$  satisfy the assumption of Section 2.2 on a finite interval  $[0, T]$ . A solution to (1.2) on the time interval  $[0, T]$ , with initial condition  $X_0(\cdot) \in L^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^d)$ , is a random controlled path  $X(\cdot)$ , such that for  $\mathbb{P}$ -a.e.  $\omega$  the paths  $X(\omega)$  and  $X_0(\omega) + \int_0^\cdot F(X_s(\omega), Y_s(\cdot)) d\mathbf{W}_s(\omega)$  coincide.

We formulate here the regularity assumptions on  $F(x, \mu)$  used in [1], in addition to **Regularity assumptions 1**, to show the well-posed character of Equation (1.2). Below, we denote by  $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$  a copy of  $(\Omega, \mathcal{F}, \mathbb{P})$ , and given a random variable  $Z$  on  $(\Omega, \mathcal{F}, \mathbb{P})$ , write  $\tilde{Z}$  for its copy on  $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ .

**Regularity assumptions 2.**

- The function  $\partial_x F$  is differentiable in  $(x, \mu)$ .
- For each  $(x, \mu) \in \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d)$ , there exists a version of  $D_\mu F(x, \mu)(\cdot) \in L^2_\mu(\mathbb{R}^d; \mathbb{R}^d \otimes \mathbb{R}^m)$  such that the map  $(x, \mu, z) \mapsto D_\mu F(x, \mu)(z)$  from  $\mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d) \times \mathbb{R}^d$  to  $\mathbb{R}^d \otimes \mathbb{R}^m \otimes \mathbb{R}^d$  is of class  $C^1$ , the derivative in the direction  $\mu$  being understood as before.
- The function  $(x, Z) \mapsto \partial_x^2 F(x, \mathcal{L}(Z))$  from  $\mathbb{R}^d \times \mathbb{L}^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^d)$  to  $\mathbb{R}^d \otimes \mathbb{R}^m \otimes \mathbb{R}^d \otimes \mathbb{R}^d$  is bounded by  $\Lambda$  and  $\Lambda$ -Lipschitz continuous.
- The three derivative functions  $(x, Z) \mapsto \partial_x D_\mu F(x, \mathcal{L}(Z))(Z(\cdot))$ ,  $(x, Z) \mapsto D_\mu \partial_x F(x, \mathcal{L}(Z))(Z(\cdot))$ ,  $(x, Z) \mapsto \partial_z D_\mu F(x, \mathcal{L}(Z))(Z(\cdot))$  are bounded by  $\Lambda$  and  $\Lambda$ -Lipschitz continuous from  $\mathbb{R}^d \times \mathbb{L}^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^d)$  to  $\mathbb{L}^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^d \otimes \mathbb{R}^m \otimes \mathbb{R}^d \otimes \mathbb{R}^d)$ .
- For each  $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ , we denote by  $D_\mu^2 F(x, \mu)(z, \cdot)$ , the derivative of  $D_\mu F(x, \mu)(z)$  with respect to  $\mu$  – which is indeed given by a function. For  $z' \in \mathbb{R}^d$ ,  $D_\mu^2 F(x, \mu)(z, z')$  is an element of  $\mathbb{R}^d \otimes \mathbb{R}^m \otimes \mathbb{R}^d \otimes \mathbb{R}^d$ . We assume that

$$(x, Z) \mapsto D_\mu^2 F(x, \mathcal{L}(Z))(Z(\cdot), \tilde{Z}(\cdot)),$$

from  $\mathbb{R}^d \times \mathbb{L}^2(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^d)$  to  $\mathbb{L}^2(\Omega \times \tilde{\Omega}, \mathcal{F} \otimes \tilde{\mathcal{F}}, \mathbb{P} \otimes \tilde{\mathbb{P}}; \mathbb{R}^d \otimes \mathbb{R}^m \otimes \mathbb{R}^d \otimes \mathbb{R}^d)$ , is bounded by  $\Lambda$  and  $\Lambda$ -Lipschitz continuous.

The two functions  $F(x, \mu) = \int f(x, y)\mu(dy)$  and  $F(x, \mu) = g(x, \int y\mu(dy))$ , for functions  $f, g \in C_b^3$  (meaning that  $f$  and  $g$  are bounded and have bounded derivatives of order 1, 2 and 3), satisfy the **Regularity assumptions 1 and 2**. The following property is taken from [1, Proposition 4.3 and (4.21)].

**Proposition 3.2.** Let  $F$  satisfy **Regularity assumptions 1 and 2** and  $w$  be a control satisfying (2.8) and (2.9). Consider two  $\omega$ -controlled paths  $X(\omega)$  and  $X'(\omega)$  with possibly different initial conditions  $(X_0(\omega), \delta_x X_0(\omega))$  and  $(X'_0(\omega), \delta_x X'_0(\omega))$ , defined on a time interval  $[0, T]$ , together with two random controlled paths  $Y(\cdot)$  and  $Y'(\cdot)$ , with possibly different initial conditions  $(Y_0(\omega), \delta_x Y_0(\omega))$  and  $(Y'_0(\omega), \delta_x Y'_0(\omega))$ , all of them satisfying  $\delta_\mu X(\omega) \equiv \delta_\mu X'(\omega) \equiv 0$  and  $\delta_\mu Y(\cdot) \equiv$

$\delta_\mu Y'(\cdot) \equiv 0$  together with

$$|\delta_x X(\omega)| \vee |\delta_x X'(\omega)| \vee \langle \delta_x Y(\cdot) \rangle_\infty \vee \langle \delta_x Y'(\cdot) \rangle_\infty \leq \Lambda, \quad (3.1)$$

and the size estimates

$$\langle \|Y(\cdot)\|_{[0,T],w,p} \rangle_8^2 \leq L_0, \quad \langle \|Y'(\cdot)\|_{[0,T],w,p} \rangle_8^2 \leq L_0, \quad (3.2)$$

$$\|X(\omega)\|_{[t_i^0, t_{i+1}^0], w, p}^2 \leq L_0, \quad \|X'(\omega)\|_{[t_i^0, t_{i+1}^0], w, p}^2 \leq L_0, \quad (3.3)$$

for  $i \in \{0, \dots, N^0\}$ , for some  $L_0 \geq 1$ , and  $N^0 = N([0, T], \omega, 1/(4L_0))$  given by (2.17), and for the sequence  $(t_i^0 = \tau_i(0, T, \omega, 1/(4L_0)))_{i=0, \dots, N^0+1}$  given by (2.16).

Then, we can find a constant  $\gamma$  depending on  $L_0$  and  $\Lambda$  such that, for any partition  $(t_i)_{i=0, \dots, N}$  included in  $(t_i^0)_{i=0, \dots, N^0}$  and satisfying  $w(t_i, t_{i+1}, \omega)^{1/p} \leq 1/(4L)$  for some  $L \geq L_0$ , we have

$$\begin{aligned} & \left\| \int_{t_i}^{\cdot} \left( F(X_r(\omega), Y_r(\cdot)) - F(X'_r(\omega), Y'_r(\cdot)) \right) d\mathbf{W}_r(\omega) \right\|_{[t_i, t_{i+1}], w, p} \\ & \leq \gamma \left( |\Delta X_0(\omega)| + |\delta_x \Delta X_0(\omega)| \right) + \langle \Delta Y_0(\cdot) \rangle_4 + \langle \delta_x \Delta Y_0(\cdot) \rangle_4 \\ & \quad + \gamma w(0, t_i, \omega)^{1/p} \left( \| \Delta X(\omega) \|_{[0, t_i], w, p} + \langle \| \Delta Y(\cdot) \|_{[0, T], w, p} \rangle_8 \right) \\ & \quad + \frac{\gamma}{4L} \left( \| \Delta X(\omega) \|_{[t_i, t_{i+1}], w, p} + \langle \| \Delta Y(\cdot) \|_{[0, T], w, p} \rangle_8 \right), \end{aligned} \quad (3.4)$$

where  $\Delta X_t(\omega) := X_t(\omega) - X'_t(\omega)$ ,  $\Delta Y_t(\cdot) := Y_t(\cdot) - Y'_t(\cdot)$ ,  $t \in [0, T]$ .

In [1], Proposition 3.2 is used to prove the following existence and uniqueness result, see Theorems 1.1 and 4.4 therein, to which we add the final estimate in the statement.

**Theorem 3.3.** *Let  $F$  satisfy **Regularity assumptions 1 and 2** and  $w$  be a control satisfying (2.8) and (2.9). Assume there exists a positive time horizon  $T$  such that the random variables  $w(0, T, \cdot)$  and  $(N([0, T], \cdot, \alpha))_{\alpha > 0}$  have sub and super exponential tails respectively, in the sense that*

$$\begin{aligned} \mathbb{P}(w(0, T, \cdot) \geq t) & \leq c_1 \exp(-t^{\varepsilon_1}), \\ \mathbb{P}(N([0, T], \cdot, \alpha) \geq t) & \leq c_2(\alpha) \exp(-t^{1+\varepsilon_2(\alpha)}), \end{aligned} \quad (3.5)$$

for some positive constants  $c_1$  and  $\varepsilon_1$ , and possibly  $\alpha$ -dependent positive constants  $c_2(\alpha)$  and  $\varepsilon_2(\alpha)$ . Then, for any  $d$ -dimensional square-integrable random variable  $X_0$ , the mean field rough differential equation (1.2) has a unique solution defined on the whole interval  $[0, T]$ . Moreover, there exist four positive real numbers  $\gamma$ ,  $L_0$ ,  $L$  and  $\eta$  (with  $\gamma, \eta > 1$ ), only depending on  $\Lambda$  and  $T$ , such that, for any subinterval  $[S_1, S_2] \subset [0, T]$  for which

$$\left\langle N([S_1, S_2], \cdot, 1/(4L_0)) \right\rangle_8 \leq 1,$$

and

$$\left\langle \left[ \gamma \left( 1 + w(0, T, \cdot)^{1/p} \right) \right]^{N([S_1, S_2], \cdot, 1/(4L))} \right\rangle_{32} \leq \eta,$$

it holds, for any  $\omega \in \Omega$

$$\|X(\omega)\|_{[S_1, S_2], w, p} \leq \left[ C \left( 1 + w(0, T, \omega)^{1/p} \right) \right]^{2N([0, T], \omega, 1/(4L))},$$

for a constant  $C$  depending only on  $\Lambda$  and  $T$ .

*Proof.* We just address the derivation of the last inequality since the latter is not given in [1, Theorem 4.4]. The key point is to sum over  $n \geq 1$  in [1, (4.30)], replacing  $[0, S]$  therein by  $[S_1, S_2]$ , which is indeed licit provided that  $\langle N([S_1, S_2], \cdot, 1/(4L_0)) \rangle_8 \leq 1$ , see for instance [1, (4.23)], and  $\left\langle \left[ \gamma \left( 1 + w(0, T, \cdot)^{1/p} \right) \right]^{N([S_1, S_2], \cdot, 1/(4L))} \right\rangle_{32} \leq \eta$  for  $\eta$  small enough, see [1, (4.29)].  $\square$

### 3.2. Strong Rough Set-Ups and Continuity of the Itô-Lyons solution Map

Uniqueness in law of the solutions to (1.2) is proven in [1, Theorem 5.3] under the additional assumption that the set-up satisfies the following definition.

**Definition 3.4.** A rough set-up is called strong if there exists a measurable mapping  $\mathcal{I}$  from  $\mathcal{C}([0, T]; \mathbb{R}^m)^2$  into  $\mathcal{C}(\mathcal{S}_2^T; \mathbb{R}^m \otimes \mathbb{R}^m)$  such that

$$\mathbb{P}^{\otimes 2} \left( \left\{ (\omega, \omega') \in \Omega^2 : \mathbb{W}^\perp(\omega, \omega') = \mathcal{I}(W(\omega), W(\omega')) \right\} \right) = 1. \quad (3.6)$$

For our prospect, the following continuity theorem is of crucial interest; see [1, Theorem 5.4].

**Theorem 3.5.** Let  $F$  satisfy the same assumptions as in Theorem 3.3. Given a time interval  $[0, T]$  and a sequence of probability spaces  $(\Omega_n, \mathcal{F}_n, \mathbb{P}_n)$ , indexed by  $n \in \mathbb{N}$ , let, for any  $n$ ,  $X_0^n(\cdot) := (X_0^n(\omega_n))_{\omega_n \in \Omega_n}$  be an  $\mathbb{R}^d$ -valued square-integrable initial condition and

$$\mathbf{W}^n(\cdot) := \left( W^n(\omega_n), \mathbb{W}^n(\omega_n), \mathbb{W}^{n, \perp}(\omega_n, \omega'_n) \right)_{\omega_n, \omega'_n \in \Omega_n}$$

be an  $m$ -dimensional rough set-up with corresponding control  $w^n$ , as given by (2.10), and local accumulated variation  $N^n$ , for fixed values of  $p \in [2, 3]$  and  $q > 8$ . Assume that

- the collection  $(\mathbb{P}_n \circ (|X_0^n(\cdot)|^2)^{-1})_{n \geq 0}$  is uniformly integrable;
- for positive constants  $\varepsilon_1, c_1$  and  $(\varepsilon_2(\alpha), c_2(\alpha))_{\alpha > 0}$ , the tail assumption (3.5) hold for  $w^n$  and  $N^n$ , for all  $n \geq 0$ ;
- associating  $v^n$  with each  $\mathbf{W}^n(\cdot)$  as in (2.6), the functions  $(\mathcal{S}_2^T \ni (s, t) \mapsto \langle v^n(s, t, \cdot) \rangle_{2q})_{n \geq 0}$  are uniformly Lipschitz continuous.

Assume also that

- there exist, on another probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , a square integrable initial condition  $X_0(\cdot)$  with values in  $\mathbb{R}^d$  and a strong rough set-up

$$\mathbf{W}(\cdot) := \left( W(\omega), \mathbb{W}(\omega), \mathbb{W}^\perp(\omega, \omega') \right)_{\omega, \omega' \in \Omega}$$

with values in  $\mathbb{R}^m$ , such that the law under the probability measure  $\mathbb{P}_n^{\otimes 2}$  of the random variable

$$\Omega_n^2 \ni (\omega_n, \omega'_n) \mapsto (X_0^n(\omega_n), W^n(\omega_n), \mathbb{W}_n(\omega_n), \mathbb{W}_n^\perp(\omega_n, \omega'_n)),$$

seen as a random variable with values in the space  $\mathbb{R}^d \times \mathcal{C}([0, T]; \mathbb{R}^m) \times \{\mathcal{C}(\mathcal{S}_2^T; \mathbb{R}^m \otimes \mathbb{R}^m)\}^2$ , converges in the weak sense to the law of

$$\Omega^2 \ni (\omega, \omega') \mapsto (X_0(\omega), W(\omega), \mathbb{W}(\omega, \omega'), \mathbb{W}^\perp(\omega, \omega')).$$

Then,  $\mathbf{W}(\cdot)$  satisfies the requirements of Theorem 3.3 for some  $p' \in (p, 3)$  and  $q' \in [8, q)$ , with control  $w$  therein given by (2.10). Moreover, if  $X^n(\cdot)$ , resp.  $X(\cdot)$ , is the solution of the mean field rough differential equation driven by  $\mathbf{W}^n(\cdot)$ , resp.  $\mathbf{W}(\cdot)$ , then  $X^n(\cdot)$  converges in law to  $X(\cdot)$  on  $\mathcal{C}([0, T]; \mathbb{R}^d)$ .

#### 4. Particle System and Propagation of Chaos

We now have all the ingredients to write down the limiting mean field rough differential equation (1.2) as the limit of a system of particles driven by rough signals (1.3).

##### 4.1. Empirical Rough Set-Up

We recall the framework used to address (1.3). The initial conditions  $(X_0^i(\cdot))_{1 \leq i \leq n}$  are  $\mathbb{R}^d$ -valued variables with the same distribution as  $X_0$  (in the statement of Theorem 3.3) and the enhanced signals  $(W^i(\cdot), \mathbb{W}^i(\cdot))_{1 \leq i \leq n}$  are  $\mathbb{R}^m \oplus \mathbb{R}^m \otimes \mathbb{R}^m$ -valued with the same distribution as  $(W(\cdot), \mathbb{W}(\cdot))$  on the space of continuous functions. Moreover, the variables  $(X_0^i(\cdot), W^i(\cdot), \mathbb{W}^i(\cdot))_{1 \leq i \leq n}$  are independent and identically distributed. All of them are constructed on a single probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . Assuming the rough set-up in Theorem 3.3 to be strong, see Definition 3.4, we let

$$\mathbb{W}^{i,j}(\omega) = \mathcal{I}(W^i(\omega), W^j(\omega)), \quad i \neq j, \quad 1 \leq i, j \leq n.$$

Obviously, equation (1.3) must be understood as a rough differential equation driven by an  $(n \times m)$ -dimensional signal  $(W^1(\omega), \dots, W^n(\omega))$ , and with  $(X^1(\omega), \dots, X^n(\omega))$  as  $(n \times d)$ -dimensional output. Our first task is to prove that (1.3) may be also understood as a mean field rough differential equation on a suitable rough set-up and that the two interpretations coincide. If we require

$\mathbb{P}^{\otimes 2}(\{(\omega, \omega') : \|\mathbb{W}^\perp(\omega, \omega')\|_{[0, T], p/2-\nu} < \infty\}) = 1$  in Definition 3.4, then it is pretty clear that, for almost every  $\omega \in \Omega$ ,

$$\mathbf{W}^{(n)}(\omega) = \left( (W^i(\omega))_{1 \leq i \leq n}, (\mathbb{W}^{i,j}(\omega))_{1 \leq i, j \leq n} \right) =: \left( W^{(n)}(\omega), \mathbb{W}^{(n)}(\omega) \right),$$

is a rough path of finite  $p$ -variation, with the convention that  $\mathbb{W}^{i,i}(\omega) = \mathbb{W}^i(\omega)$ , for  $i \in \{1, \dots, n\}$ . As explained in [1, Proposition 2.4], we may change the definition of  $((W^i(\omega))_{1 \leq i \leq n}, (\mathbb{W}^{i,j}(\omega))_{1 \leq i, j \leq n})$  on a  $\mathbb{P}$ -null set so that  $\mathbf{W}^{(n)}(\omega)$  is in fact a rough path for any  $\omega \in \Omega$ .

As mentioned in Introduction, the striking fact of the analysis was first introduced by Tanaka in [34] and used by Cass and Lyons in their seminal work [12]. The quantity  $\mathbb{W}^{(n)}(\omega)$  may be seen as a rough set-up defined on a finite probability space for any fixed  $\omega \in \Omega$ ; we call it the **empirical rough set-up**. To make it clear, observe that, throughout Section 2, the rough structure is supported by the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  itself. Here,  $\omega$  is fixed, and we see the probability space as

$$\left( \{1, \dots, n\}, \mathcal{P}(\{1, \dots, n\}), \frac{1}{n} \sum_{i=1}^n \delta_i \right), \quad (4.1)$$

where  $\mathcal{P}(\{1, \dots, n\})$  denotes the collection of subsets of  $\{1, \dots, n\}$ . The reader may object that such a probability space is not atomless whilst we explicitly assumed  $(\Omega, \mathcal{F}, \mathbb{P})$  to be atomless in the introduction (see also [1, Section 2]); actually, the reader must realize that, in [1], the atomless property is just used to guarantee that, for any probability measure  $\mu$  on a given Polish space  $S$ , the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  carries an  $S$ -valued random variable with  $\mu$  as distribution. So, it is not a hindrance that  $\{1, \dots, n\}$  is finite.

Hence, in order to draw a parallel with (2.2), the role played by  $\omega \in \Omega$  is here played by  $i \in \{1, \dots, n\}$  and the matrix (2.2) must read

$$\begin{pmatrix} \mathbb{W}_{s,t}^{i,i}(\omega) & \mathbb{W}_{s,t}^{i,\bullet}(\omega) \\ \mathbb{W}_{s,t}^{\bullet,i}(\omega) & \mathbb{W}_{s,t}^{\bullet,\bullet}(\omega) \end{pmatrix}_{0 \leq s \leq t \leq T}, \quad (4.2)$$

where  $\mathbb{W}_{s,t}^{i,\bullet}(\omega)$  is seen as  $\{1, \dots, n\} \ni j \mapsto \mathbb{W}_{s,t}^{i,j}(\omega)$ ,  $\mathbb{W}_{s,t}^{\bullet,i}(\omega)$  as  $\{1, \dots, n\} \ni j \mapsto \mathbb{W}_{s,t}^{j,i}(\omega)$  and  $\mathbb{W}_{s,t}^{\bullet,\bullet}(\omega)$  as  $\{1, \dots, n\} \ni (i, j) \mapsto \mathbb{W}_{s,t}^{i,j}(\omega)$ .

In the same spirit, the variation function  $v$  in (2.6) is

$$\begin{aligned} v^{i,n}(s, t, \omega) &:= \|W^i(\omega)\|_{[s,t], p-\nu}^p + {}^{(n)}(W^\bullet(\omega))_{q; [s,t], p-\nu}^p \\ &\quad + \|\mathbb{W}^i(\omega)\|_{[s,t], p/2-\nu}^{p/2} + {}^{(n)}(\mathbb{W}^{i,\bullet}(\omega))_{q; [s,t], p/2-\nu}^{p/2} \\ &\quad + {}^{(n)}(\mathbb{W}^{\bullet,i}(\omega))_{q; [s,t], p/2-\nu}^{p/2} + {}^{(n)}(\mathbb{W}^{\bullet,\bullet}(\omega))_{q; [s,t], p/2-\nu}^{p/2} \end{aligned} \quad (4.3)$$

where we used the notations

$${}^{(n)}(X^\bullet)_q = \left( \frac{1}{n} \sum_{j=1}^n |X^j|^q \right)^{1/q}, \quad {}^{(n)}(\mathbb{W}^{\bullet,\bullet})_q = \left( \frac{1}{n^2} \sum_{j,k=1}^n |X^{j,k}|^q \right)^{1/q},$$

the corresponding  $p$ -variation being defined as in (2.4) and (2.5). Obviously,  $v^{i,n}(0, T, \omega)$  is almost surely finite. Hence, in order to check that  $\mathbf{W}^{(n)}(\omega)$  defines a rough set-up, it remains to check that it satisfies (2.7). To do so, we strengthen the assumptions on the signal and assume that, for the same parameter  $q$  as in Section 2, it holds

$$\begin{aligned} & \mathbb{E} \left[ \left\| W(\cdot) \right\|_{[0, T], (1/p)\text{-H}}^{pq} + \left\| \mathbb{W}(\cdot) \right\|_{[0, T], (2/p)\text{-H}}^{pq/2} \right] \\ & + \mathbb{E}^{\otimes 2} \left[ \left\| \mathbb{W}^\perp(\cdot, \cdot) \right\|_{[0, T], (2/p)\text{-H}}^{pq/2} \right] < \infty, \end{aligned} \quad (4.4)$$

where

$$\begin{aligned} \left\| W(\omega) \right\|_{[s, t], (1/p)\text{-H}} &= \sup_{\emptyset \neq (s', t') \subset [s, t]} \frac{|W_{t'}(\omega) - W_{s'}(\omega)|}{|t' - s'|^{1/p}} \\ \left\| \mathbb{W}(\omega) \right\|_{[s, t], (2/p)\text{-H}} &= \sup_{\emptyset \neq (s', t') \subset [s, t]} \frac{|\mathbb{W}_{s', t'}(\omega)|}{|t' - s'|^{2/p}}, \end{aligned}$$

and similarly for  $\left\| \mathbb{W}^\perp(\omega, \omega') \right\|_{[s, t], (2/p)\text{-H}}$ , stand for the standard Hölder semi-norms of the rough path. Then, back to (4.3), we can find a universal positive constant  $c$  such that

$$\begin{aligned} v_p^{i,n}(s, t, \omega) &\leq c \left\{ \left\| W^i(\omega) \right\|_{[s, t], (1/p)\text{-H}}^p + \left\| \mathbb{W}^i(\omega) \right\|_{[s, t], (2/p)\text{-H}}^{p/2} \right. \\ &+ \binom{n}{i} \left( \left\| W^{\bullet, i}(\omega) \right\|_{[s, t], (1/p)\text{-H}}^p \right)_q + \binom{n}{i} \left( \left\| \mathbb{W}^{i, \bullet}(\omega) \right\|_{[s, t], (2/p)\text{-H}}^{p/2} \right)_q \\ &\left. + \binom{n}{i} \left( \left\| \mathbb{W}^{\bullet, i}(\omega) \right\|_{[s, t], (2/p)\text{-H}}^{p/2} \right)_q + \binom{n}{i} \left( \left\| \mathbb{W}^{\bullet, \bullet}(\omega) \right\|_{[s, t], (2/p)\text{-H}}^{p/2} \right)_q \right\} (t - s). \end{aligned} \quad (4.5)$$

Taking the mean over  $i \in \{1, \dots, n\}$  and invoking the law of large numbers (see Lemma A.3 in Appendix A.3 for a version of the law of large numbers with second order interactions), we deduce that, for almost every  $\omega \in \Omega$ ,

$$\begin{aligned} & \limsup_{n \geq 1} \sup_{0 \leq s < t \leq T} \frac{\binom{n}{i} \left( v_p^{\bullet, n}(s, t, \omega) \right)_q}{t - s} \\ & \leq c \left( \left\langle \left\| W(\cdot) \right\|_{[0, T], (1/p)\text{-H}}^{pq} + \left\| \mathbb{W}(\cdot) \right\|_{[0, T], (2/p)\text{-H}}^{pq/2} \right\rangle^{1/q} \right. \\ & \quad \left. + \left\langle \left\| \mathbb{W}^\perp(\cdot, \cdot) \right\|_{[0, T], (2/p)\text{-H}}^{pq/2} \right\rangle^{1/q} \right), \end{aligned} \quad (4.6)$$

for a new value of the constant  $c$ . Observe that, in order to derive (4.6), the law of large numbers can be directly applied to each of the first three terms in the right-hand side of (4.5), since each of them can be put in the form  $\mathcal{J}(W^i(\omega))$ , for a suitable form of the functional  $\mathcal{J}$ . Differently, the last three terms in (4.5) require a modicum of care as they lead to empirical means of the form

$$\frac{1}{n^2} \sum_{j, k=1, j \neq k}^n \left\| \mathcal{I}(W^j(\omega), W^k(\omega)) \right\|_{[0, T], (2/p)\text{-H}}^{pq/2} + \frac{1}{n^2} \sum_{j=1}^n \left\| \mathbb{W}^j(\omega) \right\|_{[0, T], (2/p)\text{-H}}^{pq/2}$$

with  $\mathcal{I}$  as in (3.6). Still, if the summands in the two sums are integrable, the limit is  $\langle \|\mathcal{I}(W^1(\cdot), W^2(\cdot))\|_{[0,T],(2/p)\text{-H}}^{pq/2} \rangle$ , see Lemma A.3 in Appendix A.3. Hence (4.6). Now, the fact that the right-hand side of (4.6) is finite guarantees that the 1-variation in the mean in (2.7) is uniformly controlled in  $n \geq 1$ , the mean therein being understood as the mean on the probability space  $(\{1, \dots, n\}, \mathcal{P}(\{1, \dots, n\}), \frac{1}{n} \sum_{i=1}^n \delta_i)$ . Here are two examples under which (4.5) holds true.

**Example 4.1.** Assume that the regularity index  $q$  used in (2.6) satisfies the inequality  $q > 1/(1 - p/3)$ , and that, for some constant  $K_T \geq 0$ ,  $\langle v(s, t, \cdot) \rangle_q \leq K_T(t - s)$  for  $(s, t) \in \mathcal{S}_2^T$ . Then, we get the bounds

$$\begin{aligned} \langle |(W_t - W_s)(\cdot)|^{pq} \rangle &\leq K_T^q |t - s|^q, \\ \langle |W_{s,t}(\cdot)|^{pq/2} \rangle &\leq K_T^q |t - s|^q, \quad \langle \langle |W_{s,t}^\perp(\cdot, \cdot)|^{pq/2} \rangle \rangle \leq K_T^q |t - s|^q. \end{aligned}$$

By Kolmogorov's criterion for rough paths, Theorem 3.1 in [19], we deduce that  $W$  has paths that are  $p' := (1 - 1/q)/p > 1/3$ -Hölder continuous. Similarly,  $\mathbb{W}$  and  $\mathbb{W}^\perp$  have paths that are  $2p' = 2(1 - 1/q)/p > 2/3$ -Hölder continuous and (4.4) holds true with  $p'$  instead of  $p$ . So, the empirical rough set-up satisfies the required conditions provided we replace  $p$  by  $p'$ .

**Example 4.2.** Assume that  $W := (W^1, \dots, W^m)$  is a tuple of independent and centred continuous Gaussian processes, defined on  $[0, T]$ , for which there exists a constant  $K$  such that, for any subinterval  $[s, t] \subset [0, T]$  and any  $k = 1, \dots, m$ , it holds

$$\sup_{i,j} \sum \left| \mathbb{E} \left[ (W_{t_{i+1}}^k - W_{t_i}^k) (W_{s_{j+1}}^k - W_{s_j}^k) \right] \right|^p \leq K |t - s|, \quad (4.7)$$

the sup being over divisions  $(t_i)_i$  and  $(s_j)_j$  of  $[s, t]$ . Then,  $\|W(\cdot)\|_{[0,T],(1/p)\text{-H}}$  has Gaussian tail and  $\|\mathbb{W}(\cdot)\|_{[0,T],(2/p)\text{-H}}$  and  $\|\mathbb{W}^\perp(\cdot, \cdot)\|_{[0,T],(2/p)\text{-H}}$  have exponential tails; see Theorem 11.9 in [19].

Now that we have defined the empirical rough set-up, we must make clear the meaning given to the rough differential equation (1.2) in Definition 3.1 when the rough set-up therein is precisely the empirical rough set-up. We call the corresponding rough differential equation the *empirical rough differential equation*.

For a given  $\omega \in \Omega$ , the probability space that carries the empirical rough set up is given by (4.1). Despite the fact it is not atomless, whilst  $(\Omega, \mathcal{F}, \mathbb{P})$  is, Theorem 3.3 applies and guarantees existence and uniqueness of a solution to the empirical rough differential equation. In this regard, observe that the square integrability requirement on the initial condition here writes  $\frac{1}{n} \sum_{i=1}^n |X_0^i(\omega)|^2 < \infty$ , which is indeed satisfied for  $\omega$  in a full event. The solution reads in the form of a  $n$ -tuple  $X^{(n)}(\omega) = (X^i(\omega))_{1 \leq i \leq n}$  in  $\mathcal{C}([0, T]; \mathbb{R}^d)^n$ . The coefficient driving the equation for  $X^i(\omega)$  reads

$$F \left( X_t^i(\omega), X_t^{\theta_n(\cdot)}(\omega) \right), \quad t \in [0, T],$$

where  $\theta_n(\cdot) : \{1, \dots, n\} \ni i \mapsto i$  is the canonical random variable on  $\{1, \dots, n\}$ . Here the dot in the notation  $X_t^{\theta_n(\cdot)}(\omega)$  refers to the current element in  $\{1, \dots, n\}$ . With this notation, the law of  $X_t^{\theta_n(\cdot)}(\omega)$  (on  $\{1, \dots, n\}$ ) must be understood as the empirical distribution  $\mu_t^n(\omega)$ . Moreover, each  $X^i(\omega)$  is controlled, in standard Gubinelli's sense, by the enhanced rough path  $(W^i(\omega), \mathbb{W}^i(\omega))$  (the remainder in the expansion being controlled by  $v^{i,n}$ ). In particular,  $X^i(\omega)$  may be seen as an  $i$ -controlled path on the empirical rough set-up: If we use  $\delta_x^{(n)}$  and  $\delta_\mu^{(n)}$  as symbols for the Gubinelli derivatives in Definition 2.1 but on the empirical rough set-up, then  $\delta_x^{(n)} X^i(\omega)$  identifies with the standard Gubinelli derivative in the expansion of  $X^i(\omega)$  along the variations of  $(W^i(\omega), \mathbb{W}^i(\omega))$  and  $\delta_\mu^{(n)} X^i(\omega) \equiv 0$ .

The key fact in our analysis lies in the interpretation of the two derivatives

$$\delta_x^{(n)} \left[ F(X^i(\omega), X^{\theta_n(\cdot)}(\omega)) \right] \quad \text{and} \quad \delta_\mu^{(n)} \left[ F(X^i(\omega), X^{\theta_n(\cdot)}(\omega)) \right]$$

in Proposition 2.4. First, it is elementary to check that

$$\begin{aligned} \delta_x^{(n)} \left( F(X^i(\omega), X^{\theta_n(\cdot)}(\omega)) \right)_t &= \partial_x F(X_t^i(\omega), X_t^{\theta_n(\cdot)}(\omega)) \delta_x^{(n)} X_t^i(\omega) \\ &= \partial_x F(X_t^i(\omega), \mu_t^n(\omega)) \delta_x^{(n)} X_t^i(\omega). \end{aligned} \quad (4.8)$$

More interestingly, we have

$$\begin{aligned} \delta_\mu^{(n)} \left( F(X^i(\omega), X^{\theta_n(\cdot)}(\omega)) \right)_t & \\ &= D_\mu F(X_t^i(\omega), \mu_t^n(\omega)) (X_t^{\theta_n(\cdot)}(\omega)) \delta_x^{(n)} X_t^{\theta_n(\cdot)}(\omega), \end{aligned} \quad (4.9)$$

both the left- and the right-hand sides being seen as random variables on  $\{1, \dots, n\}$ . The realizations of the random variable in the right-hand side may be computed by replacing the symbol  $\cdot$  by  $j \in \{1, \dots, n\}$ .

So, applying (2.13) with  $F(X^i(\omega), \mu^n(\omega))$  as integrand, the third term on the first line of (2.13) here reads

$$\frac{1}{n} \sum_{j=1}^n D_\mu F(X_t^j(\omega), \mu_t^n(\omega)) (X_t^j(\omega)) \delta_x^{(n)} X_t^j(\omega) \mathbb{W}_t^{j,i}(\omega).$$

This shows that the integral  $\int_0^t F(X_s^i(\omega), X_s^{\theta_n(\cdot)}(\omega)) d\mathbf{W}_s^{(n)}(\omega)$ , as defined by Theorem 2.3, is the limit of the compensated Riemann sums

$$\begin{aligned} &\sum_{k=0}^{K-1} \left( F(X_{t_k}^i(\omega), X_{t_k}^{\theta_n(\cdot)}(\omega)) W_{t_k, t_{k+1}}^i(\omega) \right. \\ &+ \partial_x F(X_{t_k}^i(\omega), X_{t_k}^{\theta_n(\cdot)}(\omega)) F(X_{t_k}^i(\omega), X_{t_k}^{\theta_n(\cdot)}(\omega)) \mathbb{W}_{t_k, t_{k+1}}^i(\omega) \\ &\left. + \frac{1}{n} \sum_{j=1}^n D_\mu F(X_{t_k}^j(\omega), \mu_{t_k}^n(\omega)) (X_{t_k}^j(\omega)) F(X_{t_k}^j(\omega), X_{t_k}^{\theta_n(\cdot)}(\omega)) \mathbb{W}_{t_k, t_{k+1}}^{j,i}(\omega) \right), \end{aligned} \quad (4.10)$$

as the mesh of the dissection  $0 = t_0 < \dots < t_K = t$  tends to 0<sup>3</sup>. This allows to compare the latter quantity with (1.3) if we interpret the integral with respect to  $W^i(\omega)$  therein as a rough integral with respect to the enhanced setting above  $(W^1(\omega), \dots, W^n(\omega))$ , and consider the leading coefficient  $F(X_t^i(\omega), \mu_t^n(\omega))$  as a *standard* Euclidean function of the tuple  $X_t^{(n)}(\omega) = (X_t^1(\omega), \dots, X_t^n(\omega))$ . Indeed, under the standing **Regularity assumptions 1** and **2**, the function

$$f^i : (\mathbb{R}^d)^n \ni (x^1, \dots, x^n) \mapsto F\left(x^i, \frac{1}{n} \sum_{k=1}^n \delta_{x^k}\right)$$

is  $\mathcal{C}^2$  with Lipschitz derivatives and

$$\partial_{x^j} f^i(x^1, \dots, x^n) = \delta_{i,j} \partial_x F\left(x^i, \frac{1}{n} \sum_{k=1}^n \delta_{x^k}\right) + \frac{1}{n} D_\mu F\left(x^i, \frac{1}{n} \sum_{k=1}^n \delta_{x^k}\right)(x^i),$$

with  $\delta_{i,j} = 1$  if  $i = j$  and 0 otherwise, see Chapter 5 in [9]. Therefore, (1.3) is uniquely solvable in the classical sense and the above formulas for the derivatives show that the rough integral therein may be approximated by the same Riemann sum as in (4.10). Namely, (1.3) may be rewritten as

$$\begin{aligned} & \sum_{k=0}^{K-1} \left( f^i(X_{t_k}^1(\omega), \dots, X_{t_k}^n(\omega)) W_{t_k, t_{k+1}}^i(\omega) \right. \\ & \quad \left. + \sum_{j=1}^n \partial_{x^j} f^i(X_{t_k}^1(\omega), \dots, X_{t_k}^n(\omega)) \mathbb{W}_{t_k, t_{k+1}}^{j,i}(\omega) \right). \end{aligned}$$

This proves that the solution to (1.3), when the latter is seen as a rough differential equation driven by the enhanced setting above  $(W^1(\omega), \dots, W^n(\omega))$ , coincides with the solution of the empirical version of (1.2), when the latter is understood as a mean field rough differential equation driven by the empirical rough set up.

## 4.2. Propagation of Chaos

We now have all the ingredients to prove that the empirical measure of the solution to the particle system (1.3) converges, in some sense, to the solution of the rough mean field equation (1.2), when the rough set-up therein is interpreted as originally explained in Section 2. This is what we call *propagation of chaos*. The statement takes the following form.

**Theorem 4.3.** *We make the following assumptions.*

- (a) *Let F satisfy **Regularity assumptions 1** and **2**.*

<sup>3</sup>In the second line,  $\partial_x F(X_s^i(\omega), X_s^{\theta_n(\cdot)}(\omega)) (F(X_s^i(\omega), X_s^{\theta_n(\cdot)}(\omega)) \mathbb{W}_{s,t}^i(\omega))$  is understood as  $(\sum_{\ell=1}^d \sum_{j,k=1}^m \partial_{x^\ell} F^{\ell,j}(X_s^i(\omega), X_s^{\theta_n(\cdot)}(\omega)) (F^{\ell,k}(X_s^i(\omega), X_s^{\theta_n(\cdot)}(\omega)) (\mathbb{W}_{s,t}^i)^{k,j}(\omega)))_{i=1, \dots, d}$  and similarly for the term on the third line.

- (b) Let  $w$  be a control satisfying (2.8) and (2.9). Assume there exists a positive time horizon  $T$  such that the random variables  $w(0, T, \cdot)$  and  $(N([0, T], \cdot, \alpha))_{\alpha > 0}$ , see (2.17), have sub and super exponential tails, see (3.5).
- (c) Assume that the rough set-up  $\mathbf{W}$  is strong.
- (d) Assume also that there exists a positive constant  $\varepsilon_1$  such that

$$\begin{aligned} & \mathbb{E} \left[ \exp \left( \|W(\cdot)\|_{[0, T], (1/p)\text{-H}}^{\varepsilon_1} \right) \right] + \mathbb{E} \left[ \exp \left( \|\mathbb{W}(\cdot)\|_{[0, T], (2/p)\text{-H}}^{\varepsilon_1/2} \right) \right] \\ & + \mathbb{E}^{\otimes 2} \left[ \exp \left( \|\mathbb{W}^\perp(\cdot, \cdot)\|_{[0, T], (2/p)\text{-H}}^{\varepsilon_1/2} \right) \right] < \infty. \end{aligned} \quad (4.11)$$

Then, for almost every  $\omega \in \Omega$ ,

$$\frac{1}{n} \sum_{i=1}^n \delta_{X^{i, (n)}(\omega)} \rightarrow \mathcal{L}(X(\cdot)), \quad (4.12)$$

where  $X^{(n)}(\omega) = (X^{i, (n)}(\omega))_{i=1, \dots, n}$  is the solution to (1.3) and  $X(\cdot)$  is the solution to (1.2), the convergence being the convergence in law on  $\mathcal{C}([0, T]; \mathbb{R}^d)$ . Moreover, for any fixed  $k \geq 1$ , the law of  $(X^{1, (n)}(\cdot), \dots, X^{k, (n)}(\cdot))$  converges to  $\mathcal{L}(X(\cdot))^{\otimes k}$ .

Following [1, Theorem 2.6], the above assumptions hold true for Gaussian rough paths subject to the classical conditions of Friz-Victoir [21], see Example 4.2, and the related Example 2.5 in [1].

*Proof.* The key tool for passing to the limit is the continuity Theorem 3.5, but with  $p$  therein replaced by some  $p' \in (p, 3)$ . The main difficulty is in controlling the accumulated local variation of the empirical rough set-up. To make the notations clear, we write  $X_0^{i, (n)}$  for  $X^i$ ,  $W^{i, (n)}$  for  $W^i$ ,  $\mathbb{W}^{i, (n)}$  for  $\mathbb{W}^i$  and  $\mathbb{W}^{i, j, (n)}$  for  $\mathbb{W}^{i, j}$ .

**Step 1.** As a starting point, we want to prove that, for almost every  $\omega \in \Omega$ , for any  $\alpha > 0$ , there exists a constant  $\varepsilon_2 > 0$  such that, for all  $n \geq 1$ ,

$$\sup_{n \geq 1} \frac{1}{n} \sum_{i=1}^n \exp \left( N^{i, n}(0, T, \omega, \alpha)^{1+\varepsilon_2} \right) < \infty, \quad (4.13)$$

where  $N^{i, n}(0, T, \omega, \alpha)$  is defined as the local accumulation

$$N^{i, n}([0, T], \omega, \alpha) := N_\varpi([0, T], \alpha), \quad (4.14)$$

when  $\varpi(s, t) = v_{p'}^{i, n}(s, t, \omega)^{1/p'}$ , see (2.17). Following (A.1) in appendix (see also the longer discussion in the introduction of the appendix in [1]), it suffices to prove (4.13) when  $\varpi$  in the definition of  $N^{i, n}$  is equal to each of the terms in the right-hand side of (4.3).

When  $\varpi(s, t) = \|W^i(\omega)\|_{[s, t], p'\text{-v}}$  or  $\varpi(s, t) = \|\mathbb{W}^i(\omega)\|_{[s, t], p'/2\text{-v}}^{1/2}$ , the resulting variables  $(N^{i, n}([0, T], \omega, \alpha))_{i=1, \dots, n}$  in (4.14) are independent and identically

distributed, their common law being independent of  $n$ . Then, (4.13) follows from assumption (b) in the statement and from the law of large numbers.

If  $\varpi(s, t) = {}^{(n)}(W^\bullet(\omega))_{q;[s,t],p'-v}$  or  $\varpi(s, t) = {}^{(n)}(\|W^\bullet(\omega)\|_{q;[s,t],p'/2-v}^{1/2})$ , the resulting variables  $(N^{i,n}([0, T], \omega, \alpha))_{i=1, \dots, n}$  in (4.14) only depend on  $n$ . We may denote them by  $N^n([0, T], \omega, \alpha)$ . Then, it suffices to prove that, for any  $\alpha > 0$ ,  $\limsup_{n \rightarrow \infty} N^n([0, T], \omega, \alpha)$  is almost surely finite. By (4.5), we may easily control  $N^n([0, T], \omega, \alpha)$  from above by noticing that

$$\begin{aligned} & \alpha^p N^n([0, T], \omega, \alpha) \\ & \leq c \left( {}^{(n)}(\|W^\bullet(\omega)\|_{[0,T],(1/p)-H}^p)_q + {}^{(n)}(\|W^\bullet(\omega)\|_{[0,T],(2/p)-H}^{p/2})_q \right), \end{aligned}$$

for a constant  $c$  that is independent of  $n$  and  $\omega$ . Proceeding as in (4.6), the result follows again from the law of large numbers and from assumption (b).

In fact, the most difficult cases are  $\varpi(s, t) = {}^{(n)}(\mathbb{W}^{i,\bullet}(\omega))_{q;[s,t],p'/2-v}^{1/2}$  or  $\varpi(s, t) = {}^{(n)}(\mathbb{W}^{\bullet,i}(\omega))_{q;[s,t],p'/2-v}^{1/2}$ . By symmetry, it suffices to treat the first one. And, by an obvious change of parameter, we may just focus on  $\varpi(s, t) = {}^{(n)}(\mathbb{W}^{i,\bullet}(\omega))_{q;[s,t],p'/2-v}^q$ . Then,

$${}^{(n)}(\mathbb{W}_{s,t}^{i,\bullet}(\omega))_q^q = \frac{1}{n} \sum_{j=1}^n \left( |\mathbb{W}_{s,t}^{i,j}(\omega)|^q - \langle \mathbb{W}_{s,t}^{i,\perp}(\omega, \cdot) \rangle_q^q \right) + \langle \mathbb{W}_{s,t}^{i,\perp}(\omega, \cdot) \rangle_q^q. \quad (4.15)$$

Now, Rosenthal's inequality (see [31]) together with (4.11) say that, for any  $a \geq 2$  and any  $i \in \{1, \dots, n\}$ ,

$$\int_{\Omega} \left| \frac{1}{n} \sum_{j=1}^n \left( |\mathbb{W}_{s,t}^{i,j}(\omega)|^q - \langle \mathbb{W}_{s,t}^{i,\perp}(\omega, \cdot) \rangle_q^q \right) \right|^a d\mathbb{P}(\omega) \leq C_a n^{-a/2} |t-s|^{2aq/p},$$

for a constant  $C_a$  depending on  $a$  and on the upper bound for the left-hand side in (4.11), but independent of  $i$  and  $n$ . Letting  $(t_k^{(n)} = kT/n)_{k=0, \dots, n}$  and allowing the constant  $C_a$  to vary from line to line, we deduce that

$$\sum_{1 \leq k < \ell \leq n} \sum_{i=1}^n \int_{\Omega} \left| \frac{1}{n} \sum_{j=1}^n \left( |\mathbb{W}_{t_k^{(n)}, t_\ell^{(n)}}^{i,j}(\omega)|^q - \langle \mathbb{W}_{t_k^{(n)}, t_\ell^{(n)}}^{i,\perp}(\omega, \cdot) \rangle_q^q \right) \right|^a d\mathbb{P}(\omega) \leq C_a n^{3-a/2}.$$

Choosing  $a$  large enough, we deduce from Markov inequality and from Borel-Cantelli lemma that, for  $\mathbb{P}$ -almost every  $\omega \in \Omega$ , for  $n$  large enough,

$$\begin{aligned} & \forall i \in \{1, \dots, n\}, \quad \forall 1 \leq k < \ell \leq n, \\ & \left| \frac{1}{n} \sum_{j=1}^n \left( |\mathbb{W}_{t_k^{(n)}, t_\ell^{(n)}}^{i,j}(\omega)|^q - \langle \mathbb{W}_{t_k^{(n)}, t_\ell^{(n)}}^{i,\perp}(\omega, \cdot) \rangle_q^q \right) \right| \leq n^{-1/4}. \end{aligned} \quad (4.16)$$

By assumption (d) in the statement, we recall that, for any  $\delta > 0$ ,

$$\sum_{n \geq 1} \sum_{1 \leq i, j \leq n} \mathbb{P} \left( \|W^i\|_{[0,T],(1/p)-H}^2 + \|\mathbb{W}^{i,j}\|_{[0,T],(2/p)-H} \geq n^\delta \right) < \infty.$$

Therefore, for any  $\delta > 0$ , for  $\mathbb{P}$ -almost every  $\omega \in \Omega$ , for  $n$  large enough,

$$\sup_{1 \leq i, j \leq n} \left( \|W^i(\omega)\|_{[0, T], (1/p)\text{-H}}^2 + \|W^{i, j}(\omega)\|_{[0, T], (2/p)\text{-H}} \right) \leq n^\delta, \quad (4.17)$$

which implies

$$\begin{aligned} \sup_{1 \leq i, j \leq n} \sup_{|s-t| \leq 1/n} |\mathbb{W}_{s, t}^{i, j}(\omega)| &\leq n^{\delta-2/p}, \\ \sup_{1 \leq i, j \leq n} \sup_{\min(|s-t|, |s'-t'|) \leq 1/n} \left( |W_{s, t}^i(\omega)| |W_{s', t'}^j(\omega)| \right) &\leq n^{\delta-1/p}. \end{aligned} \quad (4.18)$$

Similarly, we have for  $\mathbb{P}$ -almost every  $\omega \in \Omega$ , for  $n$  large enough,

$$\sup_{1 \leq i \leq n} \|\langle W^{i, \perp}(\omega, \cdot) \rangle_q\|_{[0, T], (2/p)\text{-H}} \leq n^\delta, \quad (4.19)$$

which implies

$$\begin{aligned} \sup_{1 \leq i \leq n} \sup_{|s-t| \leq 1/n} \langle W_{s, t}^{i, \perp}(\omega, \cdot) \rangle_q &\leq n^{\delta-2/p}, \\ \sup_{1 \leq i \leq n} \sup_{\min(|s-t|, |s'-t'|) \leq 1/n} \left( |W_{s, t}^i(\omega)| \langle W_{s', t'}(\cdot) \rangle_q \right) &\leq n^{\delta-1/p}. \end{aligned} \quad (4.20)$$

Choosing  $\delta < 1/(2p)$ , combining (4.16) with (4.18) and (4.20), and using Chen's relations (2.3) to write  $\mathbb{W}_{s, t}^{i, j} := -\mathbb{W}_{\{s\}, s}^{i, j} + \mathbb{W}_{\{s\}, \{t\}}^{i, j} + \mathbb{W}_{\{t\}, t}^{i, j} + W_{s, \{t\}}^i \otimes W_{\{t\}, t}^j - W_{\{s\}, s}^i \otimes W_{s, \{t\}}^j$  (with  $\{s\} := \lfloor ns/T \rfloor T/n$ ), and similarly for  $\langle W^{i, \perp}(\omega, \cdot) \rangle_q$ , we can find a constant  $c_q$  only depending on  $q$  such that, for almost every  $\omega \in \Omega$ , for  $n$  large enough,

$$\sup_{i=1, \dots, n} \sup_{(s, t) \in \mathcal{S}_2^T} \left| \frac{1}{n} \sum_{j=1}^n \left( |\mathbb{W}_{s, t}^{i, j}(\omega)|^q - \langle W_{s, t}^{i, \perp}(\omega, \cdot) \rangle_q^q \right) \right| \leq c_q n^{-[1/4 \wedge q/(2p)]}. \quad (4.21)$$

Meanwhile, we also have, for  $\mathbb{P}$ -almost every  $\omega \in \Omega$ ,

$$\begin{aligned} &\left| \frac{1}{n} \sum_{j=1}^n \left( |\mathbb{W}_{s, t}^{i, j}(\omega)|^q - \langle W_{s, t}^{i, \perp}(\omega, \cdot) \rangle_q^q \right) \right| \\ &\leq \sup_{1 \leq i, j \leq n} \left( \|W^{i, j}(\omega)\|_{[0, T], (2/p)\text{-H}}^q + \|\langle W^{i, \perp}(\omega, \cdot) \rangle_q\|_{[0, T], (2/p)\text{-H}}^q \right) (t-s)^{2q/p}. \end{aligned}$$

By (4.17) and (4.19), we deduce that, for  $\mathbb{P}$ -almost every  $\omega \in \Omega$ , for  $n$  large enough, for all  $(s, t) \in \mathcal{S}_2^T$ ,

$$\left| \frac{1}{n} \sum_{j=1}^n \left( |\mathbb{W}_{s, t}^{i, j}(\omega)|^q - \langle W_{s, t}^{i, \perp}(\omega, \cdot) \rangle_q^q \right) \right| \leq n^{\delta q} (t-s)^{2q/p}. \quad (4.22)$$

Taking the power  $1-p/p'$  in (4.21) and the power  $p/p'$  in (4.22), cross-multiplying both of them and then choosing  $\delta$  small enough, we get, for  $\mathbb{P}$ -almost every  $\omega \in \Omega$ , for  $n$  large enough, for all  $(s, t) \in \mathcal{S}_2^T$ ,

$$\left| \frac{1}{n} \sum_{j=1}^n \left( |\mathbb{W}_{s, t}^{i, j}(\omega)|^q - \langle W_{s, t}^{i, \perp}(\omega, \cdot) \rangle_q^q \right) \right| \leq c (t-s)^{2q/p'},$$

where  $c$  only depends on  $p$ ,  $p'$  and  $q$ . Back to (4.15), we deduce that, for  $\mathbb{P}$ -almost every  $\omega \in \Omega$ , for  $n$  large enough,

$${}^{(n)}\left(\mathbb{W}_{s,t}^{i,\bullet}(\omega)\right)_{q;[s,t],p'/2-\nu} \leq c^{1/q}(t-s)^{2/p'} + \langle \mathbb{W}^{i,\perp}(\omega, \cdot) \rangle_{q;[s,t],p/2-\nu}.$$

Since the variables  $(\mathbb{W}^{i,\perp})_{i \geq 1}$  are independent, the local accumulation associated with the second term in the right-hand side may be handled like the local accumulation associated to  $\varpi(s, t) = \|W^i(\omega)\|_{[s,t],p-\nu}$ . The local accumulation associated with the first term is easily handled.

**Step 2.** Now, from the law of large numbers (see Lemma A.3 for the law of large numbers with second order interaction terms) and from [4, Theorem 2.3, Problem 3.1], we deduce that there exists a full subset  $E \subset \Omega$  (the definition of which may vary from line to line in the rest of the proof as long as  $\mathbb{P}(E)$  remains equal to 1) such that, for any  $\omega \in E$ ,

$$\pi_n(\omega) = \left( \frac{1}{n^2} \sum_{i,j=1}^n \delta_{(X_0^{i,(n)}(\omega), W^{i,(n)}(\omega), \mathbb{W}^{i,(n)}(\omega), \mathbb{W}^{i,j,(n)}(\omega))} \right)_{n \geq 1}$$

converges in the weak sense to  $(X_0(\cdot), W(\cdot), \mathbb{W}(\cdot), \mathbb{W}^\perp(\cdot, \cdot))$  on the space  $\mathbb{R}^d \times \mathcal{C}([0, T]; \mathbb{R}^m) \times \{\mathcal{C}(\mathcal{S}_2^T; \mathbb{R}^m \otimes \mathbb{R}^m)\}^2$ .

**Step 3.** Back to the statement of Theorem 3.5, the first item in the statement is a consequence of the law of large numbers. As for the fourth item, it follows directly from the previous step. In order to check the second and third items, we now have a look at  $v_{p'}^{i,n}(s, t, \omega)$  in (4.3). Following (4.6), we already know that

$$\limsup_{n \geq 1} \sup_{0 \leq s < t \leq T} \frac{{}^{(n)}(v_{p'}^{\bullet,n}(s, t, \omega))_{2q}}{t-s} < \infty,$$

which proves the third item in the statement of Theorem 3.5. We end up with the proof of the second item. Following (4.5), there exists a constant  $c'$  such that, for any  $\varepsilon > 0$ , the quantity

$$\sup_{n \geq 1} {}^{(n)} \left( \exp \left( [v_{p'}^{\bullet,n}(0, T, \omega)]^\varepsilon \right) \right)_1 \quad (4.23)$$

is finite if

$$\begin{aligned} \sup_{n \geq 1} \frac{1}{n} \sum_{i=1}^n \exp \left( c' \|W^i(\omega)\|_{[0,T],(1/p)-H}^{p'\varepsilon} + c' \|\mathbb{W}^i(\omega)\|_{[0,T],(2/p)-H}^{p'\varepsilon/2} \right) &< \infty, \\ \sup_{n \geq 1} \frac{1}{n} \sum_{i=1}^n \exp \left( c' {}^{(n)} \left( \|\mathbb{W}^{\bullet,i}(\omega)\|_{[0,T],(2/p)-H}^{p'/2} \right)_q^\varepsilon \right) &< \infty, \end{aligned}$$

and similarly on the second line with  $\mathbb{W}^{\bullet,i}(\omega)$  replaced by  $\mathbb{W}^{i,\bullet}(\omega)$ . By the law of large numbers, the first line holds true on a full event if  $p'\varepsilon < \varepsilon_1$ . As for the second one, we use the following trick. Notice that the function

$$(0, +\infty) \ni x \mapsto \exp(x^{\varepsilon/q}), \quad (4.24)$$

is convex on  $[A_\varepsilon, \infty)$ , for some  $A_\varepsilon > 0$ . Therefore, Jensen's inequality says that, in order to check the third line, it suffices to prove that

$$\sup_{n \geq 1} \frac{1}{n^2} \sum_{i,j=1}^n \exp\left[\left(A_\varepsilon^{\varepsilon/q} \vee \|\mathbb{W}^{i,j}(\omega)\|_{[0,T],(2/p)\text{-H}}^{p'\varepsilon/2}\right)\right] < \infty, \quad (4.25)$$

and similarly for the last line. Obviously, under the standing assumption, the latter holds true with probability 1 provided  $p'\varepsilon < \varepsilon_1$ . This proves (4.23). In the statement of Theorem 3.5, this proves the condition related to the tails of  $w^n$  by a standard application of Markov inequality.

The bound on the local accumulation in the second item of Theorem 3.5 follows from the first step of the proof.

**Step 4.** By Theorem 3.5, we get (4.12) on a set of full measure. By Proposition 2.2 in [32], we deduce that, for any fixed  $k \geq 1$ , the law of  $(X^{1,(n)}, \dots, X^{k,(n)})$  converges to  $\mathcal{L}(X(\cdot))^{\otimes k}$ .  $\square$

**Remark 4.4.** Recently, the authors in [13] obtained a quantified propagation of chaos result for mean field stochastic equations with additive noise

$$dx_t = b(x_t, \mathcal{L}(x_t))dt + dw_t, \quad x_0 = \zeta, \quad (4.26)$$

for a random path  $w \in \mathcal{C}([0, T], \mathbb{R}^d)$  subject to mild integrability condition, and random initial condition  $\zeta$ . There is no need of rough paths theory to make sense of this equation and solve it by elementary means, under proper regularity assumptions on the drift  $b$ . Its distribution is even a Lipschitz function of the distribution of  $(\zeta, w)$ , in  $p$ -Wasserstein metric. Using Tanaka's trick, this continuity result entails a propagation of chaos result. The global Lipschitz continuity of the solution map  $\mathcal{L}(w, \zeta) \mapsto \mathcal{L}(x)$  ensures in particular a quantitative convergence rate for the particle system no greater than the corresponding convergence rate for the sample empirical mean of the driving noises, which is optimal. We get back such a sharp estimate in the present, much more complicated, setting in the next section. Note that the global Lipschitz bound satisfied by the natural map  $\Phi$  giving the solution to equation (4.26) as a fixed point of  $\Phi$  actually allows to deal with reflected dynamics, as the bounded variation part needed for the reflection happens to be a Lipschitz function of the non-reflected path, in Skorokhod formulation of the problem. We do not have such a strong continuity result for our solution map; see Theorem 3.5. See also the previous work [17] of the authors.

## 5. Rate of Convergence

The goal of this section is to elucidate the rate of convergence in the convergence result stated in Theorem 4.3.

The analysis is based upon a variation of Sznitman's original coupling argument, see [32]. To make its principle clear, we recall that, on the space  $(\Omega, \mathcal{F}, \mathbb{P})$ , the triples  $(X_0^1(\cdot), W^1(\cdot), \mathbb{W}^1(\cdot)), \dots, (X_0^n(\cdot), W^n(\cdot), \mathbb{W}^n(\cdot))$  are  $n$  independent copies of the original triple  $(X_0(\cdot), W(\cdot), \mathbb{W}(\cdot))$ . For each  $i \in \{1, \dots, n\}$ , the pair  $(W^i(\cdot), \mathbb{W}^i(\cdot))$  is completed into a rough set-up

$$\begin{aligned} \overline{\mathbf{W}}^i(\cdot) &:= (W^i(\cdot), \mathbb{W}^i(\cdot), \mathbb{W}^{i,\perp}(\cdot, \cdot)), \\ \mathbb{W}^{i,\perp}(\omega, \omega') &= \mathcal{I}(W^i(\omega), W^i(\omega')), \quad (\omega, \omega') \in \Omega^2. \end{aligned} \quad (5.1)$$

Here we put a bar on the symbol  $\overline{\mathbf{W}}^i$  in order to distinguish it from the finite-dimensional rough set-up  $\mathbf{W}^{(n)}(\omega)$  that lies above  $(W^1(\omega), \dots, W^n(\omega))$ . In comparison, the second-order level of  $\mathbf{W}^{(n)}$  is made of  $(\mathbb{W}^i)_{1 \leq i \leq n}$  and of  $(\mathbb{W}^{i,j} = \mathcal{I}(W^i, W^j))_{1 \leq i \neq j \leq n}$ , see (4.2). To make the notations more homogeneous, we sometimes write  $\mathbb{W}^{i,i}(\omega)$  for  $\mathbb{W}^i(\omega)$ .

With each  $(X_0^i(\cdot), \overline{\mathbf{W}}^i(\cdot))$ , we associate the corresponding solution  $\overline{X}^i(\cdot)$  to the mean field equation (1.2). The 5-tuples

$$\Omega \ni \omega \mapsto \left( X_0^i(\omega), W^i(\omega), \mathbb{W}^i(\omega), \mathbb{W}^{i,\perp}(\cdot, \omega), \overline{X}^i(\omega) \right)_{1 \leq i \leq n}$$

are independent and identically distributed,  $\Omega \ni \omega \mapsto (\mathbb{W}_t^{i,\perp}(\cdot, \omega))_{0 \leq t \leq T}$  being regarded as a process with values in  $\mathbb{L}^q(\Omega, \mathcal{F}, \mathbb{P}; \mathbb{R}^d)$ . Recalling that  $X^{(n)}(\omega) = (X^{1,(n)}(\cdot), \dots, X^{n,(n)}(\cdot))$  is the solution to (1.3), we then let

$$\mu_t^n(\omega) = \frac{1}{n} \sum_{i=1}^n \delta_{X_t^{i,(n)}(\omega)}, \quad \overline{\mu}_t^n(\omega) = \frac{1}{n} \sum_{i=1}^n \delta_{\overline{X}_t^i(\omega)}, \quad t \in [0, T], \quad \omega \in \Omega. \quad (5.2)$$

Note the use of the  $\mathbf{d}_1$ -distance (see (2.14)) in the assumption required from  $F$  in the statement below,  $\mathbf{d}_1$ -continuity being stronger than  $\mathbf{d}_2$ -continuity.

**Theorem 5.1.** *We make the following assumptions.*

- (a) *Assumptions (a)–(d) in the statement of Theorem 4.3 are satisfied.*
- (b) *The first and second derivatives of  $F$ ,  $(x, \mu) \mapsto \partial_x F(x, \mu)$ ,  $(x, \mu, z) \mapsto (D_\mu F(x, \mu)(z), \partial_x D_\mu F(x, \mu, z))$ , and  $(x, \mu, z, z') \mapsto D_\mu^2 F(x, \mu, z, z')$ , are bounded on the whole space and are Lipschitz continuous with respect to all the variables, the Lipschitz property in the direction  $\mu$  being understood with respect to  $\mathbf{d}_1$ .*
- (c) *Last, for any  $\alpha > 0$ , there exists a constant  $\varepsilon_2 > 0$  such that, for any  $n \geq 1$ , for any  $p' \in (p, 3)$ , and any random variables  $\tau, \tau' : \Omega \rightarrow [0, T]$ , with  $\mathbb{P}(\tau < \tau') = 1$ , we have*

$$\sup_{n \geq 1} \sup_{1 \leq i \leq n} \mathbb{E} \left[ \exp \left[ \left( \frac{\widehat{N}^{i,n}([\tau, \tau'], \omega, \alpha)}{\sqrt{\tau' - \tau}} \right)^{1+\varepsilon_2} \right] \right] < \infty,$$

where  $\widehat{N}^{i,n}([\tau, \tau'], \omega, \alpha)$  is defined as the accumulation  $N_{\varpi}([\tau, \tau'], \alpha)$  when  $\varpi = (\widehat{w}_{p'}^{i,n}(\omega))^{1/p'}$  with

$$\begin{aligned}\widehat{w}_{p'}^{i,n}(s, t, \omega) &:= (w_{p'}^{i,n} + \widehat{v}_{p'}^{i,n})(s, t, \omega) + {}^{(n)}(\widehat{v}_{p'}^{\bullet,n}(\omega))_{q;[s,t],1-\nu} + (t - s), \\ w_{p'}^{i,n}(s, t, \omega) &:= v_{p'}^{i,n}(s, t, \omega) + {}^{(n)}(v_{p'}^{\bullet,n}(\omega))_{q;[s,t],1-\nu}, \\ \widehat{v}_{p'}^{i,n}(s, t, \omega) &:= \langle \mathbb{W}^{i,\perp}(\omega, \cdot) \rangle_{q;[s,t],p'/2-\nu}^{p'/2} + \langle \mathbb{W}^{i,\perp}(\cdot, \omega) \rangle_{q;[s,t],p'/2-\nu}^{p'/2}.\end{aligned}\tag{5.3}$$

Then, for any  $r \geq 1$ , there exists an exponent  $q(r) \geq 8$  such that, if  $q \geq q(r)$ , with  $q$  as in Section 2, and  $X_0(\cdot)$  is in  $\mathbb{L}^{q(r)}$ , then

$$\sup_{1 \leq i \leq n} \mathbb{E} \left[ \sup_{0 \leq t \leq T} |\overline{X}_t^i - X_t^{i,(n)}|^r \right]^{1/r} + \mathbb{E} \left[ \sup_{0 \leq t \leq T} \mathbf{d}_1(\mu_t^n, \overline{\mu}_t^n)^r \right]^{1/r} \leq C \varsigma_n, \tag{5.4}$$

for a constant  $C$  independent of  $n$ , and  $\varsigma_n = n^{-1/2}$  if  $d = 1$ ,  $\varsigma_n = n^{-1/2} \ln(1+n)$  if  $d = 2$  and  $\varsigma_n = n^{-1/d}$  if  $d \geq 3$ .

**Remark 5.2.** Let us make a few remarks on this statement before embarking on its proof.

- The convergence rate  $\varsigma_n$  in (5.4) corresponds to the usual rate for the convergence in the 1-Wasserstein distance of an empirical sample of independent, identically distributed, random variables toward the limiting common distribution; see [18] together with Lemma A.2.
- Theorem 5.1 applies when  $W$  is a continuous centred Gaussian process defined over some finite interval  $[0, T]$ , with independent components, and with a covariance function that is of finite two dimensional  $\rho$ -variation, for some  $\rho \in [1, 3/2)$ , see (4.7). The proof is given in Appendix A.1.
- We refer to [9, Chapter 5] for examples of a function  $F$  satisfying item (a) in the assumptions of the statement. Importantly, we recall that a function  $G : \mathcal{P}_2(\mathbb{R}^d) \ni \mu \mapsto G(\mu) \in \mathbb{R}$ , whose derivative  $D_\mu G : \mathcal{P}_2(\mathbb{R}^d) \times \mathbb{R}^d \ni (\mu, z) \mapsto D_\mu G(\mu)(z) \in \mathbb{R}^d$  is uniformly bounded on the whole  $\mathcal{P}_2(\mathbb{R}^d) \times \mathbb{R}^d$ , is Lipschitz continuous with respect to the  $\mathbf{d}_1$ -Wasserstein distance. In particular, under the assumptions of the statement,  $F$  itself is Lipschitz continuous on  $\mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d)$ , the Lipschitz property in the direction  $\mu$  being understood with respect to  $\mathbf{d}_1$ .
- By inspecting the proof of Theorem 5.1, we could make explicit the value of  $q(r)$ , but we feel that it would not be so useful.

*Proof.* Observe that, for each  $i \in \{1, \dots, n\}$  and any  $\omega \in \Omega$ , we can define the integral process  $(\int_0^t F(\overline{X}_s^i(\omega), \overline{\mu}_s^n(\omega)) d\mathbf{W}_s^{i,(n)}(\omega))_{0 \leq t \leq T}$  using usual rough paths theory, where the label  $i$  in the notation  $\mathbf{W}^{i,(n)}(\omega)$  is here to indicate that the integral only involves  $(W^i(\omega), (\mathbb{W}^{j,i}(\omega))_{1 \leq j \leq n})$ . Equivalently,  $\mathbf{W}^{i,(n)}(\omega)$  must be seen as  $(W^i(\omega), (\mathbb{W}^{j,i}(\omega))_{1 \leq j \leq n})$ . The fact that the integral may be defined with respect to  $(W^i(\omega), (\mathbb{W}^{j,i}(\omega))_{1 \leq j \leq n})$  follows from the fact that  $\overline{X}^j(\omega)$ , for

each  $j \in \{1, \dots, n\}$  and each  $\omega \in \Omega$ , is controlled by the variations of the sole  $W^j(\omega)$ .

**Step 1.** The first step is to compare

$$\int_0^t \mathbb{F}(\overline{X}_s^i(\omega), \mathcal{L}(X_s)) d\overline{\mathbf{W}}_s^i(\omega) \quad \text{and} \quad \int_0^t \mathbb{F}(\overline{X}_s^i(\omega), \overline{\mu}_s^n(\omega)) d\mathbf{W}_s^{i,(n)}(\omega), \quad (5.5)$$

for  $t \in [0, T]$ . What makes the proof non-trivial is the fact that the rough sets-ups used in the first and the second integrals are not the same. So, in order to compare the two of them, we need to come back to the original constructions of the two integrals. To simplify notations, and for  $0 \leq t \leq T$ , set

$$\overline{F}_t^i(\omega) := \mathbb{F}(\overline{X}_t^i(\omega), \mathcal{L}(X_t)), \quad F_t^{i,n}(\omega) := \mathbb{F}(\overline{X}_t^i(\omega), \overline{\mu}_t^n(\omega)). \quad (5.6)$$

For sure,  $(\overline{F}_t^i(\omega))_{0 \leq t \leq T}$  is  $\omega$ -controlled by  $\overline{\mathbf{W}}^i(\omega)$ , see Definition 2.1, and the collection indexed by  $\omega \in \Omega$  is a random path controlled by  $\overline{\mathbf{W}}^i$ , see Definition 2.2 for a reminder. The corresponding Gubinelli derivatives are denoted by  $(\delta_x \overline{F}_t^i(\omega), \delta_\mu \overline{F}_t^i(\omega, \cdot))_{0 \leq t \leq T}$ , see Proposition 2.4. Similarly,  $(F_t^{i,n}(\omega))_{0 \leq t \leq T}$  is controlled by  $\mathbf{W}^{i,(n)}(\omega)$  and  $\mathbf{W}^{\bullet,(n)}(\omega)$  and Gubinelli derivatives are encoded in the form of a collection  $(\delta_x F_t^{i,n}(\omega), (\delta_\mu F_t^{i,j,n}(\omega))_{1 \leq j \leq n})_{0 \leq t \leq T}$ , see (4.8–4.9). To make it clear, set

$$\begin{aligned} \delta_x \overline{F}_t^i(\omega) &:= \partial_x \mathbb{F}(\overline{X}_t^i(\omega), \mathcal{L}(X_t)) \mathbb{F}(\overline{X}_t^i(\omega), \mathcal{L}(X_t)), \\ \delta_\mu \overline{F}_t^i(\omega, \cdot) &:= D_\mu \mathbb{F}(\overline{X}_t^i(\omega), \mathcal{L}(X_t)) (X_t(\cdot)) \mathbb{F}(X_t(\cdot), \mathcal{L}(X_t)), \end{aligned} \quad (5.7)$$

where  $X(\cdot)$  is the solution to (1.2) with  $\mathbf{W}(\cdot) = (W(\cdot), \mathbb{W}(\cdot), \mathbb{W}^\perp(\cdot, \cdot))$ . We also let

$$\begin{aligned} \delta_x F_t^{i,n}(\omega) &:= \partial_x \mathbb{F}(\overline{X}_t^i(\omega), \overline{\mu}_t^n(\omega)) \mathbb{F}(\overline{X}_t^i(\omega), \overline{\mu}_t^n(\omega)), \\ \delta_\mu F_t^{i,j,n}(\omega) &:= D_\mu \mathbb{F}(\overline{X}_t^i(\omega), \overline{\mu}_t^n(\omega)) (\overline{X}_t^j(\omega)) \mathbb{F}(\overline{X}_t^i(\omega), \overline{\mu}_t^n(\omega)). \end{aligned} \quad (5.8)$$

For a subdivision  $\Delta = \{s = t_0 < t_1 < \dots < t_K = t\}$ , set

$$\begin{aligned} \overline{\mathcal{I}}_{s,t}^{i,\Delta}(\omega) &:= \sum_{k=0}^{K-1} \left\{ \overline{F}_{t_k}^i(\omega) W_{t_k, t_{k+1}}^i(\omega) + \delta_x \overline{F}_{t_k}^i(\omega) \mathbb{W}_{t_k, t_{k+1}}^i(\omega) \right. \\ &\quad \left. + \mathbb{E}[\delta_\mu \overline{F}_{t_k}^i(\omega, \cdot) \mathbb{W}_{t_k, t_{k+1}}^{i,\perp}(\cdot, \omega)] \right\}, \\ \mathcal{I}_{s,t}^{i,n,\Delta}(\omega) &:= \sum_{k=0}^{K-1} \left\{ F_{t_k}^{i,n}(\omega) W_{t_k, t_{k+1}}^i(\omega) + \delta_x F_{t_k}^{i,n}(\omega) \mathbb{W}_{t_k, t_{k+1}}^i(\omega) \right. \\ &\quad \left. + \frac{1}{n} \sum_{j=1}^n \delta_\mu F_{t_k}^{i,j,n}(\omega) \mathbb{W}_{t_k, t_{k+1}}^{j,i}(\omega) \right\}. \end{aligned} \quad (5.9)$$

The two integrals in (5.5) should be understood as the respective limits of the two Riemann sums right above as  $K$  tends to  $\infty$ . In the sequel, we denote the summand in the first sum by  $\overline{\mathcal{I}}_{\{t_k, t_{k+1}\}}^{i, \partial}(\omega)$  and the summand in the second sum by  $\mathcal{I}_{\{t_k, t_{k+1}\}}^{i, n, \partial}(\omega)$ . By Lemma A.1 proved in Appendix A.2, we can find, for any  $\varrho \geq 8$ , a constant  $C$  and an exponent  $\varrho' \geq \varrho$  independent of  $n$  and  $K$  such that, when  $X_0(\cdot) \in \mathbb{L}^{\varrho'}$ , it holds for any  $k \in \{1, \dots, K-1\}$  (provided  $K \geq 2$ ),

$$\left\langle \left\{ \mathcal{I}_{s,t}^{i, n, \Delta}(\cdot) - \mathcal{I}_{s,t}^{i, n, \Delta'}(\cdot) \right\} - \left\{ \overline{\mathcal{I}}_{s,t}^{i, \Delta}(\cdot) - \overline{\mathcal{I}}_{s,t}^{i, \Delta'}(\cdot) \right\} \right\rangle_{\varrho} \leq C \varsigma_n \langle w^+(t_{k-1}, t_{k+1}, \cdot, \cdot) \rangle_{\varrho'}^{3/p},$$

where  $\Delta' := \Delta \setminus \{t_k\}$  and  $w^+(s, t, \omega, \omega') := w(s, t, \omega) + \|\mathbb{W}^\perp(\omega, \omega')\|_{[s,t], p/2-\nu}^{p/2}$ .

Formulating (4.5) and (4.6) but for the limit (instead of empirical) rough set-up, we know that the right hand side in the above inequality is less than

$$C \varsigma_n \left[ \langle \|W(\cdot)\|_{[0,T], (1/p)-\mathbb{H}} \rangle_{p\varrho'} + \langle \|\mathbb{W}(\cdot)\|_{[0,T], (2/p)-\mathbb{H}} \rangle_{p\varrho'}^{1/2} + \langle \|\mathbb{W}^\perp(\cdot, \cdot)\|_{[0,T], (2/p)-\mathbb{H}} \rangle_{p\varrho'}^{1/2} \right]^3 (t_{k+1} - t_k)^{3/p},$$

but by assumption all the expectations are finite. Now we can choose  $t_k$  such that  $|t_{k+1} - t_{k-1}| \leq 3|t-s|/K$  (if not, it means that  $3(t-s)(K-1)/K < \sum_{k=1}^{K-1} |t_{k+1} - t_{k-1}| = \sum_{k=1}^{K-1} (t_{k+1} - t_k + t_k - t_{k-1}) = 2(t-s) - (t_K - t_{K-1} + t_1 - t_0) \leq t-s$ , which is a contradiction). We get

$$\left\langle \left\{ \mathcal{I}_{s,t}^{i, n, \Delta}(\cdot) - \mathcal{I}_{s,t}^{i, n, \Delta'}(\cdot) \right\} - \left\{ \overline{\mathcal{I}}_{s,t}^{i, \Delta}(\cdot) - \overline{\mathcal{I}}_{s,t}^{i, \Delta'}(\cdot) \right\} \right\rangle_{\varrho} \leq C \varsigma_n \left( \frac{t-s}{K} \right)^{3/p},$$

the constant  $C$  being allowed to increase from line to line as long as it remains independent of  $n$  and  $K$ . Letting  $t^{(1)} = t_k$  and applying iteratively the above bound to a sequence of meshes of the form  $\Delta \setminus \{t^{(1)}\}$ ,  $\Delta \setminus \{t^{(1)}, t^{(2)}\}$ ,  $\dots$ , and then letting  $K$  tend to  $\infty$ , we deduce that

$$\left\langle \int_s^t F_r^{i, n}(\cdot) d\mathbb{W}_r^{i, (n)}(\cdot) - \int_s^t \overline{F}_r^i(\cdot) d\overline{\mathbb{W}}_r^i(\cdot) - \left\{ \mathcal{I}_{\{s,t\}}^{i, n, \partial} - \overline{\mathcal{I}}_{\{s,t\}}^{i, \partial} \right\} \right\rangle_{\varrho} \leq C \varsigma_n (t-s)^{3/p}. \quad (5.10)$$

By Lemma A.1, we also have  $\langle \mathcal{I}_{\{s,t\}}^{i, n, \partial} - \overline{\mathcal{I}}_{\{s,t\}}^{i, \partial} \rangle_{\varrho} \leq C \varsigma_n (t-s)^{1/p}$ , from which we deduce that

$$\left\langle \int_s^t F_r^{i, n}(\cdot) d\mathbb{W}_r^{i, (n)}(\cdot) - \int_s^t \overline{F}_r^i(\cdot) d\overline{\mathbb{W}}_r^i(\cdot) \right\rangle_{\varrho} \leq C \varsigma_n (t-s)^{1/p}.$$

Similarly, Lemma A.1 says that  $\langle [F^{i, n}(\cdot) - \overline{F}^i(\cdot)]_{s,t} \rangle_{\varrho} \leq C \varsigma_n (t-s)^{1/p}$ , and,

noting that

$$\begin{aligned} R_{s,t}^{\int F_r^{i,n} d\mathbf{W}_r^{i,(n)}}(\omega) &= \int_s^t F_r^{i,n}(\omega) d\mathbf{W}_r^{i,(n)}(\omega) - \mathcal{I}_{s,t}^{i,n,\partial}(\omega) \\ &\quad + \delta_x F_s^{i,n}(\omega) \mathbb{W}_{s,t}^i(\omega) + \frac{1}{n} \sum_{j=1}^n \delta_\mu F_s^{i,j,n}(\omega) \mathbb{W}_{s,t}^{j,i}(\omega), \\ R_{s,t}^{\int \overline{F}_r^i d\overline{\mathbf{W}}_r^i}(\omega) &= \int_s^t \overline{F}_r^i(\omega) d\overline{\mathbf{W}}_r^i(\omega) - \overline{\mathcal{I}}_{s,t}^{i,\partial}(\omega) \\ &\quad + \delta_x \overline{F}_s^i(\omega) \mathbb{W}_{s,t}^i(\omega) + \mathbb{E} \left[ \delta_\mu \overline{F}_s^i(\omega, \cdot) \mathbb{W}_{s,t}^{i,\perp}(\cdot, \omega) \right], \end{aligned}$$

we deduce in a similar manner, using (5.10) and Lemma A.1 once again, that

$$\left\langle R_{s,t}^{\int F_r^{i,n} d\mathbf{W}_r^{i,(n)}}(\cdot) - R_{s,t}^{\int \overline{F}_r^i d\overline{\mathbf{W}}_r^i}(\cdot) \right\rangle_\varrho \leq C_{\zeta_n} (t-s)^{2/p}.$$

So, fixing  $i \in \{1, \dots, n\}$ , choosing  $\varrho$  large enough and applying a suitable version of Kolmogorov's theorem (see for instance Theorem 3.1 in [19]), we can find  $p' \in (p, 3)$  such that

$$\begin{aligned} \left| \int_s^t F_r^{i,n}(\omega) d\mathbf{W}_r^{i,(n)} - \int_s^t \overline{F}_r^i(\omega) d\overline{\mathbf{W}}_r^i(\omega) \right| &\leq \theta^{i,n}(\omega) (t-s)^{1/p'}, \\ \left| \left[ F_r^{i,n}(\omega) - \overline{F}_r^i(\omega) \right]_{s,t} \right| &\leq \theta^{i,n}(\omega) (t-s)^{1/p'}, \\ \left| R_{s,t}^{\int F_r^{i,n} d\mathbf{W}_r^{i,(n)}}(\omega) - R_{s,t}^{\int \overline{F}_r^i d\overline{\mathbf{W}}_r^i}(\omega) \right| &\leq \theta^{i,n}(\omega) (t-s)^{2/p'}, \end{aligned} \quad (5.11)$$

with  $\langle \theta^{i,n}(\cdot) \rangle_\varrho \leq C_{\zeta_n}$ , for a new value of the constant  $C$ .

Observe now that the empirical control associated with our empirical rough set-up and with the exponent  $p'$  reads (compare with (2.10))

$$w_{p'}^{i,n}(s, t, \omega) := v_{p'}^{i,n}(s, t, \omega) + {}^{(n)}(v_{p'}^{\bullet,n}(\omega))_{q;[s,t],1-v},$$

where we used the same notation as in (4.3). In fact, there is no loss of generality in changing the definition of  $w_{p'}^{i,n}$  into

$$w_{p'}^{i,n}(s, t, \omega) := v_{p'}^{i,n}(s, t, \omega) + {}^{(n)}(v_{p'}^{\bullet,n}(\omega))_{q;[s,t],1-v} + (t-s), \quad (5.12)$$

which permits to replace  $(t-s)^{1/p'}$  by  $w_{p'}^{i,n}(s, t, \omega)^{1/p'}$  in the inequalities (5.11). Hence,

$$\left\| \int_0^t F_r^{i,n}(\omega) d\mathbf{W}_r^{i,(n)} - \int_0^t \overline{F}_r^i(\omega) d\overline{\mathbf{W}}_r^i(\omega) \right\|_{[0,T], w_{p'}^{i,n}, p'} \leq \theta^{i,n}(\omega).$$

**Step 2.** We now make use of Proposition 3.2 to compare

$$\int_0^t F(X_s^{i,(n)}(\omega), \mu_s^n(\omega)) d\mathbf{W}_s^{i,(n)}(\omega) \quad \text{and} \quad \int_0^t F(\overline{X}_s^i(\omega), \overline{\mu}_s^n(\omega)) d\overline{\mathbf{W}}_s^i(\omega),$$

see (5.2). To simplify the notations, we just write  $X^i$  for  $X^{i,(n)}$  and  $\mathbf{W}^i$  for  $\mathbf{W}^{i,(n)}$ . We then apply Proposition 3.2 with

$$(X(\omega), Y(\cdot)) = (X^i(\omega), X^\bullet(\omega)), \quad (X'(\omega), Y'(\cdot)) = (\overline{X}^i(\omega), \overline{X}^\bullet(\omega)), \quad (5.13)$$

the underlying set-up being understood as the empirical rough set-up for a given realization  $\omega$ . The difficulty here is that the variations of these two solutions are controlled by two different functionals  $w$ , see (2.11). This is the rationale for introducing  $\widehat{w}_{p'}^{i,n}$  in (5.3). Obviously,  $\widehat{w}^{i,n}(\cdot, \cdot, \omega)$  (we remove the index  $p'$  for simplicity) is not the *natural* control functional associated with  $\mathbf{W}^i(\omega)$ , but it is greater than  $w_{p'}^{i,n}(s, t, \omega)$  and it satisfies

$${}^{(n)}\langle \widehat{w}^{\bullet,n}(s, t, \omega) \rangle_q \leq 2\widehat{w}^{i,n}(s, t, \omega), \quad (5.14)$$

which suffices to apply Proposition 3.2, see also [1, Proposition 4.3], with  $w_{p'}^{i,n}(s, t, \omega)$  replaced by  $\widehat{w}^{i,n}(s, t, \omega)$ . The resulting semi-norm that must be used to control the difference  $(X(\omega) - X'(\omega), Y(\cdot) - Y'(\cdot)) = (X^i(\omega) - \overline{X}^i(\omega), X^\bullet(\omega) - \overline{X}^\bullet(\omega))$  on a given interval  $[s, t]$  is  $\|\cdot\|_{[s,t], \widehat{w}^{i,n}, p'}$ . We use the corresponding local accumulation, which we denote by  $\widehat{N}^{i,n}([0, T], \omega, \alpha)$ .

By construction of the processes  $(X^i(\omega))_{i=1, \dots, n}$  as the solution of the empirical rough equation, the pair  $(X(\omega), Y(\cdot)) = (X^i(\omega), X^\bullet(\omega))$  in (5.13) automatically satisfies the first bound in (3.3) with  $w = \widehat{w}^{i,n}$  and for some large enough deterministic  $L_0$ ; implicitly, this means that we perform the same construction as in the proof of Theorem 3.3, see [1, Theorem 4.4], using therein the empirical rough-set up and the control functionals  $(\widehat{w}^{i,n})_{i=1, \dots, n}$ . In particular, the points  $(t_\ell^0 = \tau_\ell(0, T, \omega, 1/(4L_0)))_{\ell=0, \dots, N_0+1}$  in the statement of Proposition 3.2 are understood as with respect to  $\widehat{w}^{i,n}$ . Also, by the last part in the statement of [1, Proposition 4.2], we know that  $Y(\cdot) = X^\bullet(\omega)$  satisfies condition (3.2) with respect to  ${}^{(n)}\langle \cdot \rangle_8$  if we assume that  $T$  satisfies

$${}^{(n)}\left[ \widehat{N}^{\bullet,n}([0, T], \omega, 1/(4L_0)) \right]_8 \leq c, \quad (5.15)$$

for a deterministic constant  $c$ , independent of  $n$ ,  $L_0$  and  $T$ .

In fact, using the Hölder regularity of the paths, see (4.5) for a similar use, and using the additional  $t - s$  in the definition (5.3),  $\widehat{w}^{i,n}$  dominates (up to a multiplicative constant) the control  $\overline{w}^i$  associated to  $\overline{\mathbf{W}}^i$  through (2.10) (see (5.1) for the definition of  $\overline{\mathbf{W}}^i$ ; in short, the variations of  $W^i(\omega)$  and  $\mathbb{W}^i(\omega)$  are already included in  $v_{p'}^{i,n}(\omega)$ , the variations of  $\mathbb{W}^{i,\perp}(\omega, \cdot)$  and  $\mathbb{W}^{i,\perp}(\cdot, \omega)$  are precisely included in the definition of  $\widehat{v}_{p'}^{i,n}(\omega)$  and, using the Hölder regularity of the paths, the variations of  $W(\cdot)$  and  $\mathbb{W}^\perp(\cdot, \cdot)$  (in  $L^q$ ) are dominated by the additional  $t - s$ ). Moreover, we have

$$\langle \widehat{w}^{i,n}(s, t, \cdot) \rangle_q \leq C(t - s) \leq C\widehat{w}^{i,n}(s, t, \omega),$$

for a constant  $C$  independent of  $i, n, s$  and  $t$ . Although  $C \geq 2$  (compare with (2.8)), this permits to use  $\hat{w}^{i,n}(s, t, \cdot)$  as control functional when working with the rough set-up  $\overline{\mathbf{W}}^i$  and, in particular, when invoking the solvability Theorem 3.3 – the proof would be the same. This is an important point: the path  $\overline{X}^i(\omega)$ , defined right after (5.1), is the solution of a mean-field rough equation driven by a signal that is controlled by  $\hat{w}^{i,n}(\cdot)$ . Hence,  $X^i(\omega) = \overline{X}^i(\omega)$  in (5.13) satisfies the second bound in (3.3) with  $w = \hat{w}^{i,n}$ . Also, invoking the first line in [1, Proposition 4.2, (4.4)] for each  $i \in \{1, \dots, n\}$ , we deduce that  $Y^i(\cdot) = \overline{X}^i(\omega)$  satisfies condition (3.2) with respect to  ${}^{(n)}[\cdot]_8$  provided (5.15) holds true. Due to the form of the constant in the right-hand side of [1, Proposition 4.2, (4.4)], this may require to work with a larger value of the threshold  $L_0$  in the statement of Proposition 3.2, but, as made clear in the statement of Proposition 3.2, this is not a hindrance. Then, by Proposition 3.2, we obtain, for a given  $L \geq L_0$ ,

$$\begin{aligned} & \left\| \int_{t_k}^{\cdot} F(X_r^i(\omega), \mu^n(\omega)) d\mathbf{W}_r^i(\omega) - \int_{t_k}^{\cdot} F(\overline{X}_r^i(\omega), \overline{\mu}^n(\omega)) d\mathbf{W}_r^i(\omega) \right\|_{[t_k, t_{k+1}], \hat{w}^{i,n}, p'} \\ & \leq \gamma \hat{w}^{i,n}(0, t_k, \omega)^{1/p'} \left( \left\| (X^i - \overline{X}^i)(\omega) \right\|_{[0, t_k], \hat{w}^{i,n}, p'} \right. \\ & \quad \left. + {}^{(n)} \left[ \left\| (X^\bullet - \overline{X}^\bullet)(\omega) \right\|_{[0, T], \hat{w}^{\bullet,n}, p'} \right]_8 \right) \\ & \quad + \frac{\gamma}{4L} \left( \left\| (X^i - \overline{X}^i)(\omega) \right\|_{[t_k, t_{k+1}], \hat{w}^{i,n}, p'} \right. \\ & \quad \left. + {}^{(n)} \left[ \left\| (X^\bullet - \overline{X}^\bullet)(\omega) \right\|_{[t_k, t_{k+1}], \hat{w}^{\bullet,n}, p'} \right]_8 \right), \end{aligned}$$

where  $\hat{w}^{i,n}(t_k, t_{k+1}, \omega)^{1/p'} \leq 1/(4L)$  as long as  $k \leq 2\hat{N}^{i,n}([0, T], \omega, 1/(4L))$  (since the sequence  $(t_i)_i$  must refine the sequence  $(t_j^0)_j$ , we may assume that the collection  $(t_i)_i$  counts  $2\hat{N}^{i,n}([0, T], \omega, 1/(4L)) + 2$  points, including  $t_0 = 0$ ) and where we dropped the leading coefficient  $(1 + 1/(4L))$  in the main equality of Proposition 3.2 (which is always possible by modifying  $\gamma$  accordingly,  $\gamma$  here depending on  $L_0$ ). The point now is to insert the conclusion of the first step (replacing for free  $w^{i,n}$  by  $\hat{w}^{i,n}$  therein). We get

$$\begin{aligned} & \left\| (X^i - \overline{X}^i)(\omega) \right\|_{[t_k, t_{k+1}], \hat{w}^{i,n}, p'} \\ & \leq \gamma \hat{w}^{i,n}(0, t_k, \omega)^{1/p'} \left( \left\| (X^i - \overline{X}^i)(\omega) \right\|_{[0, t_k], \hat{w}^{i,n}, p'} \right. \\ & \quad \left. + {}^{(n)} \left[ \left\| (X^\bullet - \overline{X}^\bullet)(\omega) \right\|_{[0, T], \hat{w}^{\bullet,n}, p'} \right]_8 \right) + \theta^{i,n}(\omega) \\ & \quad + \frac{\gamma}{4L} \left( \left\| (X^i - \overline{X}^i)(\omega) \right\|_{[t_k, t_{k+1}], \hat{w}^{i,n}, p'} \right. \\ & \quad \left. + {}^{(n)} \left[ \left\| (X^\bullet - \overline{X}^\bullet)(\omega) \right\|_{[t_k, t_{k+1}], \hat{w}^{\bullet,n}, p'} \right]_8 \right). \end{aligned}$$

If  $\gamma/(4L) \leq 1/2$ , we get

$$\begin{aligned} & \left\| (X^i - \bar{X}^i)(\omega) \right\|_{[t_k, t_{k+1}], \hat{w}^{i,n}, p'} \\ & \leq 2\gamma \left( \frac{1}{L} + \hat{w}^{i,n}(0, t_k, \omega)^{1/p'} \right) \left( \left\| (X^i - \bar{X}^i)(\omega) \right\|_{[0, t_k], \hat{w}^{i,n}, p'} \right. \\ & \quad \left. + {}^{(n)} \left( \left\| (X^\bullet - \bar{X}^\bullet)(\omega) \right\|_{[0, T], \hat{w}^{\bullet,n}, p'} \right)_8 \right) + 2\theta^{i,n}(\omega), \end{aligned} \quad (5.16)$$

and, allowing the value of the constant  $c$  below to increase from line to line, as long as it remains independent of  $n$ ,  $L$  and  $T$  (but  $c$  now possibly depending on  $L_0$  since  $\gamma$  does), we get (see for instance [1, footnote 7] for the concatenation of two intervals)

$$\begin{aligned} & \left\| (X^i - \bar{X}^i)(\omega) \right\|_{[0, t_{k+1}], \hat{w}^{i,n}, p'} \\ & \leq c \left( \left\| (X^i - \bar{X}^i)(\omega) \right\|_{[0, t_k], \hat{w}^{i,n}, p'} + \left\| (X^i - \bar{X}^i)(\omega) \right\|_{[t_k, t_{k+1}], \hat{w}^{i,n}, p'} \right) \\ & \leq c \left( 1 + \zeta_T^{i,n}(\omega) \right) \left\| (X^i - \bar{X}^i)(\omega) \right\|_{[0, t_k], \hat{w}^{i,n}, p'} \\ & \quad + c \zeta_T^{i,n}(\omega) {}^{(n)} \left( \left\| (X^\bullet - \bar{X}^\bullet)(\omega) \right\|_{[0, T], \hat{w}^{\bullet,n}, p'} \right)_8 + c\theta^{i,n}(\omega), \end{aligned}$$

with  $\zeta_T^{i,n}(\omega) := \frac{1}{L} + w_{p'}^{i,n}(0, T, \omega)^{1/p'}$ . So, by induction,

$$\begin{aligned} & \left\| (X^i - \bar{X}^i)(\omega) \right\|_{[0, t_{k+1}], \hat{w}^{i,n}, p'} \leq c \left( \sum_{\ell=0}^k [c(1 + \zeta_T^{i,n}(\omega))]^\ell \right) \\ & \quad \times \left( \zeta_T^{i,n}(\omega) {}^{(n)} \left( \left\| (X^\bullet - \bar{X}^\bullet)(\omega) \right\|_{[0, T], \hat{w}^{\bullet,n}, p'} \right)_8 + \theta^{i,n}(\omega) \right). \end{aligned}$$

In the end,

$$\begin{aligned} & \left\| (X^i - \bar{X}^i)(\omega) \right\|_{[0, T], \hat{w}^{i,n}, p'} \\ & \leq c \left[ c(1 + \zeta_T^{i,n}(\omega)) \right]^{2\hat{N}^{i,n}([0, T], \omega, 1/(4L))+1} \\ & \quad \times \left( \zeta_T^{i,n}(\omega) {}^{(n)} \left( \left\| (X^\bullet - \bar{X}^\bullet)(\omega) \right\|_{[0, T], \hat{w}^{\bullet,n}, p'} \right)_8 + \theta^{i,n}(\omega) \right). \end{aligned} \quad (5.17)$$

Hence, using the shorten notation  $\hat{N}_T^{i,n}(\omega)$  for  $\hat{N}^{i,n}([0, T], \omega, 1/(4L))$  and assuming  $c \geq 1$ , we obtain

$$\begin{aligned} & {}^{(n)} \left( \left\| (X^\bullet - \bar{X}^\bullet)(\omega) \right\|_{[0, T], \hat{w}^{\bullet,n}, p'} \right)_8 \\ & \leq {}^{(n)} \left[ [c^2 (1 + \zeta_T^{\bullet,n}(\omega))]^{2\hat{N}_T^{\bullet,n}(\omega)+1} \zeta_T^{\bullet,n}(\omega) \right]_8 \\ & \quad \times {}^{(n)} \left( \left\| (X^\bullet - \bar{X}^\bullet)(\omega) \right\|_{[0, T], \hat{w}^{\bullet,n}, p'} \right)_8 \\ & \quad + {}^{(n)} \left[ [c^2 (1 + \zeta_T^{\bullet,n}(\omega))]^{2\hat{N}_T^{\bullet,n}(\omega)+1} \theta^{\bullet,n}(\omega) \right]_8. \end{aligned} \quad (5.18)$$

**Step 3.** The key quantity of interest in (5.18) is the multiplicative factor in the second line, which we denote by

$$\Psi_T^n(\omega) := {}^{(n)}\left[ [c^2 (1 + \zeta_T^{\bullet,n}(\omega))]^{2\widehat{N}_T^{\bullet,n}(\omega)+1} \zeta_T^{\bullet,n}(\omega) \right]_8.$$

In particular, letting

$$\Theta_T^n(\omega) := {}^{(n)}\left[ [c^2 (1 + \zeta_T^{\bullet,n}(\omega))]^{2\widehat{N}_T^{\bullet,n}(\omega)+1} \theta^{\bullet,n}(\omega) \right]_8,$$

we rewrite (5.18) in the form

$$\begin{aligned} & {}^{(n)}\left( \left\| (X^\bullet - \overline{X}^\bullet)(\omega) \right\|_{[0,T],\widehat{w}^{\bullet,n,p'}} \right)_8 \\ & \leq \Psi_T^n(\omega) {}^{(n)}\left( \left\| (X^\bullet - \overline{X}^\bullet)(\omega) \right\|_{[0,T],\widehat{w}^{\bullet,n,p'}} \right)_8 + \Theta_T^n(\omega). \end{aligned} \quad (5.19)$$

*Here comes the key point.* The variable  $\omega$  being frozen, we can choose deterministically the time horizon  $T$  small enough, depending on  $\omega$ , and  $L \geq L_0$  large enough, such that  $\Psi_T^n(\omega) \leq 1/2$  and (5.15) holds true. The proof is made clear below. Take it for granted for a while and deduce that

$${}^{(n)}\left( \left\| (X^\bullet - \overline{X}^\bullet)(\omega) \right\|_{[0,T],\widehat{w}^{\bullet,n,p'}} \right)_8 \leq 2\Theta_T^n(\omega).$$

The above inequality sounds really close to the desired result, but it is on a small interval  $[0, T]$  only. The purpose is thus to iterate it in order to cover any given time interval.

**Step 4.** In order to iterate in a proper way, we change our notation. While we keep the notation  $T$  for the deterministic time horizon given in the statement, we use the latter  $\tau$  instead of  $T$  in the previous analysis. Put differently,  $\tau$  will stand for the random time horizon such that  $\Psi_\tau$  is small enough. More precisely, we let  $\tau_0 = \tau$  and then consider a possibly random dissection  $0 = \tau_0 < \tau_1 < \dots < \tau_M = T$  of the interval  $[0, T]$  into  $M$  subintervals. *The goal of this step is to clarify the construction of the dissection in order to iterate the previous steps to any interval  $[\tau_\ell, \tau_{\ell+1}]$ ,  $\ell = 0, \dots, M - 1$ .*

To do so, we need to revisit the statement of Proposition 3.2. Assume indeed that we have a bound for

$$\mathcal{E}_{\tau_\ell}^{i,n}(\omega) := \left( 1 + \widehat{w}^{i,n}(0, T, \omega)^{1/p'} \right) \left\| (X^i - \overline{X}^i)(\omega) \right\|_{[0,\tau_\ell],\widehat{w}^{i,n,p'}},$$

for some  $\ell \leq M$ . Then, in order to duplicate the previous two steps, we must consider a new dissection  $\tau_\ell = t_0 < t_1 < \dots < t_K = \tau_{\ell+1}$  of the interval  $[\tau_\ell, \tau_{\ell+1}]$  with the property that  $K = 2\widehat{N}^{i,n}([\tau_\ell, \tau_{\ell+1}], \omega, 1/(4L)) + 1$  and that  $\widehat{w}^{i,n}(t_k, t_{k+1}, \omega) \leq 1/(4L)$  if  $k < K$ . The key point is to apply the first inequality in (3.4) on  $[t_k, t_{k+1}]$  with  $(X(\omega), Y(\cdot)) = (X^i(\omega), X^i(\cdot))$  and  $(X'(\omega), Y'(\cdot)) =$

$(\overline{X}^i(\omega), \overline{X}^i(\cdot))$ , but with  $\tau_\ell$  instead of 0 as initial time. Upper bounding the second line in (3.4) by  $\mathcal{E}_{\tau_\ell}^{i,n}(\omega) + {}^{(n)}(\mathcal{E}_{\tau_\ell}^{i,n}(\omega))_8$ , we obtain

$$\begin{aligned} & \left\| \int_{t_k}^{\cdot} F(X_r^i(\omega), \mu_r^n(\omega)) d\mathbf{W}_r^i(\omega) - \int_{t_k}^{\cdot} F(\overline{X}_r^i(\omega), \overline{\mu}_r^n(\omega)) d\mathbf{W}_r^i(\omega) \right\|_{[t_k, t_{k+1}], \widehat{w}^{i,n}, p'} \\ & \leq \gamma \widehat{w}^{i,n}(\tau_\ell, \tau_{\ell+1}, \omega)^{1/p'} \left\{ \left\| (X^i - \overline{X}^i)(\omega) \right\|_{[\tau_\ell, t_k], \widehat{w}^{i,n}, p'} \right. \\ & \quad \left. + {}^{(n)} \left( \left\| (X^\bullet - \overline{X}^\bullet)(\omega) \right\|_{[\tau_\ell, \tau_{\ell+1}], \widehat{w}^{\bullet,n}, p'} \right)_8 \right\} \\ & + \frac{\gamma}{4L} \left\{ \left\| (X^i - \overline{X}^i)(\omega) \right\|_{[t_k, t_{k+1}], \widehat{w}^{i,n}, p'} \right. \\ & \quad \left. + {}^{(n)} \left( \left\| (X^\bullet - \overline{X}^\bullet)(\omega) \right\|_{[t_k, t_{k+1}], \widehat{w}^{\bullet,n}, p'} \right)_8 \right\} \\ & + \gamma \left[ \mathcal{E}_{\tau_\ell}^{i,n}(\omega) + {}^{(n)}(\mathcal{E}_{\tau_\ell}^{\bullet,n}(\omega))_8 \right], \end{aligned}$$

provided the analogues of (3.2) and (3.3) hold true. As for (3.3), we may argue as in the second step: It is a consequence of the proof of Theorem 3.3. As for (3.2), it is again a consequence of the statement of [1, Proposition 4.2], provided the analogue of (5.15) holds true, namely

$${}^{(n)} \left( \widehat{N}^{\bullet,n}([\tau_\ell, \tau_{\ell+1}], \omega, 1/(4L_0)) \right)_8 \leq c. \quad (5.20)$$

Then, proceeding as in the second step,

$$\begin{aligned} & \left\| (X^i - \overline{X}^i)(\omega) \right\|_{[t_k, t_{k+1}], \widehat{w}^{i,n}, p'} \\ & \leq c \widehat{w}^{i,n}(\tau_\ell, \tau_{\ell+1}, \omega)^{1/p'} \left\{ \left\| (X^i - \overline{X}^i)(\omega) \right\|_{[\tau_\ell, t_k], \widehat{w}^{i,n}, p'} \right. \\ & \quad \left. + {}^{(n)} \left( \left\| (X^\bullet - \overline{X}^\bullet)(\omega) \right\|_{[\tau_\ell, \tau_{\ell+1}], \widehat{w}^{\bullet,n}, p'} \right)_8 \right\} \\ & + c \left\{ \mathcal{E}_{\tau_\ell}^{i,n}(\omega) + {}^{(n)}(\mathcal{E}_{\tau_\ell}^{\bullet,n}(\omega))_8 + \theta^{i,n}(\omega) \right\}. \end{aligned}$$

In the end, we are in the same situation as in (5.16), but with new  $\zeta_T^{i,n}$  and  $\widehat{N}_T^{i,n}$ . Here, we let (pay attention that, to be consistent with the notations  $\zeta_T^{i,n}$  and  $\widehat{N}_T^{i,n}$ , we should use  $[\tau_\ell, \tau_{\ell+1}]$  instead of  $\ell$  as subscript below, but, for simplicity, we prefer to use  $\ell$  only)

$$\zeta_\ell^{i,n}(\omega) := \widehat{w}^{i,n}(\tau_\ell, \tau_{\ell+1}, \omega)^{1/p'}, \quad \widehat{N}_\ell^{i,n}(\omega) := \widehat{N}^{i,n} \left( [\tau_\ell, \tau_{\ell+1}], \omega, \frac{1}{4L} \right).$$

Following (5.17), we obtain

$$\begin{aligned}
& \left\| (X^i - \bar{X}^i)(\omega) \right\|_{[\tau_\ell, \tau_{\ell+1}], \hat{w}^{i,n}, p'} \\
& \leq c \left[ c(1 + \zeta_\ell^{i,n}(\omega)) \right]^{2\widehat{N}_\ell^{i,n}(\omega)+1} \\
& \quad \times \left\{ \zeta_\ell^{i,n}(\omega)^{(n)} \left( \left\| (X^\bullet - \bar{X}^\bullet)(\omega) \right\|_{[\tau_\ell, \tau_{\ell+1}], \hat{w}^{\bullet,n}, p'} \right)_8 \right. \\
& \quad \left. + \theta^{i,n}(\omega) + \mathcal{E}_{\tau_\ell}^{i,n} + {}^{(n)}(\mathcal{E}_{\tau_\ell}^{\bullet,n}(\omega))_8 \right\}. \tag{5.21}
\end{aligned}$$

Hence,

$$\begin{aligned}
& {}^{(n)} \left( \left\| (X^\bullet - \bar{X}^\bullet)(\omega) \right\|_{[\tau_\ell, \tau_{\ell+1}], \hat{w}^{\bullet,n}, p'} \right)_8 \\
& \leq \Psi_\ell^n(\omega) \times {}^{(n)} \left( \left\| (X^\bullet - \bar{X}^\bullet)(\omega) \right\|_{[\tau_\ell, \tau_{\ell+1}], \hat{w}^{\bullet,n}, p'} \right)_8 + \Theta_\ell^n(\omega),
\end{aligned}$$

with

$$\begin{aligned}
\Psi_\ell^n(\omega) & := {}^{(n)} \left[ \left[ c^2 (1 + \zeta_\ell^{\bullet,n}(\omega)) \right]^{2\widehat{N}_\ell^{\bullet,n}(\omega)+1} \zeta_\ell^{\bullet,n}(\omega) \right]_8, \\
\Theta_\ell^n(\omega) & := {}^{(n)} \left[ \left[ c^2 (1 + \zeta_\ell^{\bullet,n}(\omega)) \right]^{2\widehat{N}_\ell^{\bullet,n}(\omega)+1} \left( \theta^{\bullet,n}(\omega) + \mathcal{E}_{\tau_\ell}^{\bullet,n}(\omega) + {}^{(n)}(\mathcal{E}_{\tau_\ell}^{\bullet,n}(\omega))_8 \right) \right]_8.
\end{aligned}$$

If we can choose  $\tau_{\ell+1} - \tau_\ell$  such that  $\Psi_\ell^n(\omega) \leq 1/2$ , then we get

$${}^{(n)} \left( \left\| (X^\bullet - \bar{X}^\bullet)(\omega) \right\|_{[\tau_\ell, \tau_{\ell+1}], \hat{w}^{\bullet,n}, p'} \right)_8 \leq 2 \Theta_\ell^n(\omega).$$

Eventually, returning to (5.21) and modifying the value of the constant  $c$ , we deduce

$$\begin{aligned}
& \left\| (X^i - \bar{X}^i)(\omega) \right\|_{[\tau_\ell, \tau_{\ell+1}], \hat{w}^{i,n}, p'} \\
& \leq c \left[ c(1 + \zeta_\ell^{i,n}(\omega)) \right]^{2\widehat{N}_\ell^{i,n}(\omega)+1} \\
& \quad \times \left( \zeta_\ell^{i,n}(\omega) \Theta_\ell^n(\omega) + \theta^{i,n}(\omega) + \mathcal{E}_{\tau_\ell}^{i,n} + {}^{(n)}(\mathcal{E}_{\tau_\ell}^{\bullet,n}(\omega))_8 \right),
\end{aligned}$$

and then

$$\mathcal{E}_{\tau_{\ell+1}}^{i,n}(\omega) \leq \kappa_\ell^{i,n}(\omega) \left( \zeta_\ell^{i,n}(\omega) \Theta_\ell^n(\omega) + \theta^{i,n}(\omega) + \mathcal{E}_{\tau_\ell}^{i,n}(\omega) + {}^{(n)}(\mathcal{E}_{\tau_\ell}^{\bullet,n}(\omega))_8 \right),$$

with

$$\kappa_\ell^{i,n}(\omega) := c^2 \left( 1 + \widehat{w}^{i,n}(0, T, \omega)^{1/p'} \right) \left[ c^2 (1 + \zeta_\ell^{i,n}(\omega)) \right]^{2\widehat{N}_\ell^{i,n}(\omega)+1}, \tag{5.22}$$

using the fact that  $c \geq 1$ . By induction, we get the following global bound:

$$\mathcal{E}_{\tau_{\ell+1}}^{i,n}(\omega) \leq \sum_{k=0}^{\ell} \mathcal{K}_{k,\ell}^{i,n}(\omega) \left[ \zeta_k^{i,n}(\omega) \Theta_k^n(\omega) + \theta^{i,n}(\omega) + {}^{(n)}(\mathcal{E}_{\tau_k}^{\bullet,n}(\omega))_8 \right], \quad (5.23)$$

with

$$\mathcal{K}_{k,\ell}^{i,n}(\omega) := \prod_{j=k}^{\ell} \kappa_j^{i,n}(\omega). \quad (5.24)$$

We deduce that for any  $r > 8$ , we can find a constant  $q(r)$  such that

$$\begin{aligned} {}^{(n)}(\mathcal{E}_{\tau_{\ell+1}}^{\bullet,n}(\omega))_r &\leq \sum_{k=0}^{\ell} \left\{ {}^{(n)}(\mathcal{K}_{k,\ell}^{\bullet,n})_{q(r)} \times \left( 1 + {}^{(n)}(\widehat{w}^{\bullet,n}(0, T, \omega))^{1/p'} \right)_{q(r)} \right. \\ &\quad \times \left( 1 + {}^{(n)}\left[ c^2 (1 + \zeta_k^{\bullet,n}(\omega)) \right]^{2\widehat{N}_k^{\bullet,n}(\omega)+1} \right)_{q(r)} \\ &\quad \left. \times \left( {}^{(n)}(\theta^{\bullet,n}(\omega))_{q(r)} + {}^{(n)}(\mathcal{E}_{\tau_k}^{\bullet,n}(\omega))_r \right) \right\}. \end{aligned}$$

Using the fact that

$$\begin{aligned} &{}^{(n)}(\mathcal{K}_{k,\ell}^{\bullet,n})_{q(r)} \\ &\geq \max \left( 1, {}^{(n)}(\widehat{w}^{\bullet,n}(0, T, \omega))^{1/p'} \right)_{q(r)}, \quad {}^{(n)}\left[ c^2 (1 + \zeta_k^{\bullet,n}(\omega)) \right]^{2\widehat{N}_k^{\bullet,n}(\omega)+1} \right)_{q(r)}, \end{aligned}$$

we obtain a bound of the form  $a_{\ell+1} \leq \sum_{k=0}^{\ell} g_{k,\ell} (b + a_k)$ , with

$$a_{\ell} := {}^{(n)}(\mathcal{E}_{\tau_{\ell}}^{\bullet,n}(\omega))_r, \quad g_{k,\ell} := 4 \times \left( {}^{(n)}(\mathcal{K}_{k,\ell}^{\bullet,n})_{q(r)} \right)^3, \quad b := {}^{(n)}(\theta^{\bullet,n}(\omega))_{q(r)}.$$

Hence,

$$a_{\ell} \leq b \sum_{j=1}^{\ell} \sum_{0 \leq k_1 \leq \dots \leq k_j \leq k_{j+1} = \ell} \prod_{h=1}^j g_{k_h, k_{h+1}}. \quad (5.25)$$

Back to (5.24), we will use below the bound

$$\begin{aligned} &\mathcal{K}_{k,\ell}^{i,n}(\omega) \\ &\leq c^{2(\ell+1-k)} \prod_{j=k}^{\ell} \left\{ (1 + \widehat{w}^{i,n}(0, T, \omega))^{1/p'} \left[ c^2 (1 + \zeta_j^{i,n}(\omega)) \right]^{2\widehat{N}_j^{i,n}(\omega)+1} \right\} \\ &\leq c^{4(\ell+1-k) + 4\widehat{N}_{k,\ell+1}^{i,n}(\omega)} (1 + \widehat{w}^{i,n}(0, T, \omega))^{1/p'}{}^{\ell+1-k+2\widehat{N}_{k,\ell+1}^{i,n}(\omega)}, \end{aligned} \quad (5.26)$$

with the shortened notation  $\widehat{N}_{k,\ell}^{i,n}(\omega) := \widehat{N}^{i,n}([\tau_k, \tau_{\ell}], \omega, 1/(4L))$ .

**Step 5.** We now recall that the sequence of times  $0 = \tau_0 < \tau_1 < \dots < \tau_M$  must satisfy (5.20). A priori, the sequence may be random, but we are free to

take it deterministic, which is what we do below. To make it clear, we let

$$M_1 = \left\langle \|W(\cdot)\|_{[0,T],(1/p)-H}^p \right\rangle_q + \left\langle \|\mathbb{W}(\cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q \\ + \left\langle \|\mathbb{W}^\perp(\cdot, \cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q,$$

and we consider the events

$$A_1^n = \left\{ \omega \in \Omega : \binom{(n)}{\|W^\bullet(\omega)\|_{[0,T],(1/p')-H}^{p'}} + \binom{(n)}{\|\mathbb{W}^\bullet(\omega)\|_{[0,T],(2/p')-H}^{p'/2}} \right. \\ \left. + \binom{(n)}{\|\mathbb{W}^{\bullet,\bullet}(\omega)\|_{[0,T],(2/p')-H}^{p'/2}} \right\rangle_q \leq M_1 + 1 \Big\}, \\ A_2^{i,n} = \left\{ \omega \in \Omega : \binom{(n)}{\|\mathbb{W}^{\bullet,i}(\omega)\|_{[0,T],(2/p')-H}^{p'/2}} + \binom{(n)}{\|\mathbb{W}^{i,\bullet}(\omega)\|_{[0,T],(2/p')-H}^{p'/2}} \right. \\ \left. \leq \left\langle \|\mathbb{W}^{i,\perp}(\omega, \cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q + \left\langle \|\mathbb{W}^{i,\perp}(\cdot, \omega)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q + 1 \right\}.$$

On the event  $A_1^n \cap A_2^{i,n}$ , we have (compare with (4.3) and (4.5)) for  $s, t \in [0, T]^2$ ,  $s < t$ ,

$$v_{p'}^{i,n}(s, t, \omega) \leq \|W^i(\omega)\|_{[s,t],p'-v}^{p'} + \|\mathbb{W}^i(\omega)\|_{[s,t],p'/2-v}^{p'/2} \\ + \left( 2 + M_1 + \left\langle \|\mathbb{W}^{i,\perp}(\omega, \cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q \right. \\ \left. + \left\langle \|\mathbb{W}^{i,\perp}(\cdot, \omega)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q \right) (t - s).$$

Therefore, introducing the new event

$$A_3^n = \left\{ \binom{(n)}{\left\langle \|\mathbb{W}^{\bullet,\perp}(\omega, \cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q} \right. \\ \left. + \binom{(n)}{\left\langle \|\mathbb{W}^{\perp,\bullet}(\omega, \cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q} \right\} \leq 2(1 + M_1),$$

we get, on  $A_1^n \cap \left(\bigcap_{i=1}^n A_2^{i,n}\right) \cap A_3^n$ ,

$$\binom{(n)}{v_{p'}^{\bullet,n}(\omega)}_{q;[s,t],1-v} \leq (5 + 4M_1)(t - s).$$

Recall now the definition of  $\widehat{v}_{p'}^{i,n}$  in (5.3). We have

$$\widehat{v}_{p'}^{i,n}(s, t, \omega) \\ = \left\langle \|\mathbb{W}^{i,\perp}(\omega, \cdot)\|_{q;[s,t],p'/2-v}^{p'/2} + \left\langle \|\mathbb{W}^{i,\perp}(\cdot, \omega)\|_{q;[s,t],p'/2-v}^{p'/2} \right\rangle_q \right. \\ \left. \leq \left[ \left\langle \|\mathbb{W}^{i,\perp}(\omega, \cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q + \left\langle \|\mathbb{W}^{i,\perp}(\cdot, \omega)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q \right] (t - s).$$

And then,

$$\begin{aligned} {}^{(n)}(\widehat{v}_{p'}^{\bullet,n}(\omega))_{q:[s,t],1-v} &\leq \left( {}^{(n)}\left[ \left\langle \|\mathbb{W}^{\bullet,\perp}(\omega, \cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q \right]_q \right. \\ &\quad \left. + {}^{(n)}\left[ \left\langle \|\mathbb{W}^{\perp,\bullet}(\omega, \cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q \right]_q \right) (t-s). \end{aligned}$$

Therefore, on the event  $A_1^n \cap \left(\bigcap_{i=1}^n A_2^{i,n}\right) \cap A_3^n$ , we have

$${}^{(n)}(\widehat{v}_{p'}^{\bullet,n}(\omega))_{q:[s,t],1-v} \leq 2(1 + M_1)(t-s).$$

Using the same notations as in (5.3), we end-up with

$$\widehat{w}_{p'}^{i,n}(s, t, \omega) \leq \widetilde{w}_{p'}^{i,n}(s, t, \omega), \quad (5.27)$$

for  $\omega \in A_1^n \cap \left(\bigcap_{i=1}^n A_2^{i,n}\right) \cap A_3^n$ , where we let

$$\begin{aligned} \widetilde{w}_{p'}^{i,n}(s, t, \omega) &:= \|W^i(\omega)\|_{[s,t],p'-v}^{p'} + \|\mathbb{W}^i(\omega)\|_{[s,t],p'/2-v}^{p'/2} + (10 + 7M_1)(t-s) \\ &\quad + 2\langle \mathbb{W}^{i,\perp}(\omega, \cdot) \rangle_{q:[s,t],p'/2-v}^{p'/2} + 2\langle \mathbb{W}^{i,\perp}(\cdot, \omega) \rangle_{q:[s,t],p'/2-v}^{p'/2}. \end{aligned}$$

Using the notation (2.17), we also let  $\widetilde{N}^{i,n}([\tau, \tau'], \omega, \alpha) := N_{\varpi}([\tau, \tau'], \alpha)$ , with  $\varpi := (\widetilde{w}_{p'}^{i,n}(\omega))^{1/p'}$ . By (5.27),

$$\widehat{N}^{i,n}([\tau, \tau'], \omega, \alpha) \leq \widetilde{N}^{i,n}([\tau, \tau'], \omega, \alpha),$$

for  $\omega \in A_1^n \cap \left(\bigcap_{i=1}^n A_2^{i,n}\right) \cap A_3^n$ . The good point here is that the variables  $(\widetilde{w}_{p'}^{i,n})_{1 \leq i \leq n}$  are independent whilst the variables  $(\widehat{w}_{p'}^{i,n})_{1 \leq i \leq n}$  are not. Similarly, whenever  $\tau$  and  $\tau'$  are deterministic, the variables  $(\widetilde{N}^{i,n}([\tau, \tau'], \cdot, \alpha))_{1 \leq i \leq n}$  are independent. Moreover, it is not difficult to see that

$$\widetilde{w}_{p'}^{i,n}(s, t, \omega) \leq 2\widehat{w}_{p'}^{i,n}(s, t, \omega) + 8(1 + M_1)(t-s),$$

from which we deduce, see for instance (A.1), that, for any  $\alpha > 0$ ,

$$\widetilde{N}^{i,n}([\tau, \tau'], \omega, \alpha) \leq \widehat{N}^{i,n}\left([\tau, \tau'], \omega, \frac{\alpha}{4}\right) + C_\alpha,$$

for a constant  $C_\alpha$  only depending on  $\alpha$  and on  $M_1$ . In particular, we can easily replace  $\widehat{N}^{i,n}$  by  $\widetilde{N}^{i,n}$  in the third item of the assumption of the statement. Moreover, by (4.11), we deduce that each  $\widetilde{w}_{p'}^{i,n}$  satisfies the first bound in (3.5), uniformly in  $i$  and  $n$ .

We now have all the ingredients to choose the sequence  $0 = \tau_0 < \dots < \tau_M = T$ . We may take it, independently of  $n$ , such that

$$\left\langle \widetilde{N}^{1,n}([\tau_\ell, \tau_{\ell+1}], \cdot, 1/(4L_0)) \right\rangle_8 \leq \frac{c}{2},$$

for  $c$  as in (5.15). Existence of the dissection follows from the third item in the assumption of the statement as it says that the left-hand side can be bounded by  $C\sqrt{\tau_{\ell+1} - \tau_\ell}$ . Importantly, the latter bound says that  $M$  can be chosen independently of  $n$ . For sure, the index 1 in the left-hand side may be replaced by any  $i \in \{1, \dots, n\}$ . We then consider the family of events

$$A_4^{\ell,n} = \left\{ \binom{(n)}{\left[ \tilde{N}^{\bullet,n}([\tau_\ell, \tau_{\ell+1}], \cdot, 1/(4L_0)) \right]}_8 \leq c \right\}, \quad \ell = 0, \dots, M-1.$$

On  $A_1^n \cap \left( \bigcap_{i=1}^n A_2^{i,n} \right) \cap A_3^n \cap \left( \bigcap_{\ell=0}^{M-1} A_4^{\ell,n} \right)$ , the upper bound (5.20) is satisfied and then the conclusion of the fourth step holds true. Following (5.22), this prompts us to set:

$$\tilde{\kappa}_\ell^{i,n}(\omega) := c^2 \left( 1 + \tilde{w}^{i,n}(0, T, \omega)^{1/p'} \right) \left[ c^2 \left( 1 + \tilde{w}^{i,n}(\tau_\ell, \tau_{\ell+1}, \omega)^{1/p'} \right) \right]^{2\tilde{N}_\ell^{i,n}(\omega)+1},$$

and then  $\tilde{\mathcal{K}}_{k,\ell}^{i,n}(\omega) := \prod_{j=k}^\ell \tilde{\kappa}_j^{i,n}(\omega)$ . Returning to the conclusion of the fourth step, we get, for  $\omega \in A^n := A_1^n \cap \left( \bigcap_{i=1}^n A_2^{i,n} \right) \cap A_3^n \cap \left( \bigcap_{\ell=0}^{M-1} A_4^{\ell,n} \right)$ ,

$$\begin{aligned} & \binom{(n)}{\left( \mathcal{E}_{\tau_\ell}^{\bullet,n}(\omega) \right)_r} \\ & \leq \binom{(n)}{\left( \theta^{\bullet,n}(\omega) \right)_{q(r)}} \sum_{j=1}^\ell \sum_{0 \leq k_1 \leq \dots \leq k_j \leq k_{j+1} = \ell} \prod_{h=1}^j 4 \times \binom{(n)}{\left( \tilde{\mathcal{K}}_{k_h, k_{h+1}}^{\bullet,n}(\omega) \right)_{q(r)}}^3 \\ & \leq \ell 2^{2\ell+1} \times 4^\ell \times \binom{(n)}{\left( \theta^{\bullet,n}(\omega) \right)_{q(r)}} \times \binom{(n)}{\left( \tilde{\mathcal{K}}_{0,\ell}^{\bullet,n}(\omega) \right)_{q(r)}}^{3\ell}. \end{aligned}$$

The key fact here is that  $\tilde{\mathcal{K}}_{0,M}^{i,n}(\omega)$ , for any  $i \in \{1, \dots, n\}$ , has finite moments of any order, independently of  $i$  and  $n$ . The proof follows from (5.26), from (4.11) and from the third item in the assumption of the statement of Theorem 5.1, the last two properties implying that  $\left( 1 + \tilde{w}^{i,n}(\tau_\ell, \tau_{\ell+1}, \omega)^{1/p'} \right)^{\tilde{N}_\ell^{i,n}(\omega)}$  has finite moments of any order, independently of  $i$  and  $n$ , see for instance (5.32) below. Hence, for a constant  $C$ , independent of  $n$  but possibly depending on  $M$ , we get

$$\left\langle \mathbf{1}_{A^n}(\cdot) \binom{(n)}{\left( \mathcal{E}_T^{\bullet,n}(\cdot) \right)_r} \right\rangle_r \leq C \left\langle \binom{(n)}{\left( \theta^{\bullet,n}(\cdot) \right)_{q(r)}} \right\rangle_{2r} \leq C \left\langle \theta^{1,n}(\cdot) \right\rangle_{q(r)},$$

where we took, without any loss of generality,  $q(r) \geq 2r$ . Taking  $\varrho = q(r)$  in (5.11), we get that, for a constant  $C$  independent of  $n$ , but depending on  $r$ ,

$$\sup_{1 \leq i \leq n} \left\langle \mathbf{1}_{A^n}(\cdot) \left| \left( X^i - \bar{X}^i \right) (\cdot) \right| \right\rangle_r \leq C \zeta_n. \quad (5.28)$$

**Step 6.** From the law of large numbers and from (4.11), we claim that  $\mathbb{P}((A_1^n)^c)$  decays faster than any  $n^{-s}$ , for  $s > 0$ . The first step of the proof is to

notice that

$$\begin{aligned}
 & \mathbb{P}\left(\left(A_1^n\right)^c\right) \\
 & \leq \mathbb{P}\left(\omega : {}^{(n)}\left(\|W^\bullet(\omega)\|_{[0,T],(1/p')-H}^{p'}\right)_q - \left\langle \|W(\cdot)\|_{[0,T],(1/p')-H}^{p'} \right\rangle_q \geq \frac{1}{3}\right) \\
 & \quad + \mathbb{P}\left(\omega : {}^{(n)}\left(\|W^\bullet(\omega)\|_{[0,T],(2/p')-H}^{p'/2}\right)_q - \left\langle \|W(\cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q \geq \frac{1}{3}\right) \\
 & \quad + \mathbb{P}\left(\omega : {}^{(n)}\left(\|W^{\bullet,\bullet}(\omega)\|_{[0,T],(2/p')-H}^{p'/2}\right)_q - \left\langle \|W^\perp(\cdot, \cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q \geq \frac{1}{3}\right) \\
 & =: \pi_{1,1}^n + \pi_{1,2}^n + \pi_{1,3}^n.
 \end{aligned}$$

Since the most difficult term is the last one, we just explain how to handle it. The other two terms may be treated in the same way. Since  $q \geq 1$ , we first observe that

$$\begin{aligned}
 & \pi_{1,3}^n \\
 & \leq \mathbb{P}\left(\omega : {}^{(n)}\left(\|W^{\bullet,\bullet}(\omega)\|_{[0,T],(2/p')-H}^{p'/2}\right)_q \geq \left\langle \|W^\perp(\cdot, \cdot)\|_{[0,T],(2/p')-H}^{p'/2} \right\rangle_q^q + \frac{1}{3^q}\right) \\
 & \leq \mathbb{P}\left(\omega : \frac{1}{n^2} \sum_{i \neq j} \left(\|W^{i,j}(\omega)\|_{[0,T],(2/p')-H}^{qp'/2} - \left\langle \|W^\perp(\cdot, \cdot)\|_{[0,T],(2/p')-H}^{qp'/2} \right\rangle_1\right) \geq \frac{1}{3^{q+1}}\right) \\
 & \quad + \mathbb{P}\left(\omega : \frac{1}{n^2} \sum_{i=1}^n \|W^i(\omega)\|_{[0,T],(2/p')-H}^{qp'/2} \geq \frac{1}{3^{q+1}}\right).
 \end{aligned}$$

By (4.11), the last term in the right-hand side is easily handled. As for the first one, Markov's inequality yields, for any  $s > 1$ ,

$$\begin{aligned}
 & \mathbb{P}\left(\omega : \frac{1}{n^2} \sum_{j \neq i} \left(\|W^{i,j}(\omega)\|_{[0,T],(2/p')-H}^{qp'/2} - \left\langle \|W^\perp(\cdot, \cdot)\|_{[0,T],(2/p')-H}^{qp'/2} \right\rangle_1\right) \geq \frac{1}{3^{q+1}}\right) \\
 & \leq \frac{3^{s(q+1)}}{n^{s+1}} \sum_{i=1}^n \mathbb{E}\left[\left|\sum_{j:j \neq i} \left(\|W^{i,j}(\cdot)\|_{[0,T],(2/p')-H}^{qp'/2} - \left\langle \|W^\perp(\cdot, \cdot)\|_{[0,T],(2/p')-H}^{qp'/2} \right\rangle_1\right)\right|^s\right].
 \end{aligned}$$

By (4.11) again and by Rosenthal's inequality, see [31], we deduce that the right-hand side is less than  $Cn^{-s/2}$ , for a constant  $C$  independent of  $n$ . This completes the proof of our claim.

The same result holds for  $\mathbb{P}(A_3^n)$ . Also, since  $(\tilde{N}^{i,n}([\tau_\ell, \tau_{\ell+1}], \cdot, \alpha))_{1 \leq i \leq n}$ , are independent for any  $\ell = 0, \dots, M-1$ , we also have that  $\mathbb{P}(A_4^{\ell,n})$  decays faster than any  $n^{-s}$ , for any  $\ell = 0, \dots, M-1$ .

We finally check that the same is true for  $A_2^{i,n}$ . To make it clear, we prove that, for any  $s > 0$ ,  $\sup_{i=1, \dots, n} \mathbb{P}\left(\left(A_2^{i,n}\right)^c\right) \leq Cn^{-s}$  for a constant  $C$  independent of  $n$ , but the result is less obvious. In fact, it suffices to upper bound the first

term below, for which we have, as before,

$$\begin{aligned} & \mathbb{P}\left(\omega : {}^{(n)}(\|\mathbb{W}^{i,\bullet}(\omega)\|_{[0,T],(2/p')\text{-H}}^{p'/2})_q \geq \left\langle \|\mathbb{W}^{i,\perp}(\omega, \cdot)\|_{[0,T],(2/p')\text{-H}}^{p'/2} \right\rangle_q + \frac{1}{2}\right) \\ & \leq \mathbb{P}\left(\omega : \frac{1}{n} \sum_{j:j \neq i} \left( \|\mathbb{W}^{i,j}(\omega)\|_{[0,T],(2/p')\text{-H}}^{qp'/2} - \left\langle \|\mathbb{W}^{i,\perp}(\omega, \cdot)\|_{[0,T],(2/p')\text{-H}}^{qp'/2} \right\rangle_1 \right) \geq \frac{1}{2q+1}\right) \\ & \quad + \mathbb{P}\left(\omega : \|\mathbb{W}^{i,i}(\omega)\|_{[0,T],(2/p')\text{-H}}^{qp'/2} \geq \frac{n}{2q+1}\right). \end{aligned}$$

The last term in the right-hand side is easily handled. As for the penultimate one, we proceed as before, recalling that  $\mathbb{W}^{i,j} = \mathcal{I}(W^i, W^j)$ . By Markov's inequality,

$$\begin{aligned} & \mathbb{P}\left(\omega : \frac{1}{n} \sum_{j:j \neq i} \left( \|\mathbb{W}^{i,j}(\omega)\|_{[0,T],(2/p')\text{-H}}^{qp'/2} - \left\langle \|\mathbb{W}^{i,\perp}(\omega, \cdot)\|_{[0,T],(2/p')\text{-H}}^{qp'/2} \right\rangle_1 \right) \geq \frac{1}{2q+1}\right) \\ & \leq \frac{2^{(q+1)}}{n^s} \mathbb{E}\left[\mathbb{E}\left[\left(\sum_{j=1, j \neq i}^n \left( \|\mathcal{I}(W^i(\omega), W^j(\omega))\|_{[0,T],(2/p')\text{-H}}^{qp'/2} \right. \right. \right. \right. \\ & \quad \left. \left. \left. - \left\langle \|\mathcal{I}(W^i(\omega), W^j(\cdot))\|_{[0,T],(2/p')\text{-H}}^{qp'/2} \right\rangle_1 \right) \right)^s \mid \sigma(W^i(\cdot))\right]\right]. \end{aligned}$$

We apply Rosenthal's inequality once again, but conditional on  $\sigma(W^i(\cdot))$ . Assuming without any loss of generality that  $s \geq 2$ , we obtain, for a constant  $C$  independent of  $n$ ,

$$\begin{aligned} & \mathbb{P}\left(\omega : \frac{1}{n} \sum_{j:j \neq i} \left( \|\mathbb{W}^{i,j}(\omega)\|_{[0,T],(2/p')\text{-H}}^{qp'/2} - \left\langle \|\mathbb{W}^{i,\perp}(\omega, \cdot)\|_{[0,T],(2/p')\text{-H}}^{qp'/2} \right\rangle_1 \right) \geq \frac{1}{2q+1}\right) \\ & \leq \frac{C}{n^{s/2}} \mathbb{E}\left[\mathbb{E}\left[\|\mathcal{I}(W^i(\omega), W^j(\omega))\|_{[0,T],(2/p')\text{-H}}^{sqp'/2} \mid \sigma(W^i(\cdot))\right]\right], \end{aligned}$$

where  $j$  right above is any arbitrary integer in  $\{1, \dots, n\} \setminus \{i\}$ . We easily deduce that, for any  $s > 0$ ,  $\sup_{i=1, \dots, n} \mathbb{P}((A_2^{i,n})^c) \leq Cn^{-s}$  for a constant  $C$  independent of  $n$ .

All and all, back to the definition of  $A^n$  at the end of the fifth step, see (5.28), we deduce that, for any  $s > 0$ ,  $\mathbb{P}((A^n)^c) \leq Cn^{-s}$ . Therefore, in order to conclude, it suffices to prove that, for any  $r \geq 1$ , we can choose  $q(r) \geq 8$  such that, if  $q \geq q(r)$  and  $X_0(\cdot)$  is in  $L^q$ , then

$$\sup_{1 \leq i \leq n} \mathbb{E}\left[\sup_{0 \leq t \leq T} |X_t^i|^r\right] \leq C(r), \quad (5.29)$$

for a constant  $C(r)$  depending on  $r$  but independent of  $n$ .

The proof of (5.29) relies on the final estimate in the statement of Theorem 3.3. To make it clear, we consider a new random dissection  $0 = \tau_0 < \tau_1 < \dots < \tau_M = T$  of  $[0, T]$  (for simplicity, we use the same notation as in the previous step, but the new dissection has in fact nothing to do with the first one; in

particular, it is random) such that

$$\begin{aligned} \Gamma_1^{(n)}(\omega, [\tau_\ell, \tau_{\ell+1}]) &:= {}^{(n)}\left(\widehat{N}^{\bullet, n}([\tau_\ell, \tau_{\ell+1}], \omega, 1/(4L_0))\right)_8 \leq 1, \\ \Gamma_2^{(n)}(\omega, [\tau_\ell, \tau_{\ell+1}]) & \\ &:= {}^{(n)}\left([\gamma^2 (1 + \widehat{w}^{\bullet, n}(0, T, \omega)^{1/p'})]\widehat{N}^{\bullet, n}([\tau_\ell, \tau_{\ell+1}], \omega, 1/(4L))\right)_{32} \leq \eta, \end{aligned} \quad (5.30)$$

for the same constants as in the statement of Theorem 3.3. We deduce from Theorem 3.3 that there exists a constant  $C$  (independent of  $n$ ) such that, for any  $i \in \{1, \dots, n\}$  and  $\ell \in \{0, \dots, M-1\}$ ,

$$\|X^i(\omega)\|_{[\tau_\ell, \tau_{\ell+1}], \widehat{w}^{i, n}, p'} \leq \left[C\left(1 + \widehat{w}^{i, n}(0, T, \omega)^{1/p'}\right)\right]^{2\widehat{N}^{i, n}([0, T], \omega, 1/(4L))}.$$

Observe now that, for any  $i \in \{1, \dots, n\}$ ,

$$\begin{aligned} \sup_{0 \leq t \leq T} |X_t^i - X_0^i| &\leq \sum_{\ell=0}^{M-1} \left(\|X^i(\omega)\|_{[\tau_\ell, \tau_{\ell+1}], \widehat{w}^{i, n}, p'} \widehat{w}^{i, n}(\tau_\ell, \tau_{\ell+1}, \omega)^{1/p'}\right) \\ &\leq M \left[C\left(1 + \widehat{w}^{i, n}(0, T, \omega)^{1/p'}\right)\right]^{2\widehat{N}^{i, n}([0, T], \omega, 1/(4L))+1}. \end{aligned}$$

The second factor in the right-hand side has finite moments of any order, see (5.33) below, replacing therein  $\widehat{N}_\ell^{i, n}/\sqrt{\delta_\ell}$  by  $N([0, T], \omega, 1/(4L))/\sqrt{T}$ . Moreover, we prove below that  $M$  has sub-exponential tails, *i.e.*,  $\mathbb{P}(M > a) \leq c \exp(-a^\varepsilon)$ , for  $c, \varepsilon > 0$ . This suffices to prove (5.29).

We now prove that  $(\tau_\ell)_{\ell=0, \dots, M}$  in (5.30) may be constructed in such a way that  $M$  has sub-exponential tails. Obviously, see for instance (A.1), it suffices to construct, for each constraint in (5.30), a subdivision  $(\tau_\ell)_{\ell=0, \dots, M}$  of  $[0, T]$ , for which the corresponding constraint in (5.30) (and only this one) holds true and the number of points  $M$  has sub-exponential tails.

We start with the second constraint in (5.30). By induction, we define the sequence  $(\tau'_\ell)_{\ell=0, \dots, M'}$ , letting  $\tau'_0 := 0$  and  $\tau'_{\ell+1} := \inf\{t \geq \tau'_\ell : \Gamma_2^{(n)}(\omega, [\tau'_\ell, t]) \geq \eta\} \wedge T^4$ , with  $M' := \inf_{\ell \in \mathbb{N}}\{\ell \in \mathbb{N} : \tau'_\ell = T\}$ . We claim that we may choose  $M = 2M'$ . Indeed, since the counter  $\widehat{N}^{i, n}$  appearing in (5.30) is the local accumulation of a continuous function on  $\mathcal{S}_2^T$ , there exists  $\delta_L > 0$  such that, for any  $t \in [0, T]$  and any  $i \in \{1, \dots, n\}$ ,  $\widehat{N}^{i, n}([t, (t + \delta_L) \wedge T], \omega, 1/(4L)) = 0$ . (Of course,  $\delta_L$  depends on  $n$  and  $\omega$ , but this is not a problem in the rest of the proof.) Then, for any point  $t \in [\tau'_\ell, \tau'_{\ell+1})$ , we have, by definition of  $\tau'_{\ell+1}$ ,  $\Gamma_2^{(n)}(\omega, [\tau'_\ell, t]) < \eta$ . Moreover, if  $|\tau'_{\ell+1} - t| < \delta_L$ , then  $\Gamma_2^{(n)}(\omega, [t, \tau'_{\ell+1}]) = 1 \leq \eta$ . Therefore, we may choose  $\tau_{2\ell} = \tau'_\ell$  for  $\ell \in \{0, \dots, M'\}$  and then  $|\tau_{2\ell+1} - \tau'_{\ell+1}| \leq \delta_L$ . The sequence  $(\tau_\ell)_{\ell=0, \dots, 2M}$  satisfies the second constraint in (5.30).

<sup>4</sup>The reader may compare with (2.16), paying attention to the fact that, here,  $t \mapsto \Gamma_2^{(n)}(\omega, [\tau'_\ell, t])$  is not continuous.

We now prove that  $M'$  has sub-exponential tails (which implies that  $M = 2M'$  also has sub-exponential tails). Letting  $\delta_\ell := \tau'_{\ell+1} - \tau'_\ell$ , for any  $\ell \in \mathbb{N}$ , we have, for any  $A > 1$  (recalling  $\gamma, \eta > 1$ ),

$$\begin{aligned} \pi' &:= \mathbb{P} \left( \delta_\ell < \frac{1}{A}, \ell < M' - 1 \right) \\ &\leq \mathbb{P} \left( {}^{(n)} \left[ \gamma^2 \left( 1 + \widehat{w}^{\bullet, n}(0, T, \omega)^{1/p'} \right) \right]_{\widehat{N}_\ell^{\bullet, n}(\omega)/\sqrt{\delta_\ell}}^{1/\sqrt{A}} \geq \eta \right) \\ &= \mathbb{P} \left( {}^{(n)} \left[ \gamma^2 \left( 1 + \widehat{w}^{\bullet, n}(0, T, \omega)^{1/p'} \right) \right]_{\widehat{N}_\ell^{\bullet, n}(\omega)/\sqrt{\delta_\ell}} \geq \eta \sqrt{A} \right), \end{aligned}$$

with the shorten notation  $\widehat{N}_\ell^{\bullet, n}(\omega) = \widehat{N}^{\bullet, n}([\tau'_\ell, \tau'_{\ell+1}], \omega, 1/(4L))$ . We now introduce the function  $f(x) = \exp(\ln(x)^{1+\varepsilon})$ ,  $x > 1$ ; it is non-decreasing on  $[1, \infty)$  and convex on  $[e, \infty)$ . By Markov inequality,

$$\begin{aligned} \pi' &\leq e^{-(\ln[\eta^{32\sqrt{A}}])^{1+\varepsilon}} \mathbb{E} \left[ f \left( \frac{1}{n} \sum_{i=1}^n e \left[ \gamma^2 \left( 1 + \widehat{w}^{i, n}(0, T, \cdot)^{1/p'} \right) \right]^{32\widehat{N}_\ell^{i, n}/\sqrt{\delta_\ell}} \right) \right] \\ &\leq e^{-(\ln[\eta^{32\sqrt{A}}])^{1+\varepsilon}} \frac{1}{n} \sum_{i=1}^n \mathbb{E} \left[ f \left( e \left[ \gamma^2 \left( 1 + \widehat{w}^{i, n}(0, T, \cdot)^{1/p'} \right) \right]^{32\widehat{N}_\ell^{i, n}/\sqrt{\delta_\ell}} \right) \right], \end{aligned}$$

with  $e = \exp(1)$ . We prove in (5.32) below that, for  $\varepsilon$  small enough,

$$\sup_{i=1, \dots, n} \mathbb{E} \left[ f \left( e \left[ \gamma^2 \left( 1 + \widehat{w}^{i, n}(0, T, \cdot)^{1/p'} \right) \right]^{32\widehat{N}_\ell^{i, n}/\sqrt{\delta_\ell}} \right) \right] \leq C, \quad (5.31)$$

for  $C$  independent of  $n$ . As a result,  $\pi' \leq C \exp(-(32 \ln(\eta))^{1+\varepsilon} A^{(1+\varepsilon)/2})$ , and then,

$$\begin{aligned} \mathbb{P}(M' > \ell + 1) &= \mathbb{P}(\delta_1 + \dots + \delta_\ell < T, \ell + 1 < M') \\ &\leq \sum_{i=1}^{\ell} \mathbb{P} \left( \delta_i < \frac{T}{\ell}, i + 1 < M' \right) \leq C \ell e^{-(32 \ln(\eta))^{1+\varepsilon} (\ell/T)^{(1+\varepsilon)/2}}, \end{aligned}$$

which shows that  $M'$  has sub-exponential tails.

We now check what happens when handling the first constraint in (5.30). We may define  $M'$  as before, that is  $M' := \inf_{\ell \in \mathbb{N}} \{\ell \in \mathbb{N} : \tau'_\ell = T\}$  with  $\tau'_0 := 0$  and  $\tau'_{\ell+1} := \inf\{t \geq \tau'_\ell : \Gamma_1^{(n)}(\omega, [\tau'_\ell, t]) \geq 1\} \wedge T$ . Then, we can repeat the same proof as above by using the fact that

$$\left\{ \delta_\ell < \frac{1}{A}, \ell < M' - 1 \right\} \subset \left\{ {}^{(n)} \left( \frac{\widehat{N}^{\bullet, n}([\tau'_\ell, \tau'_{\ell+1}], \omega, 1/(4L_0))}{\sqrt{\delta_\ell}} \right)_8 \geq \sqrt{A} \right\}$$

and by recalling that

$${}^{(n)} \left( \frac{\widehat{N}^{\bullet, n}([\tau'_\ell, \tau'_{\ell+1}], \omega, 1/(4L_0))}{\sqrt{\delta_\ell}} \right)_8$$

has Weibull tails with shape parameter strictly greater than 1<sup>5</sup>, uniformly in the choice of the dissection  $0 = \tau_0 < \dots < \tau_{M'} = T$ , which follows from the third item in the assumption of Theorem 5.1 together with the convexity of the function  $[0, +\infty) \ni x \mapsto \exp(x^{1+\varepsilon})$ , for  $\varepsilon > 0$ . This permits to provide an upper bound for  $\mathbb{P}(\delta_\ell < 1/A)$  and then to deduce as before that  $M'$  has sub-exponential tails.

It now remains to prove (5.31). By (4.24) and (4.25), we can find a real  $\varepsilon_1 > 0$ , independent of  $n$ , such that  $\sup_{i=1, \dots, n} \mathbb{E}[\exp(\widehat{w}^{i,n}(0, T, \cdot)^{\varepsilon_1})] \leq C$ , for  $C$  independent of  $n$ . Hence, combining with the third item in the assumption of the statement, we get, for any  $n \geq 1$ ,  $i \in \{1, \dots, n\}$ ,  $a > 1$  and  $K > 0$ ,

$$\begin{aligned} & \mathbb{P}\left(\left(1 + \widehat{w}^{i,n}(0, T, \cdot)^{1/p'}\right)^{\widehat{N}_\ell^{i,n}/\sqrt{\delta_\ell}} \geq a\right) \\ & \leq \mathbb{P}\left(\frac{\widehat{N}_\ell^{i,n}}{\sqrt{\delta_\ell}} \geq K\right) + \mathbb{P}\left(1 + \widehat{w}^{i,n}(0, T, \cdot)^{1/p'} \geq a^{1/K}\right) \\ & \leq ce^{-K^{1+\varepsilon_2}} + ce^{-a^{\varepsilon_1 p'/K}}, \end{aligned} \quad (5.32)$$

for a new constant  $c$  independent of  $n$  and  $i$ . Choosing  $K = (\ln a)^{1/(1+\varepsilon_2/2)}$ , we deduce that there exist a constant  $c > 1$  and an exponent  $\varepsilon > 0$  such that, for any  $a > 0$ ,

$$\mathbb{P}\left(\left(1 + \widehat{w}^{i,n}(0, T, \cdot)^{1/p'}\right)^{\widehat{N}_\ell^{i,n}/\sqrt{\delta_\ell}} \geq a\right) \leq ce^{-c^{-1} \ln(a)^{1+2\varepsilon}}, \quad (5.33)$$

from which we obtain (5.31).  $\square$

## Appendix A: Integrability and Auxiliary Estimates

We prove in this appendix auxiliary results that we left aside in the body of the text to keep focused on the main problems at hand. In Appendix A.1, we show that assumption (c) in Theorem 5.1 holds true for interacting particle system driven by Gaussian rough paths satisfying, see Example 4.2 together with Remark 5.2. Appendix A.2 is dedicated to proving a crucial moment estimate for some quantity of interest in the proof of the convergence rate in the propagation of chaos result, Theorem 5.1. *This is where the convergence rate  $\zeta_n$  appears.* In the last Appendix A.3, we elaborate on the versions of law of large numbers used in the text.

### A.1. Gaussian Case

Remark 5.2 asserts that the assumptions of Theorem 4.3 are satisfied in the Gaussian framework specified in Example 4.2. Since the derivation of (4.11) is

<sup>5</sup>Recall that a positive random variable  $A$  has a Weibull tail with shape parameter  $2/\varrho$  if  $A^{1/\varrho}$  has a Gaussian tail.

already justified in the latter example, we only prove (here that we can control the empirical local accumulation as the other requirements (c) in the statement of Theorem 5.1. Following the proof of [1, Theorem 2.6], we may focus on the local accumulation of each of the various terms in (5.3). To make it clear, we have the following property: For a given threshold  $\alpha > 0$  and for any two continuous functions  $v_1 : \mathcal{S}_2^T \rightarrow \mathbb{R}_+$  and  $v_2 : \mathcal{S}_2^T \rightarrow \mathbb{R}_+$ , set  $N_i(\alpha) := N_{v_i}([0, T], \alpha)$ , for  $1 \leq i \leq 2$ ; see (2.17) for the original definition, then

$$\max\left(N_1\left(\frac{\alpha}{2}\right), N_2\left(\frac{\alpha}{2}\right)\right) \geq N(\alpha). \quad (\text{A.1})$$

Throughout the proof, we choose  $\Omega$  as the space  $\mathcal{W} = \mathcal{C}([0, T]; \mathbb{R}^d)$ . We call  $\mathcal{H}$  the corresponding Cameron-Martin space and we regard  $(\mathcal{W}, \mathcal{H}, \mathbb{P})$  as an abstract Wiener space. We then regard  $(W^1, \dots, W^n)$  as the canonical process on  $\Omega^n$  equipped with the product measure  $\mathbb{P}^{\otimes n}$ . We recall from [19, Theorem 10.4] that the processes  $(\mathbb{W}^i)_{1 \leq i \leq n}$  and  $(\mathbb{W}^{i,j})_{1 \leq i, j \leq n}$  may be regarded as random variables on  $\Omega^n$ .

**Step 1.** The first step is to consider, for a given  $\alpha > 0$ , the accumulation  $\tilde{N}^i([0, T], \omega, \alpha)$  associated with  $\|W^i(\omega)\|_{[s,t], p-v}^p + \|\mathbb{W}^i(\omega)\|_{[s,t], p/2-v}^{p/2}$ , see (4.3), namely

$$\tilde{N}^i([0, T], \omega, \alpha) := N_{\varpi}([0, T], \alpha),$$

when

$$\varpi(s, t)^p = \|W^i(\omega)\|_{[s,t], p-v}^p + \|\mathbb{W}^i(\omega)\|_{[s,t], p/2-v}^{p/2},$$

but this follows from the proof of [1, Theorem 2.6]. The term  $\hat{v}_{p'}^{i,n}(s, t, \omega)$  in (5.3) is handled by the same argument.

**Step 2.** We now focus on the local accumulation of the fourth and fifth terms in (4.3). For simplicity, we just explain what happens for the fourth term. The fifth term may be handled in the same way.

We use the same notation as in Subsection 4.1 and proceed as in the proof of [1, Theorem 2.6]. The Gaussian process  $(W^1, \dots, W^n)$  has  $(\mathcal{W}^n, \mathcal{H}^{\oplus n}, \mathbb{P}^{\otimes n})$  as abstract Wiener space. For  $\omega = (\omega_i)_{i=1}^n \in \Omega^n$  and for  $\mathbf{h} = \bigoplus_{i=1}^n h_i \in \mathcal{H}^{\oplus n}$ , we let

$$T_{\mathbf{h}} \mathbf{W}^{(n)}(\omega) = T_{\bigoplus_{i=1}^n h_i} \mathbf{W}^{(n)}(\omega)$$

for the translated rough path along  $\mathbf{h}$  (see [19, (11.5)]). By [19, Lemma 11.4], with probability 1 under  $\mathbb{P}^{\otimes n}$ , for all  $\mathbf{h} \in \mathcal{H}^{\oplus n}$ ,

$$\begin{aligned} \|\mathbb{W}^{i,j}(\omega)\|_{[s,t], (p/2)-v}^{p/2} &\leq c \left( \|(T_{\mathbf{h}} \mathbb{W})^{i,j}(\omega)\|_{[s,t], (p/2)-v}^{p/2} + \|(T_{\mathbf{h}} W)^i(\omega)\|_{[s,t], p-v}^p \right. \\ &\quad \left. + \|(T_{\mathbf{h}} W)^j(\omega)\|_{[s,t], p-v}^p + \|h_i\|_{[s,t], \varrho-v}^p + \|h_j\|_{[s,t], \varrho-v}^p \right). \end{aligned}$$

Importantly, the constant  $c$  is independent of  $n$ . Below, it is allowed to increase from line to line as long as it remains independent of  $n$ . So,

$$\begin{aligned}
 & \binom{(n)}{\left\| \mathbb{W}^{i,\bullet}(\omega) \right\|_{[s,t],(p/2)-v}^{p/2}}_q \\
 & \leq c \left\{ \binom{(n)}{\left\| (T\mathbf{h}\mathbb{W})^{i,\bullet}(\omega) \right\|_{[s,t],(p/2)-v}^{p/2}}_q + \binom{(n)}{\left\| (T\mathbf{h}\mathbb{W})^\bullet(\omega) \right\|_{[s,t],p-v}^p}_q \right. \\
 & \quad \left. + \left\| (T\mathbf{h}\mathbb{W})^i(\omega) \right\|_{[s,t],p-v}^p + \|h_i\|_{[s,t],\varrho-v}^p + \binom{(n)}{\left\| h_\bullet \right\|_{[s,t],\varrho-v}^p}_q \right\} \quad (\text{A.2}) \\
 & \leq c \left\{ \binom{(n)}{\left\| (T\mathbf{h}\mathbb{W})^{i,\bullet}(\omega) \right\|_{[0,T],(1/p)-H}^p}_q (t-s) + \|h_i\|_{[s,t],\varrho-v}^p \right. \\
 & \quad \left. + \binom{(n)}{\left\| h_\bullet \right\|_{[s,t],\varrho-v}^p}_q \right\},
 \end{aligned}$$

where for any  $i, j \in \{1, \dots, n\}$ , we let

$$\left\| \mathbf{W}^{i,j}(\omega) \right\|_{[s,t],(1/p)-H} := \left\| (W^i, W^j)(\omega) \right\|_{[s,t],(1/p)-H} + \sqrt{\left\| \mathbb{W}^{i,j}(\omega) \right\|_{[s,t],(2/p)-H}},$$

and similarly for  $\left\| (T\mathbf{h}\mathbb{W})^{i,j}(\omega) \right\|_{[0,T],(1/p)-H}$ .

The tricky term in (A.2) is the last one on the last line. The key point is to notice that, for a given  $\varepsilon \in (0, 2 - \rho)$ ,

$$\begin{aligned}
 & \binom{(n)}{\left\| h_\bullet \right\|_{[s,t],\varrho-v}^p}_q = \left[ \frac{1}{n} \sum_{j=1}^n \|h_j\|_{[s,t],\varrho-v}^{pq} \right]^{1/q} \\
 & = n^{-1/q} \left\{ \left[ \sum_{j=1}^n \|h_j\|_{[s,t],\varrho-v}^{pq} \right]^{(2-\varepsilon)/(pq)} \right\}^{p/(2-\varepsilon)} \leq n^{-1/q} \left[ \sum_{j=1}^n \|h_j\|_{[s,t],\varrho-v}^{2-\varepsilon} \right]^{p/(2-\varepsilon)},
 \end{aligned}$$

where we used the fact that  $2 - \varepsilon < pq$ . Observe in particular that, whenever  $\sum_{j=1}^n \|h_j\|_{[s,t],\varrho-v}^{2-\varepsilon} \leq n^{(2-\varepsilon)/(pq)}$ , it holds

$$\begin{aligned}
 & \binom{(n)}{\left\| h_\bullet \right\|_{[s,t],\varrho-v}^p}_q \leq n^{-1/q} \left[ \sum_{j=1}^n \|h_j\|_{[s,t],\varrho-v}^{2-\varepsilon} \right]^{p/(2-\varepsilon)} \\
 & \leq n^{-1/q} \left( n^{(2-\varepsilon)/(pq)} \right)^{p/(2-\varepsilon)-1} \sum_{j=1}^n \|h_j\|_{[s,t],\varrho-v}^{2-\varepsilon} = n^{-(2-\varepsilon)/(pq)} \sum_{j=1}^n \|h_j\|_{[s,t],\varrho-v}^{2-\varepsilon},
 \end{aligned}$$

where, in the second line, we used the fact that  $p/(2-\varepsilon) > 1$ . Returning to (A.2), we deduce that, whenever  $\|h_i\|_{[s,t],\varrho-v} \leq 1$  and  $\sum_{j=1}^n \|h_j\|_{[s,t],\varrho-v}^{2-\varepsilon} \leq n^{(2-\varepsilon)/(pq)}$ ,

$$\begin{aligned}
 & \binom{(n)}{\left\| \mathbb{W}^{i,\bullet}(\omega) \right\|_{[s,t],(p/2)-v}^{p/2}}_q \leq c \left\{ \binom{(n)}{\left\| (T\mathbf{h}\mathbb{W})^{i,\bullet}(\omega) \right\|_{[0,T],(1/p)-H}^p}_q (t-s) \right. \\
 & \quad \left. + \|h_i\|_{[s,t],\varrho-v}^{2-\varepsilon} + n^{-(2-\varepsilon)/(pq)} \sum_{j=1}^n \|h_j\|_{[s,t],\varrho-v}^{2-\varepsilon} \right\}. \quad (\text{A.3})
 \end{aligned}$$

When the left-hand side is less than or equal to  $\alpha^p$ , we can modify the constant  $c$  in such a way that the inequality remains true when  $\|h_i\|_{[s,t],\varrho-\nu} \geq 1$  or  $\sum_{j=1}^n \|h_j\|_{[s,t],\varrho-\nu}^{2-\varepsilon} \geq n^{(2-\varepsilon)/(pq)}$ . Noticing that  $2 - \varepsilon > \rho$ , (A.3) remains true with  $({}^{(n)}\mathbb{W}^{i,\bullet}(\omega))_{q;[s,t],(p/2)-\nu}^{p/2}$  in the left-hand side.

Define now  $N^{i,n,\perp}([0, T], \omega, \alpha) := N_{\varpi}([0, T], \alpha)$ , when

$$\varpi(s, t)^p = ({}^{(n)}\mathbb{W}^{i,\bullet}(\omega))_{q;[s,t],(p/2)-\nu}^{p/2}.$$

Then, (A.3) (together with  $2 - \varepsilon > \rho$ ) yields

$$\begin{aligned} N^{i,n,\perp}([0, T], \omega, \alpha)\alpha^p &\leq c \left\{ ({}^{(n)}\left( \left\| (T\mathbf{h}\mathbf{W})^{i,\bullet}(\omega) \right\|_{[0,T],(1/p)-\mathbb{H}}^p \right)_q T \right. \\ &\quad \left. + \|h_i\|_{[0,T],\varrho-\nu}^{2-\varepsilon} + n^{-(2-\varepsilon)/(pq)} \sum_{j=1}^n \|h_j\|_{[0,T],\varrho-\nu}^{2-\varepsilon} \right\} \\ &\leq c \left\{ ({}^{(n)}\left( \left\| (T\mathbf{h}\mathbf{W})^{i,\bullet}(\omega) \right\|_{[0,T],(1/p)-\mathbb{H}}^p \right)_q T + \|h_i\|_{[0,T],\varrho-\nu}^{2-\varepsilon} \right. \\ &\quad \left. + n^{-(2-\varepsilon)/(pq)+\varepsilon/2} \left[ \sum_{j=1}^n \|h_j\|_{[0,T],\varrho-\nu}^2 \right]^{(2-\varepsilon)/2} \right\}, \end{aligned}$$

where we applied Hölder's inequality to handle the last term. By choosing  $\varepsilon$  small enough such that  $(2 - \varepsilon)/(pq) - \varepsilon/2 > 0$  and by applying Proposition 11.2 in [19], we get, for a possibly new value of the constant  $c$ ,

$$\begin{aligned} N^{i,n,\perp}([0, T], \omega, \alpha)\alpha^p &\leq c \left\{ ({}^{(n)}\left( \left\| (T\mathbf{h}\mathbf{W})^{i,\bullet}(\omega) \right\|_{[0,T],(1/p)-\mathbb{H}}^p \right)_q T + \|\mathbf{h}\|_{\mathcal{H}^{\oplus n}}^{2-\varepsilon} T^{(2-\varepsilon)/(2\rho)} \right\}, \quad (\text{A.4}) \end{aligned}$$

with  $\|\mathbf{h}\|_{\mathcal{H}^{\oplus n}}^2 = \sum_{i=1}^n \|h_i\|_{\mathcal{H}}^2$ . We then notice that  $(2 - \varepsilon)/(2\rho) > 1/2$  since  $2 - \varepsilon > \rho$ . We deduce that  $T^{(2-\varepsilon)/(2\rho)} \leq cT^{1/2}$  for a possibly new value of the constant  $c$ . We then apply Theorem 11.7 in [19] but on the space  $(\mathcal{W}^{\otimes n}, \mathcal{H}^{\oplus n}, \mathbb{P}^{\otimes n})$ . Importantly, we observe that

$$\mathbb{E} \left[ ({}^{(n)}\left( \left\| \mathbf{W}^{i,\bullet}(\omega) \right\|_{[0,T],(1/p)-\mathbb{H}}^p \right)_q \right]$$

is bounded by a constant  $c$ , independent of  $i$  and  $n$ , which proves that  $N^{i,n,\perp}([0, T], \cdot, \alpha)/\sqrt{T}$  has a Weibull distribution with shape parameter  $2/(2 - \varepsilon)$ , independently of  $n$ .

**Step 3.** We now turn to the local accumulation of the sixth term in (4.3). Taking the norm  $({}^{(n)}\cdot)_q$  in (A.2), we get, with probability 1 under  $\mathbb{P}^{\otimes n}$ , for all  $\mathbf{h} \in \mathcal{H}^{\oplus n}$ ,

$$\begin{aligned} &({}^{(n)}\left( \left\| \mathbb{W}^{\bullet,\bullet}(\omega) \right\|_{[s,t],(p/2)-\nu}^{p/2} \right)_q \\ &\leq c \left\{ ({}^{(n)}\left( \left\| (T\mathbf{h}\mathbf{W})^{\bullet,\bullet}(\omega) \right\|_{[0,T],(1/p)-\mathbb{H}}^p \right)_q (t - s) + ({}^{(n)}\left( \|h_{\bullet}\|_{[s,t],\varrho-\nu}^p \right)_q \right\}. \end{aligned}$$

Following the proof of (A.3), we deduce that

$$\begin{aligned} {}^{(n)}\left(\left[\mathbb{W}^{\bullet,\bullet}(\omega)\right]_{q;[s,t],p/2-\nu}^{p/2}\right) &\leq c \left\{ {}^{(n)}\left(\left[\left[\left(T_{\mathbf{h}}\mathbf{W}\right)^{\bullet,\bullet}(\omega)\right]_{[0,T],(1/p)-\mathbf{H}}^p\right]\right)_q (t-s) \right. \\ &\quad \left. + n^{-(2-\varepsilon)/(pq)} \sum_{j=1}^n \|h_j\|_{[s,t],\varrho-\nu}^{2-\varepsilon} \right\}, \end{aligned}$$

at least when the left-hand side is less than or equal to  $\alpha^p$ . Importantly, there is no need to distinguish the coordinate  $i$  of  $\mathbf{h}$  from the other coordinates  $j \neq i$  since the coefficient in front of any  $\|h_j\|_{[s,t],\varrho-\nu}$ ,  $j = 1, \dots, n$ , has the same power decay as  $n$  tends to  $\infty$ . So, the context is simpler than in the previous step and we may conclude in the same way.

Local accumulations associated to the second term in (4.3) and to  $(s, t) \mapsto {}^{(n)}\left(\widehat{v}_{p'}^{\bullet,n}(\omega)\right)_{q;[s,t],1-\nu}$  and  $(s, t) \mapsto {}^{(n)}\left(\widehat{v}_{p'}^{\bullet,n}(\omega)\right)_{q;[s,t],1-\nu}$  in (5.3) are handled in the same way. (As for the latter one, the reader may refer to the proof of [1, Theorem 2.6].)  $\square$

## A.2. An Auxiliary Estimate

We prove in this appendix some auxiliary estimates that were used in Step 1 of the proof of Theorem 5.1. This is where the convergence rate  $\varsigma_n$  in Theorem 5.1 appears. Recall we set  $\varsigma_n = n^{-1/2}$  if  $d = 1$ , and  $\varsigma_n = n^{-1/2} \ln(1+n)$ , if  $d = 2$ , and  $\varsigma_n = n^{-1/d}$ , if  $d \geq 3$ . Recall also definitions (5.6), (5.7), (5.8) and (5.9).

**Lemma A.1.** *Fix  $\varrho \geq 8$ . There exist an exponent  $\varrho'$  and a positive constant  $C$  such that, if  $X_0(\cdot)$  is  $\varrho'$ -integrable, then, for any integers  $1 \leq i \leq n$ , and  $0 \leq r \leq s \leq t \leq T$ , one has*

$$\begin{aligned} \left\langle [F^{i,n}(\cdot) - \overline{F}^i(\cdot)]_{s,t} \right\rangle_{\varrho} &\leq C \varsigma_n \left\langle w^+(s, t, \cdot, \cdot) \right\rangle_{\varrho'}^{1/p}, \\ \left\langle \mathcal{I}_{\{s,t\}}^{i,n,\partial}(\cdot) - \overline{\mathcal{I}}_{\{s,t\}}^{i,\partial}(\cdot) \right\rangle_{\varrho} &\leq C \varsigma_n \left( \left\langle w^+(s, t, \cdot, \cdot) \right\rangle_{\varrho'}^{1/p} + \left\langle w^+(s, t, \cdot, \cdot) \right\rangle_{\varrho'}^{2/p} \right), \\ \left( \int_{\Omega} \left| \frac{1}{n} \sum_{j=1}^n \delta_{\mu} F_s^{i,j,n}(\omega) \mathbb{W}_{s,t}^{j,i}(\omega) - \mathbb{E}[\delta_{\mu} \overline{F}_s^i(\omega, \cdot) \mathbb{W}_{s,t}^{i,\perp}(\cdot, \omega)] \right|^{\rho} d\mathbb{P}(\omega) \right)^{1/\rho} \\ &\quad + \left\langle (\delta_x F_s^{i,n}(\cdot) - \delta_x \overline{F}_s^i(\cdot)) \mathbb{W}_{s,t}^i(\cdot) \right\rangle_{\rho} \leq C \varsigma_n \left\langle w^+(s, t, \cdot, \cdot) \right\rangle_{\varrho'}^{2/p}, \\ \left\langle \left\{ \mathcal{I}_{\{r,s\}}^{i,n,\partial}(\cdot) + \mathcal{I}_{\{s,t\}}^{i,n,\partial}(\cdot) - \mathcal{I}_{\{r,t\}}^{i,n,\partial}(\cdot) \right\} - \left\{ \overline{\mathcal{I}}_{\{r,s\}}^{i,\partial}(\cdot) + \overline{\mathcal{I}}_{\{s,t\}}^{i,\partial}(\cdot) - \overline{\mathcal{I}}_{\{r,t\}}^{i,\partial}(\cdot) \right\} \right\rangle_{\varrho} \\ &\leq C \varsigma_n \left\langle w^+(r, t, \cdot, \cdot) \right\rangle_{\varrho'}^{3/p}, \end{aligned}$$

where  $w^+(r, t, \omega, \omega') := w(r, t, \omega) + \|\mathbb{W}^{\perp}(\omega, \omega')\|_{[r,t],p/2-\nu}^{p/2}$ .

The reason the appearance of the quantity  $w^+$  instead of  $w$ , in the above upper bounds, will appear at the beginning of **Step 2** in the proof.

*Proof.* We directly prove the last inequality in the statement; the first three inequalities follow from the computations. Throughout the proof, we use the following notations. For each  $i \in \{1, \dots, n\}$ , we call  $\bar{w}^i$  the control associated with  $\bar{W}^i(\cdot)$  through identity (2.10). For  $j \in \{1, \dots, n\}$ , we also let

$$\bar{w}^{i,j}(s, t, \omega) := \|\mathbb{W}^{i,j}(\omega)\|_{[s,t],p-v}^p.$$

We make in the course of the proof an intense use of Lemma A.2 below, giving the convergence rate of the empirical measure of a sample of independent and identically distributed random variables towards their common law. By Theorem 3.3 and following (5.33), we know that, under the standing assumption,  $\sup_{0 \leq t \leq T} |X_t(\cdot)|$  and  $\|X(\cdot)\|_{[0,T],w,p}$  are in  $\mathbb{L}^{\rho'}$  as soon as  $X_0(\cdot)$  is in  $\mathbb{L}^{\rho'}$ . We then compute

$$\begin{aligned} & \left\{ \mathcal{I}_{\{r,s\}}^{i,n,\partial}(\omega) + \mathcal{I}_{\{s,t\}}^{i,n,\partial}(\omega) - \mathcal{I}_{\{r,t\}}^{i,n,\partial}(\omega) \right\} - \left\{ \bar{\mathcal{I}}_{\{r,s\}}^{i,\partial}(\omega) + \bar{\mathcal{I}}_{\{s,t\}}^{i,\partial}(\omega) - \bar{\mathcal{I}}_{\{r,t\}}^{i,\partial}(\omega) \right\} \\ &= \left( R_{r,s}^{F^{i,n}}(\omega) - R_{r,s}^{\bar{F}^i}(\omega) \right) W_{s,t}^i(\omega) + \left( \delta_x F_{r,s}^{i,n}(\omega) - \delta_x \bar{F}_{r,s}^i(\omega) \right) \mathbb{W}_{s,t}^i(\omega) \\ &+ \left( \frac{1}{n} \sum_{j=1}^n \delta_\mu F_{r,s}^{i,j,n}(\omega) \mathbb{W}_{s,t}^{j,i}(\omega) - \mathbb{E} \left[ \delta_\mu \bar{F}_{r,s}^i(\omega, \cdot) \mathbb{W}_{s,t}^{i,\perp}(\cdot, \omega) \right] \right), \end{aligned}$$

where

$$\begin{aligned} R_{r,s}^{F^{i,n}}(\omega) &:= F_s^{i,n}(\omega) - F_r^{i,n}(\omega) - \delta_x F_{r,s}^{i,n}(\omega) W_{r,s}^i(\omega) \\ &\quad - \frac{1}{n} \sum_{j=1}^n \delta_\mu F_r^{i,j,n}(\omega) W_{r,s}^j(\omega), \\ R_{r,s}^{\bar{F}^i}(\omega) &:= \bar{F}_s^i(\omega) - \bar{F}_r^i(\omega) - \delta_x \bar{F}_{r,s}^i(\omega) W_{r,s}^i(\omega) \\ &\quad - \mathbb{E} \left[ \delta_\mu \bar{F}_{r,s}^i(\omega, \cdot) W_{r,s}(\cdot) \right]. \end{aligned} \tag{A.5}$$

Following (5.7) and (5.8), we define differentiable functions  $G_x$  and  $G_\mu$  of their arguments setting

$$\begin{aligned} \delta_x F_t^{i,n}(\omega) &:= G_x(\bar{X}_t^i(\omega), \bar{\mu}_t^n(\omega)), \quad \delta_x \bar{F}_t^i(\omega) := G_x(\bar{X}_t^i(\omega), \mathcal{L}(X_t)), \\ \delta_\mu F_t^{i,j,n}(\omega) &:= G_\mu(\bar{X}_t^i(\omega), \bar{\mu}_t^n(\omega))(\bar{X}_t^j(\omega)), \\ \delta_\mu \bar{F}_t^i(\omega, \omega') &:= G_\mu(\bar{X}_t^i(\omega), \mathcal{L}(X_t))(X_t(\omega')). \end{aligned}$$

Finally, we can write the whole difference in the form

$$\begin{aligned}
 & \left\{ \mathcal{I}_{\{r,s\}}^\partial(\omega) + \mathcal{I}_{\{s,t\}}^\partial(\omega) - \mathcal{I}_{\{r,t\}}^\partial(\omega) \right\} - \left\{ \overline{\mathcal{I}}_{\{r,s\}}^\partial(\omega) + \overline{\mathcal{I}}_{\{s,t\}}^\partial(\omega) - \overline{\mathcal{I}}_{\{r,t\}}^\partial(\omega) \right\} \\
 &= (R_{r,s}^{F^{i,n}}(\omega) - R_{r,s}^{\overline{F}^i}(\omega)) W_{s,t}^i(\omega) \\
 &+ \left[ G_x(\overline{X}^i(\omega), \overline{\mu}^n(\omega)) - G_x(\overline{X}^i(\omega), \mathcal{L}(X)) \right]_{r,s} \mathbb{W}_{s,t}^i(\omega) \\
 &+ \frac{1}{n} \sum_{j=1}^n \left[ G_\mu(\overline{X}^i(\omega), \overline{\mu}^n(\omega))(\overline{X}^j(\omega)) - G_\mu(\overline{X}^i(\omega), \mathcal{L}(X))(\overline{X}^j(\omega)) \right]_{r,s} \mathbb{W}_{s,t}^{j,i}(\omega) \\
 &+ \frac{1}{n} \sum_{j=1}^n \left[ G_\mu(\overline{X}^i(\omega), \mathcal{L}(X))(\overline{X}^j(\omega)) \right]_{r,s} \mathbb{W}_{s,t}^{j,i}(\omega) - \mathbb{E} \left[ \delta_\mu \overline{F}_{r,s}^i(\omega, \cdot) \mathbb{W}_{s,t}^{i,\perp}(\cdot, \omega) \right].
 \end{aligned} \tag{A.6}$$

A key fact is that  $G_x$  and  $G_\mu$  are Lipschitz continuous in all the entries, the Lipschitz property in  $\mu$  being understood with respect to  $\mathbf{d}_1$ . Moreover, similar to  $F$  itself, they are jointly continuously differentiable in all the arguments and the derivatives are Lipschitz continuous, the Lipschitz property in  $\mu$  being again understood with respect to  $\mathbf{d}_1$ .

**Step 1.** Observe that

$$\begin{aligned}
 & \left[ G_x(\overline{X}^i(\omega), \overline{\mu}^n(\omega)) \right]_{r,s} \\
 &= \int_0^1 \partial_x G_x \left( \overline{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \overline{\mu}_{r;(r,s)}^{n,\lambda}(\omega) \right) \overline{X}_{r,s}^i(\omega) d\lambda \\
 &+ \frac{1}{n} \sum_{j=1}^n \int_0^1 D_\mu G_x \left( \overline{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \overline{\mu}_{r;(r,s)}^{n,\lambda}(\omega) \right) (\overline{X}_{r;(r,s)}^{j,(\lambda)}(\omega)) \overline{X}_{r,s}^j(\omega) d\lambda \quad (\text{A.7}) \\
 &= \int_0^1 \partial_x G_x \left( \overline{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \overline{\mu}_{r;(r,s)}^{n,\lambda}(\omega) \right) \overline{X}_{r,s}^i(\omega) d\lambda \\
 &+ \int_{\mathbb{R}^{2d}} \left[ \int_0^1 D_\mu G_x \left( \overline{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \overline{\mu}_{r;(r,s)}^{n,\lambda}(\omega) \right) (y) z d\lambda \right] d\overline{\nu}_{r;(r,s)}^{n,\lambda}(\omega; y, z)
 \end{aligned}$$

where

$$\overline{\mu}_{r;(r,s)}^{n,(\lambda)}(\omega) := \frac{1}{n} \sum_{j=1}^n \delta_{\overline{X}_{r;(r,s)}^{j,(\lambda)}(\omega)}, \quad \overline{\nu}_{s;(s,t)}^{n,(\lambda)}(\omega) := \frac{1}{n} \sum_{j=1}^n \delta_{(\overline{X}_{r;(r,s)}^{j,(\lambda)}(\omega), \overline{X}_{r,s}^j(\omega))},$$

with

$$\overline{X}_{r;(r,s)}^{j,(\lambda)}(\omega) := \overline{X}_r^j(\omega) + \lambda \overline{X}_{r,s}^j(\omega).$$

Proceeding similarly with  $[G_x(\bar{X}^i(\omega), \mathcal{L}(X))]_{r,s}$ , we get

$$\begin{aligned}
 & \left[ G_x(\bar{X}^i(\omega), \bar{\mu}^n(\omega)) - G_x(\bar{X}^i(\omega), \mathcal{L}(X)) \right]_{r,s} \\
 &= \int_0^1 \left[ \partial_x G_x \left( \bar{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \bar{\mu}_{r;(r,s)}^{n,(\lambda)}(\omega) \right) \right. \\
 & \quad \left. - \partial_x G_x \left( \bar{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \mathcal{L}(X_{r;(r,s)}^{(\lambda)}) \right) \right] \bar{X}_{r,s}^i(\omega) d\lambda \\
 & + \int_{\mathbb{R}^{2d}} \left[ \int_0^1 D_\mu G_x \left( \bar{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \bar{\mu}_{r;(r,s)}^{n,(\lambda)}(\omega) \right) (y) z d\lambda \right] d\bar{\nu}_{r;(r,s)}^{n,(\lambda)}(\omega; y, z) \\
 & - \int_{\mathbb{R}^{2d}} \left[ \int_0^1 D_\mu G_x \left( \bar{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \mathcal{L}(X_{r;(r,s)}^{(\lambda)}) \right) (y) z d\lambda \right] d\mathcal{L}(X_{r;(r,s)}^{(\lambda)}, X_{r,s})(y, z),
 \end{aligned}$$

where, as before,  $X_{r;(r,s)}^{(\lambda)}(\omega) = X_r(\omega) + \lambda X_{r,s}(\omega)$ . Splitting the last two terms in the above expansion into

$$\begin{aligned}
 & \int_{\mathbb{R}^{2d}} \left[ \int_0^1 D_\mu G_x \left( \bar{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \bar{\mu}_{r;(r,s)}^{n,(\lambda)}(\omega) \right) (y) z d\lambda \right] d\bar{\nu}_{r;(r,s)}^{n,(\lambda)}(\omega; y, z) \\
 & - \int_{\mathbb{R}^{2d}} \left[ \int_0^1 D_\mu G_x \left( \bar{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \mathcal{L}(X_{r;(r,s)}^{(\lambda)}) \right) (y) z d\lambda \right] d\bar{\nu}_{r;(r,s)}^{n,(\lambda)}(\omega; y, z) \\
 & + \int_{\mathbb{R}^{2d}} \left[ \int_0^1 D_\mu G_x \left( \bar{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \mathcal{L}(X_{r;(r,s)}^{(\lambda)}) \right) (y) z d\lambda \right] d\bar{\nu}_{r;(r,s)}^{n,(\lambda)}(\omega; y, z) \\
 & - \int_{\mathbb{R}^{2d}} \left[ \int_0^1 D_\mu G_x \left( \bar{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \mathcal{L}(X_{r;(r,s)}^{(\lambda)}) \right) (y) z d\lambda \right] d\mathcal{L}(X_{r;(r,s)}^{(\lambda)}, X_{r,s})(y, z),
 \end{aligned}$$

we get

$$\begin{aligned}
 & \left| \left[ G_x(\bar{X}^i(\omega), \bar{\mu}^n(\omega)) - G_x(\bar{X}^i(\omega), \mathcal{L}(X)) \right]_{r,s} \right| \\
 & \leq c \int_0^1 \mathbf{d}_1 \left( \bar{\mu}_{r;(r,s)}^{n,(\lambda)}(\omega), \mathcal{L}(X_{r;(r,s)}^{(\lambda)}) \right) d\lambda \\
 & \quad \times \left( \|\bar{X}^i(\omega)\|_{[0,T], \bar{w}^i, p} \bar{w}^i(r, s, \omega)^{1/p} + \frac{1}{n} \sum_{k=1}^n \|\bar{X}^k(\omega)\|_{[0,T], \bar{w}^k, p} \bar{w}^k(r, s, \omega)^{1/p} \right) \\
 & + c \left| \mathcal{S}_{r,s}^{i,n}(\omega, |\bar{X}_{r,s}^\bullet(\omega)|) \right|,
 \end{aligned}$$

where  $\mathcal{S}_{r,s}^{i,n}(\omega, |\bar{X}_{r,s}^\bullet(\omega)|)$  is the  $n$ -empirical mean of  $n$  variables that are dominated by  $(|\bar{X}_{r,s}^j(\omega)|)_{j=1, \dots, n}$  and  $n-1$  of which are conditionally centered and conditionally independent given the realization of the path  $(\bar{X}^i, W^i, \mathbb{W}^i)$ . Recalling (2.9) and allowing the value of the constant  $c$  to increase from line to

line, we obtain

$$\begin{aligned}
 & \left| \left[ G_x \left( \bar{X}^i(\omega), \bar{\mu}^n(\omega) \right) - G_x \left( \bar{X}^i(\omega), \mathcal{L}(X) \right) \right]_{r,s} \mathbb{W}_{s,t}^i(\omega) \right| \\
 & \leq c \int_0^1 \mathbf{d}_1 \left( \bar{\mu}_{r;(r,s)}^{n,(\lambda)}(\omega), \mathcal{L}(X_{r;(r,s)}^{(\lambda)}) \right) d\lambda \\
 & \quad \times \left[ \|\bar{X}^i(\omega)\|_{[0,T],\bar{w}^i,p} + \left( \frac{1}{n} \sum_{k=1}^n \|\bar{X}^k(\omega)\|_{[0,T],\bar{w}^k,p}^2 \right)^{1/2} \right] \\
 & \quad \times \left[ \bar{w}^i(r,t,\omega)^{3/p} + \frac{1}{n} \sum_{k=1}^n \bar{w}^k(r,t,\omega)^{3/p} \right] \\
 & \quad + c \left| \mathcal{S}_{r,s}^{i,n}(\omega, |\bar{X}_{r,s}^\bullet(\omega)|) \right| \bar{w}^i(r,t,\omega)^{2/p}.
 \end{aligned}$$

In order to conclude for the second term in the right-hand side of (A.6), it suffices to recall from Rosenthal's inequality (applied under the conditional probability given the realization of the path  $(\bar{X}^i, W^i, \mathbb{W}^i)$ ) that

$$\begin{aligned}
 \left\langle \mathcal{S}_{r,s}^{i,n}(\cdot, |\bar{X}_{r,s}^\bullet(\cdot)|) \right\rangle_{3\varrho/2} & \leq c n^{-1/2} \left\langle \|\mathcal{X}(\cdot)\|_{[0,T],w,p} w(r,s,\cdot)^{1/p} \right\rangle_{3\varrho/2} \\
 & \leq c n^{-1/2} \left\langle \|\mathcal{X}(\cdot)\|_{[0,T],w,p} \right\rangle_{\chi\varrho} \left\langle w(r,t,\cdot) \right\rangle_{\chi\varrho}^{1/p},
 \end{aligned}$$

where  $\chi \geq 1$  is a universal constant whose value may change from line to line. If  $\rho$  is large enough, we deduce from Lemma A.2 that

$$\begin{aligned}
 & \left\langle \left[ G_x \left( \bar{X}^i(\cdot), \bar{\mu}^n(\cdot) \right) - G_x \left( \bar{X}^i(\cdot), \mathcal{L}(X) \right) \right]_{r,s} \mathbb{W}_{s,t}^i(\cdot) \right\rangle_{\varrho} \\
 & \leq c \left( \int_0^1 \left\langle \mathbf{d}_1 \left( \bar{\mu}_{r;(r,s)}^{n,(\lambda)}(\cdot), \mathcal{L}(X_{r;(r,s)}^{(\lambda)}) \right) \right\rangle_{\chi\varrho} d\lambda \right) \left\langle \|\mathcal{X}(\cdot)\|_{[0,T],w,p} \right\rangle_{\chi\varrho} \\
 & \quad \times \left\langle w(r,t,\cdot) \right\rangle_{\chi\varrho}^{3/p} \\
 & \quad + c n^{-1/2} \left\langle \|\mathcal{X}(\cdot)\|_{[0,T],w,p} \right\rangle_{\chi\varrho} \left\langle w(r,t,\cdot) \right\rangle_{\chi\varrho}^{3/p} \\
 & \leq c \varsigma_n \left( 1 + \left\langle \sup_{0 \leq u \leq T} |\mathcal{X}_u(\cdot)| \right\rangle_{\chi\varrho} \right) \left\langle \|\mathcal{X}(\cdot)\|_{[0,T],w,p} \right\rangle_{\chi\varrho} \left\langle w^+(r,t,\cdot) \right\rangle_{\chi\varrho}^{3/p}.
 \end{aligned}$$

**Step 2.** By the same argument, we have

$$\begin{aligned}
 & \left| \left[ G_\mu \left( \bar{X}^i(\omega), \bar{\mu}^n(\omega) \right) \left( \bar{X}^j(\omega) \right) - G_\mu \left( \bar{X}^i(\omega), \mathcal{L}(X) \right) \left( \bar{X}^j(\omega) \right) \right]_{r,s} \mathbb{W}_{s,t}^{j,i}(\omega) \right| \\
 & \leq c \left( \int_0^1 \mathbf{d}_1 \left( \bar{\mu}_{r;(r,s)}^{n,(\lambda)}(\omega), \mathcal{L}(X_{r;(r,s)}^{(\lambda)}) \right) d\lambda \right) \bar{w}^{j,i}(s, t, \omega)^{2/p} \\
 & \quad \times \left[ \left\| \bar{X}^i(\omega) \right\|_{[0,T], \bar{w}^i, p} + \left\| \bar{X}^j(\omega) \right\|_{[0,T], \bar{w}^j, p} + \left( \frac{1}{n} \sum_{k=1}^n \left\| \bar{X}^k(\omega) \right\|_{[0,T], \bar{w}^k, p}^2 \right)^{1/2} \right] \\
 & \quad \times \left[ \bar{w}^i(r, s, \omega)^{1/p} + \bar{w}^j(r, s, \omega)^{1/p} + \left( \frac{1}{n} \sum_{k=1}^n \bar{w}^k(r, s, \omega)^{2/p} \right)^{1/2} \right] \\
 & \quad + c \left| \mathcal{S}_{r,s}^{i,j,n}(\omega, |\bar{X}_{r,s}^\bullet(\omega)|) \right| \bar{w}^{j,i}(s, t, \omega)^{2/p},
 \end{aligned}$$

where

$$\begin{aligned}
 \left\langle \mathcal{S}_{r,s}^{i,j,n}(\cdot, |\bar{X}_{r,s}^\bullet(\cdot)|) \right\rangle_{3\varrho/2} & \leq c n^{-1/2} \langle \|X\|_{[0,T], w, p} w(r, s, \cdot)^{1/p} \rangle_{3\varrho/2} \\
 & \leq c n^{-1/2} \langle \|X\|_{[0,T], w, p} \rangle_{\chi\varrho} \langle w(r, t, \cdot) \rangle_{\chi\varrho}^{1/p}.
 \end{aligned}$$

Observing that  $\langle \bar{w}^{j,i}(s, t, \cdot)^{2/p} \rangle_{\chi\varrho} \leq \langle w^+(r, t, \cdot) \rangle_{\chi\varrho}^{2/p}$  – this is the rationale for introducing  $w^+$ , and taking expectation, we get

$$\begin{aligned}
 & \left\langle \left[ G_\mu \left( \bar{X}^i(\cdot), \bar{\mu}^n(\cdot) \right) \left( \bar{X}^j(\cdot) \right) - G_\mu \left( \bar{X}^i(\cdot), \mathcal{L}(X) \right) \left( \bar{X}^j(\cdot) \right) \right]_{r,s} \mathbb{W}_{s,t}^{j,i}(\omega) \right\rangle_{\varrho} \\
 & \leq c \left( \int_0^1 \left\langle \mathbf{d}_1 \left( \bar{\mu}_{r;(r,s)}^{n,(\lambda)}(\cdot), \mathcal{L}(X_{r;(r,s)}^{(\lambda)}) \right) \right\rangle_{\chi\varrho} d\lambda \right) \langle \|X(\cdot)\|_{[0,T], w, p} \rangle_{\chi\varrho} \\
 & \quad \times \langle w^+(r, t, \cdot, \cdot) \rangle_{\chi\varrho}^{3/p} \\
 & \quad + c n^{-1/2} \langle \|X(\cdot)\|_{[0,T], w, p} \rangle_{\chi\varrho} \langle w^+(r, t, \cdot, \cdot) \rangle_{\chi\varrho}^{3/p}.
 \end{aligned}$$

Taking the mean over  $j$ , we obtain for upper bound for the third term in the right-hand side of (A.6) the quantity

$$\begin{aligned}
 & \left\langle \frac{1}{n} \sum_{j=1}^n \left[ G_\mu \left( \bar{X}^i(\cdot), \bar{\mu}^n(\cdot) \right) \left( \bar{X}^j(\cdot) \right) - G_\mu \left( \bar{X}^i(\cdot), \mathcal{L}(X) \right) \left( \bar{X}^j(\cdot) \right) \right]_{r,s} \mathbb{W}_{s,t}^{j,i}(\omega) \right\rangle_{\varrho} \\
 & \leq c \left( \int_0^1 \left\langle \mathbf{d}_1 \left( \bar{\mu}_{r;(r,s)}^{n,(\lambda)}(\cdot), \mathcal{L}(X_{r;(r,s)}^{(\lambda)}) \right) \right\rangle_{\chi\varrho} d\lambda \right) \langle \|X(\cdot)\|_{[0,T], w, p} \rangle_{\chi\varrho} \\
 & \quad \times \langle w^+(r, t, \cdot, \cdot) \rangle_{\chi\varrho}^{3/p} \\
 & \quad + c n^{-1/2} \langle \|X(\cdot)\|_{[0,T], w, p} \rangle_{\chi\varrho} \langle w^+(r, t, \cdot, \cdot) \rangle_{\chi\varrho}^{3/p}.
 \end{aligned}$$

By Lemma A.2, we get the same bound as in the first step.

**Step 3.** We now turn to the last term in the right-hand side of (A.6). It reads as the empirical mean of  $n$  random variables,  $n - 1$  of which are conditionally centered and conditionally independent given the realization of the

paths  $(\bar{X}^i, W^i, \mathbb{W}^i)$ , namely

$$\frac{1}{n} \sum_{j=1}^n [G_\mu(\bar{X}^i(\omega), \mathcal{L}(X))(\bar{X}^j(\omega))]_{r,s} \mathbb{W}_{s,t}^{j,i}(\omega) - \mathbb{E}[\delta_\mu \bar{F}_{r,s}^i(\omega, \cdot) \mathbb{W}_{s,t}^{i,\perp}(\cdot, \omega)].$$

Invoking Rosenthal's inequality once again (in a conditional form), it suffices to compute the  $L^p$  norm of  $[G_\mu(\bar{X}^i(\omega), \mathcal{L}(X))(\bar{X}^j(\omega))]_{r,s} \mathbb{W}_{s,t}^{j,i}(\omega)$ . Doing as before (see (A.7)), it is less than  $c \langle \|\bar{X}(\cdot)\|_{[0,T],w,p} \rangle_{\chi_\varrho} \langle \langle w^+(r, t, \cdot, \cdot) \rangle \rangle_{\chi_\varrho}^{3/p}$ . So,

$$\begin{aligned} & \left\langle \frac{1}{n} \sum_{j=1}^n [G_\mu(\bar{X}^i(\omega), \mathcal{L}(X))(\bar{X}^j(\omega))]_{r,s} \mathbb{W}_{s,t}^{j,i}(\omega) - \mathbb{E}[\delta_\mu \bar{F}_{r,s}^i(\omega, \cdot) \mathbb{W}_{s,t}^{i,\perp}(\cdot, \omega)] \right\rangle_e \\ & \leq c n^{-1/2} \langle \|\bar{X}(\cdot)\|_{[0,T],w,p} \rangle_{\chi_\varrho} \langle \langle w^+(r, t, \cdot, \cdot) \rangle \rangle_{\chi_\varrho}^{3/p}, \end{aligned}$$

which suffices to conclude.

**Step 4.** We now handle the remainders in (A.6). By expanding (A.5) and by using similar notations for the remainders in the expansion of each  $(\bar{X}^j)_{j=1, \dots, n}$ , we have (see for instance the proof of [1, Proposition 3.5])

$$\begin{aligned} R_{r,s}^{F^{i,n}}(\omega) &= \partial_x F(\bar{X}_r^i(\omega), \bar{\mu}_r^n(\omega)) R_{r,s}^{\bar{X}^i}(\omega) \\ &+ \frac{1}{n} \sum_{j=1}^n D_\mu F(\bar{X}_r^i(\omega), \bar{\mu}_r^n(\omega))(\bar{X}_r^j(\omega)) R_{r,s}^{\bar{X}^j}(\omega) \\ &+ \int_0^1 [\partial_x F(\bar{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \bar{\mu}_{r;(r,s)}^{n,(\lambda)}(\omega)) - \partial_x F(\bar{X}_r^i(\omega), \bar{\mu}_r^n(\omega))] \bar{X}_{r,s}^i(\omega) d\lambda \quad (\text{A.8}) \\ &+ \frac{1}{n} \sum_{j=1}^n \int_0^1 [D_\mu F(\bar{X}_{r;(r,s)}^{i,(\lambda)}(\omega), \bar{\mu}_{r;(r,s)}^{n,(\lambda)}(\omega))(\bar{X}_{r;(r,s)}^{j,(\lambda)}(\omega)) \\ &\quad - D_\mu F(\bar{X}_r^i(\omega), \bar{\mu}_r^n(\omega))(\bar{X}_r^j(\omega))] \bar{X}_{r,s}^j(\omega) d\lambda. \end{aligned}$$

Expanding  $R_{r,s}^{\bar{X}^i}(\omega)$  in a similar way, we have to investigate four difference terms in order to estimate the difference  $R_{r,s}^{F^{i,n}}(\omega) - R_{r,s}^{\bar{F}^i}(\omega)$ . The first difference term corresponds to the first term in the right-hand side of (A.8)

$$\begin{aligned} & \left| \left[ \partial_x F(\bar{X}_r^i(\omega), \bar{\mu}_r^n(\omega)) - \partial_x F(\bar{X}_r^i(\omega), \mathcal{L}(X_r)) \right] R_{r,s}^{\bar{X}^i}(\omega) \right| \\ & \leq c \mathbf{d}_1(\bar{\mu}_r^n(\omega), \mathcal{L}(X_r)) \|\bar{X}^i(\cdot)\|_{[0,T],\bar{w}^i,p} \bar{w}^i(r, s, \omega)^{2/p}. \end{aligned}$$

Then, we must recall that, in the first line of the right-hand side in (A.6), the difference  $R_{r,s}^{F^{i,n}}(\omega) - R_{r,s}^{\bar{F}^i}(\omega)$  is multiplied by  $W_{s,t}^i(\omega)$ , which is less than  $\bar{w}^i(s, t, \omega)^{1/p}$ . In other words, we must multiply both sides in the above inequality

by  $\bar{w}^i(r, t, \omega)^{1/p}$ . By Cauchy Schwarz inequality, the  $\mathbb{L}^\ell$  norm of the resulting bound is less than

$$c \langle \mathbf{d}_1(\bar{\mu}_r^n(\cdot), \mathcal{L}(X_r)) \rangle_{\chi_\ell} \langle \|X(\cdot)\|_{[0, T], w, p} \rangle_{\chi_\ell} \langle w(r, t, \cdot) \rangle_{\chi_\ell}^{3/p}.$$

The second difference term that we have to handle corresponds to the second term in the right-hand side of (A.8). With an obvious definition for  $R^X(\cdot)$ , it reads

$$\left| \frac{1}{n} \sum_{j=1}^n D_\mu \mathbb{F}(\bar{X}_r^i(\omega), \bar{\mu}_r^n(\omega))(\bar{X}_r^j(\omega)) R_{r,s}^{\bar{X}^j}(\omega) - \langle D_\mu \mathbb{F}(\bar{X}_r^i(\omega), \mathcal{L}(X_r))(X_r(\cdot)) R_{r,s}^X(\cdot) \rangle \right|.$$

Proceeding exactly as in the first step, the latter is bounded by

$$c \mathbf{d}_1(\bar{\mu}_r^n(\omega), \mathcal{L}(X_r)) \left( \frac{1}{n} \sum_{j=1}^n |R_{r,s}^{\bar{X}^j}(\omega)| \right) + c \left| \mathcal{S}_{r,s}^{i,n}(\omega, |R_{r,s}^{\bar{X}^\bullet}(\omega)|) \right|,$$

where  $\mathcal{S}_{r,s}^{i,n}(\omega, |R_{r,s}^{\bar{X}^\bullet}(\omega)|)$  is the  $n$ -empirical mean of  $n$  variables that are dominated by  $(|R_{r,s}^{\bar{X}^j}(\omega)|)_{j=1, \dots, n}$  and  $n-1$  of which are conditionally centered and conditionally independent given the realization of the path  $(\bar{X}^i, W^i, \mathbb{W}^i)$ . Hence, the  $\mathbb{L}^\ell$  norm of the right-hand side, after multiplication as before by  $\bar{w}^i(s, t, \omega)^{1/p}$ , is less than

$$c \left( \langle \mathbf{d}_1(\bar{\mu}_r^n(\cdot), \mathcal{L}(X_r)) \rangle_{\chi_\ell} + n^{-1/2} \right) \langle \|X(\cdot)\|_{[0, T], w, p} \rangle_{\chi_\ell} \langle w(r, t, \cdot) \rangle_{\chi_\ell}^{3/p}.$$

As for the third term in the right-hand side of (A.8), it fits, up to the additional factor  $\bar{X}_{r,s}^i(\omega)$ , the analysis in the first step. So we get as an upper bound for its  $\mathbb{L}^\ell$  norm, after multiplication by  $\bar{w}^i(s, t, \omega)^{1/p}$ , the quantity

$$\begin{aligned} & c \left( \int_0^1 \int_0^1 \langle \mathbf{d}_1(\bar{\mu}_{r;(r,s)}^{n,(\lambda\lambda')}(\cdot), \mathcal{L}(X_{r;(r,s)}^{(\lambda\lambda')})) \rangle_{\chi_\ell} d\lambda d\lambda' \right) \\ & \quad \times \langle \|X(\cdot)\|_{[0, T], w, p} \rangle_{\chi_\ell}^2 \langle w(r, t, \cdot) \rangle_{\chi_\ell}^{3/p} \\ & + c n^{-1/2} \langle \|X(\cdot)\|_{[0, T], w, p} \rangle_{\chi_\ell}^2 \langle w(r, t, \cdot) \rangle_{\chi_\ell}^{3/p}. \end{aligned}$$

Following Step 2, we get exactly a similar bound for the fourth term in the right-hand side of (A.8). Applying once again Lemma A.2 completes the proof.  $\square$

### A.3. About Law of Large Numbers

**Lemma A.2.** *There exists a real  $q_d \geq 1$  such that, for any  $q \geq q_d$  and any probability measure  $\mu$  on  $\mathbb{R}^d$  satisfying  $M_q(\mu) := (\int_{\mathbb{R}^d} |x|^q \mu(dx))^{1/q} < \infty$ , it*

holds

$$\mathbb{E} \left[ \mathbf{d}_1(\mu^n(\cdot), \mu)^{q/3} \right]^{3/q} \leq c_{q,d} M_q(\mu) \varsigma_n,$$

for a constant  $c_{q,d}$  depending on  $q$  and  $d$ , where  $\mu^n(\cdot)$  is the empirical distribution of  $n$  independent identically distributed random variables.

*Proof.* Without any loss of generality, we can assume that  $M_q(\mu) = 1$ , see the argument in [9, Chapter 5]. Then Theorem 2 in [18] gives us the following results. For  $d \geq 3$ , we have

$$\mathbb{P}(\mathbf{d}_1(\mu^n(\cdot), \mu) \geq A\varsigma_n) \leq C \exp(-cn\varsigma_n^d A^d) + Cn(nA\varsigma_n)^{-q/2},$$

in which case the result easily follows. When  $d = 1$ , we have

$$\mathbb{P}(\mathbf{d}_1(\mu^n(\cdot), \mu) \geq A\varsigma_n) \leq C \exp(-cn\varsigma_n^2 A^2) + Cn(nA\varsigma_n)^{-q/2},$$

and the result follows as well by our choice of  $\varsigma_n$ . Finally, when  $d = 2$ ,

$$\mathbb{P}(\mathbf{d}_1(\mu^n(\cdot), \mu) \geq A\varsigma_n) \leq C \exp\left(-\frac{cn\varsigma_n^2 A^2}{(\ln(2 + A^{-1}\varsigma_n^{-1}))^2}\right) + Cn(nA\varsigma_n)^{-q/2}.$$

Assuming without any loss of generality that  $A \geq 1$ , we have

$$\ln(2 + A^{-1}\varsigma_n^{-1}) \leq \ln(2 + \varsigma_n^{-1}) = \ln(1 + 2\varsigma_n) - \ln(\varsigma_n),$$

which is less than  $-2\ln(\varsigma_n)$  for  $n$  large enough. Given our choice of  $\varsigma_n$ , we have  $-\ln(\varsigma_n) = \ln(n)/2 - \ln(\ln(1 + n))$ , which is less than  $\ln(n)/2$ . Hence, modifying the value of the constant  $c$ , we get, for  $A \geq 1$  and for  $n$  large enough, independently of the value of  $A$ , we get the bound

$$\mathbb{P}(\mathbf{d}_1(\mu^n(\cdot), \mu) \geq A\varsigma_n) \leq C \exp\left(-\frac{cA^2 \ln(1 + n)^2}{(\ln(n))^2}\right) + Cn(nA\varsigma_n)^{-q/2},$$

which suffices to complete the proof.  $\square$

**Lemma A.3.** *Let  $(X_n)_{n \geq 1}$  be a collection of independent and identically distributed random variables with values in a Polish space  $S$  and let  $f$  be a real-valued Borel function on  $S^2$  such that  $\mathbb{E}[|f(X_1, X_2)|]$  and  $\mathbb{E}[|f(X_1, X_2)|^2]$  are both finite. Then, with probability 1,*

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} \sum_{i,j=1}^n f(X_i, X_j) = \mathbb{E}[f(X_1, X_2)].$$

*Proof.* By the standard version of the law of large numbers, it suffices to prove that, with probability 1,

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} \sum_{i,j=1, i \neq j}^n f(X_i, X_j) = \mathbb{E}[f(X_1, X_2)].$$

Letting  $S_n = \sum_{i,j=1, i \neq j}^n f(X_i, X_j)$ , for  $n \geq 1$ , we then define the  $\sigma$ -field  $\mathcal{G}_n = \sigma(S_k, k \geq n)$ . By independence of the variables  $(X_k)_{k \geq 1}$ , we have, for any  $(i, j) \in \{1, \dots, n\}^2$  with  $i \neq j$ ,  $\mathbb{E}[f(X_i, X_j)|\mathcal{G}_n] = \mathbb{E}[f(X_i, X_j)|S_n]$ . By exchangeability, this is also equal to  $\mathbb{E}[f(X_1, X_2)|S_n]$ . We get

$$\mathbb{E}[f(X_1, X_2)|\mathcal{G}_n] = \frac{1}{n^2 - n} \sum_{i,j=1, i \neq j}^n \mathbb{E}[f(X_i, X_j)|S_n] = \frac{S_n}{n^2 - n}.$$

By Lévy's downward theorem and by Kolmogorov zero-one law, the left-hand side converges almost surely to  $\mathbb{E}[f(X_1, X_2)]$ .  $\square$

## References

- [1] Bailleul, I., Catellier, R., Delarue, F., Solving mean field rough differential equations. *Submitted*, 2019.
- [2] Bailleul, I., Catellier, R., Delarue, F., Mean field rough differential equations. [arXiv:1802.05882](https://arxiv.org/abs/1802.05882), 2018.
- [3] Bailleul, I., Flows driven by rough paths. *Revista Mat. Iberoamericana*, 31(3):901–934, 2015.
- [4] Billingsley, P., *Convergence of probability measures*. Second Edition. Wiley Series in Probability and Statistics: Probability and Statistics, John Wiley & Sons Inc., 1999.
- [5] Budhiraja, A. and Dupuis, P. and Fischer, M., Large deviation properties of weakly interacting processes via weak convergence methods. *Ann. Probab.*, **40**(1):74–102, 2012.
- [6] Budhiraja, A. and Wu, R., Moderate Deviation Principles for Weakly Interacting Particle Systems. *Probab. Th. Rel. Fields*, **168**(2-4):721–771, 2017.
- [7] Cardaliaguet, P., Notes on mean field games. <https://www.ceremade.dauphine.fr/~cardaliaguet/MFG20130420.pdf>, 2013.
- [8] Carmona, R. and Delarue, F., Forward-backward Stochastic Differential Equations and Controlled McKean Vlasov Dynamics. *Ann. Probab.*, **44**(6):3740–3803, 2016.
- [9] Carmona, R. and Delarue, F. *Probabilistic Theory of Mean Field Games: vol. I, Mean Field FBSDEs, Control, and Games*. Probability Theory and Stochastic Modelling, Springer Verlag, 2018.
- [10] Carmona, R. and Delarue, F. *Probabilistic Theory of Mean Field Games: vol. II, Mean Field Games with Common Noise and Master Equations*. Probability Theory and Stochastic Modelling, Springer Verlag, 2018.
- [11] Cass, T., Litterer, C. and Lyons, T., Integrability and tail estimates for Gaussian rough differential equations. *Ann. Probab.*, **41**(4):3026–3050, 2013.
- [12] Cass, T. and Lyons, T., Evolving communities with individual preferences. *Proc. Lond. Math. Soc. (3)*, **110**(1):83–107, 2015.

- [13] Coghi, M., Deuschel, J.-D., Friz, P. and Maurelli, M., Pathwise McKean-Vlasov theory. [arXiv:1812.11773](https://arxiv.org/abs/1812.11773), 2018.
- [14] Coghi, M. and Flandoli, F., Propagation of chaos for interacting particles subject to environmental noise. *Ann. Appl. Probab.*, **26**(3):1407–1442, 2016.
- [15] Dawson, D. and Gärtner, J., Large deviations from the McKean-Vlasov limit for weakly interacting diffusions. *Stochastics*, **20**:247–308, 1987.
- [16] Dawson, D. and Vaillancourt, J., Stochastic McKean-Vlasov equations. *Nonlinear Diff. Eq. Appl.*, **2**,199–229, 1995.
- [17] Deuschel, J.-D., Friz, P., Maurelli, M. and Slowik, M., The enhanced Sanov theorem and propagation of chaos. *Stoch. Proc. Appl.*, **128**(7):2228–2269, 2018.
- [18] Fournier, N. and Guillin, A., On the rate of convergence in the Wasserstein distance of the empirical measure. *Probab. Theory Related Fields*, **162**:707–738, 2015.
- [19] Friz, P. and Hairer, M., A course on rough paths, with an introduction to regularity structures. *Universitext, Springer*, 2014.
- [20] Friz, P. and Victoir, N., Multidimensional stochastic processes as rough paths. *Cambridge studies in advanced Mathematics*, **120**, 2010.
- [21] Friz, P. and Victoir, N., Differential equations driven by Gaussian signals. *Ann. Inst. H. Poincaré, Probab. Statist*, **46**(2):369–413, 2010.
- [22] Gärtner, J., On the McKean-Vlasov limit for interacting diffusions. *Math. Nachr.*, **137**:197–248, 1988.
- [23] Gubinelli, M., Controlling rough paths. *J. Funct. Anal.*, **216**(1):86–140, 2004.
- [24] Jourdain, B., and Méléard, S., *Propagation of chaos and fluctuations for a moderate model with smooth initial data*, **34** (1998), 727–766.
- [25] Kac, M., Foundations of kinetic theory. *Third Berkeley Symp. on Math. Stat. and Probab.*, **3**:171–197, 1956.
- [26] Kolokoltsov, V.N., and Troeva, M., On the mean field games with common noise and the McKean-Vlasov SPDEs. [arXiv:1506.04594](https://arxiv.org/abs/1506.04594), 2015.
- [27] Lions, P.-L., Théorie des jeux à champs moyen et applications, *Lectures at the Collège de France*. [http://www.college-de-france.fr/default/EN/all/equ\\_der/cours\\_et\\_seminaires.htm](http://www.college-de-france.fr/default/EN/all/equ_der/cours_et_seminaires.htm)”, 2007-2008
- [28] Lyons, T., Differential equations driven by rough paths. *Rev. Mat. Iberoamericana*, **14**(2):215–310, 1998.
- [29] McKean, H.P., A class of markov processes associated with nonlinear parabolic equations. *Prov. Nat. Acad. Sci.*, **56**:1907–1911, 1966.
- [30] Méléard, S., *Asymptotic behaviour of some interacting particle systems; McKean-Vlasov and Boltzmann models*, Probabilistic models for nonlinear partial differential equations (Montecatini Terme, 1995), Lecture Notes in Math., vol. 1627, Springer, 1996, pp. 42–95.
- [31] Rosenthal, H.P., *On the span in  $L^p$  of sequences of independent random variables. II*. Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Volume 2: Probability Theory,

University of California Press, 1972, pp. 149–167.

- [32] Sznitman, A.-S., Topics in propagation of chaos. *Lect. Notes Math.*, **1464**, 1991.
- [33] Tanaka, H., Probabilistic treatment of the Boltzman equation of Maxwellian molecules. *Probab. Th. Rel. Fields*, **46**:67–105, 1978.
- [34] Tanaka, H., Limit theorems for certain diffusion processes with interaction. *Stochastic analysis (Katata/Kyoto, 1982)*:469–488, North-Holland Math. Library, 32, North-Holland, Amsterdam, 1984.
- [35] Wu, C. and Zhang, J., An elementary proof for the structure of derivatives in probability measures. [arXiv:1705.08046](https://arxiv.org/abs/1705.08046), 2017.