



HAL
open science

Universality of Noise Reinforced Brownian Motions

Jean Bertoin

► **To cite this version:**

| Jean Bertoin. Universality of Noise Reinforced Brownian Motions. 2019. hal-02341310v1

HAL Id: hal-02341310

<https://hal.science/hal-02341310v1>

Preprint submitted on 31 Oct 2019 (v1), last revised 9 Apr 2020 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Universality of Noise Reinforced Brownian Motions

Jean Bertoin*

Dedicated to the memory of Vladas Sidoravicius

Abstract

A noise reinforced Brownian motion is a centered Gaussian process $\hat{B} = (\hat{B}(t))_{t \geq 0}$ with covariance

$$\mathbb{E}(\hat{B}(t)\hat{B}(s)) = (1 - 2p)^{-1}t^p s^{1-p} \quad \text{for } 0 \leq s \leq t,$$

where $p \in (0, 1/2)$ is a reinforcement parameter. Our main purpose is to establish a version of Donsker's invariance principle. Specifically, \hat{B} arises as the universal scaling limit for a large family of step-reinforced random walks in the diffusive regime. This extends known results on the asymptotic behavior of the so-called elephant random walk.

Keywords: Reinforcement, Brownian motion, Invariance Principle, Elephant Random Walk.

Mathematics Subject Classification: 60G50; 60G51; 60K35.

1 Introduction

This work concerns a rather simple real-valued and centered Gaussian process $\hat{B} = (\hat{B}(t))_{t \geq 0}$ with covariance function

$$\mathbb{E}(\hat{B}(t)\hat{B}(s)) = \frac{t^p s^{1-p}}{1 - 2p} \quad \text{for } 0 \leq s \leq t, \quad (1)$$

where $p \in (0, 1/2)$ is a fixed parameter. Recently, this process has notably appeared as the scaling limit for diffusive regimes of the so-called elephant random walk, a simple random walk with memory that has been introduced by Schütz and Trimper [21]. Just as the standard Brownian motion B corresponds to the integral of a white noise, \hat{B} can be thought of as the integral of a reinforced

*Institute of Mathematics, University of Zurich, Switzerland, jean.bertoin@math.uzh.ch

version of the white noise, hence the name *noise reinforced Brownian motion*. Here, reinforcement means that the noise tends to repeat itself infinitesimally as time passes; we refer to [19] for a survey of various models of stochastic processes with reinforcement and their applications, and to [16] and works cited therein for more recent contributions in this area. The parameter p should be interpreted as the strength of the reinforcement; specifically, it represents the probability that an infinitesimal portion of the noise is a repetition. In the limiting case $p = 0$ without reinforcement, one just recovers the standard Brownian motion.

Our purpose here is twofold. We will first present several basic properties of \hat{B} that mirror well-known facts for the standard Brownian motion, even though the laws of B and \hat{B} are mutually singular. We will then establish a version of Donsker's invariance principle. That is, we will show that any so-called step-reinforced random walk with reinforcement parameter p , whose typical step has a finite moment of order r for some $r > 1/p$, converges after a proper rescaling to a noise reinforced Brownian motion. In the special case when the typical step has the Rademacher law, this encompasses known results on the large time asymptotic behavior for the elephant random walk in diffusive regimes.

Our proof of this invariant principle uses an embedding of a step-reinforced random walk in a branching process with types in \mathbb{R} , which is based on a time-substitution via an independent Yule process. This yields a remarkable martingale, whose quadratic variation can then be estimated using well-known properties of Yule processes. In turn, this enables the application of the martingale functional central limit theorem (later on referred to as martingale FCLT).

2 Some basic properties

We start by observing from (1) that the noise reinforced Brownian motion admits a simple representation as a Wiener integral, namely

$$\hat{B}(t) = t^p \int_0^t s^{-p} dB(s), \quad t \geq 0, \quad (2)$$

where $B = (B(s))_{s \geq 0}$ is a standard Brownian motion. Note that equivalently,

$$\hat{B} \text{ has the same law as } \left(\frac{t^p}{\sqrt{1-2p}} B(t^{1-2p}) \right)_{t \geq 0}. \quad (3)$$

It follows immediately from (3) and the classical law of the iterated logarithm for B that

$$\limsup_{t \rightarrow \infty} \frac{\hat{B}(t)}{\sqrt{2t \ln \ln t}} = \limsup_{t \rightarrow 0^+} \frac{\hat{B}(t)}{\sqrt{2t \ln \ln(1/t)}} = \frac{1}{\sqrt{1-2p}} \quad \text{a.s.} \quad (4)$$

In particular, we see that for different reinforcement parameters p , the distributions of noise reinforced Brownian motions, say on the time interval $[0, 1]$, yield

laws on $\mathcal{C}([0, 1])$ which are mutually singular, and are also singular with respect to the Wiener measure.

We deduce from (2) by stochastic calculus that \hat{B} is a semi-martingale which solves the stochastic differential equation

$$d\hat{B}(t) = dB(t) + \frac{p}{t}\hat{B}(t)dt, \quad \hat{B}(0) = 0. \quad (5)$$

This shows that \hat{B} is actually a time-inhomogeneous diffusion process with quadratic variation $\langle \hat{B} \rangle(t) = t$. In this direction, we also infer from (1) that the process

$$\hat{b}(t) = \hat{B}(t) - t^{1-p}\hat{B}(1) \quad \text{for } 0 \leq t \leq 1,$$

is independent of $\hat{B}(1)$. Hence, for any $x \in \mathbb{R}$,

$$\hat{b}(t) + t^{1-p}x = \hat{B}(t) + t^{1-p}(x - \hat{B}(1)) \quad \text{for } 0 \leq t \leq 1,$$

is a version of the bridge of \hat{B} from 0 to x with unit duration.

We further see from (1) that the scaling property,

$$\text{for every } c > 0, \quad \left(c^{-1}\hat{B}(c^2t)\right)_{t \geq 0} \text{ has the same law as } \hat{B}, \quad (6)$$

as well as the time-inversion property,

$$\left(t\hat{B}(1/t)\right)_{t > 0} \text{ has the same law as } \left(\hat{B}(t)\right)_{t > 0},$$

both hold. In this vein, we also point at a remarkable connection with Ornstein-Uhlenbeck processes: the process

$$\hat{U}(t) = e^{-t/2}\hat{B}(e^t), \quad t \in \mathbb{R}$$

is a stationary Ornstein-Uhlenbeck process with infinitesimal generator

$$\mathcal{G}f(x) = \frac{1}{2}f''(x) + (p - 1/2)xf'(x),$$

where $f \in \mathcal{C}^2(\mathbb{R})$. This can be checked by stochastic calculus from (5), or directly by observing from (1) and (6) that \hat{U} is a stationary Gaussian process with covariance

$$\mathbb{E}(\hat{U}(t)\hat{U}(0)) = \frac{e^{(p-1/2)t}}{1-2p} \quad \text{for } t \geq 0.$$

On the other hand, several classical results for the standard Brownian motion plainly fail for its reinforced version. For instance, the increments of \hat{B} are clearly not independent, and the time-reversal property (e.g. Exercise 1.11 in Chapter I of [20]) also fails.

3 An invariance principle with reinforcement

The main purpose of this section is to point out that noise reinforced Brownian motions arise as universal scaling limits of a large class of random walks with step reinforcement. We first recall some features on the so-called elephant random walk, where the story begins.

The *elephant random walk* has been introduced by Schütz and Trimper [21] as a discrete-time nearest neighbor process with memory on \mathbb{Z} ; it can be depicted as follows. Fix some $q \in (0, 1)$ and call q the memory parameter. Imagine that a walker (an elephant) makes a first step in $\{-1, +1\}$ at time 1; then at each time $n \geq 2$, it selects randomly a step from its past. With probability q , the elephant repeats this step, and with complementary probability $1 - q$, it makes the opposite step. Note that for $q = 1/2$, the elephant merely follows the path of a simple symmetric random walk. The elephant random walk has generated much interest in the recent years, we refer notably to [2, 3, 7, 8, 9, 17, 18], see also [1, 4, 5, 6, 12] for variations, and references therein for further related works. A remarkable feature is that its large time asymptotic behavior of an elephant random walk is diffusive when the memory parameter q is less than $3/4$ and super-diffusive when $q > 3/4$.

Remark 3.1. The laws of the iterated logarithm (4) for a noise reinforced Brownian motion bear the same relation to that for the elephant random walk in the diffusive regime (Corollary 1 in [8] and Theorem 3.2 in [3]; beware that the memory parameter denoted by p there corresponds to $q = (1 + p)/2$ in the present notation as it will be stressed below), as the law of the iterated logarithm for the standard Brownian motion due to P. Lévy does to that for the simple random walk due to A.Y. Khintchine.

We are interested in a generalization where the distribution of a typical step of the walk is arbitrary, that we call *step reinforced random walk*. Fix a parameter $p \in (0, 1)$, called the *reinforcement parameter*¹; at each discrete time, with probability p , a step reinforced random walk repeats one of its preceding steps chosen uniformly at random, and otherwise, i.e. with probability $1 - p$, it has an independent increment with a fixed distribution. More precisely, consider a sequence X_1, X_2, \dots of i.i.d. copies of a random variable X in \mathbb{R} and define recursively $\hat{X}_1, \hat{X}_2, \dots$ as follows. Let $(\varepsilon_i : i \geq 2)$ be an independent sequence of Bernoulli variables with parameter p . We set first $\hat{X}_1 = X_1$, and next for $i \geq 2$, we let $\hat{X}_i = X_i$ if $\varepsilon_i = 0$, whereas we define \hat{X}_i as a uniform random sample from $\hat{X}_1, \dots, \hat{X}_{i-1}$ if $\varepsilon_i = 1$. Finally, the sequence of the partial sums

$$\hat{S}(n) = \hat{X}_1 + \dots + \hat{X}_n, \quad n \in \mathbb{N},$$

is referred to as a step reinforced random walk.

¹Beware that we assumed $p < 1/2$ in the preceding sections. The first part of the present section (super-diffusive regime) does not involve any noise reinforced Brownian motion, and the case $p \geq 1/2$ is allowed. In the second part of this section (diffusive regime), we shall again focus on the case $p < 1/2$ and noise reinforced Brownian motions will then re-appear.

When the typical step X has the Rademacher law, i.e. $\mathbb{P}(X = 1) = \mathbb{P}(X = -1) = 1/2$, Kürsten [18] (see also [10]) pointed out that \hat{S} is a version of the elephant random walk with memory parameter $q = (p + 1)/2$ in the present notation. We also note that \hat{S} is a time-inhomogeneous Markov chain when X can take at most two values, but otherwise the Markov property fails for \hat{S} . When X has a symmetric stable distribution, \hat{S} is the so-called shark random swim which has been studied in depth by Businger [6]. More general versions when the distribution of X is infinitely divisible have been considered in [5].

The main result of Businger [6] is that the large time asymptotic behavior of a shark random swim exhibits a phase transition similar to that for the elephant random walk, but for a different critical parameter. We shall now extend this to a large class of step reinforced random walks. In the sequel we implicitly rule out the degenerate case when the typical step variable X is a constant. We start with the super-diffusive regime for which all the ingredients are already in Section 3.1 in [6].

Theorem 3.2. *Let $p \in (1/2, 1)$, and suppose that $X \in L^2(\mathbb{P})$. Then*

$$\lim_{n \rightarrow \infty} \frac{\hat{S}(n) - n\mathbb{E}(X)}{n^p} = L \quad \text{in } L^2(\mathbb{P}),$$

where L is some non-degenerate random variable.

Proof. By centering and normalizing, we may assume without loss of generality that $\mathbb{E}(X) = 0$ and $\text{Var}(X) = 1$. One first introduces for every $i, n \in \mathbb{N}$, the number $r_{i,n}$ of repetitions of the variable X_i in the reinforced sequence $\hat{X}_1, \dots, \hat{X}_n$ (in particular, $r_{i,n} = 0$ when either $\varepsilon_i = 1$ or $i > n$). We stress that repetition numbers only depend on the Bernoulli variables ε_j and are hence independent of the variables X_j . We know from Lemmas 3 and 5 in [6] that for each $i \in \mathbb{N}$

$$\lim_{n \rightarrow \infty} n^{-p} r_{i,n} = R_i, \quad \text{a.s.}$$

where R_i is some non-degenerated random variable with $\sum_i \mathbb{E}(R_i^2) < \infty$.

In particular, $\sum_i R_i^2 < \infty$ a.s., and since the variables X_i are i.i.d. centered and with unit variance, the sum $\sum_i R_i X_i = L$ is well-defined a.s. (as a martingale limit, conditionally on the R_i). More precisely, $\mathbb{E}(L) = 0$ and $\mathbb{E}(L^2) = \sum_i \mathbb{E}(R_i^2)$.

To conclude, we recall from Equation (6) in [6] that

$$\lim_{n \rightarrow \infty} \sum_i \mathbb{E} \left(|n^{-p} r_{i,n} - R_i|^2 \right) = 0. \quad (7)$$

By the construction of the step reinforced random walk, we have

$$\hat{S}(n) = \sum_i r_{i,n} X_i.$$

Since the variables $r_{i,n}$ and R_i are independent of the X_i , and further $\mathbb{E}(X) = 0$ and $\mathbb{E}(X^2) = 1$, there is the identity

$$\mathbb{E} \left(|n^{-p} \hat{S}(n) - L|^2 \right) = \mathbb{E} \left(\sum_i |n^{-p} r_{i,n} - R_i|^2 \right).$$

An appeal to (7) completes the proof. \square

We then turn our attention to the diffusive regime, and obtain a version of Donsker's invariance principle for step reinforced random walks.

Theorem 3.3. *Let $p \in (0, 1/2)$, and suppose that $X \in L^r(\mathbb{P})$ for some $r > 1/p$. Then as $n \rightarrow \infty$, the sequence of processes*

$$\frac{\hat{S}(\lfloor tn \rfloor) - tn\mathbb{E}(X)}{\sqrt{n\text{Var}(X)}}, \quad t \geq 0$$

converges in distribution in the sense of Skorohod towards a noise reinforced Brownian motion \hat{B} with reinforcement parameter p .

Remark 3.4. It is of course tempting to conjecture that Theorem 3.3 holds under the weaker assumption $X \in L^2(\mathbb{P})$. Note in particular that our requirement $X \in L^r(\mathbb{P})$ for some $r > 1/p$ gets stronger as the reinforcement parameter p decreases to $0+$, which is clearly contrary to the intuition. This requirement is essential for establishing the forthcoming Lemma 3.8, which however is much stronger than the asymptotic negligibility of the maximal jumps that is needed for applying the martingale FCLT.

The method used in [2] to establish Theorem 3.3 for the elephant random walk relied crucially on limit theorems for generalized Pólya urns due to Janson [14]. Unfortunately this approach is not available for arbitrary step distributions, notably as one would need to work with urn models having types in an infinite space (the real line \mathbb{R} to be more specific), and how to deal with this kind of urns is still an open problem; see Remark 4.1 in [14]. Our approach relies essentially on the martingale FCLT, which also plays a key role for urn models (see [11, 14]), as well in the works of Coletti *et al.* [7, 8] and of Bercu [3] on the elephant random walk. For the sake of simplicity, we only consider here the one-dimensional setting only, however the argument could be readily adapted to \mathbb{R}^d ; see also [4].

The rest of this section is devoted to the proof of Theorem 3.3; without loss of generality, we henceforth assume that $X \in L^r(\mathbb{P})$ for some $r > 1/p > 2$ with $\mathbb{E}(X) = 0$ and $\text{Var}(X) = 1$. These assumptions are implicit in the forthcoming statements and their proofs.

We first embed the step reinforced random walk in a branching process as follows. Let $Y = (Y_t)_{t \geq 0}$ be a standard Yule process, that is Y is a pure birth process started from $Y_0 = 1$, with birth rate n from any state $n \in \mathbb{N}$. We

further assume that Y and \hat{S} are independent, and will mainly work with the time-changed process $\hat{S}(Y)$.

Before tackling the proof of Theorem 3.3, it may be worth discussing a bit our motivation for introducing the time-substitution based on an independent Yule process. Even though this discussion will not be used in the proof, it nonetheless provides a useful guiding line for our approach. Roughly speaking, Y describes a population model started from a single ancestor, where individuals are eternal, and each begets a child at unit rate, independently of the other individuals. Those individuals are naturally enumerated in the increasing order of their birth time, in particular the ancestor is the first individual. We decide to assign the type \hat{X}_n to the n -th individual, and write

$$\mathbf{Z}_t(dx) = \sum_{n=1}^{Y_t} \delta_{\hat{X}(n)}(dx), \quad x \in \mathbb{R},$$

for the point process of the types of individuals alive at time t . We see from basic properties of independent exponential clocks that $\mathbf{Z} = (\mathbf{Z}_t)_{t \geq 0}$ is a (multitype) branching process, in which each individual begets a child at unit rate, independently of the other individuals. A child is either a clone of its parent, an event which occurs with probability p , or a mutant, an event which occurs with probability $1 - p$. If a child is a clone, then its type is the same as that of its parent, whereas if it is a mutant, its type is given by an independent copy of X . In this setting, there are the identities

$$Y_t = \mathbf{Z}_t(\mathbf{1}) \quad \text{and} \quad \hat{S}(Y_t) = \mathbf{Z}_t(\text{Id}), \quad (8)$$

with the notation $\mathbf{1}(x) = 1$, $\text{Id}(x) = x$ and $\mathbf{Z}_t(f) = \int_{\mathbb{R}} f(x) \mathbf{Z}_t(dx)$. The key point is that the functions $\mathbf{z} \mapsto \mathbf{z}(\mathbf{1})$ and $\mathbf{z} \mapsto \mathbf{z}(\text{Id})$ on the space of point measures \mathbf{z} are eigenfunctions for the infinitesimal generator of the branching process \mathbf{Z} , for the eigenvalues 1 and p , respectively.

Lemma 3.5. *The process*

$$M(t) = e^{-pt} \hat{S}(Y_t), \quad \text{for } t \geq 0,$$

is a square integrable purely discontinuous martingale.

Proof. Plainly, $\mathbb{E}(\hat{S}(n)^2) \leq n^2 \mathbb{E}(X^2) = n^2$, and since \hat{S} and Y are independent with $\mathbb{E}(Y_t^2) < \infty$, $M(t)$ is indeed square integrable. We next point out that for any $n \in \mathbb{N}$,

$$\begin{aligned} \mathbb{E}(\hat{X}(n+1) \mid \hat{X}(1), \dots, \hat{X}(n)) &= (1-p)\mathbb{E}(X) + p \frac{\hat{X}(1) + \dots + \hat{X}(n)}{n} \\ &= p \frac{\hat{S}(n)}{n} \end{aligned}$$

(this observation is also the starting point of the analysis of the elephant random walk in [7, 8, 3]). Since the Yule process has precisely jump rate n from the state n , this entails that

$$\hat{S}(Y_t) - p \int_0^t \hat{S}(Y_s) ds, \quad \text{for } t \geq 0$$

is a martingale, and our statement then follows from elementary stochastic calculus. \square

We will need to estimate the angle-bracket $\langle M \rangle$ of M , and in this direction, we first introduce

$$\hat{V}(n) = \hat{X}(1)^2 + \dots + \hat{X}(n)^2, \quad n \in \mathbb{N}.$$

Lemma 3.6. *We have that, as $t \rightarrow \infty$,*

$$\mathbb{E} \left(\left| e^{-pt} \hat{V}(Y_t) - (1-p) \int_0^t e^{-ps} Y_s ds \right| \right) = o(e^{(1-p)t}).$$

Proof. Just as in the proof of Lemma 3.5, we note the identity

$$\mathbb{E}(\hat{X}(n+1)^2 \mid \hat{X}(1), \dots, \hat{X}(n)) = p \frac{\hat{V}(n)}{n} + 1 - p,$$

and get by stochastic calculus that the process

$$M'(t) = e^{-pt} \hat{V}(Y_t) - (1-p) \int_0^t e^{-ps} Y_s ds, \quad t \geq 0$$

is a purely discontinuous martingale in $L^{r/2}(\mathbb{P})$.

Suppose first that $r \geq 4$. Then the square bracket of M' can be expressed as

$$[M'](t) = \int_{(0,t]} e^{-2ps} |\hat{X}(Y_s)|^4 dY_s, \quad t \geq 0.$$

We compute its expected value and get

$$\begin{aligned} \mathbb{E}(X^4) \mathbb{E} \left(\int_{(0,t]} e^{-2ps} dY_s \right) &= \mathbb{E}(X^4) \int_0^t e^{(1-2p)s} ds \\ &= o(e^{2(1-p)t}), \end{aligned}$$

where for the first equality, we used that the instantaneous jump rate of the Yule process at time s equals Y_s and $\mathbb{E}(Y_s) = e^s$. We conclude by the Cauchy-Schwarz inequality that

$$\mathbb{E}(|M'(t) - M'(0)|) \leq \mathbb{E}((M'(t) - M'(0))^2)^{1/2} = \mathbb{E}([M'](t))^{1/2} = o(e^{(1-p)t}).$$

Suppose then that $2 < r < 4$. Since $r/4 < 1$ and dY_s is a point measure, there is the inequality

$$[M']^{r/4}(t) \leq \int_{(0,t]} e^{-prs/2} |\hat{X}(Y_s)|^r dY_s, \quad t \geq 0.$$

We compute the expected value of the right-hand side just as above, and get

$$\mathbb{E} \left([M']^{r/4}(t) \right) \leq \mathbb{E}(|X|^r) \int_0^t e^{(1-pr/2)s} ds = o(e^{(1-p)rt/2}).$$

Since $r/2 > 1$, we conclude from Jensen's inequality at the first line and the Burkholder-Davis-Gundy inequality (see, e.g. Theorem 26.12 in [15]) at the second line that

$$\begin{aligned} \mathbb{E}(|M'(t) - M'(0)|) &\leq \mathbb{E}(|M'(t) - M'(0)|^{r/2})^{2/r} \\ &\leq c(r) \mathbb{E}([M']^{r/4}(t))^{2/r} \\ &= o(e^{(1-p)t}), \end{aligned}$$

where $c(r)$ denotes some constant depending on r only. \square

We next recall that the asymptotic behavior of the Yule is described by

$$\lim_{t \rightarrow \infty} e^{-t} Y_t = \epsilon \quad \text{a.s.} \quad (9)$$

where ϵ is a standard exponential variable. Needless to say, ϵ is independent of \hat{S} .

Corollary 3.7. *The angle-bracket $\langle M \rangle$ of the martingale M in Lemma 3.5 fulfills*

$$\lim_{t \rightarrow \infty} e^{-(1-2p)t} \langle M \rangle(t) = \frac{\epsilon}{1-2p} \quad \text{in probability.}$$

Proof. The square-bracket process of M can be expressed in the form

$$[M](t) = \int_{(0,t]} e^{-2ps} |\hat{X}(Y_s)|^2 dY_s, \quad t \geq 0,$$

and it follows readily that its predictable compensator $\langle M \rangle$ is given by

$$\langle M \rangle(t) = \int_{(0,t]} e^{-2ps} \left(p \hat{V}(Y_s) + (1-p) Y_s \right) ds, \quad t \geq 0. \quad (10)$$

Since $2p < 1$, our statement now derives from (9), (10) and Lemma 3.6. \square

Next, we point out that the maximal jump size of M has a finite moment of order r .

Lemma 3.8. *The random variable $\Delta^* = \sup_{t \geq 0} |M_t - M_{t-}|$ belongs to $L^r(\mathbb{P})$.*

We stress that the initial value M_0 is viewed as an initial jump (i.e. we use with the convention $M_{0-} = 0$) and therefore is taken into account in the definition of Δ^* .

Proof. Let $\tau(n) = \inf\{t > 0 : Y_t = n\}$ denote the first hitting time of $n \in \mathbb{N}$ by the Yule process. From the definition of M , there is the identity

$$\Delta^* = \sup_{n \in \mathbb{N}} e^{-p\tau(n)} |\hat{X}(n)|.$$

Since $\tau(n)$ and $\hat{X}(n)$ are independent and each $\hat{X}(n)$ is a copy of X , there is the inequality

$$\mathbb{E}((\Delta^*)^r) \leq \mathbb{E}(|X|^r) \sum_{n=1}^{\infty} \mathbb{E}(e^{-rp\tau(n)}),$$

and we just need to check that the series in the right-hand side converges.

We express $\mathbb{E}(e^{-rp\tau(n)}) = \mathbb{P}(Y_{\sigma(rp)} > n)$, where $\sigma(rp)$ denotes an exponential random variable with parameter rp , which is independent of the Yule process. In this setting, it is well-known that $Y_{\sigma(rp)}$ has the so-called Yule-Simon distribution with parameter rp ,

$$\mathbb{P}(Y_{\sigma(rp)} = k) = rpB(k, rp + 1) \sim rp\Gamma(rp + 1)k^{-(rp+1)} \quad \text{as } k \rightarrow \infty,$$

where B (respectively, Γ) stands for the beta (respectively, gamma) function. Since $rp > 1$, this ensures the finiteness of $\sum_{n=1}^{\infty} \mathbb{E}(e^{-rp\tau(n)})$, and hence completes the proof. □

The angle-bracket $\langle M \rangle$ is a continuous strictly increasing bijection from \mathbb{R}_+ to \mathbb{R}_+ a.s., see (10), and we write T for the inverse bijection. We introduce for each $n \in \mathbb{N}$ the process

$$N_n(t) = n^{-1/2} M_{T(nt)} = n^{-1/2} e^{-pT(nt)} \hat{S}(Y_{T(nt)}), \quad t \geq 0.$$

We are in position of applying the martingale FCLT.

Proposition 3.9. *As $n \rightarrow \infty$, the sequence of processes N_n converges in distribution in the sense of Skorohod to a standard Brownian motion.*

Proof. Plainly, each N_n is a square-integrable martingale with angle-bracket $\langle N_n \rangle(t) = t$, and Lemma 3.8 ensures that the maximum jump in N_n is asymptotically negligible in $L^r(\mathbb{P})$. This enables us to apply the martingale FCLT; see e.g. Theorem 2.1 in [22], and also Section VIII.3 in [13] for more general versions. □

We shall actually need a slightly stronger version of Proposition 3.9 in the which the convergence holds conditionally on the variable ϵ in (9).

Corollary 3.10. *As $n \rightarrow \infty$, the sequence of pairs (ϵ, N_n) converges in distribution in the sense of Skorohod to (ϵ, B) where B is a standard Brownian motion independent of ϵ .*

Proof. First fix $t > 0$ and consider a random variable $A(t)$ which is measurable with respect to the sigma-algebra $\sigma(M_{T(s)} : 0 \leq s \leq t)$. The statement with $A(t)$ replacing ϵ follows from the martingale FCTL applied to the sequence of processes $(N_n(s + t/n))_{s \geq 0}$, just as in Proposition 3.9. Corollary 3.10 can then be deduced, provided that we can choose $A(t)$ such that

$$\lim_{t \rightarrow \infty} A(t) = \epsilon \quad \text{a.s.} \quad (11)$$

Specifically, consider any continuous functional F on the Skorohod space with $|F| \leq 1$, and any continuous and bounded function f on \mathbb{R} . Taking (11) for granted, for any $\eta > 0$ arbitrarily small, we can first choose $t > 0$ sufficiently large so that

$$\mathbb{E}(|f(A(t)) - f(\epsilon)|) \leq \eta,$$

and then $n(t) \in \mathbb{N}$ such that

$$|\mathbb{E}(f(A(t))F(N_n)) - \mathbb{E}(f(A(t)))\mathbb{E}(F(B))| \leq \eta \quad \text{for all } n \geq n(t).$$

We can conclude from the triangle inequality that

$$|\mathbb{E}(f(\epsilon)F(N_n)) - \mathbb{E}(f(\epsilon))\mathbb{E}(F(B))| \leq 3\eta \quad \text{for all } n \geq n(t).$$

We now have to construct variables $A(t)$ so that (11) holds, and in this direction, we first assume that $\mathbb{P}(X = 0) = 0$. All the steps of \hat{S} are non-zero a.s., and $Y_{T(t)}$ is then the total number of jumps of the process $M_{T(\cdot)}$ on the time interval $[0, t]$ (recall that this takes the initial jump at time 0 into account), and hence is measurable with respect to $\sigma(M_{T(s)} : 0 \leq s \leq t)$. On the other hand, we deduce from Corollary 3.7 that

$$T(t) = \frac{1}{1-2p} \ln \left(\frac{(1-2p)t}{\epsilon} \right) + o(1) \quad \text{as } t \rightarrow \infty, \quad (12)$$

and then from (9) that

$$Y_{T(t)} \sim ((1-2p)t)^{1/(1-2p)} \epsilon^{-2p/(1-2p)} \quad \text{as } t \rightarrow \infty. \quad (13)$$

The construction of the sought variables $A(t)$ is plain from (13).

The case when $\mathbb{P}(X = 0) = a \in (0, 1)$ only requires a minor modification. We can no longer identify $Y_{T(t)}$ with the total number of jumps of the process $M_{T(\cdot)}$ on the time interval $[0, t]$. Nonetheless the latter is now close to $(1-a)Y_{T(t)}$ for $t \gg 1$, and we can conclude just as above. \square

We now have all the ingredients needed for the proof of the invariance principle.

Proof of Theorem 3.3. Set

$$s = s(t) = (\epsilon^{-2pt})^{1/(1-2p)} \quad \text{and} \quad k = k(n) = ((1-2p)n)^{1/(1-2p)},$$

so (13) and (12) yield respectively

$$Y_{T(nt)} \sim ks \quad \text{and} \quad e^{-pT(nt)} \sim (ks/\epsilon)^{-p}.$$

Since $n = k^{1-2p}/(1-2p)$ and $t = \epsilon^{2p}s^{1-2p}$, we deduce from Corollary 3.10 and the very definition of the Skorohod's topology involving changes of time (see, e.g. Section VI.1 in [13]) that as $k \rightarrow \infty$, the sequence of processes $k^{-1/2}(\hat{S}(\lfloor ks \rfloor))_{s \geq 0}$ converges in distribution in the sense of Skorohod towards

$$\left(\frac{\epsilon^{-p}}{\sqrt{1-2p}} s^p B(\epsilon^{2p}s^{1-2p}) \right)_{s \geq 0},$$

where B is a standard Brownian motion independent of ϵ . By the scaling property of Brownian motion, the process displayed above has the same distribution as

$$\left(\frac{s^p}{\sqrt{1-2p}} B(s^{1-2p}) \right)_{s \geq 0},$$

and we complete the proof with (3). □

References

- [1] BAUR, E. On a class of random walks with reinforced memory. arXiv:1909.04633.
- [2] BAUR, E., AND BERTOIN, J. Elephant random walks and their connection to Pólya-type urns. *Phys. Rev. E* 94 (Nov 2016), 052134.
- [3] BERCU, B. A martingale approach for the elephant random walk. *J. Phys. A* 51, 1 (2018), 015201, 16.
- [4] BERCU, B., AND LAULIN, L. On the multi-dimensional elephant random walk. *Journal of Statistical Physics* 175, 6 (Jun 2019), 1146–1163.
- [5] BERTOIN, J. Noise reinforcement for Lévy processes. arXiv:1810.08364.
- [6] BUSINGER, S. The shark random swim (Lévy flight with memory). *J. Stat. Phys.* 172, 3 (2018), 701–717.
- [7] COLETTI, C. F., GAVA, R., AND SCHÜTZ, G. M. Central limit theorem and related results for the elephant random walk. *J. Math. Phys.* 58, 5 (2017), 053303, 8.

- [8] COLETTI, C. F., GAVA, R., AND SCHÜTZ, G. M. A strong invariance principle for the elephant random walk. *J. Stat. Mech. Theory Exp.*, 12 (2017), 123207, 8.
- [9] COLETTI, C. F., AND PAPAGEORGIOU, I. Asymptotic analysis of the elephant random walk. arXiv:1910.03142.
- [10] GONZÁLEZ-NAVARRETE, M., AND LAMBERT, R. Non-Markovian random walks with memory lapses. *J. Math. Phys.* 59, 11 (2018), 113301, 11.
- [11] GOUET, R. Martingale functional central limit theorems for a generalized Pólya urn. *The Annals of Probability* 21, 3 (1993), 1624–1639.
- [12] GUT, A., AND STADTMUELLER, U. Elephant random walks with delays. arXiv:1906.04930.
- [13] JACOD, J., AND SHIRYAEV, A. N. *Limit Theorems for Stochastic Processes*. Grundlehren der mathematischen Wissenschaften. Springer Berlin Heidelberg, 2002.
- [14] JANSON, S. Functional limit theorems for multitype branching processes and generalized pólya urns. *Stochastic Processes and their Applications* 110, 2 (2004), 177 – 245.
- [15] KALLENBERG, O. *Foundations of modern probability*, second ed. Probability and its Applications (New York). Springer-Verlag, New York, 2002.
- [16] KIOUS, D., AND SIDORAVICIUS, V. Phase transition for the once-reinforced random walk on \mathbb{Z}^d -like trees. *Ann. Probab.* 46, 4 (07 2018), 2121–2133.
- [17] KUBOTA, N., AND TAKEI, M. Gaussian fluctuation for superdiffusive elephant random walks. arXiv:1909.02834.
- [18] KÜRSTEN, R. Random recursive trees and the elephant random walk. *Phys. Rev. E* 93, 3 (2016), 032111, 11.
- [19] PEMANTLE, R. A survey of random processes with reinforcement. *Probab. Surveys* 4 (2007), 1–79.
- [20] REVUZ, D., AND YOR, M. *Continuous martingales and Brownian motion*, third ed., vol. 293 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [21] SCHÜTZ, G. M., AND TRIMPER, S. Elephants can always remember: Exact long-range memory effects in a non-markovian random walk. *Phys. Rev. E* 70 (Oct 2004), 045101.
- [22] WHITT, W. Proofs of the martingale FCLT. *Probab. Surv.* 4 (2007), 268–302.