



The Bacterial and Fungal Diversity of an Aged PAH- and Heavy Metal-Contaminated Soil is Affected by Plant Cover and Edaphic Parameters

Amélia Bourceret, Aurélie Cebron, Emilie E. Tisserant, Pascal Poupin, Pascale Bauda, Thierry Beguiristain, Corinne Leyval

► To cite this version:

Amélia Bourceret, Aurélie Cebron, Emilie E. Tisserant, Pascal Poupin, Pascale Bauda, et al.. The Bacterial and Fungal Diversity of an Aged PAH- and Heavy Metal-Contaminated Soil is Affected by Plant Cover and Edaphic Parameters. *Microbial ecology*, 2016, 71 (3), pp.711-724. 10.1007/s00248-015-0682-8 . hal-02341848

HAL Id: hal-02341848

<https://hal.science/hal-02341848>

Submitted on 13 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The bacterial and fungal diversity of an aged PAH- and heavy metal-contaminated soil is affected by plant cover and edaphic parameters

BOURCERET Amélie^{1,2}, CÉBRON Aurélie^{1,2,†}, TISSERANT Emilie³, POUPIN Pascal^{4,5},
BAUDA Pascale^{4,5}, BEGUIRISTAIN Thierry^{1,2}, LEYVAL Corinne^{1,2}

¹ CNRS, LIEC UMR7360, Faculté des Sciences et Technologies, BP70239, 54506 Vandoeuvre-lès-Nancy, France.

² Université de Lorraine, LIEC UMR7360, Faculté des Sciences et Technologies, BP 70239, 54506 Vandoeuvre-lès-Nancy, France.

³ INRA, IAM UMR1136, Centre INRA de Nancy, 54280 Champenoux, France.

⁴ Université de Lorraine, LIEC UMR7360, Campus Bridoux - Rue du Général Delestraint, 57070 Metz Borny, France.

⁵ CNRS, LIEC UMR7360, Campus Bridoux - Rue du Général Delestraint, 57070 Metz Borny, France.

[†] Corresponding author: aurelie.cebron@univ-lorraine.fr; Phone: (+33) 3 83 68 42 96

ABSTRACT

Industrial wasteland soils with aged PAH and heavy metal contaminations are environments where pollutant toxicity has been maintained for decades. Although the communities may be well adapted to the presence of stressors, knowledge about microbial diversity in such soils is scarce. Soil microbial community dynamics can be driven by the presence of plants, but the impact of plant development on selection or diversification of microorganisms in these soils has not been established yet. To test these hypotheses, aged-contaminated soil samples from a field trial were collected. Plots planted with alfalfa were compared to bare soil plots, and bacterial and fungal diversity and abundance were assessed after 2 and 6 years. Using pyrosequencing of 16S rRNA gene and ITS amplicons, we showed that the bacterial community was dominated by *Proteobacteria*, *Actinobacteria* and *Bacteroidetes* and was characterized by low *Acidobacteria* abundance, while the fungal community was mainly represented by members of the *Ascomycota*. The short-term toxic impact of pollutants usually reduces the microbial diversity, yet in our samples bacterial and fungal species richness and diversity was high suggesting that the community structure and diversity adapted to the contaminated soil over decades. The presence of plants induced higher bacterial and fungal diversity than in bare soil. It also increased the relative abundance of bacterial members of the

Actinomycetales, *Rhizobiales* and *Xanthomonadales* orders and of most fungal orders. Multivariate analysis showed correlations between microbial community structure and heavy metal and PAH concentrations over time, but also with edaphic parameters (C/N, pH, phosphorus and nitrogen concentrations).

Key words: bacteria, fungi, PAH, heavy metals, rhizosphere, 454-pyrosequencing

1. Introduction

The decline of coal-mining activities has left behind large areas of industrial wasteland soils that display high levels of multi-contamination by polycyclic aromatic hydrocarbons (PAHs) and heavy metals (HMs) [1]. During the 20th century, industrial facilities such as gas or coking plants caused chronic pollution of soils. After an ageing period, both PAHs and HMs persisted in the environment due to their low availability/mobility and low degradability/reactivity. Thus the high concentrations of pollutants and their toxicity have been exerting a selection pressure on microorganisms for decades. These particular environmental conditions potentially enabled the microbial community to adapt, and favored microorganisms able of using pollutant as carbon source or tolerate pollutant by detoxification mechanisms [2]; [3]. This long-term process should lead to the development of a new microbial community diversity after a few decades [4]; [5]. Yet, little is known about the impact of ancient contamination on microbial diversity in such aged multi-polluted soils. Most studies focus on the short-term modifications of microbial diversity following recent contamination events, mainly in laboratory experiments (*i.e.* spiking of unpolluted soils) [8];[9];[10]. Environmental conditions and edaphic parameters are components that shape the soil microbial communities [11]. Similarly, chronic pollution could also be an important parameter that can select soil microbial diversity. While the short-term impact of contamination tends to decrease microbial abundance, richness and diversity, we hypothesized that in an aged multi-contaminated soil the microbial community would adapt over time toward a unique and rich diversity pattern [4]; [5].

Some studies show that vegetation can help to restore contaminated environments. Phytoremediation strategies have been developed based on plants' contribution to the stabilization, extraction and degradation of pollutants [12]. While shaping microbial diversity by stimulating some microbial populations, plant root exudates could favor PAH degradation

through rhizodegradation processes [13];[14];[15] and modify HM speciation, mobility and availability [12] in the rhizosphere. Among different plant species used in rhizoremediation assays, alfalfa is described as tolerant to pollution and interesting for PAH remediation [16];[17], similarly to other legumes such as red lentil [18]. A contribution of mycorrhizal colonization of plant roots to PAH dissipation [19] and HM tolerance [20] has also been shown. There is an effect of plants on soil microbial biomass and diversity because vegetal litter or residues bring complex organic matter into soils [21] and also pose the question of selection or diversification of microorganisms. As most studies focus on one particular microbial function or population, little is known about the impact of plants and the effect of the rhizosphere on bacterial and fungal richness and diversity in multi-contaminated soils. A dramatic shift in the bacterial community structure in the rhizosphere of sunflower in a creosote-contaminated soil was demonstrated in a pot experiment after 90 days in the greenhouse during PAH-degradation [22]. Although fungal community dynamic is mainly driven by plants [23], the diversity of this microbial community needs to be explored. Overall, as the rhizosphere effect could modify soil parameters and fate of pollutants, there is still a need to study the long-term impact of plants in comparison to bare soil, on shaping the microbial communities during natural attenuation processes of aged multi-contaminated soils. It is well established that environmental parameters impact microbial community composition [11]. Although bacteria and fungi are in constant interaction in soils [24], these two components of the microbial community have a different ecology (e.g. nutrient and carbon source requirement, growth rate...) and respond differently to environmental conditions and stressors. To study the system as a whole, bacterial and fungal diversity must be studied together. To our knowledge, no study has yet evaluated how plants and edaphic parameters drive both fungal and bacterial diversity of a multi-contaminated soil on the long term.

In this context, the aims of our study were to (1) describe the diversity of bacterial and fungal communities in an aged multi-contaminated soil from a former coking plant site (Neuves-Maisons, France), (2) determine the impact of plant development on microbial richness and diversity, (3) characterize the temporal modifications of microbial diversity according to evolution of edaphic parameters over 6 years. To address this issue, we collected soil samples from a field trial where a long-term plant-assisted natural attenuation experiment was being conducted [25]. We studied three replicate plots of planted and non-planted soil, sampled at three time-points (0, after 2 and 6 years). We analyzed bacterial and fungal community abundance using real-time quantitative PCR, and diversity using tag-encoded pyrosequencing

of the 16S rRNA gene and ITS amplicons, respectively.

2. Materials and methods

2.1. Site description and sample collection

The field trial, previously described by Ouvrard et al. [25], was set up in September 2005. Among the 24 plots filled with an aged PAH- and HM-contaminated soil (NM) from a coking plant wasteland (Neuves-Maisons, Meurthe et Moselle, France), only three replicates of two treatments were studied. The bare soil plots (NM-BS) were kept clear of vegetation by hand weeding and the planted plots (NM-Msm) were sown with alfalfa (40 g of seeds of *Medicago sativa* var. Europe *per* plot) in September 2005 and co-inoculated with two mycorrhizal fungal strains (*Glomus mosseae* and *Glomus intra-radices*, recently renamed *Funneliformis mosseae* and *Rhizophagus intraradices* [26], commercial inoculum from the Institut für Pflanzenkultur, Germany). The plots were monitored twice a year, in May and September, for 6 years. In September, the alfalfa biomass was cut and dry weight was measured (Table 1). Mycorrhizal colonization of roots was determined by Trypan blue staining as described by [27] and notation was performed using the method described in [28]. Six soil subsamples were collected *per* plot and mixed to obtain one composite soil sample *per* plot. After sieving to 5 mm, 500 g were dried and dedicated to soil characterization, and 50 g were stored at -80°C until DNA extraction. Three sampling dates were studied: samples T0 (September 2005), T4 (September 2007) and T12 (September 2011), i.e. 18 soil samples altogether. Sample characteristics (humidity level, physico-chemical properties, total and extractable heavy metals, and PAH concentrations) were measured at the LAS-INRA laboratory (Arras, France) and are listed in Table 1.

2.2. DNA extraction

Total genomic DNA was extracted from 0.5 g wet weight of each of the 18 soil samples with the FastDNA SPIN Kit for Soil (MP Biomedicals, Illkirch, France), and resuspended in 100 µl of DES (Dnase-Free water). DNA concentrations and quality (purity ratio A_{260}/A_{280}) were measured using a spectrophotometer (UV1800, Shimadzu) equipped with a TrayCell unit (Hellma).

2.3. Assessment of fungal and bacterial abundance by real-time quantitative PCR

Real-time quantitative PCR was performed as previously described in Cébron et al. [29] and

Thion et al. [23]. Primer sets FF390R (5'-GAGGTCTCGTTCGTTATCG-3') / Fung5F (5'-GGGAACCAGGACTTTTAC-3') [30] and 968F (5'-AACGCGAAGAACCTTAC-3') / 1401R (5'-CGGTGTGTACAAGACCC-3') [31] were used to quantify fungi and bacteria by targeting the 18S rRNA and 16S rRNA genes, respectively. Amplification reactions (20 µl) were carried out using 1 µl of ten-fold diluted DNA (corresponding to 2.8 to 6.5 ng), 10 µl of iQ SYBR Green SuperMix (Bio-Rad), 0.8 µl of each primer (10 µM), 0.4 µl of Bovine Serum Albumin (3%), 0.2 µl of Dimethylsulfoxide, and 0.08 µl of T4gp32 (500 µg/ml) (MP Biomedicals). Quantification was performed using a CFX96 Real-Time PCR detection system (Bio-Rad) and standard plasmid dilution series from 10⁸ to 10¹ copies µl⁻¹. Data were then expressed as gene copy numbers *per* gram of dry weight soil after data correction using soil sample humidity values.

2.4. PCR and pyrosequencing

The primer set 515F (5'-GTG CCA GCM GCC GCG GTA A-3') / 907R (5'-CCG TCA ATT CMT TTR AGT TT-3') [32] was used to amplify the V4-V5 region of bacterial 16S rRNA genes and the primer set ITS1F (5'-CTTGGTCATTTAGAGGAAGTAA-3') [33] / ITS2 (5'-GCTGCGTTCTTCATCGATGC-3') [34] was used to amplify the ITS1 (Internal Transcribed Spacer) region of fungal rRNA genes. Primers were combined at the 5' end with a MID (Multiplex Identifier, Roche) corresponding to a sequence of 10 sample-specific nucleotides. The PCR mixture (50 µl) contained 5X Taq-&Go™ ready to use PCR Mix (MP Biomedicals), 10 pM of each primer, and 2 µl of ten-fold diluted template DNA (or 2 µl of sterile water for the negative control). Five independent PCR reactions were performed for each sample. The PCR conditions were 94°C for 5 min; 30 cycles of 94°C for 30 s, 55°C or 50°C for 1 min, 72°C for 1 min, followed by 72°C for 10 min for primers 515F/907R or ITS1F/ITS2, respectively. PCR efficiency was checked by agarose gel electrophoresis (1%) and visualized after ethidium bromide staining under UV light with a GelDoc XR transilluminator (Bio-Rad).

The five independent PCR products were pooled, purified using the QIAquick PCR Purification Kit (Qiagen, France) and DNA was quantified by using a spectrophotometer (UV1800, Shimadzu) equipped with a TrayCell unit (Hellma) and by visualization and comparison with a mass ladder (Low DNA Mass Ladder, Invitrogen) on 1% agarose gels as described above. Two equimolar mixes (multiplexed samples) from the 18 samples, one for 16S rRNA genes and one for ITS amplicons, were prepared in 50 µl to reach a total concentration of 40 ng/µl. Pyrosequencing was performed by Beckman Coulter Genomics

(Danvers, MA, USA) using a 454/Roche GS-FLX Titanium system.

2.5. Analysis of 16S rRNA genes and ITS pyrosequencing data

2.5.1. Bacterial community analysis

Ribosomal Database Project (RDP, [35]) pyrosequencing pipeline was used for filtering and trimming of 16S rRNA gene reads based on sff-files. The presence of both forward and reverse primers (max. edit distance of 2) was checked to recover sense and anti-sense sequences. At the end, 482,163 reads passed the length (longer than 300 bp) and quality criteria (exclusion a reads with unresolved nucleotides and a low average quality score below 20) before being distributed into 18 samples according to their MIDs, the 16S rRNA gene reads were filtered and trimmed according to different criteria. Mothur 1.29.2 program [36] was used following previously described analyses steps [37]. Alignment of bacterial sequences used the SILVA database as a template. Based on distance matrix generation, the sequences were clustered and OTU (Operational Taxonomic Unit) were formed to 97% sequence similarity. To compare all sample conditions, a sub-sampling to 18,100 sequences per sample was performed. Finally, the taxonomic affiliation was obtained using the RDP database as a template.

2.5.2. Fungal community analysis

For the fungal community, 360,477 quality reads of ITS were obtained from the Mothur-based pipeline previously described by Counce et al. [38]. The fungal ITS sequences were distributed into 18 samples according to their MIDs, the reads were cleaned in Mothur 1.20.1 with the default parameters [36]. ITS1 was extracted by using Fungal ITS extractor version 2 [39] and a filtration step by length (> 100 bp) was performed. Quality ITS reads were aligned and clustered into OTUs (97% similarity) and consensus sequences were generated by using Uclust version 3.0 (parameter iddef = 2.97% similarity). Then sub-sampling to 9,100 sequences per sample was performed. Consensus sequences were assigned against the UNITE database by using the Basic Local Alignment search Tool algorithm Blastn version 2.2.2.3 [40].

2.5.3. Diversity analysis

Rarefaction curves were obtained using the Analytic Rarefaction 2.0 tool (Hunt Mountain Software); richness and diversity indices were calculated using EstimateS 9.1.0 [41]. The sequence data generated in this study were deposited in the NCBI Sequence Read Archive

(SRA) and are available under the BioProject ID: PRJNA269565 and PRJNA269566 for bacteria and fungi, respectively

2.6. Statistical analyses

Statistical analyses were performed using XLStat2012 software (Addinsoft). A two-way analysis of variance (ANOVA) followed by a Newman-Keuls post-hoc test was performed to determine statistical differences ($p < 0.05$) in the microbial communities among the six conditions, depending on time and the presence of a plant cover. Canonical correspondence analysis (CCA) followed by a Monte Carlo permutation test (1000 permutations) in XLStat-ADA 2015 (Addinsoft) was used to evaluate the relationship between environmental factors (plant biomass, Log_{10} of bacterial and fungal abundances obtained through qPCR and edaphic characteristics that were significantly different (ANOVA, $p < 0.05$) between samples). Two-dimensional hierarchical clustering was performed on standardized data set (z-score) of the most abundant ($> 2\%$ in at least one sample) bacterial and fungal OTUs (affiliation at 97% similarity; i.e. 24 bacterial and 37 fungal OTUs) in the three replicates of the six conditions. This was performed within the XLStat2011-Omics package (Addinsoft). Both samples and OTUs were clustered with Ascending Hierarchical Classification based on distance metric (euclidian), generating dendrograms. Heat maps were reordered by dendrogram clustering to relocate covarying OTUs and samples.

3. Results

3.1. Evolution of soil characteristics with time and plant colonisation

The edaphic parameters of the NM soil evolved over time as described in Table 1. Over the six years, pH values, C/N, available phosphorous, and humidity increased (or were higher at T12), while PAH concentrations, extractable Mn and Zn decreased with time, and total metal concentration remained unchanged.

3.2. Bacterial and fungal abundance

Over time, bacterial abundance (mean value of $1.69 \times 10^9 \pm 0.58 \times 10^9$ 16S rRNA gene copies g dw soil⁻¹) was about ten-fold higher than fungal abundance (mean value of $2.14 \times 10^8 \pm 0.15 \times 10^8$ 18S rRNA gene copies g dw soil⁻¹; Figure 1). Bacterial abundance was significantly higher ($p < 0.05$) at T0 and then decreased with time. Conversely, fungal abundance remained constant over time. In the Msm plots, no colonization of roots by mycorrhizal fungi was observed at T4, while the frequency of mycorrhizal roots reached 24% at T12 and arbuscules

were observed (Table 1). The presence of vegetation had no impact on bacterial abundance, while it enhanced fungal abundance: it was significantly higher ($p<0.05$) in MsmT4 samples when the biomass of alfalfa colonizing the Msm plots reached its highest value (1693 ± 156 g plot⁻¹; Figure 1). Consequently, the ratio fungi to bacteria was higher in MsmT4 soil compared to the other samples.

3.3. Bacterial and fungal richness and diversity

After filtering steps, 88% and 82% of 16S rRNA genes and ITS reads were kept for further analyses (Table 2). The rarefaction analysis revealed that plateau levels were not reached in any of the samples (Supplementary Figure S1). Based on complete data, we found ten times more OTUs for the bacterial (1,963 to 3,057 OTUs g dw soil⁻¹) than for the fungal community (284 to 592 OTUs g dw soil⁻¹; Table 2). Similarly, the Chao1 estimator of OTU richness predicted a ten-fold higher richness for the bacterial community (mean of 6,350 estimated species at T0) as compared to the fungal community (mean of 463 estimated species at T0; Table 2). Bacterial richness was neither influenced by the presence of a plant cover or by time (Table 2 and Supplementary Figure S1). By contrast, fungal richness increased significantly ($p<0.05$) with time and was significantly higher ($p<0.05$) under the MsmT4 condition (Table 2). The Shannon and inverse of Simpson indices of bacterial and fungal diversity increased significantly ($p<0.05$) with time and with the presence of plants at MsmT4, as well as at T12 for fungi (Table 2).

Taxonomic assignment of OTUs allowed us to identify at least 17 bacterial phyla and 5 fungal phyla (Figure 2). The bacterial community was mainly composed of *Proteobacteria* ($57.4 \pm 13.4\%$), *Actinobacteria* ($13.2 \pm 7.7\%$), *Bacteroidetes* ($6.0 \pm 1.9\%$) and unclassified bacteria ($2.4 \pm 1.0\%$). *Proteobacteria* were dominated by members of the *Gamma*- ($21.3 \pm 8.8\%$), *Alpha*- ($19.0 \pm 1.5\%$) and *Beta-Proteobacteria* ($15.0 \pm 6.4\%$) classes. The fungal community was mainly composed of *Ascomycota* ($53.6 \pm 9.4\%$), unclassified fungi ($41.3 \pm 8.2\%$) and *Basidiomycota* ($4.2 \pm 5.2\%$).

We found 24 and 37 dominant bacterial and fungal OTUs, represented by at least 2% of the sequences for at least one sample (Figure 3). They accounted for $28.6 \pm 7.5\%$ and $67.9 \pm 3.3\%$ of the bacterial 16S rRNA genes and fungal ITS sequences, respectively. At T0, the bacterial community was dominated by OTU11 affiliated to the *Thiobacillus* genus (*Beta-Proteobacteria*) and the fungal community was dominated by two OTUs affiliated to the

Talaromyces and *Scedosporium* genera (Ascomycota) and by a third OTU affiliated to unclassified fungi.

3.4. Modification of bacterial and fungal diversity over time

The relative abundance of some bacterial phyla significantly ($p < 0.05$) increased (*Bacteroidetes*, *Firmicutes*, *Nitrospirae*, *OD1*, *Gemmatimonadetes*, *Verrucomicrobia*) or decreased (*Gamma-Proteobacteria*, *Beta-Proteobacteria*) over time, while others were specifically increased at T4 (*Actinobacteria*, *Acidobacteria*, *Alpha-Proteobacteria*) (Figure 2). The relative abundance of *Glomeromycota* phylum increased significantly ($p = 0.007$) at T12 (Figure 2). Moreover, the relative abundance of two orders belonging to the *Ascomycota* (*Hypocreales* and *Pleosporales*) increased significantly ($p = 0.062$ and $p = 0.009$) over time. On the contrary, the relative abundance of *Eurotiales* decreased significantly ($p < 0.001$) over time.

Clustering of samples according to the dominant OTUs (horizontal dendrogram, Figure 3) showed higher similarity between samples at T0 than between samples at the two others times. This clustering of T0 samples is mainly due to 19 OTUs, mostly belonging to bacteria (Figure 3). The relative abundance of 12 bacterial and 10 fungal OTUs increased significantly ($p < 0.05$) over time (Figure 3 and Table S1). The 12 bacterial OTUs were affiliated to one *Actinobacteria* (*Arthrobacter*), one *Bacteroidetes* (*Ohtaekwangia*), one unclassified *Firmicute*, one *Gemmatimonadetes* (*Gemmatimonas*), one *Nitrospirae* (*Nitrospira*), two *Alpha-Proteobacteria* (unclassified *Erythrobacteraceae* and unclassified *Bradyrhizobiaceae*), three *Beta-Proteobacteria* (*Rhodocyclaceae*, *Methyloversatilis* and unclassified *Burkholderiales*), and two *Gamma-Proteobacteria* (unclassified and *Thiohalophilus*). The 10 fungal OTUs were affiliated to 5 different genera: 3 were affiliated to the *Ascomycota* phylum, the *Pezizomycotina* subphylum and the *Sordariomycetes* class (*Acremonium*, *Bionectria*, *Fusarium*), one belonged to unclassified *Basidiomycota*, and one was affiliated to unclassified fungi. The relative abundance of 2 bacterial OTUs affiliated to the *Acidobacteria* Gp7 and *Ohtaekwangia* genera (*Bacteroidetes*) and of 7 fungal OTUs affiliated to *Phoma*, unclassified *Plectosphaerellaceae* and unclassified fungi, increased significantly at T4 (Figure 3 and Table S1). The relative abundance of 9 bacterial and 3 fungal OTUs decreased significantly ($p < 0.05$) over time (Table S1). Seven of these bacterial OTUs were affiliated to *Gamma-Proteobacteria* and belonged to the orders *Acidithiobacillales* (*Thermithiobacillus*), *Chromatiales* (*Thiohalophilus*, *Halothiobacillus*), *Xanthomonadales* (unclassified *Xanthomonadaceae*, *Dyella*, *Rhodanobacter*), *Methylohalomonas* or unclassified *Gamma-*

Proteobacteria, one was affiliated to *Chloroflexi* (*Sphaerobacter*) and one to *Beta-Proteobacteria* (*Thiobacillus*). The 3 fungal OTUs belonged to the *Ascomycota* phylum, *Pezizomycotina* subphylum and *Eurotiomycetes* (*Talaromyces*) or *Sordariomycetes* (*Pseudallescheria*) class, and one was affiliated to unclassified fungi (Figure 3).

3.5. Modification of bacterial and fungal diversity with the presence of plant

Clustering of samples according to the dominant OTUs (horizontal dendrogram, Figure 3) showed high similarities in planted soils and between bare soils at T4 and T12. Clustering of the dominant OTUs (vertical dendrogram, Figure 3) showed that plant development increased the relative abundance of 15 and 11 OTUs, at T4 and T12 respectively, mainly belonging to fungi. The relative abundance of other bacterial OTUs was increased in bare soil (Figure 3 and Table S1). The relative abundance of some bacterial phyla increased significantly ($p<0.05$) (*Actinobacteria*) or decreased (*Bacteroidetes*, *Nitrospira*, *Planctomycetes*, *Gamma-Proteobacteria*, *OD1*, *Gemmatimonadetes*) with the presence of plants. Concerning fungal phyla, the relative abundance of *Zygomycota* and *Glomeromycota* increased significantly in alfalfa-planted plots ($p<0.05$). Moreover, the fungal order *Microascales* was significantly ($p<0.001$) less represented in planted soils (Figure 3).

The relative abundance of 5 bacterial and 10 fungal OTUs increased significantly ($p<0.05$) with the presence of plants (Figure 3 and Table S1). These bacterial OTUs were affiliated to one *Actinobacteria* (*Arthrobacter*), one unclassified *Firmicute*, one *Alpha-Proteobacteria* (unclassified *Bradyrhizobiaceae*) and two *Gamma-Proteobacteria* belonging to the *Xanthomonadaceae* family (unclassified and *Dyella*). Among the fungal OTUs, four belonged to the *Sordariomycetes* class among the *Pezizomycotina* subphylum (*Acremonium*, *Bionectria*, and *Fusarium* and unclassified *Plectosphaerellaceae*), and the other six were unclassified. The relative abundance of 6 bacterial and 5 fungal OTUs decreased significantly ($p<0.05$) with the presence of plants (Table S1). These bacterial OTUs were affiliated to one *Bacteroidetes* (*Ohtaekwangia*), one *Gemmatimonadetes* (*Gemmatimonas*), one *Nitrospiraea* (*Nitrospira*), one *Alpha-Proteobacteria* (unclassified *Erythrobacteraceae*), and two *Beta-Proteobacteria* (unclassified *Rhodocyclaceae* and *Burkholderiales*). Three fungal OTUs belonged to the *Sordariomycetes* (*Phoma*, *Pseudallescheria*, *Scedosporium*) and two to unclassified fungi (Figure 3).

3.6. Multivariate analysis

A Canonical correspondence analysis (CCA) was used to evaluate which variables best explained the microbial community compositions (Figure 4). CCA ordination of bacterial and fungal taxons (relative abundance at the order level) and environmental variables (plant biomass, microbial abundance and edaphic characteristics of the samples) depict relationships between the different plots and sampling dates. The first 2 CCA axes explained 69.63% of the total variance in the microbial diversity data (Figure 4). The six T0 samples grouped together and were separated from the other samples on the 1st axis (F1) discriminating the samples according to the effect of time. These T0 samples were discriminated according to higher bacterial abundance (16S), higher total calcium carbonate concentration, and higher extractable zinc concentration. At T4, the three samples from the planted plots MsmT4 were clearly separated from the BST4 samples on the 2nd axis (F2) discriminating the samples according to the presence of plants, higher total N content and fungal abundance (18S). At T12, although the samples were still discriminated, the difference between the Msm and BS plots was thinner. T12 samples were discriminated according to higher total metal contents, extractable Fe and Cu, higher pH, C/N ratio and available phosphorous.

At T0, four bacterial orders belonging to *Gamma-proteobacteria* (*Chromatiales*), *Beta-proteobacteria* (*Acidithiobacillales* and *Hydrogenophilales*) and *Elusimicrobia* phyla (*Elusimicrobiales*) and one fungal order (*Sporidiobolales*) belonging to *Basidiomycota* were the most correlated with high extractable Zinc concentration, high CaCO₃ and high bacterial abundance. Over 6 years part of the microbial community was modified and more affected by changes in edaphic parameters. At T12, five bacterial orders belonging to five different phyla: *Nitrospirae* (*Nitrospirales*), *Verrucomicrobia* (*Puniceicoccales*), *Actinobacteria* (*Euzebyales*), *Acidobacteria* (Unclassified *Acidobacteria_GPI6*), and *Chloroflexi* (Unclassified *Thermomicrobia*), and two fungal orders: Unclassified *Glomeromycota*, and Unclassified *Basidiomycota*, were highly correlated with higher C/N ratio, higher pH, higher P₂O₅, higher extractable iron and copper content and higher zinc and chrome concentrations. Moreover, seven bacterial orders belonging to five different phyla: *Actinobacteria* (*Solirubrobacterales*), *Thermoleophilales*, *Chloroflexi* (Unclassified *Chloroflexi*), *Acidobacteria* (Unclassified *Acidobacteria_GPI0*), *Planctomycetes* (Unclassified), and *Firmicutes* (Unclassified), were highly correlated with higher copper and lead concentrations. At T4, most fungal orders were positively correlated to alfalfa biomass and to the higher fungal abundance. Among these seven orders, six belong to the Ascomycota phylum and *Pezizomycotina* sub-phylum and belong to four different classes: *Dothideomycetes* (*Capnodiales*), *Eurotiomycetes* (*Onygenales*), *Leotiomycetes* (*Helotiales*, *Thelebolales*), and *Sordariomycetes*

(*Chaetosphaeriales*, and Unclassified *Hypocreomycetidae*), and one to the *Basidiomycota* phylum (*Tremellales*).

4. Discussion

4.1. Microbial diversity of the aged multi-contaminated NM soil

The NM soil had a high carbon concentration, neutral pH, low nutrient concentrations and high concentrations of PAHs and heavy metals that explain its low fertility and high toxicity [25]. This soil represented a unique ecosystem: contamination dated back to ca. 100 years, so selection pressure on soil microorganisms had lasted for decades. While the short-term impact of contamination tends to decrease microbial abundance, richness and diversity, we hypothesized that in such aged multi-contaminated soil the microbial community would adapt over time toward a unique diversity as compared to unpolluted soil.

In NM soil, bacteria were approximately ten-fold more abundant than fungi regardless of time or of the presence of vegetation. The same trend was previously observed in the same soil [42], and in many uncontaminated soils but with a higher difference between bacterial and fungal abundance levels. In fact, the environments harboring such high fungal abundance and a high fungal to bacterial ratio are usually rich in carbon, like the litter horizon in forest soils [43]. The Chao1 index evidenced higher bacterial than fungal species richness in our NM soil samples, as previously shown in other soils [44]. Freshly contaminated soils usually harbor low bacterial richness, evenness and diversity after spiking [8]. Yet bacterial richness and diversity in our NM soil was relatively high and comparable to the levels found in a diesel-contaminated soil in Poland [45] and in an aged PAH-contaminated soil from a coking plant wasteland in Beijing, China [46], using a similar sequencing effort. The result from our NM soil suggests a long-term adaptation of the microbial community toward a diversified and pollutant-resistant community. In addition, fungal richness was close to the richness found in forest soils [38], but fungal diversity was relatively low and close to the level found in a soil highly contaminated by hydrocarbons [6].

The dominance of 3 bacterial phyla (*Proteobacteria*, *Actinobacteria*, and *Bacteroidetes*) and *Ascomycota* and unclassified fungi with poor representation of *Basidiomycota* is in accordance with fingerprinting analyses of dominant bacterial and fungal species inhabiting the NM soil [47];[23]. Overall, at the phylum level, the bacterial and fungal community diversities were close to other soil types. However the relative proportion of each bacterial phylum seems to be specific to polluted environments, contrary to fungal ones. *Proteobacteria* are usually the dominant phylum in contaminated soils [7]. The co-occurrence

of *Alpha*-, *Beta*- and *Gamma-Proteobacteria* in similar relative proportions (15 - 21%) seems typical of polluted soils: *Alpha*- are usually in higher proportions in non-contaminated soils, while *Beta*- and *Gamma*- subdivisions are favored in hydrocarbon-contaminated soils [48]. Furthermore, *Actinobacteria* was the second most abundant phylum in our NM soil as found in other heavy-metal contaminated soils [49];[7]. *Actinobacteria* were previously described in extremely arid environments [50], with similar properties (low nutrient and low carbon availability) to aged multi-contaminated soil. Finally, some genera belonging to the *Actinobacteria* are able to degrade PAH compounds [51]. Members of *Acidobacteria* were found scarce in our NM soil as found in other PAH-contaminated soil [46], but they were largely less represented than in non-polluted soils [52];[53]. Overall, similar fungal phylum diversity was also found in non-polluted soils [54].

4.2. Impact of edaphic parameters on soil microbial diversity over time

Although this result could be specific to this particular site under specific climatic condition, the NM soil monitoring over 6 years revealed interesting relationship between bacterial and fungal diversity and edaphic parameters. Our canonical correspondence analysis showed that a temporal shift occurred in the composition of the soil microbial community, with some taxa correlated with modifications in edaphic parameters.

Among the different soil parameters that increased simultaneously over time (C/N ratio, available phosphorous, extractable iron, copper, chrome, zinc concentrations), the pH can be an important driver of the bacterial community but not of the fungal one [55]. Most orders belonging to *Acidobacteria* were favored over time, with subgroup-16 positively correlated with pH values. Unlike the other subgroups the relative abundance of *Acidobacteria* subgroup 1 was greater at T0 and was negatively correlated with the pH level, as previously found [55]. Despite a low availability and mobility of PAH and HM, and significant decrease of these fractions over time [25], we observed a significant decrease in total PAH (c.a. 40% dissipation) and extractable Zn concentrations over the 6 years of monitoring that could also contribute to explain changes in microbial diversity. Xu et al. [46] suggested that a decrease in pollutant concentration allowed for the development of members of the *Acidobacteria* phylum in a PAH-contaminated soil. The PAH decrease could also explain the emergence of PAH-degrading microorganisms that benefited from the biodegradation of the pollutant. As previously observed [47];[23], OTUs affiliated to the *Arthrobacter* and *Fusarium* genera increased over time in NM soil samples. Members of the *Arthrobacter* and *Fusarium* genera have been found in hydrocarbon-contaminated environments [56];[57] and could be potential

PAH degraders [51];[58]. The relative abundance of the *Nitrospirae* phylum, one *Nitrospira* OTU and the *Nitrospirales* order increased over time. On CCA it was positively correlated to the C/N ratio, and negatively correlated to total nitrogen content. This bacterial group is more competitive with other nitrite-oxidizing bacteria (i.e. *Nitrobacter*) in soils with low N availability [59]. Relative abundance of the *Gemmatimonadetes* phylum and one OTU affiliated to the *Gemmatimonas* genus also increased over time. Although members of this phylum are frequently detected in soils and more prevalent in dry soils with poor nutrient availability and low microbial activity [60], their ecological properties are still unclear. On the contrary, the relative abundance of other microbial taxa decreased over time. The fungal *Eurotiales* order and one OTU affiliated to *Talaromyces* (*Ascomycota*; *Eurotiales*) were more abundant at T0. This fungal taxon seems to be well adapted to high PAH contamination levels, as similar strains were isolated from an oil-contaminated soil and were able to grow at high PAH concentrations [61]. Within the bacterial community, the relative abundance of the *Gamma-Proteobacteria* phylum and of seven OTUs affiliated to *Gamma-Proteobacteria* decreased over time. Among these, *Thiohalophilus*, *Halothiobacillus*, *Methylohalomonas*, known as halophilic bacteria, could be repressed by the leaching of extractable metals and sulfate salts over time. The relative abundance of *Thiobacillus* (*Beta-proteobacteria*; *Hydrogenophilales*) was the most abundant bacterial OTU at T0 (ca. 15%), but also decreased over time. *Thiobacillus* was previously detected in heavy metal-contaminated soils [7]. The implication of the *Thiobacillus* genus in PAH degradation has also been described [62] although our results indicate a drastic decrease over time.

4.3. Impact of vegetation on soil microbial abundance and diversity

Our results show that the plant cover influenced microbial abundance in the NM soil, especially at T4 when the plant cover was the thickest. Although the role of root exudates in regulating soil fungal community composition and diversity is well known [63], the impact of the rhizosphere on increasing soil fungal abundance, as found here, is less documented as compared to bacteria [63].

The plant cover had an unexpected impact on fungal and bacterial community structure in the NM soil. At T4, PAH dissipation was significantly lower in presence of plants than in bare soil, probably explained by the release of more easily degradable carbon compounds in root exudates [64]. At T4, the community diversity is contrasted between bare and alfalfa planted soils, and the multivariate analysis showed that plant biomass is the dominant explicative factor compared to PAH content. Bacterial and fungal diversity indices increased with the

presence of vegetation, as previously shown [47];[23]. Although the impact of plants varies depending on seasons, usually fungal and bacterial diversity usually decrease in the rhizospheric or planted soils as compared to the bulk or unplanted controls [65];[66]. Lower microbial diversity in the rhizosphere could be due to selective pressure from root exudates. In aged hydrocarbon- and heavy metal-contaminated soils that are extremely oligotrophic environments, microbial activity is limited by nutrient availability [67]. Most studies describe the impact of plants on microbial diversity in non-contaminated and/or nutrient-rich soils (e.g. agricultural or forest soils). The presence of plants on a nutrient-poor soil could increase diversity due to the input of diversified carbon and nutrient sources, such as carbohydrates from exudates and plant debris preferentially degraded by bacteria and fungi, respectively. Moreover, the development of plants and roots creates numerous novel niches favoring the establishment of more diverse microbial communities. As previously shown, the functional community was also impacted by plant covers because enzymatic activity measured (FDase, arylamidase, aryl sulfatase, cellulose, urease...) were higher in vegetalized NM soil than in bare soil [42].

Our analyses allowed us to discriminate between planted soil and bare soil plots and to identify microbial taxa whose relative abundance levels were differently impacted by the presence of plants. *Actinomycetales* and *Arthrobacter*, the most represented order and most abundant OTU of the bacterial community, were positively correlated to alfalfa biomass. Increased numbers of *Actinobacteria* in the rhizosphere of different plant species such as maize [66] and strawberry [68] are documented. In addition, the *Bradyrhizobiaceae* family and the *Rhizobiales* order were among the dominant bacterial OTUs favored by the development of alfalfa. These nitrogen-fixing bacteria are known to be abundant in the rhizosphere of legumes due to their symbiotic association [69]. However we never observed nodules throughout the monitoring period. Finally, the relative abundance of the *Nitrospirae* phylum and *Nitrospira* OTUs, a well-known nitrite-oxidizing group [70], decreased in alfalfa-planted plots. Competition for nitrogen between plants and nitrifying bacteria has already been reported in unfertilized nutrient-poor soil [71].

Plants have a higher impact on fungal than on bacterial community structure. According to our multivariate analysis, most fungal orders were positively correlated to the presence of plants (alfalfa biomass). The relative abundance of *Ascomycota* belonging to the *Pezizomycotina* sub-division and affiliated to various classes, *i.e.* *Sordariomycetes*, *Leotiomycetes*, *Dothideomycetes*, and *Eurotiomycetes*, increased. Among the *Sordariomycetes*, the *Chaetosphaeriales* and the unclassified *Hypocreales* were the most

important orders allowing for discrimination between treatments. Among these orders, the relative abundance of some dominant fungal OTUs affiliated to the *Acremonium*, *Bionectria*, and *Fusarium* genera significantly increased with the presence of alfalfa. While the *Acremonium* genus is well-known as a fungal endophyte of plants that improves mineral nutrition [72], *Bionectria* and *Fusarium* species are often plant pathogens [73];[74]. Conversely, in bare soil the *Microascales* order was the only one whose relative abundance increased. Although alfalfa was co-inoculated with two mycorrhizal *Glomus* strains, *Glomeromycota* was not a dominant phylum, but its relative abundance was higher in planted plots. Our results show that unclassified *Glomeromycota* were more correlated with planted soil after 6 years than after 2 years. Colonization of roots by mycorrhizal fungi was detected only after 6 years, suggesting potential initial toxicity of the contaminated soil towards mycorrhizal fungi, as suggested previously [25].

5. Conclusion

To conclude, the high fungal and bacterial abundance levels in the NM soil probably reflect its high carbon content due to contamination. Microbial diversity and richness in the aged-contaminated soil were surprisingly high and closer to that found in an unstressed soil than in freshly contaminated soil. A specific microbial community settled over time through modifications of edaphic parameters, pollution content and due to the presence of plants. Over time the C/N increased, due to a decrease of soil nitrogen content (no fertilization), favoring *Nitrospirae* phylum, and nitrogen-fixing bacteria in planted soil. Over time the pH increased, probably due to sulphate leaching that favoured some *Acidobacteria* groups and repressed halophilic bacteria. In 6 years the decrease of PAH content by c.a. 40% favoured the appearance of few taxa previously described as PAH degraders. The modification of pollution content and bioavailability contributed probably to reduce environment toxicity and led to adaptation in microbial community composition. The alfalfa cover structured the microbial community, probably due to a rhizosphere effect while this partly inhibited the PAH dissipation. Indeed vegetation (i) selected a specific community by releasing root exudates or plant debris that input nutrient sources more easily degraded than PAHs in the oligotrophic soil; (ii) increased bacterial diversity and (iii) favoured most fungal species compared to bare soil.

Acknowledgements

This study was funded by OTELo (Observatoire Terre et Environnement de Lorraine) and the

Agence Nationale de la Recherche (ANR-2008-CESA-010 and ANR-13-JSV7-0007_01). The authors thank the GISFI (www.gisfi.fr) for the access to the experimental site at Homécourt and Annie Buchwalter for English editing.

Tables

Table 1. Summary of the soil characteristics for the six conditions (mean and standard deviation, n=3). Different letters indicate significant differences ($p<0.05$) among treatments by two-way analysis of variance (ANOVA) followed by Newmans-Keuls multiple comparison test (nd: not determined).

Table 2. Pyrosequencing data characteristics, and richness and diversity estimators (3% cutoff) of the bacterial and fungal communities in the six different conditions (means and standard deviations, n=3): BS (Bare soil) and Msm (planted with *Medicago sativa* and inoculated with mycorrhizal fungi) at three time-points (T0: setting up, T4: after 2 years and T12: after 6 years). We created OTUs based on 97% sequence similarity (OTU₉₇) and randomly subsampled 16S rRNA genes and ITS data to 18,100 and 9,100 reads *per* sample. A two-way analysis of variance (ANOVA, $p<0.05$) followed by Newman-Keuls multiple comparison test was performed on richness and diversity estimator data to determine statistical differences (lowercase) between treatments.

Supplemental Table S1. Affiliation at 97% similarity of the most abundant (>2% in at least one sample) bacterial and fungal OTUs in the six conditions (mean and standard deviation, n=3). Two-way analysis of variance (ANOVA) followed by Neuman-Keuls multiple comparison test to study the effect of plant cover and time on the most abundant bacterial (16S) and fungal (ITS) OTUs. Sections colored in dark grey / grey / light grey and white mean respectively that OTUs were significantly ($p<0.05$) favored at all time-points / favored at T4 / inhibited or not influenced by the plant cover or by time.

Figure captions

Figure 1. Abundance of fungal and bacterial (18S and 16S rRNA gene copy numbers) communities and alfalfa biomass over time (T0: September 2005, T4: September 2007 and T12: September 2011) for the two BS (bare soil) and Msm (planted with *Medicago sativa* and

inoculated with mycorrhizal fungi) treatments. Different letters indicate significant differences among plots ($p < 0.05$) resulting from a two-way analysis of variance (ANOVA) followed by Neuman-Keuls multiple comparison test ; abc, AB and A'B' refer respectively to statistics on 16S rRNA gene copy numbers. g of dry weight soil⁻¹, 18S rRNA gene copy numbers. g of dry weight soil⁻¹ and alfalfa biomass.

Figure 2. Relative abundance of the dominant (represented by more than 10 sequences) fungal and bacterial phyla in the six conditions (means of 3 replicates).

Figure 3. Clustering analysis generating double dendrograms (vertically: OTUs dendrogram and horizontally: sample conditions dendrogram) and heat map representation reflecting data values of the most abundant (>2% in at least one sample) bacterial and fungal OTUs in the six conditions (affiliation at 97% similarity). The relative abundance of 24 bacterial and 37 fungal OTUs were normalised (z-score) to compare studied conditions (3 replicates per condition).

Figure 4. Canonical correspondence analysis (CCA) ordination of bacterial (in green) and fungal (in blue) taxons (relative abundance of the dominant orders represented by more than 10 sequences), and environmental variables (abundance of fungal 18S and bacterial 16S rRNA gene copy numbers, biomass of dry alfalfa and edaphic characteristics (in red), showing the relationships between the different plots and sampling dates (3 replicates of the 6 conditions, in black bold type). Numbers in *Italic* correspond to the quantity of OTUs that belong to the taxon. Fungal and bacterial orders: *Acidi.*: *Acidimicrobiales*, *Acidithio.*: *Acidithiobacillales*, *Actinomyc.*: *Actinomycetales*, *Altero.*: *Alteromonadales*, *Bacill.*: *Bacillales*, *Bdello.*: *Bdellovibrionales*, *Burkhol.*: *Burkholderiales*, *Caldi.*: *Caldilineale*, *Capno.*: *Capnodiales*, *Caulo.*: *Caulobacterales*, *Chaetos.*: *Chaetosphaeriales*, *Chlamy.*: *Chlamydiales*, *Chroma.*: *Chromatiales*, *Clostri.*: *Clostridiales*, *Dehalo.*: *Dehalococcoidales*, *Desulfur.*: *Desulfuromonadales*, *Elusi.*: *Elusimicrobiales*, *Euro.*: *Eurotiales*, *Euzeb.*: *Euzebyales*, *Flavo.*: *Flavobacteriales*, *Gemma.*: *Gemmatimonadales*, *Helot.*: *Helotiales*, *Hydrogeno.*: *Hydrogenophilales*, *Hypo.*: *Hypocreales*, *Lactob.*: *Lactobacillales*, *Legion.*: *Legionellales*, *Methylo.*: *Methylococcales*, *Microasc.*: *Microascales*, *Mortier.*: *Mortierellales*, *Myxoc.*: *Myxococcales*, *Nitro.*: *Nitrospirales*, *Nitroso.*: *Nitrosomonadales*, *Oceano.*: *Oceanospirillales*, *Onyg.*: *Onygenales*, *Opitu.*: *Opitutales*, *Parvu.*: *Parvularculales*, *Pseudo.*: *Pseudomonadales*, *Punicei.*: *Puniceicoccales*, *Rhizo.*: *Rhizobiales*, *Rhodo.*: *Rhodobacterales*, *Rhodocyc.*: *Rhodocyclales*, *Rhodospiri.*: *Rhodospirillales*, *Rickett.*: *Rickettsiales*, *Soliru.*: *Solirubrobacterales*, *Sphaero.*: *Sphaerobacterales*, *Sphingo.*: *Sphingomonadales*,

Sphingobacter.: *Sphingobacteriales*, *Sporidio*.: *Sporidiobolales*, *Syntro*.:
Syntrophobacterales, *Theleb*.: *Thelebolales*, *Thermo*.: *Thermoanaerobacterales*, *Thermoleo*.:
Thermoleophilales, *Thiotric*.: *Thiotrichales*, *Treme*.: *Tremellales*, *Un. Acido_Gp1*:
Unclassified Acidobacteria_GP1, *Un. Acido_Gp10*: *Unclassified Acidobacteria_GP10*, *Un.*
Acido_Gp16: *Unclassified Acidobacteria_GP16*, *Un. Acido_Gp17*: *Unclassified*
Acidobacteria_GP17, *Un. Acido_Gp3*: *Unclassified Acidobacteria_GP3*, *Un. Acido_Gp6*:
Unclassified Acidobacteria_GP6, *Un. Acido_Gp7*: *Unclassified Acidobacteria_GP7*, *Un.*
Actino.: *Unclassified Actinobacteria*, *Un. Bacteria*.: *Unclassified Bacteria*, *Un. Bacteroid*.:
Unclassified Bacteroidetes, *Un. Basidio*.: *Unclassified Basidiomycota*, *Un. Chloro*.:
Unclassified Chloroflexi, *Un. Firmi*. *Unclassified Firmicutes*, *Un. Fungi*: *Unclassified Fungi*,
Un. Glomero.: *Unclassified Glomeromycota*, *Un. Hypocre*.: *Unclassified*
Hypocreomycetidae, *Un. OD1*: *Unclassified OD1*, *Un. Plancto*.: *Unclassified*
Planctomycetes, *Un. Strepto*.: *Unclassified Streptophyta*, *Un. Thermo*.: *Unclassified*
Thermomicrobia, *Un. α -proteo*.: *Unclassified Alphaproteobacteria*, *Un. γ -proteo*.:
Unclassified Gammaproteobacteria, *Un. δ -proteo*.: *Unclassified Deltaproteobacteria*,
Xantho.: *Xanthomonadales*. Edaphic characteristics: extractable copper, iron, and zinc (Cu
ext, Fe ext, and Zn ext, respectively), 16 PAHs (mg/Kg of dry weight soil).

Supplementary Figure S1. Rarefaction curve representing the number of operational
taxonomic units (OTUs) depending on the number of 16S rRNA genes (A) and ITS (B)
sequences amplified from the six conditions. After OTU formation to 97% similarity, sub-
sampling was performed to 18,100 and 9,100 sequences *per* sample for bacterial and fungal
data, respectively.

References

1. Sandrin TR, Maier RM (2003) Impact of Metals on the Biodegradation of Organic Pollutants. *Environ Health Perspect* 111:1093.
2. Cerniglia CE, Heitkamp MA (1989) Microbial degradation of polycyclic aromatic hydrocarbons (PAH) in the aquatic environment. *Metab Polycycl Aromat Hydrocarb Aquat Environ* 41–68.
3. Peng R-H, Xiong A-S, Xue Y, et al. (2008) Microbial biodegradation of polyaromatic hydrocarbons. *FEMS Microbiol Rev* 32:927–955.
4. Schimel J, Balser TC, Wallenstein M (2007) Microbial Stress-Response Physiology and Its Implications for Ecosystem Function. *Ecology* 88:1386–1394. doi: 10.2307/27651246

5. Paissé S, Coulon F, Goñi-Urriza M, et al. (2008) Structure of bacterial communities along a hydrocarbon contamination gradient in a coastal sediment. *FEMS Microbiol Ecol* 66:295–305. doi: 10.1111/j.1574-6941.2008.00589.x
6. Bell TH, El-Din Hassan S, Lauron-Moreau A, et al. (2014) Linkage between bacterial and fungal rhizosphere communities in hydrocarbon-contaminated soils is related to plant phylogeny. *ISME J* 8:331–343.
7. Gołębiewski M, Deja-Sikora E, Cichosz M, et al. (2014) 16S rDNA Pyrosequencing Analysis of Bacterial Community in Heavy Metals Polluted Soils. *Microb Ecol* 67:635–647. doi: 10.1007/s00248-013-0344-7
8. Ding GC;Heuer H;Smalla K (2012) Dynamics of bacterial communities in two unpolluted soils after spiking with phenanthrene: soil type specific and common responders. *Front Microbiol* 3:290.
9. Zhou H, Wong AY, Yu RK, et al. (2009) Polycyclic Aromatic Hydrocarbon-Induced Structural Shift of Bacterial Communities in Mangrove Sediment. *Microb Ecol* 58:153–160. doi: 10.1007/s00248-008-9456-x
10. Gremion F, Chatzinotas A, Kaufmann K, Von Sigler W, Harms H (2004) Impacts of heavy metal contamination and phytoremediation on a microbial community during a twelve-month microcosm experiment. *FEMS Microbiol Ecol Wiley* 48:273–283.
11. Fierer N, Bradford MA, Jackson RB (2007) Toward an ecological classification of soil bacteria. *Ecology* 88:1354–1364. doi: 10.1890/05-1839
12. Wenzel W (2009) Rhizosphere processes and management in plant-assisted bioremediation (phytoremediation) of soils. *Plant Soil* 321:385–408. doi: 10.1007/s11104-008-9686-1
13. Binet P, Portal JM, Leyval C (2000) Fate of polycyclic aromatic hydrocarbons (PAH) in the rhizosphere and mycorrhizosphere of ryegrass. *Plant Soil* 227:207–213. doi: 10.1023/A:1026587418611
14. Corgié SC, Joner EJ, Leyval C (2003) Rhizospheric degradation of phenanthrene is a function of proximity to roots. *Plant Soil* 257:143–150. doi: 10.1023/A:1026278424871
15. Haritash AK, Kaushik CP (2009) Biodegradation aspects of Polycyclic Aromatic Hydrocarbons (PAHs): A review. *J Hazard Mater* 169:1–15. doi: 10.1016/j.jhazmat.2009.03.137
16. Sun M, Fu D, Teng Y, et al. (2011) In situ phytoremediation of PAH-contaminated soil by intercropping alfalfa (*Medicago sativa* L.) with tall fescue (*Festuca arundinacea* Schreb.) and associated soil microbial activity. *J Soils Sediments* 11:980–989. doi: 10.1007/s11368-011-0382-z
17. Muratova AY, Turkovskaya OV, Hübner T, Kusch P (2003) Studies of the Efficacy of Alfalfa and Reed in the Phytoremediation of Hydrocarbon-Polluted Soil. *Appl Biochem Microbiol* 39:599–605. doi: 10.1023/A:1026238720268

18. Kawasaki A, Watson E, Kertesz M (2012) Indirect effects of polycyclic aromatic hydrocarbon contamination on microbial communities in legume and grass rhizospheres. *Plant Soil* 358:169–182. doi: 10.1007/s11104-011-1089-z
19. Joner EJ, Leyval C (2001) Influence of arbuscular mycorrhiza on clover and ryegrass grown together in a soil spiked with polycyclic aromatic hydrocarbons. *Mycorrhiza* 10:155–159. doi: 10.1007/s005720000071
20. Leyval C, Turnau K, Haselwandter K (1997) Effect of heavy metal pollution on mycorrhizal colonization and function: physiological, ecological and applied aspects. *Mycorrhiza* 7:139–153. doi: 10.1007/s005720050174
21. Cébron A, Beguiristain T, Bongoua-Devisme J, et al. (2015) Impact of clay mineral, wood sawdust or root organic matter on the bacterial and fungal community structures in two aged PAH-contaminated soils. *Environ Sci Pollut Res* 1–15.
22. Tejeda-Agredano MC, Gallego S, Vila J, et al. (2013) Influence of the sunflower rhizosphere on the biodegradation of PAHs in soil. *Soil Biol Biochem* 57:830–840. doi: 10.1016/j.soilbio.2012.08.008
23. Thion C, Cébron A, Beguiristain T, Leyval C (2012) Long-term in situ dynamics of the fungal communities in a multi-contaminated soil are mainly driven by plants. *FEMS Microbiol Ecol* 82:169–181. doi: 10.1111/j.1574-6941.2012.01414.x
24. Buée M, De Boer W, Martin F, et al. (2009) The rhizosphere zoo: An overview of plant-associated communities of microorganisms, including phages, bacteria, archaea, and fungi, and of some of their structuring factors. *Plant Soil* 321:189–212. doi: 10.1007/s11104-009-9991-3
25. Ouvrard S, Barnier C, Bauda P, et al. (2011) In Situ Assessment of Phytotechnologies for Multicontaminated Soil Management. *Int J Phytoremediation* 13:245–263. doi: 10.1080/15226514.2011.568546
26. Schüßler A, Walker C (2010) The Glomeromycota: a species list with new families and new genera.
27. Koske RE, Gemma JN (1989) A modified procedure for staining roots to detect VA mycorrhizas. *Mycol Res* 92:486–488. doi: 10.1016/S0953-7562(89)80195-9
28. Trouvelot A, Kough JL, Gianinazzi-Pearson V (1986) Mesure du taux de mycorhization VA d'un système racinaire. Recherche de méthodes d'estimation ayant une signification fonctionnelle. 217–221.
29. Cébron A, Norini M-P, Beguiristain T, Leyval C (2008) Real-Time PCR quantification of PAH-ring hydroxylating dioxygenase (PAH-RHD α) genes from Gram positive and Gram negative bacteria in soil and sediment samples. *J Microbiol Methods* 73:148–159. doi: 10.1016/j.mimet.2008.01.009
30. Lueders T, Wagner B, Claus P, Friedrich MW (2004) Stable isotope probing of rRNA and DNA reveals a dynamic methylotroph community and trophic interactions with fungi and protozoa in oxic rice field soil. *Environ Microbiol* 6:60–72. doi: 10.1046/j.1462-2920.2003.00535.x

31. Felske A, Akkermans ADL, De Vos WM (1998) Quantification of 16S rRNAs in Complex Bacterial Communities by Multiple Competitive Reverse Transcription-PCR in Temperature Gradient Gel Electrophoresis Fingerprints. *Appl Environ Microbiol* 64:4581–4587.
32. Turner S, Pryer KM, Miao VP, Palmer JD (1999) Investigating deep phylogenetic relationships among cyanobacteria and plastids by small subunit rRNA sequence analysis. *J Eukaryot Microbiol* 46:327–338.
33. Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes- application to the identification of mycorrhizae and rusts. *Mol Ecol* 2:113–118.
34. White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, White JJSJ (eds) *PCR Protoc.* Academic Press, San Diego, pp 315–322
35. Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naïve Bayesian Classifier for Rapid Assignment of rRNA Sequences into the New Bacterial Taxonomy. *Appl Environ Microbiol* 73:5261–5267.
36. Schloss PD, Westcott SL, Ryabin T, et al. (2009) Introducing mothur: Open-Source, Platform-Independent, Community-Supported Software for Describing and Comparing Microbial Communities. *Appl Environ Microbiol* 75:7537–7541.
37. Sogin ML, Morrison HG, Huber JA, et al. (2006) Microbial diversity in the deep sea and the underexplored “rare biosphere.” *Proc Natl Acad Sci* 103:12115–12120.
38. Counce A, Caël O, Bach C, et al. (2013) Below-ground fine-scale distribution and soil versus fine root detection of fungal and soil oomycete communities in a French beech forest. *Fungal Ecol* 6:223–235. doi: 10.1016/j.funeco.2013.01.002
39. Nilsson RH, Veldre V, Hartmann M, et al. (2010) An open source software package for automated extraction of ITS1 and ITS2 from fungal ITS sequences for use in high-throughput community assays and molecular ecology. *Fungal Ecol* 3:284–287. doi: 10.1016/j.funeco.2010.05.002
40. Altschul SF, Gish W, Miller W, et al. (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410. doi: 10.1016/S0022-2836(05)80360-2
41. Colwell R (2013) EstimateS: Statistical estimation of species richness and shared species from samples. 2006. Consult. En [Httpvicroy Eeb Uconn Eduestimates](http://vicroy.eeb.uconn.edu/estimates)
42. Cébron A, Cortet J, Criquet S, et al. (2011) Biological functioning of PAH-polluted and thermal desorption-treated soils assessed by fauna and microbial bioindicators. *Spec Issue Environ Microbiol* 162:896–907. doi: 10.1016/j.resmic.2011.02.011
43. Baldrian P, Kolarik M, Stursova M, et al. (2012) Active and total microbial communities in forest soil are largely different and highly stratified during decomposition. *ISME J* 6:248–258.

44. Gottel NR, Castro HF, Kerley M, et al. (2011) Distinct Microbial Communities within the Endosphere and Rhizosphere of *Populus deltoides* Roots across Contrasting Soil Types. *Appl Environ Microbiol* 77:5934–5944. doi: 10.1128/AEM.05255-11
45. Sutton NB, Maphosa F, Morillo JA, et al. (2013) Impact of Long-Term Diesel Contamination on Soil Microbial Community Structure. *Appl Environ Microbiol* 79:619–630. doi: 10.1128/AEM.02747-12
46. Xu Y, Sun G-D, Jin J-H, et al. (2014) Successful bioremediation of an aged and heavily contaminated soil using a microbial/plant combination strategy. *J Hazard Mater* 264:430–438. doi: 10.1016/j.jhazmat.2013.10.071
47. Cébron A, Beguiristain T, Faure P, et al. (2009) Influence of Vegetation on the In Situ Bacterial Community and Polycyclic Aromatic Hydrocarbon (PAH) Degraders in Aged PAH-Contaminated or Thermal-Desorption-Treated Soil. *Appl Environ Microbiol* 75:6322–6330.
48. Labbé D, Margesin R, Schinner F, Whyte LG, Greer CW (2007) Comparative phylogenetic analysis of microbial communities in pristine and hydrocarbon-contaminated Alpine soils. *FEMS Microbiol Ecol Wiley* 59:466–475.
49. Gremion F, Chatzinotas A, Harms H (2003) Comparative 16S rDNA and 16S rRNA sequence analysis indicates that Actinobacteria might be a dominant part of the metabolically active bacteria in heavy metal-contaminated bulk and rhizosphere soil. *Environ Microbiol* 5:896–907. doi: 10.1046/j.1462-2920.2003.00484.x
50. Köberl M, Müller H, Ramadan EM, Berg G (2011) Desert Farming Benefits from Microbial Potential in Arid Soils and Promotes Diversity and Plant Health. *PLoS ONE* 6:e24452. doi: 10.1371/journal.pone.0024452
51. Kallimanis A, Frillingos S, Drainas C, Koukkou A (2007) Taxonomic identification, phenanthrene uptake activity, and membrane lipid alterations of the PAH degrading *Arthrobacter* sp. strain Sphe3. *Appl Microbiol Biotechnol* 76:709–717. doi: 10.1007/s00253-007-1036-3
52. Roesch LFW, Fulthorpe RR, Riva A, et al. (2007) Pyrosequencing enumerates and contrasts soil microbial diversity. *ISME J* 1:283–290.
53. Nacke H, Thürmer A, Wollherr A, et al. (2011) Pyrosequencing-Based Assessment of Bacterial Community Structure Along Different Management Types in German Forest and Grassland Soils. *PLoS ONE* 6:e17000. doi: 10.1371/journal.pone.0017000
54. Buée M, Reich M, Murat C, et al. (2009) 454 Pyrosequencing analyses of forest soils reveal an unexpectedly high fungal diversity. *New Phytol* 184:449–456. doi: 10.1111/j.1469-8137.2009.03003.x
55. Rousk J, Baath E, Brookes PC, et al. (2010) Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME J* 4:1340–1351.
56. Panov AV, Esikova TZ, Sokolov SL, et al. (2013) Influence of soil pollution on the composition of a microbial community. *Microbiology* 82:241–248. doi: 10.1134/S0026261713010116

- 791 57. Lladó S, Gràcia E, Solanas AM, Viñas M (2013) Fungal and bacterial microbial
792 community assessment during bioremediation assays in an aged creosote-polluted soil.
793 *Soil Biol Biochem* 67:114–123. doi: 10.1016/j.soilbio.2013.08.010
- 794 58. Hong JW, Park JY, Gadd GM (2010) Pyrene degradation and copper and zinc uptake
795 by *Fusarium solani* and *Hypocrea lixii* isolated from petrol station soil. *J Appl*
796 *Microbiol* 108:2030–2040. doi: 10.1111/j.1365-2672.2009.04613.x
- 797 59. Attard E, Poly F, Commeaux C, et al. (2010) Shifts between *Nitrospira*- and
798 *Nitrobacter*- like nitrite oxidizers underlie the response of soil potential nitrite
799 oxidation to changes in tillage practices. *Environ Microbiol* 12:315–326.
- 800 60. DeBruyn JM, Nixon LT, Fawaz MN, et al. (2011) Global Biogeography and
801 Quantitative Seasonal Dynamics of Gemmatimonadetes in Soil. *Appl Environ*
802 *Microbiol* 77:6295–6300.
- 803 61. Reyes-César A, Absalón Á, Fernández F, et al. (2014) Biodegradation of a mixture of
804 PAHs by non-ligninolytic fungal strains isolated from crude oil-contaminated soil.
805 *World J Microbiol Biotechnol* 30:999–1009. doi: 10.1007/s11274-013-1518-7
- 806 62. Singleton D, Jones M, Richardson S, Aitken M (2013) Pyrosequence analyses of
807 bacterial communities during simulated in situ bioremediation of polycyclic aromatic
808 hydrocarbon-contaminated soil. *Appl Microbiol Biotechnol* 97:8381–8391. doi:
809 10.1007/s00253-012-4531-0
- 810 63. Berg G, Smalla K (2009) Plant species and soil type cooperatively shape the structure
811 and function of microbial communities in the rhizosphere. *FEMS Microbiol Ecol*
812 68:1–13. doi: 10.1111/j.1574-6941.2009.00654.x
- 813 64. Tian L, Ma P, Zhong J-J (2003) Impact of the presence of salicylate or glucose on
814 enzyme activity and phenanthrene degradation by *Pseudomonas mendocina*. *Process*
815 *Biochem* 38:1125–1132.
- 816 65. Shahsavari E, Adetutu E, Anderson P, Ball A (2013) Tolerance of Selected Plant
817 Species to Petrogenic Hydrocarbons and Effect of Plant Rhizosphere on the Microbial
818 Removal of Hydrocarbons in Contaminated Soil. *Water Air Soil Pollut* 224:1–14. doi:
819 10.1007/s11270-013-1495-3
- 820 66. Li X, Rui J, Mao Y, et al. (2014) Dynamics of the bacterial community structure in the
821 rhizosphere of a maize cultivar. *Soil Biol Biochem* 68:392–401. doi:
822 10.1016/j.soilbio.2013.10.017
- 823 67. Breedveld GD, Karlsen DA (2000) Estimating the availability of polycyclic aromatic
824 hydrocarbons for bioremediation of creosote contaminated soils. *Appl Microbiol*
825 *Biotechnol* 54:255–261. doi: 10.1007/s002530000362
- 826 68. Smalla K, Wieland G, Buchner A, et al. (2001) Bulk and Rhizosphere Soil Bacterial
827 Communities Studied by Denaturing Gradient Gel Electrophoresis: Plant-Dependent
828 Enrichment and Seasonal Shifts Revealed. *Appl Environ Microbiol* 67:4742–4751.
- 829 69. McCaig AE, Glover LA, Prosser JI (1999) Molecular Analysis of Bacterial

Community Structure and Diversity in Unimproved and Improved Upland Grass Pastures. *Appl Environ Microbiol* 65:1721–1730.

70. Koops H-P, Purkhold U, Pommerening-Röser A, et al. (2006) The Lithoautotrophic Ammonia-Oxidizing Bacteria. In: Dworkin M, Falkow S, Rosenberg E, et al. (eds) *The Prokaryotes*. Springer New York, pp 778–811

71. Stienstra AW, Klein Gunnewiek P, Laanbroek HJ (1994) Repression of nitrification in soils under a climax grassland vegetation. *FEMS Microbiol Ecol* 14:45–52. doi: 10.1111/j.1574-6941.1994.tb00089.x

72. Rahman MH, Saiga S (2007) Endophyte Effects on Nutrient Acquisition in Tall Fescue Grown in Andisols. *J Plant Nutr* 30:2141–2158. doi: 10.1080/01904160701700632

73. Luo J, Zhuang W-Y (2010) New species and new Chinese records of Bionectriaceae (Hypocreales, Ascomycota). *Mycol Prog* 9:17–25.

74. Pegg GF, Parry DW (1983) Infection of lucerne (*Medicago sativa*) by *Fusarium* species. *Ann Appl Biol* 103:45–55. doi: 10.1111/j.1744-7348.1983.tb02739.x

Figure1
Click here to download Figure: Figure1.tiff




Figure2
Click here to download Figure: Figure2.tiff




Table1

Condition	MsmT0	BST0	MsmT4	BST4	MsmT12	BST12
Sampling date	September 2005		September 2007		September 2011	
Humidity (%)	48.85 ± 7.83 ^b	44.74 ± 10.12 ^b	47.40 ± 5.81 ^b	58.65 ± 6.12 ^{ab}	64.37 ± 5.18 ^a	67.91 ± 3.89 ^a
Biomass of alfalfa (g)	0	0	1693.3 ± 156.3	0	109.0 ± 106.1	0
Mycorrhizal colonization						
Frequency of colonized roots (%)	nd	nd	0	nd	24.6 ± 10.1	nd
Frequency of arbuscules (%)	nd	nd	0	nd	15.6 ± 20.3	nd
N (g/Kg)	2.76 ± 0.14 ^a	2.73 ± 0.17 ^a	2.85 ± 0.07 ^a	2.51 ± 0.09 ^{bc}	2.66 ± 0.04 ^{ab}	2.40 ± 0.02 ^c
C/N	21.5 ± 0.7 ^b	22.2 ± 0.2 ^b	22.5 ± 0.8 ^b	23.5 ± 1.1 ^{ab}	25.4 ± 1.1 ^a	25.5 ± 1.7 ^a
TOC (g/Kg)	59.23 ± 1.45	60.67 ± 4.02	64.13 ± 1.88	59.07 ± 4.68	67.47 ± 2.68	61.27 ± 3.58
Organic matter (g/Kg)	102.33 ± 2.31	105.00 ± 6.93	111.00 ± 3.46	102.17 ± 8.28	117.00 ± 4.58	106.00 ± 6.08
pH	6.72 ± 0.16 ^c	6.96 ± 0.14 ^c	7.15 ± 0.07 ^{bc}	7.26 ± 0.15 ^b	7.47 ± 0.04 ^a	7.59 ± 0.01 ^a
CaCO ₃ (g/Kg)	14.83 ± 1.35 ^{ab}	18.40 ± 1.85 ^a	10.92 ± 1.29 ^b	13.93 ± 2.46 ^{ab}	10.10 ± 0.53 ^b	14.47 ± 3.86 ^{ab}
Available P ₂ O ₅ (Olsen) (g/Kg)	0.040 ± 0.004 ^b	0.048 ± 0.003 ^{ab}	0.055 ± 0.006 ^a	0.058 ± 0.009 ^a	0.055 ± 0.006 ^a	0.061 ± 0.007 ^a
Extractable metal (mg/Kg)						
Cu	10.28 ± 1.78 ^c	9.85 ± 0.21 ^b	12.03 ± 1.10 ^b	12.70 ± 0.50 ^b	13.03 ± 0.06 ^b	15.47 ± 0.23 ^a
Fe	82.00 ± 3.76 ^b	82.30 ± 4.71 ^b	97.00 ± 8.72 ^a	101.10 ± 4.25 ^a	99.83 ± 0.29 ^a	107.00 ± 3.00 ^a
Mn	435.33 ± 19.40 ^a	359.00 ± 73.75 ^b	126.00 ± 9.54 ^c	72.20 ± 6.39 ^c	64.40 ± 2.01 ^c	73.37 ± 2.97 ^c
Zn	242.67 ± 3.79 ^a	242.67 ± 9.07 ^a	152.67 ± 16.01 ^c	170.67 ± 7.64 ^{bc}	154.33 ± 8.50 ^c	188.67 ± 16.20 ^b
Metal (mg/Kg)						
B	9.2 ± 0.2 ^a	8.5 ± 0.3 ^b	5.4 ± 0.0 ^d	6.0 ± 0.2 ^c	3.8 ± 0.3 ^e	3.6 ± 0.1 ^e
Cr	658.7 ± 51.9 ^b	648.3 ± 23.5 ^b	783.7 ± 40.5 ^b	767.7 ± 84.1 ^b	1012.3 ± 61.9 ^a	1033.3 ± 75.1 ^a
Cu	140.7 ± 14.2 ^{ab}	131.0 ± 5.6 ^b	157.3 ± 4.2 ^a	147.0 ± 2.0 ^{ab}	146.7 ± 3.5 ^{ab}	146.3 ± 6.7 ^{ab}
Zn	1926.7 ± 55.1 ^b	2000.0 ± 187.3 ^b	2306.7 ± 20.8 ^{ab}	2300.0 ± 226.1 ^{ab}	2196.7 ± 40.4 ^{ab}	2416.7 ± 223.7 ^a
Pb	441.0 ± 5.6 ^c	451.0 ± 35.3 ^c	610.3 ± 19.5 ^a	596.0 ± 32.1 ^a	491.7 ± 6.4 ^{bc}	545.3 ± 56.0 ^{ab}
Σ16 PAH (US-EPA) (mg/kg)	1870.6 ± 146.5 ^a	1839.1 ± 262.3 ^a	2002.0 ± 100.5 ^a	1281.8 ± 314.2 ^b	1333.3 ± 57.7 ^b	1133.3 ± 152.8 ^b

Table2

		Numbers of reads		Richness estimator		Diversity estimator	
	Conditions	Raw data	Filtered data	Observed OTU ₉₇	Chao1	Shannon's index (H')	Inverse of Simpson's index (1/D)
16S	MsmT0	28922 ± 549	25784 ± 637	2658 ± 735	8173 ± 3599	5.46 ± 0.25 ^b	31.13 ± 8.04 ^c
	MsmT4	25205 ± 4287	21958 ± 3832	3057 ± 435	8433 ± 2061	6.38 ± 0.19 ^a	180.74 ± 24.73 ^a
	MsmT12	32767 ± 2910	28822 ± 2354	2754 ± 760	7301 ± 3622	6.22 ± 0.23 ^a	128.61 ± 11.77 ^b
	BST0	31608 ± 5355	27725 ± 4406	1963 ± 528	4527 ± 1931	5.45 ± 0.24 ^b	38.92 ± 10.08 ^c
	BST4	30640 ± 1828	26994 ± 1301	2402 ± 803	5610 ± 3373	6.06 ± 0.26 ^a	123.89 ± 6.03 ^b
	BST12	32736 ± 4867	29439 ± 4449	2762 ± 222	7136 ± 1228	6.10 ± 0.11 ^a	104.52 ± 14.24 ^b
	Total of sequences	545631	482163				
ITS	MsmT0	23887 ± 3613	19837 ± 2666	284 ± 26 ^c	452 ± 89 ^b	2.90 ± 0.18 ^c	7.20 ± 1.23 ^b
	MsmT4	26700 ± 13721	21722 ± 10983	592 ± 69 ^a	995 ± 157 ^a	4.20 ± 0.25 ^a	35.91 ± 19.50 ^a
	MsmT12	27495 ± 3256	22867 ± 2716	563 ± 80 ^a	977 ± 115 ^a	4.01 ± 0.27 ^a	16.39 ± 3.96 ^{ab}
	BST0	21020 ± 1517	17336 ± 868	294 ± 27 ^c	474 ± 117 ^b	2.98 ± 0.05 ^c	7.79 ± 0.28 ^b
	BST4	20951 ± 6250	16889 ± 4907	441 ± 11 ^b	728 ± 9 ^a	3.22 ± 0.10 ^c	7.21 ± 0.32 ^b
	BST12	26111 ± 3932	21508 ± 3276	519 ± 28 ^{ab}	847 ± 138 ^a	3.57 ± 0.10 ^b	18.39 ± 14.15 ^{ab}
	Total of sequences	438492	360477				

FigureS1

[Click here to download Supplementary Material: FigureS1.tiff](#)

TableS1

[Click here to download Supplementary Material: tableS1_290715.doc](#)