



HAL
open science

Small Solar System Bodies as granular media

D. Hestroffer, P. Sanchez, L. Staron, A. Campo Bagatin, S. Eggl, W. Losert,
N. Murdoch, E. Opsomer, F. Radjai, D. C. Richardson, et al.

► **To cite this version:**

D. Hestroffer, P. Sanchez, L. Staron, A. Campo Bagatin, S. Eggl, et al.. Small Solar System Bodies as granular media. *Astronomy and Astrophysics Review*, 2019, 27 (1), 10.1007/s00159-019-0117-5 . hal-02342853

HAL Id: hal-02342853

<https://hal.science/hal-02342853>

Submitted on 4 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Small solar system bodies as granular media

**D. Hestroffer · P. Sánchez · L. Staron ·
A. Campo Bagatin · S. Eggl · W. Losert ·
N. Murdoch · E. Opsomer · F. Radjai ·
D. C. Richardson · M. Salazar · D. J.
Scheeres · S. Schwartz · N. Taberlet · H.
Yano**

Received: date / Accepted: date

Made possible by the International Space Science Institute (ISSI, Bern) support to the international team “Asteroids & Self Gravitating Bodies as Granular Systems”

D. Hestroffer
IMCCE, Paris Observatory, universit  PSL, CNRS, Sorbonne Universit , Univ. Lille, F-75014
Paris, France
Tel.: +33 1 4051 2260
E-mail: hestroffer@imcce.fr

P. S nchez
CCAR, University of Colorado Boulder, Boulder, CO 80309, USA

L. Staron
Inst. Jean Le Rond d’Alembert, Sorbonne Universit , CNRS, F-75005 Paris, France

A. Campo Bagatin
Departamento de F sica, Ingenier a de Sistemas y Teor a de la Se al, Universidad de Alicante,
E-03080 Alicante, Spain
Instituto Universitario de F sica Aplicada a las Ciencias y las Tecnolog as, Universidad de
Alicante, E-03080 Alicante, Spain

S. Eggl
IMCCE, Paris Observatory, PSL Research University, CNRS, Sorbonne Univ., UPMC Paris
06, Univ. Lille, F-75014 Paris, France
JPL/NASA, CalTech, Pasadena, CA, USA

W. Losert
Institute for Physical Science and Technology, and Department of Physics, University of Mary-
land, College Park, MD 20742, USA

N. Murdoch
DEOS/SSPA, Institut Sup rieur de l’A ronautique et de l’Espace (ISAE-SUPAERO), Univer-
sit  de Toulouse, 31400 Toulouse, France

E. Opsomer
GRASP, Research Unit CESAM, University of Li ge, B-4000 Li ge, Belgium

F. Radjai
LMGC, Universit  de Montpellier, CNRS, F-34090 Montpellier, France

D. C. Richardson
Department of Astronomy, University of Maryland, College Park, MD 20742, USA

M. Salazar
LICB, Univ. de Bourgogne, CNRS, F-21078 Dijon, France

Abstract Asteroids and other Small Solar System Bodies (SSSBs) are of high general and scientific interest in many aspects. The origin, formation, and evolution of our Solar System (and other planetary systems) can be better understood by analysing the constitution and physical properties of small bodies in the Solar System. Currently, two space missions (Hayabusa 2, OSIRIS-REx) are on their way to their respective targets and perform samples return; other missions have been selected or proposed to space agencies. The threat posed to our planet by Near-Earth Objects (NEOs) is also considered at the international level, and this has prompted dedicated research on possible mitigation techniques. Thus, the DART space mission will test the kinetic impact technique. Even ideas for industrial exploitation have risen during the last years, as future space mission projects will undoubtedly target some asteroids or other SSSBs. Lastly, the origin of water and life on Earth appears to be connected to asteroids.

In all these cases and research topics, specific knowledge of the structure and mechanical behaviour of the surface as well as the bulk of those celestial bodies is crucial. In contrast to large telluric planets and dwarf planets, a large proportion of such small bodies is believed to consist of gravitational aggregates ('rubble piles') with no—or low—internal cohesion, with varying macro-porosity and surface properties (from smooth regolith covered terrain, to very rough collection of boulders), and varying topography (cratering, depressions, ridges). Bodies with such structure can sustain some plastic deformation without being disrupted in contrast to the classical visco-elastic models that are generally valid for planets, dwarf planets, and large satellites. These SSSBs are hence better approached through granular mechanics theories, which have been a subject of intense theoretical, experimental, and numerical research over the last four decades.

This being the case, it has been necessary to use the theoretical, numerical and experimental tools developed within Soil Mechanics, Granular Dynamics, Celestial Mechanics, Chemistry, Condensed Matter Physics, Planetary and Computer Sciences, to name the main ones, in order to understand the data collected and analysed by observational astronomy (visible, thermal, and radio), and different space missions.

In this paper, we present a review of the multi-disciplinary research carried out by these different scientific communities in an effort to study SSSBs.

Keywords Small bodies of the Solar System SSSB, minor planets, asteroids: general · gravitational aggregates · granular media · methods: numerical, laboratory, observational · planetary formation

D. J. Scheeres

Aerospace Engineering Sciences, University of Colorado Boulder, Boulder, CO 80309, USA

S. Schwartz

Labratoire Lagrange, Univ. Nice, CNRS, Observatoire de la Cte d'Azur, F-06304 Nice, France
Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ, USA

N. Taberlet

Université de Lyon, Laboratoire de Physique, École Normale Supérieure de Lyon, CNRS, F-69364 Lyon, France

H. Yano

JAXA/Institute of Space and Astronautical Science, 3-1-1 Yoshinodai, Chuo-ku, Sagami-hara, Kanagawa 252-5210, Japan

1 Introduction

Small Solar System Bodies (SSSBs) correspond to a class of Solar System objects, all orbiting around the Sun, that are smaller than dwarf planets and are likely not internally differentiated. Exception made of **active comets**—having hence bright and large comas—their appearance in the 19th century **telescopes** was stellar-like (not resolved), which is at the origin of the ‘asteroid’ designation. This naming and classification has evolved with time, between minor planet and asteroid designation, to end up in 2006, with the definition of SSSB class, following an IAU resolution (van der Hucht 2008). So, Small Solar System Bodies currently include most of the asteroids, Trans-Neptunian Objects (TNOs) and comets. SSSBs are hence found in different regions of the Solar System: Near Earth Objects (NEOs) orbiting in the vicinity of the Earth (with perihelion $q \leq 1.3$ AU), Main Belt Asteroids (MBAs) orbiting between Mars and Jupiter, Trojan asteroids co-orbiting Jupiter or other major planets, Centaurs orbiting between Jupiter and Neptune, and in the belt of trans-Neptunian objects (TNOs) beyond the orbit of Neptune (with semi-major axis $a \geq 30$ AU). Because they share some properties with these small bodies, we shall also consider in the rest of the paper some small planetary satellites (orbiting around a planet, and hence not strictly speaking SSSBs).

These small icy or rocky bodies are of particular interest for fundamental scientific research to understand the formation and evolution of planetary systems in general, and of our Solar System in particular. SSSBs, being pristine **in general** or having experienced little geological evolution are valuable tracers of the early time of the Solar System. Asteroids, comets and planetesimals are also believed to be the source of water on Earth through past collisions (Morbidelli et al 2000; O’Brien et al 2014; Bancelin et al 2017, and references therein) and so they have been proposed to be the fundamental bricks that formed an environment apt to support life on Earth. Following the success of the Hayabusa space mission that visited the asteroid (25143) Itokawa, other sample-**return** missions (Hayabusa 2, OSIRIS-REx) are currently on their way to their respective targets, and will greatly enhance our science of asteroids in these aspects. Other mission are targeting small planetary satellites for sample return (MMX), or have been proposed to space agencies (MarcoPolo-R, Phobos-Grunt-2, ZhengHe). Space missions are of great value to deepen our understanding of SSSBs, as will be seen in Sect. 2; so missions have been recently accepted to target a metallic asteroid and several Trojans of Jupiter (**Psyche and Lucy, respectively**), while other where—or are—proposed (Castalia, DePhine, OKEANOS).

On a more societal aspect, there are a number of asteroids (potentially hazardous asteroids - PHAs) with a non-zero probability of impacting the Earth. These impacts can yield extinction level events at planetary scales on astronomical/geological time frames of several tens of million of years, or more local natural disasters on a hundred-years time-frame (Board et al 2010; Pelton and Allahdadi 2015, and references therein). Understanding our vulnerability to such PHAs (for life and goods) necessitates a detailed understanding of the results of an impact on the ground (land or water), the physics of the entry in the atmosphere, and the structure and properties of the asteroid itself. In addition to the DART space mission, several missions are proposed to assess the mitigation techniques and capabilities, including their dependence to the asteroid’s properties (Don Quijote, AIDA with DART and its complementary part HERA, NEOTwIST).

Last, our interest in asteroids has increased during the last decade for economic purpose, in exploration and exploitation. Near Earth Asteroids were the intended target for the next human exploration, or resource collection by a redirect mission (the Asteroid Redirect Mission, ARM, currently dismissed) before considering a human mission to Mars. Extraction of possible extra-terrestrial resources (In-Space or In-Situ Resource Utilization - ISRU) in space is raising interest from industries for business purposes, mostly for use in space and for sustaining habited missions in space; pushing governments to adopt laws in agreement with the Outer Space Treaty¹ of the UNO-SO. This also requires a thorough understanding of the targeted object and moreover, where and how to mine potential resources (Galache et al 2017). As a consequence of these interests, several space missions have—during the space exploration era—targeted, or are proposed to target, asteroids, comets, and satellites; with a prevalence to Near Earth asteroids - because of their proximity to Earth (and therefore reduced mission duration and costs). Main belt asteroids are mostly seen as an opportunity **fly-by** during a cruise to a planet.

Presently, more than 700,000 asteroids have been identified and discovery is still progressing with current surveys (for instance, the discovery rate is of more than 1800/year for only NEOs (Chamberlin 2018)), and more is to come as the limiting magnitude of these surveys is pushed to track fainter objects. The recent WISE/NEOWISE space mission (Masiero et al 2014) has revised downward the estimation of the total population of asteroids by size range; however, a large fraction of objects smaller than approximately 100 meters still remains to be discovered. The whole mass of TNOs is estimated to be at least one order of magnitude larger than that of asteroids, but they are very faint and few details on their structure are known – except in the case of the dwarf planet Pluto. Therefore, most of the current discussion will be **focused** on asteroids.

There is no strict classification of objects by size, in the size-range of several meters to hundreds of kilometres. Even if small in size, compared to planets or giant satellites, asteroids can no longer be considered as tiny point-like masses and have gained interest as small worlds on their own right. Moreover, asteroids are driven by a great variety of both dynamical and physical mechanisms and show a large variety and diversity of composition, size, shapes, morphology, and surface properties according to what is observed. Indeed, asteroid orbital dynamics is mainly driven by their interaction with the Sun through gravity and other gravitational or non-gravitational perturbations. In this respect, they are considered as test particles for some General Relativity tests (Will 2014). However, given the accuracy reached in many studies on their dynamics (either short- or long-term) it appears that the effects connected to their physical properties can no longer be neglected. The size and shape have a direct effect on astrometric measurements, or photocentre offset to the centre of mass, and subsequently on their orbit determination or improvement (Hestroffer 1998). Mutual perturbations between asteroids or by telluric planets are affecting their orbit propagation (e.g. Hilton 2002; Mouret et al 2007). Even without considering such secondary effects, the mass and density are fundamental parameters that characterise an asteroid. After the discovery of binary asteroids in the early 1990s, such as (243) Ida-Dactyl (Chapman et al 1995) from space, and (45) Eugenia from ground-based telescope

¹ <http://www.unoosa.org/oosa/en/ourwork/spacelaw/treaties/introouterspacetreaty.html>

(Merline et al 1999), and some space probe visits to asteroids, as the recent mission to (25143) Itokawa (Fujiwara et al 2006), mass and macroscopic porosities could be estimated, completely renewing our interpretation of these objects from a geological point of view.

Many asteroids and small bodies are supposed to be ‘rubble-piles’ (a terminology introduced by C.R. Chapman (1977) for asteroids shattered as the result of impacts process and gravitationally reaccumulated into a single body) and not monolithic bodies governed only by material strength. This term of rubble-pile is however confusing due to conflict with the different structural interpretation commonly used in Geology. For this reason, other terms have been proposed: gravitational aggregates, self-gravitating aggregates and, more recently, granular asteroids. Such low bulk density asteroids are thus gravitational aggregates held together mostly by mutual gravity (Richardson et al 2002; Bagatin et al 2001), with surfaces covered by pounds, craters, grooves, boulders, large topographic slopes, etc. For instance, the 17 km size asteroid (433) Eros shows a large crater and a smooth and deep surface of regolith, while (25143) Itokawa, approximately 500 m large, presents a shallow layer of gravel-like grains. All these observations have revealed a number of surface features that are yet to be fully understood (Murdoch et al 2015). In contrast to large gas giants or planetary bodies that are generally in hydrostatic equilibrium, these smaller bodies are able to sustain comparatively large shear stresses, yields, and plastic deformation. Their surfaces and interiors as well can be modelled as granular media formed by particles ranging from dust to boulders. So, during the last two decades, asteroids have often been modelled as granular systems, **with many works from different groups and authors; a summary can be found in a few review papers** (Richardson et al 2002; Murdoch et al 2015; Scheeres et al 2015; Hestroffer et al 2017; Campo Bagatin et al 2018b, and references therein). However, a comprehensive theory of granular media is elusive, depending on a large set of parameters and not easily scalable to planetary objects.

In the following sections we provide a review of the concept of granular systems, its application to small bodies in planetary science, and connected open questions. We start with Sect. 2 that presents the current knowledge on the bodies under study, what has been learned from observations and the models that have been developed. Section 3 gives a general introduction on the theory of grains and granular systems, and what should be relevant for planetary science of gravitational aggregate objects. Section 4 will present the general modelling techniques to study granular media as either a continuum (FEM) or as a discrete (DEM) system. In Sect. 5 we explore experiments on granular systems either on the ground (that is under Earth’s gravity), or in a micro-gravity environment. After presenting benchmark cases in Sect. 6, and applications to the study of cohesion, spin-up and YORP effect, segregation, and post-impact re-accumulation, we give a general perspective in Sect. 7.

2 Asteroids and small bodies

2.1 Observations and knowledge

Asteroids were long suspected to be monolithic, possibly differentiated bodies, dry and dense, in particular in the inner regions of the Solar System, with bare rock surfaces. However, during the last thirty years, our vision of asteroids' surfaces and interiors has considerably evolved. This is a result of the particular advancement in observational techniques (from the ground and from space) that has allowed us to remotely characterize a number of SSSBs. Over time, this has resulted in a significant progress on our knowledge and understanding of asteroids, satellites and comets, which has increased even further with the analysis of the data collected through in-situ space exploration. Since the first review book on asteroids (Chapman 1977) we have not only increased, but deepened our knowledge of their physical characteristics (sizes, shapes and masses, composition, thermal inertia, albedos, etc., and their statistics). Astronomers and planetary scientists have also discovered new features, objects or phenomena (Yarkovsky, YORP, planetary migration, binaries and multiple systems, tumblers, fast rotators, main-belt comets MBC and active asteroids, to mention a few) which were described in the subsequent books Asteroids II, III, and IV (Gehrels and Matthews 1979; Binzel et al 1989; Bottke et al 2002; Michel et al 2015). The same holds true for comets (Wilkening and Matthews 1982; Festou et al 2004), though to a lesser degree.

2.1.1 Asteroids and collisions

The population of asteroids originates from steady-state collisional dynamics (Dohnanyi 1969), so that the number density distribution, for a mass m , follows a power law, with:

$$N(m) \propto m^{-11/6} \quad (1)$$

with the smaller objects being more numerous. C. R. Chapman proposed that—as a result of the inter-collisional evolution—larger asteroids would be fractured or constitute a recollection of fragments that did not escape after the catastrophic collision; he introduced the terminology of ‘rubble pile’ (Chapman 1977) (see Sect. 1). Such bodies would hence be self-gravitating collections of blocks—or gravitational aggregates—with more or less fractured rocks, with no or little internal cohesion, sometimes highly fractured and porous bodies, with more or less coherent arrangements of the constituting blocks. This vision has been confirmed with the estimation of porosity (**Britt and Consolmagno 2001**), see Eq. (2), for some of the asteroids for which the bulk density could be measured. Space missions to small bodies (see Table 1), either as flybys or rendez-vous, have considerably improved our knowledge of these bodies, by bringing detailed measurements and observations of their interiors and surfaces. Flybys are valuable even if they are only brief encounters, unfortunately, however, they are not systematically incorporated in mission trajectories.

Asteroids were long suspected to have bare rock surfaces, but the NASA NEAR mission showed that the predictions for (433) Eros were incorrect. Indeed, this in-situ mission—followed by other space missions—revealed a substantial layer of unconsolidated rocky material and dust (regolith) covering the surface of (951)

Gaspra, (243) Ida, (433) Eros, (21) Lutetia, and (25143) Itokawa (Murdoch et al 2015). Additionally, this regolith seems to be active: there is evidence of motion in the form of landslides, particle migration, particle size sorting and regolith production. So, the geophysical characteristics of the surfaces of small planetary bodies appear to be diverse and complex (Murdoch et al 2015). Also, observations made by the Hubble Space Telescope of the disruption events of active asteroids P2013/R3 (Jewitt et al 2014) and P2013/P5 (Jewitt et al 2013) imply that not only asteroid surfaces are covered by regolith, but that their internal structure is granular and not monolithic.

The usual assumption—that prevailed for more than a century—that asteroids are single monoliths, is also incorrect. While long speculated in the end of last century (Weidenschilling et al 1989), and albeit negative result from surveys around the largest minor planets (Gehrels et al 1987), evidence has been obtained only in the last decades that “binary asteroids *do* exist” (Merline et al 2002). It is noteworthy that there are no satellites **orbiting around the largest asteroids of the main belt (the dwarf planets)**, but satellites and moons are found in different asteroid dynamical classes (Near Earth regions, main belt, Trojans, Centaurs and trans-Neptunian, ...). Furthermore, some have rings (Braga-Ribas et al 2014; Ortiz et al 2017). The number of detected double and multiple systems, and also asteroid pairs, has rapidly increased, showing also a higher proportion of systems in the NEO and TNO populations; though – due to observational bias – they are not all sampled identically. The formation scenario of these binary or multiple systems, containing a variety of mass fraction, separation, angular momentum, etc. is not fully understood. Different **mechanisms** can be required within various populations of asteroids. For examples, formation mechanism can include fission and mass shedding, post catastrophic collision, re-accumulation, capture, ... (Noll et al 2008; Margot et al 2015; Walsh and Jacobson 2015; Tardivel et al 2018).

2.1.2 Spin rate, size, and shape

There are several fundamental parameters **to characterise asteroids**, both physical and dynamical. One such fundamental parameter is the spin-rate or period of rotation, that can be obtained relatively easily together with the object’s brightness. The spin period is obtained with some confidence from the light-curve (for those objects that are not spheroidal in shape); while the size is generally estimated from the brightness, sometimes by assuming the object’s albedo (when no direct or radiometric measure is available). This provides information for a spin-rate versus size plot, which gives a good view of some of the general properties of the objects we are studying (Fig. 1). In particular, two features are apparent: 1) a spin barrier at about $P \approx 2.4$ hr for the larger objects, suggesting that internal strength is not high enough to balance the centrifugal acceleration, and 2) the existence of fast and superfast rotators (Pravec and Harris 2000; Pravec et al 2002; Scheeres et al 2015). The spin barrier is supposed to **be good evidence** that most large object in the range 1 – 100 km, in particular close to spin limit, are granular in nature and not monolithic (Bagatin et al 2001). Fast rotators on the other hand, must be either monolithic rocks with natural strength, or have additional cohesion to bound a granular structure (see Sect. 4); such differences are however not recognizable from simply the size and spin-rate parameters.




Fig. 1 Spin period for more than 5500 asteroids as function of their estimated diameter. A ‘spin barrier,’ above which no large objects will rotate, is apparent; consistent with a gravitational aggregate structure of loose conglomerations held together by mutual gravitation. Conversely, small objects can spin very fast, with full rotation within minutes; a hint towards a more monolithic structure. Near Earth objects are predominant in the population of objects smaller than approximately 1 km. (Source: the LCBD light curve database (Warner et al 2009)).

When these bodies reach their critical spin rate, which is dictated by their size, density and internal strength and strength distribution, they will undergo deformation, surface mass shedding, fission or catastrophic disruption events (Hirabayashi et al 2015; Sánchez and Scheeres 2016, 2018). Particles that are ejected near the equatorial plane can supposedly re-accumulate and form moonlet(s) (Walsh et al 2008). A more abrupt fission event could also produce a binary system (Tardivel et al 2018) or could be at the origin of the observed asteroid pairs (Pravec et al 2010; Sánchez and Scheeres 2016; Zhang et al 2018). Only the structurally strongest asteroids will reach the 2.4 h spin barrier. More generally, whatever the spin-rate is, the surface gravity on a body of several 100 m size is in any case weak (approx. $10^{-3} - 10^{-6} g$), and escape velocities are in the cm/s range.

Our modelling of asteroid shapes has dramatically evolved during the last decades. The observed periodic variation in an asteroid lightcurve over a rotation period can be explained by either a variation of the projected surface for a non-spherical or non-spheroidal shape, or by a variation in albedo of the observed surface (Russell 1906). It eventually appears that such variations are mainly driven by shape effects (Kaasalainen and Torppa 2001; Kaasalainen et al 2001; Li et al 2015). Starting from simple tri-axial ellipsoid models spinning about their shortest axis (as an outcome of collisional and dynamical processes), or more sophisticated

cellinoids (Lu et al 2016), we now have tools to derive convex and non-convex topographic shapes (Kaasalainen et al 2002). Stellar occultations and high angular resolution observations valuably complete that information and modelling (Durech et al 2015), as well as radiometric and polarimetric observations for size and albedo (Masiero et al 2017; Stansberry et al 2008; Harris and Lagerros 2002; Delbo et al 2015; Belskaya et al 2015, and references therein). Radar observations—by analysing the echoed wavefront on the asteroid’s surface—also provide information on shape and more detailed physical properties (Ostro et al 2002; Benner et al 2015). **The body’s shape and outer envelope can be related to its internal structure, and also modify its spin barrier level (Harris 1996; Holsapple 2001).**

2.1.3 Mass, bulk-density, and porosity

Binaries and multiple systems are of particular interest since, as a result of collisions, they could correspond to the case proposed by Chapman (1977) and likely be gravitational aggregates. Moreover, binaries are of particular interest here, because by deriving their mutual orbits, it is possible to derive another fundamental parameter: the (total) mass of the asteroidal system.

Measuring the mass is a difficult goal to reach which can be achieved from the careful astrometric observations of binary systems, or otherwise from the analysis of their gravitational perturbation during a **rendez-vous or fly-by** with a space probe, or a close encounter with another (small, target) asteroid. **Such close encounters between asteroids are much more frequent than space mission fly-bys, but** far less precise or accurate, and require high accuracy astrometric measurement together with a global inversion to take into account all the effects (Mouret et al 2007; Baer and Chesley 2017). In any case, estimating the mass of any small planetary object studied is of high importance as it allows an indication of its density, porosity, internal structure or internal mass distribution, and its global behaviour to stresses or impacts. Given the bulk density ρ of a small body (generally measured from the knowledge of mass and volume, with some uncertainty), and ρ_g the mean density of the material that constitutes the constituting grains (generally unknown, and estimated from the taxonomic class), the macro-porosity² p is defined by:

$$p = 1 - \rho_g/\rho \quad (2)$$

It is a scale invariant parameter that also measures the interstitial voids in the global structure of the body. The bulk density uncertainty is dominated by the error on the volume which can be large. Besides, the porosity estimation depends on our knowledge of the interior material or internal constituents, and the densities of analogue meteorites (i.e., those representing a good match to the asteroid taxonomic class). Progress in the classification, and determination of such meteorite densities and micro-porosities has been achieved in laboratories. However, the meteorites collected on Earth may not always be the best analogue for representing an asteroid as a whole, as it could also be formed of other materials, containing volatiles or fragile material, or having regions with different porosities. This means

² Micro-porosity on the other hand is in the matrix of the grains or meteorites. Micro-porosity is a porosity that will survive entry in the atmosphere.




Fig. 2 Possible internal structure arrangement of gravitational aggregates, based on their macro-porosity (adapted from Britt et al (2002); Carry (2012)). Note that the uncertainty on the porosity is generally large ($\pm 5\%$ and more).

that our knowledge on asteroids' porosity remains limited to a few objects, and with substantial uncertainty (see Fig. 2).

Porosity and material strength play crucial roles in how asteroids react to impacts of smaller debris or "projectiles" as they determine how shock-waves and damage zones caused by the disruption event propagate through the target (e.g. Asphaug et al 1998; Jutzi et al 2010; Syal et al 2013). Several studies have found, for instance, that strength dominated asteroids below a few hundred meters in size are more difficult to disrupt if they are porous (Jutzi and Michel 2014, and references therein). For larger asteroids self-gravity starts to dominate, and the role of porosity becomes less clear cut. Cases where a collision leads to an asteroid's disruption has been discussed extensively in literature (see e.g. Henych and Holsapple 2018). Comparing the imparted kinetic energy per unit mass Q to the specific energy needed to disrupt a target Q^* can provide useful first insights. Let (Henych and Holsapple 2018):

$$Q = \frac{1}{2} \frac{m u^2}{M},$$

$$Q^* = (44R^{-0.6\mu} + 68R^{3\mu}) [\cos(\phi) U]^{2-3\mu}, \quad (3)$$

where m is the mass of the projectile, M the mass of the asteroid, $R[km]$ the radius of the asteroid in kilometers, $u[m/s]$ the relative impact velocity in m/s , and U the same quantity in km/s . Furthermore, ϕ denotes the impact angle and μ is the so-called point-source scaling-law exponent that is around $\mu = 0.4$ for porous materials and $\mu = 0.55$ for non-porous rock-like material. Both, Q and Q^* are given in units of J/kg . The ratio between Q/Q^* determines whether or not enough energy has been deposited to disrupt the target. A value of $Q/Q^* \approx 1$ characterizes an impact that dismantles about 50% of the body. A value much




Fig. 3 Disruption limits for collisions as a function of impactor and target size. The relative velocity at impact is 5 km/s, typical for main belt collisions. The target asteroids are assumed to be non-porous and spherical ($\mu = 0.55$) with a density of 3500 kg/m³.

greater than unity indicates pulverization. On the other hand, $Q/Q^* \ll 1$ describes a simple cratering event. Fig. 3 shows that typical main belt collision speeds suffice for relatively small "projectiles" to affect the structural integrity of kilometer-sized asteroids. Catastrophic disruption is more likely to occur in collisions between main belt asteroids of similar size.

The result of a post-catastrophic re-accumulation should be a loosely bound gravitational aggregate (e.g. **Tanga et al 2009a; Michel and Richardson 2013**). Surprisingly, low bulk densities of the order of 1200 – 1500 kg/m³ were observed on some asteroids, either single or binaries (e.g. Mathilde, Eugenia, ...), while chondritic or silicate material densities are of the order of 2500 – 3500 kg/m³. This is supposed to be evidence of the presence of large voids in the interior of the body, that can be connected to macro porosity of the order of 40 – 50%. How force chains and friction are acting, or how the blocks are arranged inside of the body, is not clear. The mechanism that prevents the fine regolith grains at the smooth upper surface from filling the voids in the case of porous bodies could be linked to friction (Britt and Consolmagno 2001), but needs to be better understood with regard to seismic shaking and segregation mechanism. How asteroids react to collisions isn't clear neither; collisions could either increase the porosity, voids, and fragments, or otherwise reduce the porosity through compaction. The asteroid (253) Mathilde is an example of an object with very high macro-porosity with smooth surface and large craters indicating that it has survived major impacts. However, the actual internal structure, or the size distribution of its constituent particles remains unknown.

From the **knowledge of asteroids mass and porosity**, Britt et al (2002) identified three classes of objects that could reflect different internal structures: coherent, fractured, and loosely consolidated (see Fig. 2). Further, Richardson




Fig. 4 Relative tensile strength (RTS) and porosity parameter space (adapted from Richardson et al (2002)).

et al (2002) schematically characterised gravitational aggregates by relative tensile strength (RTS), in addition to macro-porosity (see Fig. 4). These are driving parameters that qualitatively identify different classes of internal structures and global mechanical behaviour. So, there could be different regimes between strength dominated and gravity dominated bodies (see **Holsapple et al 2002**), with different reactions to mechanical tensile or compressive stresses and impacts. Thus, bodies in the range of concern in this paper are in the transition regime between purely strength dominated (like stones and meteorites) and gravity dominated coherent bodies (like major planets), and can react differently to impacts and cratering (Asphaug et al 2002; Britt et al 2002). For instance, porous and structurally weak bodies may, through stress dissipation, be highly resistant to catastrophic disruption. Additionally, it modifies the cumulative size-distribution slope from Eq. (1) (Bottke et al 2005, and references therein). Lastly, all the mechanisms present in granular media can govern gravitational aggregates' shapes, strength and failure limits (Scheeres et al 2015) (impacts, landslides, fission, tides, ...).

Table 1 List of flown space missions to small bodies ((Hestroffer et al 2017, updated)). Missions to the dwarf planets Ceres, Vesta, and Pluto, or to large planetary satellites, have not been included here; while two launched and ongoing—not yet achieved—sample-return missions have been included. These are mainly NASA, JAXA, and ESA space agencies interplanetary missions, with contributions from national agencies, CNSA, and former USSR for the ‘Halley armada’ and Phobos2. Legend for mission type: F=fly-by ; O=orbit or/and hovering ; SR=sample return ; L=landing ; I=impact.

Target	Mission	Type	Date
Asteroids			
(101955) Bennu	<i>OSIRIS-REx</i>	O+SR	<i>Jul. 2020</i>
(162173) Ryugu	<i>Hayabusa2</i>	O+SR	<i>Jul. 2018</i>
(4179) Toutatis	Chang'E 2	F	Dec. 2012
(21) Lutetia	Rosetta	F	July 2010
(2867) Steins	Rosetta	F	Sep. 2008
(25143) Itokawa	Hayabusa	O+SR	June 2005
(5535) AnneFrank	Stardust	F	Nov. 2002
(433) Eros	NEAR Shoemaker	O+L	Feb. 2000
(2685) Masursky	Cassini/Huygens	F	Jan. 2000
(9969) Braille	Deep Space 1	F	Jan. 1999
(253) Mathilde	NEAR Shoemaker	F	June 1997
(243) Ida+Dactyl	Galileo	F	Aug. 1993
(951) Gaspra	Galileo	F	Oct. 1991
Comets			
67P/Churyumov-Gerasimenko	Rosetta	O+L	Aug. 2014
103P/Hartley 2	EPOXI (Deep Impact ext.)	F	Nov. 2010
9P/Tempel 1	Deep Impact, Stardust	O+I	Jul. 2005, 2011
81P/Wild 2	Stardust	O+SR	Jan. 2004
19P/Borrelly	Deep Space 1	F	Sep. 2001
Grigg-Skjellerup	Giotto	F	July 1992
1P/Halley	The Halley armada: Vega 1&2, Sakigake, Suisei, Giotto, ICE	F	March 1986
21P/Giacobini-Zinner	ICE	F	Sep. 1985
Moons			
M1 Phobos	MEX, ODY, MGS, Viking1 + Phobos2	F+O	2004, 2001, 1997, 1989, 1976
M2 Deimos	MEX, ODY, MGS, Viking2	F	2004, 2001, 1997, 1976
satellites of Saturn	Cassini	O	2000-2017

2.2 Links between asteroids, planetary satellites, and comets

Comets and asteroids show very distinctive physical and dynamical properties. On the dynamical side, first, objects with parabolic or hyperbolic orbits, i.e. eccentricities $e \geq 1$, would be categorized as comets (or interstellar objects (Meech et al 2017)). Next, the Tisserand parameter T_J for a body whose periodic orbit is characterised by a semi-major axis a , eccentricity e , and inclination i :

$$T_J = \frac{a_J}{a} + 2 \sqrt{\frac{a}{a_J} (1 - e^2)} \cos i \quad (4)$$

given here with respect to the massive Jupiter (with orbit $a_J \approx 5.2$ AU), is a well known dynamical parameter that broadly separates objects that are strongly perturbed by Jupiter, or not, as a function of their encounter velocity. This helps to separate the orbital classes between asteroids (typically $T_J > 3$), and comets

(typically $T_J < 2$ for Halley type comets - HTC, and $2 < T_J < 3$ for Jupiter family comets - JFC). On the physical and visible side, when approaching the Sun at approximately less than 3 AU, a comet begins to sublimate its volatiles, giving rise to a large and bright coma and tail, making comets very distinct in appearance to asteroids.

Now, extinct comets, after having lost their volatiles and after a dynamical evolution bringing them in the Near-Earth orbiting region, are difficult to distinguish from asteroids using remote observations: (“a comet in disguise” Kerr 1985; Weissman et al 2002). On the other hand, the old viewpoint that asteroids—as a kind of minor planet—are dry with no volatiles, possibly differentiated with iron core and mantle and hence dense, while all objects beyond the snow line at approximately 5 AU are icy bodies that have retained their volatiles, has changed. Indeed, ‘active asteroids’ (and not only the large dwarf planet Ceres) or ‘main belt comets’ have been observed in the main belt, well below the hypothesized **snow** line (“a comet among the asteroids” Hsieh and Jewitt 2006; Jewitt et al 2015). Moreover, as difficult as it is, ices have been detected on the surface of a few asteroids (Rivkin and Emery 2010; Campins et al 2010), and it has been showed that volatiles can have long lifetimes if buried under a moderate layer of regolith (Schorghofer 2008; Delbo et al 2015, and reference therein). So, volatiles can be present in asteroids as it is the case in comets, albeit in different proportions. Conversely, some Centaurs have shown activity at large distance from the Sun, invalidating the commonly accepted scenario of possible activity (Jewitt 2009). Moreover, in-situ collection of comet grains by the Stardust mission has revealed the presence of silicates, so that comets are not only made of volatiles and interstellar dust but share also some composition with—closer to Sun—asteroids. It has been confirmed by the Rosetta mission, that gives more insight in the grains and dust, composed of compact grains and fluffy aggregates (Levasseur-Regourd et al 2015), rich in carbon and non-hydrated minerals (Bardyn et al 2017). This is supported by our current understanding of the formation and dynamical evolution of the whole Solar System, showing a big mixing of the distant and inner regions (DeMeo et al 2015) with conglomerates that include refractive material and volatiles, hence erasing or fading any initial solar nebula density or composition gradient. Lastly, the derived masses of some asteroids show very low bulk densities, which is assumed to reflect a high porosity (Mathilde density of ≈ 1.3 (Yeomans et al 1997), Eugenia density of ≈ 1.2 (Merline et al 1999)), similarly to comets. The same low densities have been found for Phobos and Deimos (with values of 1.5 and 1.9, respectively (Rosenblatt et al 2016)), for Trojans and TNOs (Patroclus density of ≈ 0.8 (Marchis et al 2006)), and other extremely low density are found in comets and small Saturnian moons (Thomas 2010).

This all has pushed a change of paradigm in the last decades in our understanding of the real differences between the different classes of minor bodies, bringing forth the concept of an asteroid-comet continuum (Bockelée-Morvan et al 2016). Additionally, the irregular satellites of Jupiter, Saturn, Uranus, Neptune, the small moons of Pluto, the moons of Mars Phobos and Deimos, etc., have been hypothesized to be either captured asteroids or TNOs (Peale and Canup 2015, and references therein). Such capture origin remains **sometimes unclear, and impact scenarios are also possible as in the case of Mars**. Nevertheless—whatever their origin—the similarities in size, surface and composition remain. Thus asteroids, comets and small moons show common features. Although they have some

variation and gradients, for instance in the constituent ingredients or even with different formation mechanism, their final global structure and general mechanical behaviour could still be modelled with a similar approach.

2.3 Summary and Open questions

We have a general knowledge of the surface composition of these small bodies, from telescopic observations, and sometimes of their surface roughness through thermal inertia (Delbo et al 2015; Müller et al 2017). But detailed knowledge on the presence of regolith, craters, boulders and ‘chaotic terrain’, or other geological features is much more limited. **And in that case, they have shown** a large diversity that is still difficult to predict, or to unequivocally correlate with observable data such as size, taxonomic type and spin-rate. **We also expect to see segregation phenomena acting at the surface of these bodies. However size or density segregation and sorting in granular media, and moreover, how it really behaves on self-gravitating small granular bodies, is still a matter of research.** Exploration of these small bodies has now entered a new era with landers and sample return devices that are designed to make contact with their surfaces. This was started with the Hayabusa and Rosetta missions, and we are awaiting the event—and success—of both Hayabusa-2 and OSIRIS-REx missions. In all these cases, or in the case of the DART kinetic impactor on Didymos’ secondary, different mechanisms and scenarios have been considered and the actual reaction and behaviour of the surface will be of high interest for our general understanding. Many space missions are motivated for scientific, or exploration reasons, often requiring a contact with the surface, or landing and manoeuvring. Response of the surface during anchoring, drilling, or sampling is of the utmost importance and needs a better understanding of the physics of granular media in low gravitational conditions (Daniels 2013). **The same applies for mitigation in space of a PHA asteroid that would likely impact the Earth. In that case, the threatening asteroids’ material properties can play a fundamental role when changing its trajectory via impulsive deflection techniques, or when attempting its complete disruption (Ahrens and Harris 1992; Sanchez et al 2009; Sugimoto et al 2014; Eggl et al 2015).**

As seen before, detailed knowledge on asteroids’ surface composition and terrain has been obtained for only a small sample of targets, but **still nothing is certain about their interior.** Tomography (Hérique et al 2018), seismic investigation (Murdoch et al 2017b) (either passive or active), in addition to classical planetary geodesy (Yeomans et al 2000; Konopliv et al 2002; Abe et al 2006; Rosenblatt et al 2008; Andert et al 2010; Jacobson et al 2007; Paetzold 2017; McMahon et al 2018), and possibly with CubeSats (Walker et al 2016; Murdoch et al 2016; Hestroffer et al 2017), adapted to low-gravity small bodies and their perturbed environments have been proposed as viable techniques to probe the interior, and internal structure of relatively small asteroids. At present, most of the available evidence on the internal composition of small solar system bodies are indirect: bulk-density, rotation periods, and crater sizes. As said before, it has been observed that several asteroid classes can be defined in the asteroidal population depending on their bulk porosity, reflecting different degrees of porosity (Britt et al 2002). At the moment, the origin of such porosity is not completely understood,

yet we think that collisions could likely have played a role. However, this does alone not explain for instance the differences between asteroids Mathilde and Itokawa which have similar bulk densities, but different surface characteristics.

Given these observational facts, it is evident that we need to better understand the possible evolution of asteroids: how gravitational aggregates evolve under the YORP effect, how this depends on size/mass, on internal cohesion or other contact forces. What is the formation mechanism of binaries and multiple system: continuous mass shedding, fission, or reaccumulation? How strong are these objects to meteoritic bombardment and tides, kinetic impactor, sampling mechanism, etc.? Given the large variety and diversity among these objects we should limit our assumptions, as we often need to span large parameter sets. Most of these require theoretical and numerical modelling, benchmarking, and experiments. This is the focus of the following sections.

3 Granular systems and Granular mechanics

3.1 What are grains?

A grain is a discrete, rigid, macroscopic particle that can interact with other particles through dissipative contacts, and the motion of which can be accurately described by Newtonian dynamics³. This means that the behaviour of the grains is dominated by particle interactions and gravity forces, in comparison to which thermodynamic agitation appears negligible (Jaeger et al 1996a). The behaviour of an assembly of grains without cohesion or adhesive forces is dictated by the interactions at the individual grain scale. In dense granular systems, aside from the gravitational attraction, the interactions are essentially mechanical contact forces, which can be decomposed into normal forces (arising from the elasto-plastic behavior of the material) and tangential forces (due to the frictional properties of the grains).

Although grains found in nature and asteroids can display a variety of shapes, the case study of spherical grains has been at the focus of theoretical and experimental studies in view of shedding light on the mechanical properties of generic granular materials. A good approximation of the repulsive force involved when two rigid spherical grains—assumed to be identical—are in contact, is given by the Hertz Law (Landau and Lifshitz 1986):

$$F_{\text{Hertz}} = E \frac{\sqrt{2R}}{3(1-\nu^2)} \delta^{3/2} \quad (5)$$

where R is the radius of the grains, E is Young's modulus and ν is Poisson's ratio of the material, and $\delta = 2R - r$ (where r is the distance between the centres of the grains) is the overlap that represents the elastic deflection at the contact area (Agnolin and Roux 2007b). The details of the calculation are somewhat accessory but the striking result is the non-linearity of the interaction, since the repulsive force scales as the $3/2$ power of the overlap. This specific power arises from the spherical shape of grains, although the calculation is based on the theory of linear

³ Possibly including relativistic effects, but this is not relevant in this paper on small bodies.

elasticity. However, the nonlinear nature of the grain-grain interactions remains valid for a wide variety of particle shapes and for large contact deformations.

In addition to the elastic component of the normal force, there exist dissipative effects during particle collisions. Energy dissipation can originate from the visco-elastic properties of grains as well as their plasticity, causing irreversible deformations at the contact scale, which result in inelastic collisions and solid friction (Andreotti et al 2013). Friction is a fundamental aspect as it leads to an indeterminacy problem. Indeed, Coulomb's law for solid friction sets only bounds on the value of the tangential forces between two solid bodies. Hence, the internal mechanical state of an assembly of grains is not uniquely determined by the position and velocity of the grains, but may display residual stresses which induce strongly history-dependent mechanical responses (Toiya et al 2004; Agnolin and Roux 2007a).

Due to the rigidity **of the individual grains**, granular packings may sustain intense forces, which propagate into the system through force chains. Force chains are a manifestation of the high heterogeneity of granular systems in general, and explain why the mean stress state in a granular system does not fully describe the actual mechanical behaviour exhibited by the system (Liu et al 1995; Jaeger et al 1996b; Radjai et al 1996; Radjai 2015).

On Earth, other effects include cohesion forces caused by **capillary** bridges due to the humidity, air drag acting on particles or thermal agitation (Radjai and Richefeu 2009a; Andreotti et al 2013). Due to the absence of atmosphere, these effects appear to be negligible in small asteroids. At the same time, the extremely weak gravitational environment will make other forces, particularly van der Waals cohesion and electrostatic forces, appear comparatively stronger and therefore, important in the behaviour of the system. On asteroids, electrostatic force, which are typically of the order of 10^{-6} N, can easily exceed the gravitational forces in a micro-gravity environment. However, an appropriate modelling of such forces still remains challenging (Hartzell and Carter 2017).

3.2 What are granular systems?

A granular system is a collection of grains that interact through binary or multiple contact interactions, and whose mean behaviour results from the collective dynamics of the grains. Due to the complexity and richness of the grain-scale interactions, granular systems display a wide variety of behaviours (thixotropic solid, liquid, gas), depending on the external mechanical excitation or loading (Jaeger et al 1996b; Patrick et al 2005).

In the presence of a large input of energy in the system (for instance, large slope angles, vibrating boundary conditions), grains form dilute assemblies which are reminiscent of molecular gases. Binary collisions occur between grains whose mean free path is larger than their typical diameter. Examples of such gaseous systems include planetary rings in which individual grains interact through rare collisions. In less energetic settings, granular systems flow in dense packings dominated by long-lasting multiple contact interactions, and frictional dissipation. Rock avalanches or landslides as observed on Earth and Mars (Lucchitta 1979; Neuffer and Schultz 2006) mostly fall into this category. In this case, the mean behaviour is

reminiscent of that of a viscous fluid, however with strongly non-Newtonian properties. Finally, at equilibrium or in quasi-static flow, a collection of grains may resist stress without immediate deformation, a behaviour characteristic of a solid (Liu and Nagel 1998; Gao et al 2014). Rubble-pile asteroids fall in this category and may undergo gravitational, tidal or centrifugal forces without rearranging.

It appears therefore vain to propose a general theoretical framework for the description of such a wide range of behaviours. Instead, specific theories have been developed, or adapted, for each specific state. Kinetic theory (based on the classical thermodynamics molecular kinetic theory) has proven to be very successful in modelling and describing the physics of dilute assemblies of grains (Jenkins and Richman 1985; Brilliantov and Pöschel 2010). On the other hand, classical soil-mechanics and structural engineering theories are often applied to describe the mechanical properties of dense granular material under slow deformation (Terzaghi et al 1996; McCarthy and McCarthy 1977). Finally, hydraulic equations (or the more complex Navier-Stokes equation) are applied to describe the liquid state of granular matter, coupled with semi-empirical models for the effective viscosity (MiDi-GDR 2004). These models will be presented in greater details in Sect. 4.

3.3 Phenomenology

When a flow starts, it will do so for slope angles lower than the angle at which avalanching started, thereby implying a modification of its frictional properties, often referred to as the hysteretic behaviour of the granular pile. While flowing, the internal friction will be affected by the flow dynamics and the pressure, and thus the context of the flow itself will strongly affect the properties of the granular mass (MiDi-GDR 2004). The characterisation of the internal friction of model granular media has thus prompted many research activities. A significant amount of progress has been made, but there are still many aspects relevant to asteroids to explore, including cohesive forces and grain size segregation. A model for the granular rheology in the simple case of mono-sized non-cohesive grains will be presented in Sect. 4.

Another important aspect of granular behaviour in the context of small solar system bodies is their ability to segregate grains according to their size. This phenomenon is omnipresent on Earth, in geophysical flows (debris and rocks flows) which classically involve grain sizes covering three to four orders of magnitude, but also in human activities (food industry, civil engineering, powder technologies etc). When a granular bed is submitted to vibrations or is flowing, larger grains and smaller grains tend to separate, with the larger grains classically rising to the surface of the granular system (also the potential level on Earth). Essentially, this phenomenon results from the higher probability that smaller grains have to fill in the gaps opening in the system while flowing or shaking, a mechanism known as the kinetic sieve (Savage and Lun 1988). Size segregation results in patterns that may partly affect the properties of the system. In the case of unconfined flows for instance (like rock flows), larger boulders are segregated at the front and pushed aside by the flow, forming channels or levées which will in turn affect the flow dynamics (Félix and Thomas 2004). In the context of small solar system bodies, understanding the dynamics of segregation may allow for the interpretation of grain size distribution at the surface of asteroids in terms of formation his-

tory, structure, exposure to impacts or other sources of agitation. Although much studied, the general formulation of a lift force that would describe the segregation dynamics is still mostly lacking (Guillard et al 2014). Presenting existing models is beyond the scope of the present paper, however, interesting insights will be found in Kudrolli (2004a).

3.4 What is relevant?

The previous sections have clearly outlined an introduction to the study of granular matter, granular media, the intricacies of some puzzling phenomena observed in nature and shed some light on our understanding of their innerworkings. The following sections on the other hand, will provide the reader with a more in-depth, theoretically strict and sound explanation of the current theoretical, experimental and numerical methods that have been developed and used to study granular systems. However, as insightful as this all is, we need to answer two very fundamental questions, as otherwise, all this knowledge will simply be inapplicable:

1. are small Solar System bodies granular in nature? and if so,
2. how does all we know about granular dynamics relate to them?

The answer to the first question came as a result of the space missions that visited these bodies (see Table 1). As seen in Sect. 2, snapshot images taken of those asteroids showed them to be covered with surface regolith, to be heavily cratered, and in the case of Ida to have a binary asteroid companion which itself had a regolith covering. Since these initial observations, the space-based and ground-based observations of asteroids have grown along with the realisation that these bodies are fundamentally granular in nature and hence that their physical evolution must be described using principles of granular mechanics applied to these extreme environments, which bring us to answer the second question.

Since other planetary bodies apart from the Earth have been out of human reach for most of our history, most of our knowledge of granular systems is restricted to systems on Earth. What this means is that they were all subjected to an almost constant gravitational field with a magnitude of $\approx 9.81 \text{ m.s}^{-2}$, in which cohesive and electrostatic forces were negligible, except for fine powders ($\approx 10^{-6} \text{ m}$). The accumulated knowledge collected by early builders (Fall et al 2014), engineers, scientists (Faraday 1831) and the common farmer had these two premises that were true for any practical purpose, but not so much for asteroids, comets, or even the Moon. Additionally, on Earth grains always interact with fluids (Burtally et al 2002; Kok et al 2012), be this in the form of wind (and atmosphere) or water currents (and humidity) in most cases.

Given the size and shape differences between the Earth and small planetary bodies, the gravity vector on their surfaces is not as strong as on Earth and changes dramatically from point to point. This is compounded by the rotational state of these bodies, and the possible influence of solar wind and solar radiation pressure.

In spite of these environmental differences, phenomena such as size segregation, phase transitions and granular flows have either been hinted (Sierks et al 2011) or observed directly (Jewitt et al 2013) and this is precisely the point: these are only environmental differences. Fundamentally, each individual grain is still solid, grain-grain interactions are still governed by surface, contact forces, these interactions

are still very dissipative, and individual grains still follow the laws of Newtonian physics. Therefore, there is no fundamental difference between granular systems on Earth and those in SSSBs; not even for two solitary grains colliding in the vast emptiness of space. In that sense, a planetary ring (e.g. Wisdom and Tremaine 1988; Salo 2001; Tiscareno and Murray 2018) could be seen as a self-gravitating granular gas, whereas SSSBs could be seen as self-gravitating granular systems in a condensed, solid phase which will be the focus of the rest of this paper. That being the case, the theoretical, numerical and experimental tools and methods that have been already developed should still be adequate to study them as long as we allow for the implementation of a correct gravitational field, and cohesive and electrostatic forces at the bare minimum. As Scheeres et al (2010) point out, cohesive and electrostatic forces become important due to the greatly reduced magnitude of the gravitational field in comparison to Earth's gravity.

This is the approach that many scientists in the Planetary Sciences and Granular Dynamics community have taken. Numerical tools need to implement particle-particle cohesive and electrostatic forces as well as self-gravity in the case of discrete element methods (DEM) (Richardson et al 2000; Stadel 2001; Sánchez and Scheeres 2011; Richardson et al 2011). Whereas in the case of finite element methods (FEM) (Hirabayashi et al 2016; Hirabayashi and Scheeres 2014; Holsapple 2004), the specific geometry of the gravitational field is calculated from the specific shape of the studied body, with a given rheology and equation of state of the material. Experiments need to be carried out in droptowers (Sunday et al 2016), and in aeroplanes that perform a number of parabolic flights to emulate lower gravitational conditions (milli-g levels for tens of seconds are common) (Murdoch et al 2013a). In the best case scenario, experiments can be carried out in the International Space Station as this guarantees micro-gravity conditions for extended periods of time (Fries et al 2016; Fries et al 2018). The following sections in this paper will detail how this has been carried out, the caveats and main results.

4 Physical Models

4.1 Continuum models

Granular systems are by definition discrete, namely made of a multitude of individual macroscopic components whose state can be individually described in classical terms: position, velocities, contact and volume forces. However, when zooming out and observing the motion of a granular system as a whole (flowing for instance), a mean behaviour emerges from the multiple interactions of the individual grains: they can be described as an equivalent continuum media, with effective mean properties. Nevertheless, because grain interactions are dissipative in nature, and energy is constantly lost within the granular mass, the theoretical step necessary to derive continuum equations and properties from individual grain-scale behaviour is far from straightforward. As was described in Sect. 3, depending on the external energy available to the system, a granular mass can exist in a dilute state and behave like a gas, or flow in a dense state and resemble a viscous fluid, and eventually stop and become like a solid with specific elasto-plastic properties. Each of these three states of granular matter can borrow the theoretical tools classically used in thermodynamics, fluids mechanics and soil mechanics respectively. We will

provide a short introduction to each of them in the following. However, in nature, granular systems very quickly transit from one state to the other: a granular flow down a steep slope will transform into a gas but will soon stop solid on a gentle topography, while a static granular heap may start deforming as the result of a quake or any other perturbation, leading to shear banding (namely failure) or rapid flow. The physical understanding and modelling of these transitions remain a challenge, and little will be said here on these aspects.

4.1.1 Granular gases

When a granular mass is very dilute, grains interact through short-lived binary collisions, in between which each experiences an independent free flight over a distance comparable to their diameter. In this situation, they resemble the molecules of a gas. Because grains are macroscopic, they do not exhibit a temperature in the thermodynamical sense. However, by analogy with molecular gas, where temperature describe the fluctuating part of the kinetic energy, one may introduce a granular temperature as the mean grains velocity fluctuations (Ogawa 1978; Haff 1983; Jenkins and Savage 1983):

$$T = \langle \delta v^2 \rangle = \langle (v - V)^2 \rangle, \quad (6)$$

where v and V are the grains individual velocity and the mean velocity respectively. The definition of a granular temperature lays the basis for the derivation of a kinetic theory of granular gases, by analogy with molecular gases. Based on Boltzmann equation, it implies complex mathematical formulation that are beyond the scope of the present paper. Here, we will only introduce how transport coefficients can be derived from simple arguments about the transfer of momentum during collisions. Let's consider a collection of grains of diameter d interacting through binary collisions with a typical free-flight distance ℓ . Between two collisions, grains cover the distance ℓ with a typical velocity given by the velocity fluctuations $\delta v = T^{1/2}$ from Eq. (6), so that the typical collision rate is $T^{1/2}/\ell$. During a collision, grains exchange a momentum of the order of $m\delta v = \rho D^3 \delta v$, where m and ρ are the mass and density of a grain. The pressure in the media, given by the total amount of momentum transfer per unit of time (namely, $(\rho D^3 \delta v) \times (T^{1/2}/\ell)$) over an effective section area given by the grains diameter D^2 , will thus obey:

$$P \simeq \rho \frac{D}{\ell} T \quad (7)$$

and we recover the proportionality between pressure and temperature observed in molecular gases.

When the granular gas is sheared with a shear stress τ , it deforms with a shear rate $\dot{\gamma} = du/dz$ accordingly (where u is the velocity in the shear direction), thus defining the viscosity $\eta = \tau/\dot{\gamma}$. The transfer of momentum in the shear direction is thus $mdu = \rho D^3 du$ times the collision rate $T^{1/2}/\ell$ over the effective section area given by the grains diameter D^2 . Considering moreover that $du/dz \simeq du/D$, since D represents the minimum length scale over which gradients may develop in the media, we obtain for the viscosity:

$$\eta \simeq \rho \frac{D^2}{\ell} \sqrt{T} \quad (8)$$

Again, we recover the same proportionality between temperature and viscosity as in molecular gases. Using similar arguments, we can recover the thermal conductivity, which relates temperature gradients to the heat flux (*i.e.* kinetic energy flux), as in molecular gases.

An important aspect of granular gases however is that they lose energy during collisions which are inelastic. Introducing the coefficient of restitution during a collision e , the kinetic energy lost per collision is $\Delta E = \rho D^3 (1 - e^2) \delta v^2 = \rho D^3 (1 - e^2) T$. The rate of dissipation (per unit time and volume) is thus:

$$\Gamma = \rho \frac{(1 - e^2)}{\ell} T^{3/2} \quad (9)$$

Associated to conservation equations, these coefficients provide a simple constitutive model to describe simple granular gases. Moreover, they illustrate the problem posed by granular systems and their rigidity. Indeed, in a granular gas, the free flight distance ℓ is not much larger than the grain size and can rapidly vanish if collisions are highly dissipative. In that case, $\ell \rightarrow 0$ and all coefficients diverge. More elaborate theories will be found in Andreotti et al (2013), as well as how kinetic theory has been successfully applied to Saturn's rings, probably the most famous example of granular gas in space (Goldreich and Tremaine 1978; Spahn and Schmidt 2006; Richardson 1994).

4.1.2 Soil Mechanics

At rest, granular systems can sustain shear stresses and remain static, at equilibrium. This is for instance the case when one consider a granular heap, or the face of a dune: part of the weight of the granular material is sustained by the slope, without the latter starting to flow. This is true however only if the slope does not exceed a certain critical value θ_c , which defines the internal angle of friction of the material. Past this value, the material flows until a new equilibrium is reached. In a similar fashion, a granular material in a shear cell, where both applied pressure P and shear stress τ are controlled, will start to deform and flow only when the shear stress reaches a certain critical value τ_c . This critical value is proportional to the pressure P , and the coefficient of proportionality μ (coefficient of friction) defines the internal friction of the material:

$$\tau_c = \mu P = \tan \phi P \quad (10)$$

where ϕ is the angle of repose defined by Coulomb (1776). This provides at first order a plastic criterion for the failure of granular material: either the shear stress is below the friction threshold and the system remains static, or the shear stress reaches the friction threshold and the system undergoes plastic, irreversible, transformations.

This is the basis of the Mohr-Coulomb criteria (see Fig.5), a well-known tool of soils mechanics, which generalises the above criteria to a fully three-dimensional system for which the direction of the failure is not known a priori. In this case, the ratio τ/P must be evaluated in all possible directions. The derivation of Mohr's circle implies no specific subtleties: it is recovered by projecting the forces applied to a given surface as a function of the surface inclination, and the details of the calculation will be found in Andreotti et al (2013). It provides a graphic way of




Fig. 5 The Mohr-Coulomb failure criteria and Mohr's circle. The failure criterion represents the linear envelope that is obtained from a plot of the shear strength τ of a material versus the applied normal stress σ , where ϕ is the angle of friction, considering also cohesion c .

understanding the distance of a given stress state to the material plastic limit. The existence of cohesive forces between the grains is easily taken into account by the addition of an effective cohesion C to the critical stress τ_c , which is then defined as follows:

$$\tau_c = C + \mu P \quad (11)$$

so that, in contrast to a purely loosely bound media, the higher the cohesion the higher the stress that can be sustained before failure of the granular material. The static angle of repose is the maximum slope that can be supported before the formation of an avalanche, and a dynamic angle of repose is the slope that results after this avalanche has taken place. However, for granular matter, a resisting difficulty lies in the unambiguous definition of the frictional properties, as explained in Sect. 3.3 above. Moreover, granular matter behaves differently if densely or loosely packed, and probably also in different gravity regimes. This difference of behaviour is specifically important when a packing is submitted to a shear stress. If initially densely packed, a granular packing will need to dilate to adapt the shear. This will lead to the localisation of the deformation and the occurrence of shear banding. By contrast, an initially loose packing will adapt shear stresses by deforming in an homogeneous fashion. Hence, a given granular material will exhibit different flow rule depending on its initial state. Critical state theories address these aspects, relating the system distance to the plastic limit (*i.e.* to failure) to its volume fraction (Roux and Radjai 1998).

Finite element methods (FEM) have been used in this context to see how friction can account for the global shapes and stabilities of small bodies, or derive stresses and yields in their interior, and possible failures or surface shedding conditions. The gravitational field is computed from a specific (and approximated) shape of the body, with a given rheology and possible equation of state of the material. This has been applied to both asteroids (Hirabayashi et al 2016; Hirabayashi and Scheeres 2015; Sharma 2013; Harris et al 2009; Sharma et al 2009; Holsapple 2004, 2001), and small planetary satellites (Dobrovolskis 1982; Kay and Dombard 2018).

4.1.3 Fluid Mechanics

In many natural situations, granular matter flows in a dense state that makes it resemble a liquid. In this case, grains are interacting through simultaneous multiple (rather than binary) collisions with many neighbours, while undergoing large deformation, so that theoretical frameworks borrowed from molecular thermodynamics or soil mechanics are not relevant. Here, fluid mechanics provides the basic tools, however with many modifications needed. These tools are essentially the mass conservation, and the Navier-Stokes equation which relates the velocity gradients to the pressure gradients and most importantly, the viscous or frictional dissipation.

Considering that granular flows, such as rocks or debris flows, are often thin compared to their longitudinal extension, a handy simplification consists in assuming that vertical velocities are negligible, and that the motion of the flow is accurately described by solving the one- or two-dimensional problem along the flow path (Savage 1989). In this case, the problem essentially resumes into solving numerically the mass and energy conservation equations which relates flow height and mean flow longitudinal velocity:

$$\begin{aligned} \frac{\partial h}{\partial t} + \frac{\partial}{\partial x}(h\bar{u}) &= 0, \\ \frac{\partial h\bar{u}}{\partial t} + \frac{\partial}{\partial x}(h\bar{u}^2) &= \rho g h \cos \theta \left(\tan \theta - \mu - \frac{\partial h}{\partial x} \right), \end{aligned} \quad (12)$$

where h is the flow height, \bar{u} the flow longitudinal velocity, $\frac{\partial}{\partial t}$ and $\frac{\partial}{\partial x}$ are the gradients in time and in space, ρ the flow density, g the gravity, θ the slope of the topography. The main problem to correctly apply the model to real flows is to have an accurate description of the friction coefficient μ , which balances the two driving forces: gravity and pressure gradients. In depth averaged equations such as Eqs. (12), the coefficient of friction is a basal term, namely it encompasses all the dissipation that occurs in the bulk and at the interface with the substrate into a single scalar describing frictional interactions at the bottom of the flow. As a result, much effort has been devoted to understanding what controls the value of μ in real flows (Brian Dade and Huppert 1998; Kelfoun 2011; Lucas et al 2014), on Earth as well as on other planetary bodies.

When the flow configuration is such that the shallow-layer approximation no longer holds (namely when both longitudinal and normal velocities shape the flow), as for instance in vertical collapses or flow through apertures, the full Navier-Stokes equation must be solved, which is challenging in terms of numerical techniques. An important difficulty is to handle the fact that the flow viscosity may become infinite in some places, where the granular matter turns to a solid-like behaviour, and starts to creep slowly instead of flowing. Here, the viscosity is often defined using the frictional properties of the flow through an effective coefficient of friction μ . The numerical resolution of such complex flows is computationally expensive and its applicability to natural flows down realistic topographies is limited. However, because these more challenging flow situations provide valuable benchmarking cases to test rheological models for viscosity, they have been the subject of many research in recent years (Jop et al 2006; Pouliquen et al 2006; Lagrée et al

2011; Staron et al 2012; Kou et al 2017). The rheology of granular flows is more specifically introduced in next section.

4.1.4 The rheology of granular flows

As we have seen in previous sections, dense granular matter is in essence frictional and defining a coefficient of friction μ is both the simplest and the most efficient way to describe it. Moreover, friction is a finite scalar: it does not diverge nor fall to zero, so that constraining its value experimentally seems feasible. A great variety of configurations have been employed to the task, and the wealth of measurements gathered has eventually led to the identification of a non-dimensional number which controls the value of the friction coefficient of flowing granular matter: the inertial number I (MiDi-GDR 2004). The inertial number can be understood as the ratio of the two characteristic time-scales dominating in a granular flow: the time scale of macroscopic deformation given by the shear rate $\dot{\gamma}$, and the time scale of grain-scale rearrangements under the local pressure P . For a collection of grains of diameter d and density ρ ,

$$I = \frac{d |\dot{\gamma}|}{\sqrt{P/\rho}}. \quad (13)$$

The dependence of the coefficient of friction μ on I is phenomenological and implies the knowledge of three parameters: two extremal values μ_1 and μ_2 for the friction, and a coefficient I_0 setting how quickly μ evolves between the two:

$$\mu(I) = \mu_1 + \frac{\mu_2 - \mu_1}{I_0/I + 1}. \quad (14)$$

Although the existence of three independent parameters might be seen as limiting the predictive value of the $\mu(I)$ dependence, the latter has proven very efficient in many flow configurations. By definition, friction relates pressure and shear stress. As such, it does not describe the way the system deforms or flows in response to a solicitation, and additional operations are needed to turn it into a full rheological description. This is done by deriving an effective viscosity η_{eff} using the norm of the shear rate and the pressure:

$$\eta_{\text{eff}} = \frac{\mu(I) |\dot{\gamma}|}{P}. \quad (15)$$

This operation is valid only if deformations and shear stresses are aligned, which is satisfied for rapid flows. However, this might not be strictly the case for slow motion verging to creep, for which the $\mu(I)$ rheology may not be sufficient. In these slow regimes moreover, non-local effects may start to play a role, whereby distant shear deformations need to be taken into account to describe the local state. Non-locality is however much beyond the scope of this introduction to granular behaviour; further reading may include Bouzid et al (2015), and references therein.

4.2 Discrete, Numerical Models

In contrast to continuum models and finite element methods (FEM) in Sect. 4.1 above, discrete methods are modelling granular media as a collection particles that conserve some of the geophysical characteristics of the aggregates to be studied and the real particles. Several computer simulations of granular material, structures, and flows have been developed (Radjai and Dubois 2011; Mehta 2012, **and other references in the sections below**). A discrete element method (DEM) is any of a family of numerical methods that simulates the dynamics of an ensemble of solid, macroscopic, particles. In these methods, the particles possess not only translational, but also rotational degrees of freedom and the motion of each particle is calculated over time individually, in accordance with its interaction with the rest of the particles in the system. The particles that constitute the aggregates can be subject in general to a variety of external forces fields (long range) and contact forces (short range). In particular, for Planetary Sciences applications, these fields are self-gravity (for an asteroid size body), imposed micro- and milli-gravity fields (for asteroid surfaces). Below, we introduce three of these methods, the first two have already been used for the simulation of SSSBs with mutual gravitational attraction and the third is starting to be used with the same purpose.

4.2.1 Soft Spheres DEM - SSDEM

The soft-sphere discrete element (or molecular dynamics, SSDEM) method for simulating granular material consists in applying Newton's laws of motion to individual deformable spheres. Each individual grain can be submitted to long-distance forces (most often gravity) and interacts with its neighbours through direct, long-lasting collisions. The method is time-driven, and consists in computing the positions and velocities (both translational and rotational) of each grain at time $t + dt$ based on the state of the system at time t . The search for contacts and the integration of the equations of motion can be optimized using the appropriate numerical methods, e. g. the Verlet (Verlet 1967) or linked list (Mattson and Rice 1999) methods, and the leapfrog (Hut et al 1995) or the predictor-corrector (Butcher 2016) integration schemes, respectively.

Great care must be given to the choice of the force models (Radjai et al 1997). The normal force acting between two spheres is given by the Hertzian contact (see Eq. (5) and Sect. 3.1) but a dissipative component needs to be included to model the inelasticity of granular collisions. Most often, the material is considered either visco-elastic (adding a viscous term to the normal force) or plastic (mimicking irreversible deformations of the grains) (Shäfer et al 1996). In spite of this, many SSDEM codes have implemented a linear spring-dashpot contact law due to its analytical simplicity and the fact that the coefficient of restitution is independent of the impact velocity (Herrmann and Luding 1998). The modelling of the frictional tangential forces also deserves attention. Several methods have been proposed, among which the most widely used are the simple regularized Coulomb's law (appropriate for loose systems with no residual stress) and the history-dependent Cundall model (Cundall and Strack 1979).

The main advantage of the soft-sphere discrete element method lies in its truly physical modelling. The forces are computed from solid mechanics and integrated using fundamental laws of physics, while other methods, including event-driven

simulations (HSDEM, Sect. 4.2.2) and cellular automata, may rely on somewhat arbitrary choices. The SSDEM soft-sphere method also has the advantage of computational efficiency, which allows for simulations of a large number of grains (typically up to several millions).

4.2.2 Hard Spheres DEM - HSDEM

In the hard-sphere discrete element method (HSDEM), impacts are treated as instantaneous and point contact, so restitution and momentum conservation equations are used (rather than integrating over the contact duration). This is appropriate in the ballistic regime, where the time to cover the mean free path (t_f) is much greater than time the particles stay in contact during a collision (t_c). This being the physical reality, the approximation that the numerical method makes is to assume that the duration of a collision is exactly equal to zero ($t_c = 0$), making the collisions instantaneous and sound speed infinite. This implies that by construction, all collisions are exclusively binary, and particles cannot sustain long-lasting contacts. The **instantaneity** of the collisions is artificial, but the approximation is valid for kinetically active systems. In the context of granular matter, this method is well suited to the simulation of granular gases and granular flows in a collisional regime, though more generally (and originally) it was used for the simulation of dilute molecular gases (Alder and Wainwright 1960).

Numerically, given a set of particles that form the granular system and their dynamical state (position, velocity and forces on them, including e.g. mutual gravity) the method considers that the particles will move ballistically from collision to collision (Alder and Wainwright 1959). This implies that the order in which collisions happen must be calculated in advance, based on the known state **of the system. Every single collision marks an instant at which a pair of particles stops moving ballistically and exchanges momentum.** The method calculates the amount of time that needs to pass between the present instant and all the possible collisions in the system as the particles move ballistically. Then a list of collisions in hierarchical (chronological) order is build and all particles move for the needed amount time so that the first collision in the list takes place. At this moment, the system has changed, the collision list has to be rebuilt and the process is indefinitely repeated. A brute force approach to the building of the list would be to update the entire system and, though simple, it is very inefficient for large systems. To solve that, Lubachevsky (1991) suggested to update only the two particles involved in the last collision.

From a computation standpoint, HSDEM poses a challenge for parallel processing, since for maximum realism the particle collisions should be treated in time order, so there is a computational bottleneck as these processes only ever involve 2 particles at a time. In contrast, in SSDEM, since particle contacts are finite in duration and treated as extra forces in the equations of motion, no bookkeeping is required to treat the collisions in time order, and everything can be parallelised. Both HSDEM and SSDEM need to identify particle neighbours in order to check for potential (HSDEM) or occurring (SSDEM) collisions. The brute-force neighbour search is an intrinsically order N^2 calculation; tree codes or other data sorting methods can reduce this to order $N \log(N)$ or better (Richardson et al 2011; Rocchetti et al 2017). This however, implies that some collisions could be missed and so care must be taken to make sure that this does not happen. Finally, in HSDEM,

if the timestep is much less than the relevant dynamical time(s), the trajectories can be treated as linear and any collisions can be predicted in advance; in SSDEM, timesteps need to be small enough to resolve the restoring forces on contact. This means HSDEM methods are generally much faster than SSDEM in the ballistic regime (e.g. collision and ejection); the opposite is true in the dense/quasi-static regime, when parallelism wins out despite the tiny steps needed in SSDEM to resolve contact forces.

4.2.3 Non-Smooth Contact Dynamics CD

The contact dynamics (CD) method is based on a mathematical formulation of nonsmooth dynamics, and the subsequent algorithmic developments by J. J. Moreau and M. Jean (Moreau 1994; Jean 1999; Radjai and Richefeu 2009b; Radjaï and Dubois 2011). As in HSDEM, the particles are assumed to be perfectly rigid (non-deformable), but their contacts can persist in time as in the SSDEM case.

In the CD method, the rigid-body equations of motion are integrated for all particles using ‘contact laws’ (instead of contact force laws as in SSDEM) expressing mutual exclusion and dry friction between particles. The equations of motion for each particle are formulated as differential inclusions in which possible velocity jumps replace accelerations. The unilateral contact interactions and Coulomb friction law are treated as complementarity relations, or set-valued contact laws. The time-stepping scheme is implicit but requires explicit determination of the contact network. Due to implicit time integration, inherent in the CD method, this scheme is unconditionally stable. At a given step of evolution, all kinematic constraints implied by lasting contacts and friction are simultaneously taken into account—together with the equations of dynamics—in order to determine all velocities and contact forces in the system, by means of an iterative process pertaining to the non-linear Gauss-Seidel method. The latter consists in solving a single contact problem, with other contact forces set to their values from the previous iteration, and iteratively updating the forces and velocities until a convergence criterion is fulfilled. The iterations in a time step are stopped when the calculated contact forces are stable with respect to the update procedure. The convergence criterion is based on the variation of the mean contact force and/or velocity between two successive iterations. In this process, no distinction is made between smooth evolution of a system of rigid particles during one time step and nonsmooth evolution in time due to collisions or dry friction effects. The uniqueness of the solution at each time step is not guaranteed by the CD method, as the particles are perfectly rigid. However, by initializing each step of calculation with the forces calculated in the preceding step, the set of admissible solutions shrinks to fluctuations which are basically below the numerical resolution. In this way, the solution remains close to the present state of forces. When dealing with complex-shaped particles, the same iterative process can be applied although several contact points may occur between two neighbouring particles. The multiple contacts between two particles are treated as independent unilateral constraints.

The implicit time-stepping scheme makes the method unconditionally stable. Hence, as the small elastic response times are absent from the model, much larger time steps can be used as compared to the SSDEM. The CD method has been extensively employed for the simulation of granular materials in 2D and 3D with various particle shapes.

4.2.4 *N*-body codes, dealing with mutual attraction

Long-range forces such as inter-particle gravity are expensive to compute, since every particle in the system contributes a force on every other particle. There is a great variety of cosmological *N*-body codes (e.g. Klypin 2017), with various approximations used to reduce the force cost. The particle mesh method (Hockney and Eastwood 1988) can be used to compute the potential in Fourier space, but it is not accurate for small separations. One popular approach is to use a tree code, which assigns particles to cells and approximates forces from distant cells by expanding the gravitational potential around the cell's centre of mass to some fixed order. The computation cost is reduced to $O(N \log(N))$ – where N is the number of particles in the system – instead of $O(N^2)$ in the direct pair-wise summation. Various tree algorithms exist, but one of the most widely used is the oct-tree due to Barnes and Hut (1986). Briefly, the space occupied by particles in the system is subdivided recursively into nested cubical cells until each particle uniquely occupies its own cell. To compute the force on a particle, the largest cells are considered first; if a cell subtends an angle less than some critical value (the opening-angle parameter, θ), its contribution to the force is obtained from its moments (see below). Otherwise, its occupied sub-cells are considered recursively in turn until the θ criterion is satisfied, or the sub-cell contains only a single particle. The speed and accuracy of the method depends on both θ and the expansion order. As an example, the acceleration up to quadrupole order is given by Richardson (1993):

$$\mathbf{a} = -\frac{M}{r^3}\mathbf{r} + \frac{\mathbf{Q} \cdot \mathbf{r}}{r^5} - \frac{5}{2} \frac{(\mathbf{r} \cdot \mathbf{Q} \cdot \mathbf{r})\mathbf{r}}{r^7}, \quad (16)$$

where the gravitational constant $G = 1$, \mathbf{r} is the vector between the particle under consideration and the cell's centre of mass, M is the total mass in the cell, and \mathbf{Q} is the quadrupole moment tensor given by

$$Q_{jk} = \sum_i m_i (3x_{i,j}x_{i,k} - r_i^2 \delta_{jk}), \quad (17)$$

where $\mathbf{r}_i = (x_{i,1}, x_{i,2}, x_{i,3})$ is the position of particle i in the cell relative to the cell's centre of mass, and δ_{jk} is the Kronecker delta function. Note the quadrupole moment of a cell can be computed recursively from the moments of its sub-cells using the shift theorem (Hernquist 1987). Higher order increases the complexity but improves the accuracy for a given θ . A typical goal is to achieve an average force accuracy of 1% or better, which generally requires at least quadrupole order and $\theta < 0.7$ rad (≈ 40 deg). Other optimizations are possible to improve efficiency, such as allowing more than one particle to occupy a terminal sub-cell (see Wadsley et al (2004) for a discussion). Note that most trees are not momentum conserving (they violate Newton's third law of equal and opposite forces); fast multipole methods (FMM) can be momentum conserving, but are more complex, and still approximate the force (Greengard and Rokhlin 1987). A big advantage to using a tree for computing forces is that it can also be used to find particle neighbours in $N \log(N)$ time or better (the moments are not needed for this). Parallelising tree codes can be complex, but a basic strategy is to use a balanced spatial tree to distribute equal work among processors, then each processor has its own tree for its particles (Wadsley et al 2004).

Another approach, though following the same basic idea (details in Sánchez and Scheeres (2011, 2012)), is to divide the simulation space with a static grid and keep the distances between their geometrical centres, and their powers, in memory so that look-up tables can be used. The static-grid geometry takes advantage of the fact that the simulations of self-gravitating aggregates usually requires the particles to be concentrated in a region of space and not disperse. Whereas the data kept in memory avoids unnecessary, repeated calculations.

4.2.5 *Monodisperse, polydisperse spheres, and other shapes*

In nature, grains are found in various sizes and shapes, from round beads to complex anisotropic particles. The morphology of the individual grains impacts strongly the global behaviour of the entire granular media (Cleary and Sawley 2002; Pena et al 2007; Lu and McDowell 2007; Donev et al 2004). Moreover, phenomena such as convection and segregation may arise in polydisperse systems (Kudrolli 2004a; Dziugys and Navakas 2009; Metzger et al 2011; Rietz and Stannarius 2008; Miyamoto et al 2007), see Sect. 5.2. Since the handling of the contacts in SS-DEM relies on sphere interaction, a common method to build particles of complex shape is to agglomerate several spheres (Ferrellec and McDowell 2010; Thomas and Bray 1999). For this, spheres of different sizes are glued together allowing large overlaps in order to minimize bumpy edges along the surface. This procedure can be used to create macroscopic rugosity (Ludewig and Vandewalle 2012) as well as anisotropic particles like ellipsoids (Vu-Quoc et al 2000) and even more complex bodies.

Regarding the algorithm, only few modifications have to be made in order to simulate complex shapes. Indeed, the contact detection between two particles remains a contact detection between their respective constitutive spheres. Moreover, the mass and the inertia matrix of each agglomerate are calculated at their creation so that the different forces and momenta can be easily integrated during the simulation. However, a major drawback remains: the large number of spheres that is required to model each particle will strongly impact the computational time. Along with this, the increased complexity of the equations of motion for the aggregates and the additional memory requirements which makes parallelism very difficult.

Another approach to simulating realistic granular systems and the behaviour of non-spherical is the implementation of rolling and twisting friction (Ai et al 2011). This still relies on spherical particles, but the additional friction terms will change the way they rotate on top of one another to mimic the individual and bulk-behaviour of non-spherical particles. DEM methods were adapted to include grains of polyhedra shapes. The main difficulty when moving from a sphere-based grains to more complex shapes is the implementation of a fast contact detection algorithm. This can be done thanks to development obtained in the video game industry (Movshovitz et al 2012). Lastly, in the CD method, exact particle shapes can be used since the constraints of mutual exclusion and dry friction do not refer to particle shape. Thus the CD method allows for considering general polyhedral shapes, and other complex-shaped particles with multiple contacts (Azéma et al 2017), but with considerably increasing computational cost.

4.3 Limitations

As any other numerical method that has been developed in order to study a physical system, the methods presented here have limitations which have to be taken into account, so that results are not misinterpreted and methods are not misapplied.

First off, one of the main common limitations of DEM codes is the number of particles that can be simulated given the computational and time constraints. Though some naturally occurring granular systems, and in particular laboratory experiments, can be replicated and simulated almost grain by grain, many more consist of more particles than is possible to simulate with the current computational facilities available for research. Therefore, a compromise must be reached between the number of particles that form a real system and those that can be simulated for any practical purpose. Ideally, we want a simulation with enough particles so that some characteristic parameters such as angle of friction, angle of repose, cohesive/tensile/compressive strength, bulk density and filling fraction—among the most commonly used—match the real system. The inclusion of self-gravity in the calculations only exacerbates this problem. DEM methods however can handle a large number of particles with the aid of HPC computing.

Another common limitation is the size distribution and shapes that can be used for the particles. In general, very wide size distributions are not simulated as the number of particles required to have a representative system grows to impractical levels. As an example, for a 50/50 mixture by volume of 1 mm and 1 cm spherical particles, for every 1 cm particle there are up to ≈ 740 of the small 1 mm size particles, depending on the packing. Additionally, the contact detection algorithms rely on the search for the closest neighbours to avoid unnecessary checks and for that the simulation space is divided into cells that are large enough to contain the largest particle. A large size disparity means that any given cell will contain a great number of small particles that are too far away to collide. However, the contact detection algorithm will be forced to search for possible contacts when almost none is possible. As for the shapes, an increasing complexity in the particle shape means more sophisticated contact detection algorithms and equations of motion, which will in turn require greater computing time. Besides, Procopio and Zavaliangos (2005) analysing the case of compaction with multiple particle FEM method, showed that compaction is not only porosity driven. The mechanical response of an assembly of particles during **dense packing** with finite elements models (FEM) is softer than in discrete element models (DEM), which are generally limited to non-interacting contacts, or need to be corrected to allow densities larger than approximately 0.8 (Harthong et al 2009).

For HSDEM, the premise that the free time path is much larger than the duration of collision, $t_f \gg t_c$, is of paramount importance as this determines the systems that can accurately be simulated with this method. More specifically, this method can be used to simulate the behaviour of molecular and granular gases, systems in a collisional regime. The premise begins to be untrue for systems in frictional regimes (granular liquids) and condensed phases (granular solids). In these latter regimes, the particles sustain long-lasting contacts and the approximation of $t_c = 0$ is no longer valid. Furthermore, if applied to these systems, unless the restitution equations are modified, the particles, not being able to stay in contact, will try to collide infinitely often in a finite amount of time. This is what has been

defined as *inelastic collapse* (McNamara and Young 1994). Even with this modification however, unless in a crystalline structure, the system will have an angle of friction between $0^\circ - 5^\circ$ and will behave as a granular liquid.

For SSDEM one specific limitation is the time step (δt) needed for the integration of the equations of motion. The magnitude of δt is determined by two characteristics of the simulated system: the stiffness of the grains - given by the contact model (linear spring-dashpot, Hertzian spring-dashpot are commonly used), and the characteristic collision speed. In general, in SSDEM codes the stiffness of the grains is underestimated in order to avoid a too small δt as it is chosen as a fraction of t_c . Using a δt which is too large compared to t_c would result in missed collisions or particles overlapping excessively as the collision dynamics cannot be accurately resolved. On the other hand, the grains cannot be so soft that the overlap of the particles is too great for a typical collision (1% overlap is common). It is worth mentioning that changing the stiffness does fundamentally change the material behaviour; if it is important to the problem, this parameter can be tuned to better capture the critical physics (e.g. sound speed), at the expense of a smaller maximum overlap and therefore a smaller timestep. Thus, a compromise must be reached so that the simulation is still realistic without being impractically slow. This makes this method much slower than HSDEM for highly active, rarefied systems and much better suited for more densely packed, quasi-static systems.

As to the CD method, it has the advantage of allowing for much larger time steps than the SSDEM but needs sweeping the contact network a number of times in order to determine the contact forces and particle velocities at each time step, until the convergence criterion (precision on forces and velocities) is reached. The number of iterations increases with both the number of particles and the required precision (Radjai and Richefeu 2009b). Hence, in practice, CD simulations can be either much faster or slower than SSDEM simulations, depending on the precision used. Obviously, the CD method is more adequate than the SSDEM when the ratio of external stress to the particle stiffness (leading to small particle deformations) is small and when the real stiffness matters for the physical behaviour of a granular material. Moreover, except for spherical particles for which the Hertz force law is classical, there is no general force law for contacts between particles of arbitrary shape (for example, for face-face contacts between polyhedral particles). For this reason, in applications of the SSDEM arbitrary particle shapes are generally modelled as aggregates of spherical particles. Besides, it is also important to consider the parallelisation potential of different methods. To this respect the CD method, being based on a global determination of forces by iterations, cannot be as efficiently parallelised as the SSDEM.

5 Experiments

5.1 Lab experiments on ground

Ground-based impact experiments have provided fundamental insights into granular flows. The experiments on a wide range of flows provide a basis for understanding granular flows on asteroids, but with the strong caveat that a number of observations are linked to ground-based environmental conditions very distinct from small bodies in space. In particular, humidity and the air surrounding gran-

ular matter in most ground-based experiments affect the behaviour of granular systems. For example, analyses of ejecta flow showed the importance of interstitial air in driving very high ejecta flows (Lohse et al 2004). Moreover the gravity field of the Earth is predominant, thus experiments in vacuum and in micro-gravity would be closer to reality.

Since there are a number of possible interaction forces between grains, ground-based experiments also provide important insights into the relative importance of various physical mechanisms in granular materials. Here intuition from molecular or atomic systems often leads our intuition astray. For example, collisions between dielectric molecules tend to neutralize the system for molecules or atoms of equal type. It requires mixtures of different materials to charge molecular systems through collisions. However, Pächtz et al (2010) found that granular particles of equal types, colliding under dry conditions, can accumulate charge. The key insight is that during collisions of granular matter only the areas near a contact point neutralize. Thus in the presence of electric fields (e.g. due to charges of neighbouring particles) collisions can charge particles.

Similarly, Shinbrot et al (2004) discovered through ground-based experiments that the features of Martian gullies typically associated with fluid immersed granular flows, may also be observed in dry flows when the flow speeds are high compared to the typical settling speeds, an effect expected to be enhanced under the reduced gravity conditions of Mars.

For slow granular flows and plastic deformations of granular matter, where grains remain in contact with each other and rearrange through rolling or sliding, gravity in the bulk of the flow is small compared to contact forces between particles. Thus ground-based experiments will yield flows that are also expected under microgravity conditions. New experiments allow us to measure translations of all particles in three dimensions in slow granular flows (Dijksman et al 2012). First studies on these systems have provided insights into the important question of reversibility and ageing of granular matter at the particle level. More recent experimental studies also allow for analysis of the rotations of particles within a three dimensional granular flows (Harrington et al 2014).

Finally, earth-based experiments have revealed one of the most ubiquitous differences between granular matter and atomic or molecular matter: segregation of particles by size, weight, or shape.

5.2 Segregation

An intriguing feature of polydisperse granular matter is their tendency to segregate when submitted to external constraints. The phenomenon is omnipresent in nature but also in diverse processes implying the handling of granular matter. From an industrial point of view, segregation is often considered as a parasitic effect that hinders the homogeneous mixing of components (Poux et al 1991) especially since it can be triggered by any variation in mechanical properties of the grains. Despite most of its driving mechanisms are related to gravity, granular segregation is also observed in low gravity environment where it is held responsible for particular surface granulometry of small celestial bodies covered by regolith (Miyamoto et al 2007; Asphaug 2009; Gundlach and Blum 2013). Generally speaking, granular segregation is observed and studied in various situations. Segregation is observed

in both rapid granular flows and plastic flows for almost all conditions (Ottino and Khakhar 2000; Jaeger et al 1996a), and often accompanied by convective flows (Rognon and Einav 2010). For mixtures of large and small, or heavy and light particles the tendency of granular flows to segregate manifests itself in the Brazil Nut Effect (BNE), where large or heavy particles rise to the top. For plastic flows driven by periodic forcing, the onset of segregation with increasing forcing amplitude was seen to coincide with the onset of convective flow (Harrington et al 2013).

When grains are slowly poured onto a plate a heap forms and starts to grow. However, the slope angle between the piles surface and the plate will never exceed a limit value called the angle of repose that depends on the size, shape and rugosity of the grains. When a binary mixture flows down a heap, segregation occurs since the different mechanical properties of both species lead to different angles of repose. This phenomenon is known as granular stratification and results in the formation of layers in which grains of different species are separated (Makse et al 1998; Koeppel et al 1998; Aranson and Tsimring 2006; Fan et al 2012; Shimokawa et al 2015). The formation of strata is linked to the avalanches along the heaps surface. As the granular matter flows down the heap, voids are created along the surface. These voids are more likely to be filled by small grains with creates a downward flux of the latter while large grains remain on the top (Savage 1993; Cizeau et al 1999; Kudrolli 2004b; Gray and Thornton 2005; Schröter et al 2006; Gray and Ancy 2011). Each of the so obtained pair of layers grows, by the propagation of a kink, from the bottom to the top of the pile.

Segregation can also be studied in a rotating cylinder partially filled by a granular mixture. In this case, radial segregation occurs quite rapidly: small and rough particles migrate towards the centre while large and smooth grains rotate around them (Khakhar et al 1997; Hill et al 2004; Hajra and Khakhar 2011). Under certain conditions, the radial core develops more complex patterns (Khakhar et al 2003; Zuriguel et al 2006). If the cylinder is long and narrow, radial segregation is often followed by axial segregation where patterns of segregated bands appear along the axis of rotation (Zik et al 1994; Clement et al 1995; Cantelaube and Bideau 1995; Caps et al 2003; Fischer et al 2009). This phenomenon is well known and was observed for the first time by Oyama (1939). The mechanics of axial segregation are related to the different dynamic angles of repose of the rotated grains, i.e. the angle of the slope in the drum for continuous flow regime (Makse 1999; Orpe and Khakhar 2001; Seiden and Thomas 2011). Moreover, axial segregation has recently been observed in the case of a spherical container, rotating about its horizontal axis (Finger et al 2016).

When a granular mixture is vibrated vertically (i.e. on Earth's gravity), larger particles rise to the top of the system. This phenomenon, known as Brazil Nut Effect, has been studied for a long time (Rosato et al 1987; Möbius et al 2001; Garcimartin et al 2002; Godoy et al 2008; Metzger et al 2011; Matsumura et al 2014) and is linked to mechanisms such as percolation and granular convection (Knight et al 1993; Hong et al 2001; Huerta and Ruiz-Suarez 2004). Under the effect of vertical shaking, the particles in the system lift off. The voids that are created that way are easily filled by the small grains which leads to a ratchet-like rise of the large ones. Moreover, the vibration induced convection in the container creates a wide upward flow in the central part of the system but only a thin downward flow along its boundaries so that all sorts of grains can rise to the

surface but only small particles can dive back to the bottom. Investigations during parabolic flights (Güttler et al 2013) have shown that BNE can be observed in low gravity conditions even though its driving mechanisms are strongly linked to gravity, although sometimes in a complex manner (Staron 2016). This result consolidates the theory that certain surface structures on asteroids are created by this kind of segregation (Miyamoto et al 2007).

Segregation has also been observed in granular gases (Pöschel and Brilliantov 2003). Unlike in continuous media, thermal agitation is not enough to generate the motion of the particles composing a granular gas. Furthermore, collisions between grains are dissipative so that external energy has to be injected permanently (often through vibrations) into the system in order to maintain a stationary gas like regime. Depending on the filling properties and driving mechanism of the system, granular gases exhibit intriguing phenomena such as anomalous scaling of pressure and non Gaussian velocity distribution (Rouyer and Menon 2000; Losert et al 1999; Tatsumi et al 2009; Falcon et al 2013; Scholz and Pöschel 2017). If one stops the external energy supply, the average energy in the system decays which is known as the cooling of a granular gas. After a while, slow and dense regions called cluster form in the cold regions of the system (Goldhirsch and Zanetti 1993; McNamara and Young 1994; Maaß et al 2008; Brilliantov et al 2018). In the case of binary granular gases it has been shown that clustering can be followed by a particular kind of segregation where domains of the same granular species tend to merge together (Cattuto and Marconi 2004). In denser granular gases, clustering can occur despite of an external energy injection (Falcon et al 1999; Noirhomme et al 2017). In these systems, a large and slow domain forms in the centre of the container and acts as a liquid phase coexisting with a surrounding gas phase. Here again, in the case of a binary mixture, clustering goes hand in hand with segregation. It has been shown that the different granular species segregate within the cluster, giving rise to a layered structure of the bulk (Serero et al 2009; Opsomer et al 2014, 2017).

5.3 Experiments in microgravity

Microgravity experiments can be realized by several means. However, in all of them, the key mechanism to approach weightlessness is free falling. Indeed, experiments in droptowers, in parabolic flights, and even in the international space station are falling down towards earth all together with their measuring instruments (Pletser 2004; Von Kampen et al 2006), see Fig. 6. Granular materials have been studied in microgravity for about two decades. In particular, granular gases have been the focus of this research since the microgravity conditions allow for a more homogeneous distribution of the particles in the system (Pöschel and Brilliantov 2003; Heisselmann et al 2010; Hou et al 2008; Leconte et al 2006). In the late nineties, Falcon et al (1999) observed for the first time clustering of a continuously driven granular gas during the Mini-Texus 5 rocket experiment. Since then, the transition from granular gas to cluster has been investigated numerically (Opsomer et al 2011; Noirhomme et al 2017) and experimentally (Maaß et al 2008; Tatsumi et al 2009), and is nowadays in the focus of the VIP-Gran Topical Team of

the European Space Agency⁴. Experiments with driven rod shaped particles were realized in a REXUS flight. Thanks to a precise tracking of the anisotropic particles, non-Gaussian velocity distributions and energy non-equipartition could be highlighted (Harth et al 2013). In another experiment (Lee et al 2015), free falling charged particles have been studied. The authors report the observations of individual collide-and-capture events between particles, including Kepler-like orbits. Other parabolic flights have studied granular convection in varying gravitational conditions (Murdoch et al 2013), and shear reversal in regolith dynamics (Murdoch et al 2013b). Bouncing and cohesion through Van der Waals force on aggregates or clusters of small grains (< 1 mm) have been tested on ZARM droptower (Brisset et al 2017) at 10^{-6} g, as well as on sub-orbital rocket down to 10^{-3} g (Brisset et al 2016). Low velocity impacts (2-40 cm/s) of larger (approx. 10cm) projectile on a granular surface have been tested using an Atwood machine installed in a small droptower (Sunday et al 2016). This system, which uses a system of pulleys and counterweights, allows the effective surface acceleration of the granular material to be varied from $0.2 - 1$ m/s² (Murdoch et al 2017a). The latter experiment showed shallow penetration ($< 1/4$ of the projectile diameter) and no rebound; nevertheless further experiments using other surface materials, impactor properties, and gravity regimes could be performed to confirm this behaviour.

Segregation mechanisms of denser systems with an intruder were also studied during parabolic flights (Güttler et al 2013). A large particle is placed at the bottom of an assembly of smaller ones. Through the shaking of the cell, the intruder rises up to the surface of the pile (the Brazil Nut Effect, see Sect. 5.2 above). The uprising speed was then measured for different values of gravity (Earth, Moon and Mars gravity) and a first scaling law was proposed. In bi-disperse granular media, segregation phenomena can occur even in microgravity (Louge et al 2012; Opsomer et al 2017). The driving mechanism is no longer convection and percolation as on Earth, but rather the gradients in the fluctuation energy of the grains. Though not many experiments involving granular matter have been carried out to date in the ISS, the Strata-1 experiment (Fries et al 2016; Fries et al 2018) was the first to put different mixtures of grains in orbit for a long period of time. The samples ranged from spherical glass beads to glass shards, to crushed meteorite simulants and even a real crushed meteorite sample. At the moment, the results of this experiment are still being analysed (Dove et al 2018) and the Strata equipment is being repurposed as the Hermes facility (John et al 2018) that will be made available to other researchers in the near future. One of the advantages of running experiments in the ISS is the long duration and quality of the microgravity environment which are essential in order to observe the evolution of systems that take more than a few seconds to be finalised and therefore, could not be carried out in droptower or parabolic flights due to their short duration.

⁴ SpaceGrains ESA Topical Team from the European Space Agency. <https://spacegrains.org>




Fig. 6 Schematic summary of low- to micro-gravity experiments available, with different duration of the gravity environment.

6 Benchmarking cases

6.1 Macro and micro-scale benchmarks: what to learn from observations

Theoretical development, numerical modelling, and experiments of granular systems in low, and self-gravity regime, all concur to a better understanding of asteroids. This is often needed for studying sample return mechanisms, for estimating the outcomes of high-velocity impacts, collisions and ejecta, or for predicting low-velocity impact and bouncing of platforms on the surface, etc. (e.g. Biele et al 2017; Thuillet et al 2017; Murdoch et al 2017a; Ballouz 2017). We will focus in the following on segregation phenomenon, long term evolution spin-up and mass shedding, and analysis of reaccumulation process. Several computer simulations have been proposed to tackle the analysis of self-gravitating planetary bodies as granular systems, starting mostly with HSDM (Richardson 1993; Richardson et al 2011; Leinhardt et al 2000; Murdoch et al 2012; Michel et al 2002; Walsh and Richardson 2006; Tanga et al 2009b; Walsh et al 2012; Campo Bagatin et al

2018b), and later with SSDEM (Sánchez and Scheeres 2011; Schwartz et al 2012; Tancredi et al 2012), see Sect. 4.

6.2 Segregation

Shaking giving rise to segregation (see Sect. 5.2) can have different origins on asteroids (reaccumulation, impacts, tides, stress cycles, ...), with different frequencies and amplitudes. Seismic shaking mechanism has been proposed to resurface asteroids and account for presence of ponds on (433) Eros (Cheng et al 2002; Richardson et al 2004), or boulders on (25143) Itokawa (Saito et al 2006; Miyamoto et al 2007), or young surface on specific classes of Near-Earth asteroids (Binzel et al 2010). In addition to flows, the BNE could then be effective on asteroids and provide segregation on regolith grains. This has been simulated numerically by methods presented in Sect. 4 (Sánchez and Scheeres 2009; Tancredi et al 2012; Murdoch et al 2012; Matsumura et al 2014; Perera et al 2016; Maurel et al 2017), with hard-sphere DEM, and later soft-sphere DEM. Perera et al (2016) suggest that the mechanism of the BNE that is most relevant—in presence of a binary size-distribution—is that of percolation, and not convection, and that the innermost regions remain unsorted. Further, Maurel et al (2017) have performed numerical simulations investigating the BNE in an unconfined environment. They show that, under Earth gravity ($1g$), a void-filling mechanism is predominant, in contrast to more classical granular convection driven BNE in presence of walls. While this void-filling mechanism remains relevant in lower gravity regime ($10^{-4}g$), it is however differently influenced by the friction properties of the particles. Last, Chujo et al (2017) have analysed the BNE (as well as the reverse BNE, occurring when oscillation frequency is high, and the bulk density of the larger particles is larger than that of the smaller ones) in low-gravity environment with an intruder, under ‘less-convective’ conditions. They also point out that the amplitude and frequency of vibrations that may be induced on small bodies, and relevant to the problem, is still not well known.

6.3 Rotation, spin-up, mass shedding

6.3.1 The YORP Effect:

It has been generally accepted that solar radiation pressure, the absorption, re-emission and reflection of photons emitted by the Sun can, over the lifetime of SSSBs, produce not only an acceleration of its orbit (the Yarkovsky effect) but also a net torque on them which will change their rotation state (Rubincam 2000). For this torque to appear however, it is essential for solar photons to impact a body with an irregular shape as otherwise photons would be re-emitted or reflected in symmetric directions and their effect would average out. An asymmetric body on the other hand would be similar to a propeller; sunlight bouncing off the blades and causing it to change its rotation. This is what is defined as the Yarkovsky-O’Keefe-Radzievskii-Paddack effect (YORP for short).

6.3.2 Deformation and Disruption

Given that SSSBs are granular in nature, their structure can yield/fail under stress. Events that can cause this structural failure could be rotation at high enough spin rates Ω , collisions and planetary flybys. From these, in this section we will focus our attention on the failure that is produced due to high rotation rates.

As previously explained, the interaction of an irregularly shaped body and solar photons, over millennia, can produce a net torque that can change the rotation rate of a SSSB. Though it is possible to calculate the critical rotation period for a SSSBs of arbitrary shape numerically, it is useful to start with a simpler shape so that we can obtain some insight. For this, we could start with a sphere of radius r and bulk density ρ .

For a gravitationally bound aggregate, the critical spin period

$$P_c = 2\pi/\Omega_c$$

can be calculated by equating the acceleration of gravity at the surface of a sphere with the centrifugal acceleration at the equator, just like it was done by Pravec and Harris (2000). This would allow us to derive a criterion for the critical limit of rotation period, depending only on the density of the sphere ρ ,

$$\frac{Gm}{r^2} = \Omega_c^2 r \rightarrow P_c = \sqrt{\frac{3\pi}{G\rho}} \quad (18)$$

where G is the gravitational constant and m is the mass of the sphere. For bulk density between 2000–3000 kg.m⁻³, this expression gives us a spin period P_c between 2 – 2.3 h. Conversely, the spin barrier of the observed asteroid population is ≈ 2.4 h (see Sect. 2.1, Fig. 1). Though this simple calculation provides a first approximation, and a great insight to the make-up of asteroids, there is a feature that it could not explain. For the smaller asteroids (below $\approx 100 - 200$ m), this spin barrier did not seem to exist (Pravec and Harris 2000). Additionally, this spin barrier might more accurately be described as a deformation or a fission barrier, as asteroids reaching it would have to deform or go through a fission process as their structure would fail at such high spin rates (Jewitt et al 2013, 2014; Hirabayashi et al 2014; Walsh and Richardson 2008; Sánchez and Scheeres 2016; Hirabayashi and Scheeres 2015).

From here, we can take one step further and study what happens to tri-axial ellipsoids when subjected to high spin rates. The form in which these bodies would fail can also be solved analytically, though this requires the use of Soil Mechanics theories, elastic-plastic theories to be more precise (Holsapple 2001; Sharma et al 2009), in particular, limit analyses. Within this approach, one seeks the maximum load that a body (whatever its nature) can sustain without failing. This circumvents the need to know the past history of the body that would necessarily affect its evolution. For these theories to be applied a framework is needed, a basic principle for many granular (soil or gravel) or solid (rock) geological materials is that the shear yield stress on any plane increases with the normal pressure on that plane. The simplest criterion of that type is the MohrColoumb (Drucker-Prager is another, more sophisticated possibility) yield criterion (Holsapple 2001), determined solely by a cohesive strength and an angle of friction. Also required is some ‘flow rule’ to determine the flow: some prescription of the plastic deformations that

occur when the stresses meet or exceed the yield criteria. Then, in principle, it should be possible to trace a particular loading history and determine the resulting stress fields.

By doing this, it was possible to determine that most asteroids shapes and rotation rates were well within the equilibrium limits for low angles of friction which are typical for dry, cohesionless soils and so the observed shape and spin rate of the asteroid population could not be used to assert that these bodies had to be cohesive. This analysis however, could not be applied to comets where small cohesive forces could provide additional strength to the structure.

This, of course would mean that the source of the strength of asteroids is purely gravitational – in the case they are not monoliths – and that all observed asteroids should be below the spin barrier. However, it was observed that the small asteroid population⁵ (approximately $< 100 - 200$ m) contained not only fast, but even super-fast rotators with periods of only a few minutes possibly depending on their taxonomic type (Pravec and Harris 2000; Warner et al 2009; Taylor et al 2012; Perna et al 2016). This is what has led scientists to believe that cohesive forces could be important for these granular systems; **or that otherwise the small fast-spinning asteroids are essentially monolithic rocks** (e.g., Polishook et al 2016).

At this moment we have to make a point about nomenclature, we will use the word cohesion and cohesive force to refer to the surface-surface attractive force between any two grains in contact. The term cohesive strength will be used to refer to the parameter in the different yield criteria which **is defined as the shear stress at zero normal stress**. Complementary to this, the term tensile strength is defined as *the normal stress at zero shear stress*. These two could be related for specific materials, but there are indications that their relationship can change at very small loads (Kim et al 2009).

The deformation of ideal spherical and ellipsoidal self-gravitating aggregates can also be studied with the same theoretical tools presented above (Holsapple 2001, 2004) and extend them to allow for cohesive and therefore, tensile strength (Holsapple 2007).

By doing this, it was found that the presence of tensile and cohesive strength for a large body (> 10 km) makes no difference in the permissible spin. This is, gravitational effects dominate the strength of the body. This implies that the observed spin limit for large bodies cannot be used to infer zero-strength (cohesive/tensile) granular asteroids. On the other hand, the strength that allows the higher spins of the smaller and fast-spinning km-sized bodies is only on the order of $10 - 100$ kPa, a very small value compared to small terrestrial rocks. Additionally, it is stated that the strength needed for small granular asteroids to become fast rotators could be originated in an accumulated slight bonding between their constitutive particles.

For asteroids between approximately 0.2 and 10 km, cohesive strength does not seem to influence the maximum spin rate they can reach, but it affects the overall shape of the bodies. This could partially explain the shapes and surface morphology of asteroids Itokawa, Ryugu and Bennu for which photographic evidence is available.

⁵ Sizes, when not measured directly, are estimated from the absolute magnitude H and by assuming an albedo of 0.2

Though the theoretical results described above provided great insight into the mechanisms that shaped the observed asteroid population as well as their internal structure, questions about the origin of their cohesive strength as well as the origin of binary asteroids, asteroid pairs were still unanswered. Other questions, more related to planetary defence and space exploration, also needed answers.

One weakness of the theoretical models used in the above described research is that they are not dynamical. This is, they cannot show the reshaping of the bodies and are always constrained to ellipsoidal shapes. It is here where the use of DEM codes becomes important. The use of these kinds of codes followed the same path as the theoretical efforts, that is, start with spherical and ellipsoidal shapes, no cohesive strength and the added numerical particularity of having only mono-disperse spherical particles as the constituents of the aggregates. A drawback of this last point was that crystallization was unavoidable, but even this was used to emulate higher angles of friction (Walsh and Richardson 2008) when needed. These aggregates would reproduce the behaviour of assembles of particles with friction angles of $\approx 40^\circ$ when crystallized and between $0^\circ - 5^\circ$ when the particles were randomly packed (not naturally found for granular matter).

The influential work of Walsh et al (Walsh and Richardson 2008; Walsh et al 2012) provided evidence supporting the idea that binary asteroids could be continuously formed through the shedding of asteroid material at high enough spin rates and its subsequent reaccumulation to form the secondary of a binary system. However, there were questions about the needed time for this process (Jacobson and Scheeres 2011) to take place as well as the influence of the crystallization of the aggregate. Besides, the final axisymmetric top-shapes may be a particular outcome from the simulation of Walsh and Richardson (2008), as shown by Cotto-Figueroa et al (2015) who analysed the coupled spin-shape evolution with different initial configurations of the aggregated particles, and reducing crystallisation effects. **Including polydisperse spheres, and different block shapes adds extra complexity and realism to the simulation models (Walsh and Richardson 2008; Walsh et al 2012; Michel and Richardson 2013; Campo Bagatin et al 2018a).**

Subsequent studies carried out by Sánchez and Scheeres (2011) using a SSDEM code **did implement random packings—as in Comito et al (2011), and in contrast to Walsh and Richardson (2008)—so that crystallisation was explicitly avoided**, and they started to study the influence of surface-surface friction. Their studies showed that this type of simulation fully agreed with the theoretical models of Soil Mechanics (Sánchez and Scheeres 2012) for aggregates with angles of friction of $\approx 12^\circ$ and $\approx 25^\circ$ (these angles of friction are not naturally found in nature for gravel). That is, the aggregates followed the deformation path that was determined by the theory and at the correct spin rates. Additionally, it was found that the aggregates failed at the centre, not producing granular flows on the surface which was the failure mode showed by Walsh and Richardson (2008). One intriguing feature was however, that at times the aggregates would split into two almost symmetrical pieces and some others, it would simply eject individual particles. Besides, Tanga et al (2013) showed that particle ejection is not incompatible with splitting, as the two can occur in the same system if the spin-up phase is continued.

6.3.3 Cohesive Strength

Even though it had already been established that **a hypothetical fast rotating gravitational aggregate** would necessarily have some cohesive strength, its source had not been established. To tackle this problem, the work of Scheeres et al (2010) attributed this to the comparatively strong Van der Waals forces. If we define a *bond number* \mathbf{B} to the ratio between the cohesive/adhesive attachment a particle feels and its weight, in an asteroid environment, this ratio becomes 1 for centimetre size particles in the ideal case. **Of course small SSSBS with high spin rate can either be strong monoliths or gravitational aggregates with relatively weak cohesion; indeed, the observed spin rates of fast rotators do not require high cohesive strength of solid rock (Sánchez and Scheeres 2014).**

Supported in these findings, Sánchez and Scheeres (2014), using the same DEM code, find out that the tensile strength of an ensemble of self-gravitating spherical particles is inversely proportional to the average particle size. This result, allows them to study self-gravitating aggregates with realistic asteroid sizes, friction angles and tensile strength, but without the burden of having to simulate an impractically large number of particles (Sánchez and Scheeres 2014). **Angles of friction of $\approx 35^\circ$ were obtained through the implementation of rolling resistance (Ai et al 2011).** In essence, in their model, the larger boulders are embedded in a cohesive matrix that holds the entire aggregate together. In a way, this would be a Van der Waals cement. They calculate that a cohesive strength between 25-100 Pa would be enough to explain why asteroids approximately $< 100 - 200$ m in size could have spin periods below the 2.2 h of the spin barrier. Other works compared the hard-sphere and soft-sphere DEM simulations including cohesion as well (Schwartz et al 2013). Studies of observed fast-rotating asteroids also confirm to have cohesive strengths that are far below what would be expected for competent rocks (Rozitis et al 2014; Hirabayashi and Scheeres 2015; Hirabayashi et al 2014).

Additionally to this, a subsequent study by Sánchez and Scheeres (2016) found that the amount of deformation of a self-gravitating aggregate is greater for low angles of friction and vice-versa. Whereas the amount of cohesion was directly related to the fission process. That is, at low or no cohesion, particles would be ejected in a one-by-one-fashion, producing a tail. Whilst at high cohesion, the aggregates would eject larger, coherent groups of particles at once. This could result in the aggregate going through a catastrophic disruption process that would break the body in several coherent pieces. For the most cohesive cases, the aggregates would split in two symmetrical pieces. **Similar work by Zhang et al (2018), incorporating friction and cohesive interactions in the manner suggested by Sánchez and Scheeres (2016), confirmed most of these results.**

6.3.4 Internal Structure

Up to this point, one assumption made by most studies is that the modelled aggregates and asteroids had completely homogeneous interiors. However, this is not necessarily the case. For instance, it is also believed that in a post-catastrophic collision large blocks with lower velocities reaccumulate first, and are at the centre of the body, while smaller material could be kept closer to the surface by friction (Britt and Consolmagno 2001). In this regard, it has been showed through theory

and simulations that a spherical aggregate with a central core, which is structurally stronger than its external shell, would avoid the initial internal failure prevalent in homogeneous aggregates and fail at the surface (Walsh et al 2012; Scheeres 2015; Hirabayashi 2014; Hirabayashi et al 2015). In numerical simulations, such structure was the only one able to reproduce the surface slope of asteroid 1999 KW4 (Sánchez 2015) when the radius of the core was $\approx 0.7R_b$, where R_b is the radius of the aggregate, in agreement with Walsh et al (2012), and as predicted by the theory. Those researchers also explain the surface shedding process found by Walsh and Richardson (2008). Essentially, the crystalline packing used in Walsh’s numerical simulations had intrinsically build an interior which was stronger than the outermost shell formed by surface particles. These aggregates were originally built to fail at the surface though the crystalline packing was used only as a tool to avoid the fluid-like behaviour of the grains when simulated with an HSDEM code.

On the other hand, aggregates with weak cores and strong shells have been studied only through numerical simulations. Upon rotation, these aggregates locate the greatest stress at their centre where a severely weakened interior would fail long before the stronger shell. This means that an equatorial ridge would not be formed by granular flow—which would not happen under these conditions—but rather by the global deformation of the body. Preliminary results also show that if the core is about $\approx 0.5R_b$, it is possible to obtain a shape similar to that of asteroid Itokawa, though its reproducibility is still to be proven (Sánchez and Scheeres 2018).

A possible confirmation of this correlation between fission size, cohesion and internal structure came from a study of the equatorial cavities in asteroids 2008 EV5 and 2000 DP107 that had previously been attributed to impact events (Busch et al 2011). Tardivel et al (2018) showed that if a hypothetical original body with a filled cavity rotated at high enough spin rate, the first place to fail in tension would be exactly where the cavity appears at the moment. Kinetic sieving (Gray and Thornton 2005), produced as a result of surface flow, which should in turn be produced by a strong core, could produce a ‘rocky equator’ (size segregation with a flow of the largest rocks to the equator) and explain the low tensile strength of this specific region for this fission mechanism to work.

Therefore, all these studies suggest that the shape of granular asteroids, and the specific form towards which they evolve under rotation, are intrinsically linked to internal structure and structural strength (Sánchez and Scheeres 2018). Examples of this statement have already been mentioned, asteroids Itokawa, 1999 KW4, 2008 EV5 and 2000 DP107. Additionally, late in 2013 two more asteroids that were observed, P/2013 R3 and P/2013 P5 which were termed *active asteroids*. Both structures failed, but in very different ways (Jewitt et al 2013, 2014). The former broke apart in several coherent pieces, whereas the latter exhibited several long tails. If we assume that the aggregate and core approach can be applied to these two bodies, asteroid P/2013 R3 might have had a homogeneous interior and asteroid P/2013 P5 a strong core (Hirabayashi et al 2015; Sánchez and Scheeres 2016).

6.3.5 Scaling

As it has been proven by the authors cited above, the cohesive strength of a self-gravitating aggregate could become important for small (approximately ≤ 100 m) bodies. This would imply that the onset of deformation or disruption is the result of an interplay between the effects of material properties (cohesion, density, porosity, particle size distribution, friction) and gravitational forces which depend on the aggregate size and mass. Intuitively, increasing the size of a cohesive aggregate should be equivalent (structurally) to reducing the value of cohesive strength and vice-versa (Sánchez and Scheeres 2015). However, the scaling was not very clear. To solve this, Azéma et al (2018) analysed the stress-strain behaviour and micro-structure of a granular asteroid, modelled as a cohesive granular agglomerate of spherical particles, subjected to vertical compression. Based on this, they defined a modified inertial number that relates the particle-particle tensile strength (as defined by Sánchez and Scheeres (2014)) and the overburden pressure generated inside the aggregate by gravity alone. This newly defined scaling still needs to be tested against previous results.

6.4 Post-impact reaccumulation

The evolution of fragments resulting from catastrophic disruption determines the distribution of aggregates that will reaccumulate by self-gravity. SPH plus N-body simulations indicate that the ejecta cloud shall typically collapse into multiple gravitational aggregates, while particles with higher velocities can escape from the system. Fragments with kinetic energy smaller than gravitational binding energy shall wind up in the largest remnant—which by definition has less than 50% of the original progenitors mass in a catastrophic collision. The mass distribution and angular momenta of the resulting aggregates, and how they relate to the parent body’s mass and spin, are a heritage of the impact conditions under which the parent object was disrupted. Very energetic collisions, relative to the the parent body’s binding energy, will result in a final aggregate of much lower mass, and composed of the fragments that were ejected at the lowest speeds. These are likely the largest fragments, though it still needs some further investigation. An off-centre collision will show its signature in a high-angular-momentum aggregate body. In any case, fragments of diverse shape and size will comprise the aggregated bodies, as shown in laboratory experiments of catastrophic collisions (e.g., Durda et al 2015).

Numerical simulations oriented to post-catastrophic gravitational reaccumulation were initially focused on deriving the mass and size distribution of asteroid families (Michel et al 2001, 2015). They showed that larger family members are likely reaccumulation products rather than discrete competent fragments. Gravitational aggregates obtained as a product of less catastrophic—shattering collisions—will have voids in between fragments, largely contributing to its global (macro-)porosity. An aggregate porosity is the result of the particular packing of a polydisperse collection of its components. Recently, due to the influence of granular dynamics, granular packing configurations have been investigated both

experimentally and numerically—including studies in different gravitational environments. Attempts to understand the observed global shapes and structures of aggregates through DEM numerical simulation have been performed by several authors (Tanga et al 2009a; Comito 2012; Michel and Richardson 2013; Schwartz et al 2018; Campo Bagatin et al 2018a). Originally, Farinella et al (1981) suggested that such a reaccumulation process would produce elongated tri-axial asteroids, following equilibrium figures of incompressible fluids. This is however not strictly the case for bodies smaller than $\sim 100 - 200$ km in diameter, where shape can significantly depart from a fluid-equilibrium figure, thanks to friction and sustained shear stresses (Holsapple 2004, 2007). However, Tanga et al (2009b) showed that external mechanisms (e.g., low-energy impacts) can gradually reshape the bulk of the body, pushing it toward a minimum-energy state while remaining compatible with observed shapes and spins. Nevertheless, a careful examination of the equilibrium figures show that these are very loosely defined by flat minima in the energy potential of the self-gravitating, rotating body (Tanga et al 2009a). As such, tri-axial ellipsoids significantly far from equilibrium are possible with minimal strength. Moreover, Richardson et al (2009) analysed resulting shapes from re-accumulation and rotational disruption, including variable material strength/cohesion and irregular pieces (modelled as ‘bonded aggregates’) in the DEM `pkdgrav` code (Stadel 2001; Richardson et al 2000; Schwartz et al 2012). Other simulations have been performed to describe the outcome of a catastrophic disruption on a gravitational aggregate. Michel and Richardson (2013) show that the general shape of an asteroid like Itokawa, together with the presence of boulders on its surface, can be the natural result of the reaccumulation process, given some specific material parameters for the aggregate (strength, bouncing coefficient, ...). Ballouz et al (2015) showed the influence on the catastrophic disruption threshold when taking into account the initial spin of the parent body, and some frictional effect and shear strength; but they didn’t consider material fragmentation, heating and compaction from hyper-velocity impacts. Recent numerical simulations of the reaccumulation process by Campo Bagatin et al (2018b,a) were based on such `pkdgrav` code. Using irregular rigid fragments instead of spherical individual particles, they study the packing fraction and the shapes of the final aggregates. They find that the bulk macro-porosity of aggregate asteroids can be related to the mass fraction of the largest component of the aggregate structure (Campo Bagatin et al 2018b). This in turn may be related to the specific energy of the collision that formed the aggregate itself. Then, some relationship between the macro-porosity of an asteroid and the kind of event that produced the object itself may be assessed. Studying the shapes of the aggregates formed at the end of the reaccumulation process, Campo Bagatin et al (2018b) find a general mechanism to explain some asteroid shapes, in particular bilobated (‘contact binary’) asteroids.

The reaccumulation process, while relatively short on astronomical times scales (from a few hours to several days, depending on total mass and collision boundary conditions), can be complex and result in phenomena like shaking and segregation as seen above, re-arrangements of grains and blocks and compaction (Ben-Naim et al 1998; Patrick et al 2005; Yu et al 2017) that can be difficult to model given uncertainties in the initial conditions of the aggregate cloud. However, numerical simulations have shown at least that bilobated shapes and satellites can be formed from continuous mass shedding (Walsh and Richardson 2008; Walsh et al

2012), from fission, or—over a shorter time-scale—from the reaccumulation process (Tanga et al 2013; Schwartz et al 2018; Campo Bagatin et al 2018a).

7 Discussions and prospective

From scientific point of view, surveys and space missions will allow us to address many of the topics and questions raised here. As an example LSST and Gaia will bring deeper knowledge and insights on a much larger number of solar system objects. The Gaia mission, now that the second catalogue has been released (Gaia-Collaboration et al 2018), is promising to gather valuable data of physical and dynamical properties for hundreds of thousands of small bodies (Hestroffer et al 2010); and LSST will extend this to fainter objects, with order of magnitudes in the number of objects observed compared to the present situation (Jones et al 2009; Schwamb et al 2018). Possibly the first thing we can learn from on-going and new space missions is that their success highly depends on our ability to understand their targets, small asteroids and comets, as self-gravitating granular systems. JAXA Hayabusa and Hayabusa-2 missions, as well as NASA OSIRIS-REx, Rosetta and DART missions have already had to deal with the complexities of these systems due to their design. The Hayabusa mission had as one of its objectives to shoot a small pellet to the surface of the asteroid Itokawa and capture the ejecta material to bring it back to Earth as a sample. Unfortunately the system failed and the sample canister was sealed without the pellet being fired. At its return to Earth it was found that in spite of that, some of the dust of the surface had been luckily collected. The Hayabusa-2 mission, in its way to asteroid Ryugu, will attempt to land a pod on the surface of the asteroid. This manoeuvre has encompassed a very detailed study of the interaction of the landing pod with the surface of the asteroid of which almost nothing is known. The usual characterizing parameters such as porosity, angle of friction, strength or even surface density, or boulder abundance, or topography are not known and the science team has had to work with a large range of parameters to make sure of their success. Something similar can be said about the OSIRIS-REx mission which main science objective is to return a sample of asteroid Bennu to Earth for its subsequent analysis. The Touch-and-Go-Sample-Acquisition-Mechanism (TAGSAM) had to go through a thorough design and test process that involved not only experimentation, but also two teams carrying out numerical simulations to make sure that, regardless of the variations in the characteristics of the surface of the asteroid, the sample would be acquired and sealed safely. The Double Asteroid Redirection Test (DART, Cheng et al 2016) mission is the only one that has not been launched yet, and much of the work of the different working groups that form the science team has been focused on the geophysical characteristics of Didymos. Hopefully the complementary HERA/AIDA mission will be able to complete the picture (Michel et al 2018). These missions will give more detailed information on the surface, not only from remote imaging, but also from a gentle touch down and physical interactions. They will also show how the surface and body reacts to high speed impacts, cratering, ejecta generating, and transfer of linear momentum. We still need to learn more for future applications, on the possible anchoring processes, bouncing at the surface, and how the regolith can react to different tools for exploration or excavation, depending for instance on its compaction or particle size distribution.

Unfortunately, the search for knowledge is not the only reason to obtain a better understanding about asteroid Geophysics. A precise understanding of the composition and mechanics of asteroids is essential to guarantee that humankind can protect itself from potential impactors and other hazardous asteroids. Whether an asteroid produces a blast in the atmosphere or an impact on the ground depends—for a given size or mass—on the asteroid global properties. Deflection demonstration missions can help resolve some of the challenges encountered in asteroid deflection by providing ground truths. The DART mission, for instance, is a NASA mission concept intended to demonstrate the change of the state of motion of an asteroid through a kinetic impact. Targeting the moonlet of the near-Earth asteroid (65803) Didymos, DART would alter the orbital period of the binary asteroid system. Measurements of this alteration would then allow us to draw conclusions on the composition and dynamics of the binary asteroid system on the one hand and kinetic impactor-based deflection techniques, on the other hand. The European space agency (ESA) is currently investigating a concept for an observer spacecraft, that would allow for a more detailed in situ assessment of the effects of the DART impact. Another deflection demonstration mission concept named NEOTwIST has been developed in the framework of EUs NEOShield project (Harris et al 2013; Drube et al 2016). NEOTwIST would use a kinetic impactor to spin up the well characterized asteroid (25143) Itokawa (Eggl et al 2016). The resulting change in the spin state, would provide clues as to the magnitude and direction of the momentum enhancement vector. The data acquired during deflection tests would be invaluable to gauge simulations and structure models that will be used to predict the outcomes of future deflection mission.

Regardless of the final application of acquired knowledge, furthering our understanding of the interior, internal structure and evolution of asteroids is needed. This understanding can be obtained through the determination of their gravity fields, tomography, and through seismic experiments and such experiments can only be obtained from future space missions. On the other hand, laboratory experiments need to approach the conditions of low gravity, vacuum, without specific confinement, particles size dispersion, etc., in order to better understand and model the many phenomena at play. Segregation, clustering, and internal cohesion are some of the mechanism and properties that need further studies. Numerical simulations have reached a high level of fidelity to model Earth based experiments and can predict phenomenon on low-gravity surfaces as well as on self-gravitating bodies, but would greatly benefit from experimental validation.

Having said all this, there are definitely a few things that we have been able to learn from the successes and failures in these missions: first off, that small asteroids are not only monoliths with bare surfaces. If we start with that, other things have come to light as a consequence. As the gravitational fields in asteroids are very weak, cohesive, adhesive and electrostatic forces are as, if not, more important than particle weight. This facilitates the formation of very porous interiors. As asteroids can sustain shear stress, their shapes are not going to be regular. Due to their formation and evolution process, they are likely to be formed by particles with sizes that range from microns to tens of meters. They are indeed affected by the YORP effect and so, apart from the collisional evolution, we need to take into account their rotational evolution, strength, internal heterogeneities and individual shapes and surface features if we really want to understand them.

We have seen that our objects of study span a large space of physical and dynamical parameters, with orders of magnitudes in size and mass range, with orders of magnitude in spin-rate range, different taxonomic class which could mean different constituent materials, large difference in their bulk densities, different distance from the Sun and different thermal environments. Each asteroid is indeed a small world that can provide much insight about asteroids as granular systems, but not as much as to paint a complete picture. Space missions and observations of specific targets remain circumstantial which means that theoretical, experimental and modelling efforts are needed to bridge the knowledge gap.

All these examples show the need for a great interdisciplinary effort. Really, it is not only Granular Dynamics, Soil Mechanics or Aerospace Engineering, but all of them together that have some of the tools to study Small Solar System Bodies, and only a true interdisciplinary effort will bring further understanding.

Notes added in proof

HAYABUSA-2.

OSIRIS-REx. In late 2018 the OSIRIS-REx spacecraft arrived at asteroid (101955) Bennu and began to characterize that body (Lauretta et al. 2019, Nature 568, 55). The overall bulk density of the asteroid was determined to be 1190 kg/m^3 , and its macroporosity was estimated to be up to 50%, consistent with the body being a rubble pile (Scheeres et al. 2019, Nature astronomy 3, 352). The surface of the body was seen to be uniformly covered by boulders across a large size distribution, with very few regions covered by finer regolith (Walsh et al. 2019, Nature Geoscience 12, 242). At a morphological level, the asteroid Bennu has many similarities with the asteroid Ryugu (Watanabe et al. 2019, Science 263, 268), target of the Hayabusa2 mission, and thus future comparisons between the bodies will be of great interest.

Acknowledgements This work is a direct result of support by the International Space Science Institute, ISSI Bern, Switzerland, through the hosting and provision of financial support for the international team “Asteroids & Self Gravitating Bodies as Granular Systems” led by DH. The authors would like to thank the ISSI Institute and staff for their support, and the Paris observatory for financial support. Thanks to MIAPP, Munich Institute for Astro and Particle Physics of the DFG cluster of excellence “Origin and Structure of the Universe” and participants of the the workshop on NEOS for fruitful discussions. EO thanks Prodex (Belpo) and ESA (Topical Team No. 4000103461) for financial support. DCR was supported in part by NASA grant NNX15AH90G awarded by the Solar System Workings program.

We are grateful to all the other members of the ISSI international team for discussions, exchanges, inputs, and contributions. We are grateful to Brian Warner for kindly providing us an up-to-date ‘spin-rate versus diameter’ figure. Special acknowledgment to the editor F. Schultz for his patience.

This work has made use of Wm R. Johnston archive data <http://www.johnstonsarchive.net>, and intensive use of NASA’s Astrophysics Data System.

References

- Abe S, Mukai T, Hirata N, Barnouin-Jha OS, Cheng AF, Demura H, Gaskell RW, Hashimoto T, Hiraoka K, Honda T, et al (2006) Mass and local topography measurements of Itokawa by Hayabusa. Science 312(5778):1344–1347

- Agnolin I, Roux JN (2007a) Internal states of model isotropic granular packings. I. Assembling process, geometry, and contact networks. *Phys Rev E Stat Nonlin Soft Matter Phys* 76(6-1):061,302
- Agnolin I, Roux JN (2007b) Internal states of model isotropic granular packings. III. Elastic properties. *Phys Rev E Stat Nonlin Soft Matter Phys* 76:061,304
- Ahrens TJ, Harris AW (1992) Deflection and fragmentation of near-Earth asteroids. *Nature* 360(6403):429–433
- Ai J, Chen JF, Rotter JM, Ooi JY (2011) Assessment of rolling resistance models in discrete element simulations. *Powder Technology* 206(3):269–282, DOI 10.1016/j.powtec.2010.09.030
- Alder BJ, Wainwright TE (1959) Studies in Molecular Dynamics. I. General Method. *The Journal of Chemical Physics* 31(2):459–466, DOI 10.1063/1.1730376
- Alder BJ, Wainwright TE (1960) Studies in Molecular Dynamics. II. Behavior of a Small Number of Elastic Spheres. *The Journal of Chemical Physics* 33(5):1439–1451, DOI 10.1063/1.1731425
- Andert T, Rosenblatt P, Pätzold M, Häusler B, Dehant V, Tyler G, Marty J (2010) Precise mass determination and the nature of Phobos. *Geophysical Research Letters* 37(9)
- Andreotti B, Forterre Y, Pouliquen O (2013) *Granular media: between fluid and solid*. Cambridge University Press
- Aranson IS, Tsimring LS (2006) Patterns and collective behavior in granular media: Theoretical concepts. *Review of Modern Physics* 78:641
- Asphaug E (2009) Growth and evolution of asteroids. *Annual Review of Earth and Planetary Sciences* 37:413–448
- Asphaug E, Ostro SJ, Hudson R, Scheeres DJ, Benz W (1998) Disruption of kilometre-sized asteroids by energetic collisions. *Nature* 393(6684):437
- Asphaug E, Ryan EV, Zuber MT (2002) Asteroid interiors. In: Bottke W, Cellino A, Paolicchi P, Binzel R (eds) *Asteroids III*, pp 463–484
- Azéma E, Estrada N, Preechawuttipong I, Delenne JY, Radjai F (2017) Systematic description of the effect of particle shape on the strength properties of granular media. In: *EPJ Web of Conferences*, EDP Sciences, vol 140, p 06026
- Azéma E, Sánchez P, Scheeres DJ (2018) Scaling behavior of cohesive self-gravitating aggregates. *Phys Rev E* 98:030,901, DOI 10.1103/PhysRevE.98.030901, URL <https://link.aps.org/doi/10.1103/PhysRevE.98.030901>
- Baer J, Chesley SR (2017) Simultaneous mass determination for gravitationally coupled asteroids. *The Astronomical Journal* 154(2):76
- Bagatin AC, Petit JM, Farinella P (2001) How many rubble piles are in the asteroid belt? *Icarus* 149(1):198–209
- Ballouz R (2017) Numerical simulations of granular physics in the solar system. PhD thesis, University of Maryland, College Park
- Ballouz RL, Richardson DC, Michel P, Schwartz SR, Yu Y (2015) Numerical simulations of collisional disruption of rotating gravitational aggregates: Dependence on material properties. *Planetary and Space Science* 107:29–35
- Bancelin D, Pilat-Lohinger E, Maindl TI, Ragossnig F, Schäfer C (2017) The influence of orbital resonances on the water transport to objects in the circumprimary habitable zone of binary star systems. *The Astronomical Journal* 153(6):269
- Bardyn A, Baklouti D, Cottin H, Fray N, Briois C, Paquette J, Stenzel O, Engrand C, Fischer H, Hornung K, et al (2017) Carbon-rich dust in comet 67P/Churyumov-Gerasimenko measured by COSIMA/Rosetta. *Monthly Notices of the Royal Astronomical Society* 469(Suppl.2):S712–S722
- Barnes J, Hut P (1986) A hierarchical $O(N \log N)$ force-calculation algorithm. *Nature* 324:446–449, DOI 10.1038/324446a0
- Belskaya I, Cellino A, Gil-Hutton R, Muinonen K, Shkuratov Y (2015) Asteroid polarimetry. In: Michel P, DeMeo F, Bottke W (eds) *Asteroids IV*, University of Arizona Press, Tucson, AZ, pp 151–163
- Ben-Naim E, Knight J, Nowak E, Jaeger H, Nagel S (1998) Slow relaxation in granular compaction. *Physica D: Nonlinear Phenomena* 123(1-4):380–385
- Benner LAM, Busch MW, Giorgini JD, Taylor PA, Margot JL (2015) Radar Observations of Near-Earth and Main-Belt Asteroids. In: *Asteroids IV* (P Michel, F DeMeo, W Bottke, eds) pp 165–182, DOI 10.2458/azu_uapress_9780816532131-ch009

- Biele J, Kessler L, Grimm CD, Schröder S, Mierheim O, Lange M, Ho TM (2017) Experimental Determination of the Structural Coefficient of Restitution of a Bouncing Asteroid Lander. ArXiv e-prints 1705.00701
- Binzel RP, Gehrels T, Matthews MS (eds) (1989) Asteroids II. Proceedings of the Conference, Tucson, AZ, Mar. 8–11, 1988
- Binzel RP, Morbidelli A, Merouane S, DeMeo FE, Birlan M, Vernazza P, Thomas CA, Rivkin AS, Bus SJ, Tokunaga AT (2010) Earth encounters as the origin of fresh surfaces on near-Earth asteroids. *Nature* 463(7279):331
- Board SS, Council NR, et al (2010) Defending planet earth: Near-Earth-Object surveys and hazard mitigation strategies. National Academies Press
- Bockelée-Morvan D, Gil-Hutton R, Hestroffer D, Belskaya IN, Davidsson BJ, Dotto E, Fitzsimmons A, Kawakita H, Mothe-Diniz T, Licandro J, et al (2016) Division F Commission 15: Physical Study of Comets and Minor Planets. Proceedings of the International Astronomical Union 11(T29A):316–339
- Bottke WF Jr, Cellino A, Paolicchi P, Binzel RP (eds) (2002) Asteroids III
- Bottke WF Jr, Durda DD, Nesvorný D, Jedicke R, Morbidelli A, Vokrouhlický D, Levison H (2005) The fossilized size distribution of the main asteroid belt. *Icarus* 175(1):111–140
- Bouzid M, Izzet A, Trulsson M, Clément E, Claudin P, Andreotti B (2015) Non-local rheology in dense granular flows. *The European Physical Journal E* 38(11):125
- Braga-Ribas F, Sicardy B, Ortiz J, Snodgrass C, Roques F, Vieira-Martins R, Camargo J, Assafin M, Duffard R, Jehin E, et al (2014) A ring system detected around the Centaur (10199) Chariklo. *Nature* 508(7494):72
- Brian Dade W, Huppert HE (1998) Long-runout rockfalls. *Geology* 26(9):803–806
- Brilliantov NV, Pöschel T (2010) Kinetic theory of granular gases. Oxford University Press
- Brilliantov NV, Formella A, Pöschel T (2018) Increasing temperature of cooling granular gases. *Nature Communications* 9:797
- Brisset J, Heißelmann D, Kothe S, Weidling R, Blum J (2016) Submillimetre-sized dust aggregate collision and growth properties-experimental study of a multi-particle system on a suborbital rocket. *Astron Astrophys* 593:A3
- Brisset J, Heißelmann D, Kothe S, Weidling R, Blum J (2017) Low-velocity collision behaviour of clusters composed of sub-millimetre sized dust aggregates. *Astron Astrophys* 603:A66
- Britt DT, Consolmagno GJ (2001) Modeling the structure of high porosity asteroids. *Icarus* 152(1):134–139
- Britt DT, Yeomans D, Housen K, Consolmagno G (2002) Asteroid density, porosity, and structure. In: Bottke W, Cellino A, Paolicchi P, Binzel R (eds) Asteroids III, pp 485–500
- Burtally N, King PJ, Swift MR (2002) Spontaneous air-driven separation in vertically vibrated fine granular mixtures. *Science* 295(5561):1877–1879, DOI 10.1126/science.1066850
- Busch MW, Ostro SJ, Benner LAM, Brozovic M, Giorgini JD, Jao JS, Scheeres DJ, Magri C, Nolan MC, Howell ES, Taylor PA, Margot JL, Brisken W (2011) Radar observations and the shape of near-Earth ASTEROID 2008 EV5. *Icarus* 212:649–660, DOI 10.1016/j.icarus.2011.01.013, 1101.3794
- Butcher JC (2016) Numerical methods for ordinary differential equations. John Wiley & Sons
- Campins H, Hargrove K, Pinilla-Alonso N, Howell ES, Kelley MS, Licandro J, Mothé-Diniz T, Fernández Y, Ziffer J (2010) Water ice and organics on the surface of the asteroid 24 Themis. *Nature* 464(7293):1320
- Campo Bagatin A, Alemañ RA, Benavidez PG, Richardson DC (2018a) Gravitational re-accumulation as the origin of most contact binaries and other small body shapes. *Icarus* submitted
- Campo Bagatin A, Alemañ RA, Benavidez PG, Richardson DC (2018b) Internal structure of asteroid gravitational aggregates. *Icarus* 302:343–359
- Cantelaube F, Bideau D (1995) Radial segregation in a 2d drum: An experimental analysis. *Europhysics Letters* 30:3
- Caps H, Michel R, Lecoq N, Vandewalle N (2003) Long lasting instabilities in granular mixtures. *Physica A* 326:313
- Carry B (2012) Density of asteroids. *Planetary and Space Science* 73(1):98–118
- Cattuto C, Marconi UMB (2004) Ordering phenomena in cooling granular mixtures. *Physical Review Letters* 92:174,502
- Chamberlin A (2018) Neo discovery statistics. URL <http://neo.jpl.nasa.gov/stats>
- Chapman C (1977) The evolution of asteroids as meteorite parent-bodies. In: IAU Colloquium 39, Cambridge University Press, vol 39, pp 265–275

- Chapman C, Veverka J, Thomas P, Klaasen K, Belton M, Harch A, McEwen A, Johnson T, Helfenstein P, Davies M, et al (1995) Discovery and physical properties of Dactyl, a satellite of asteroid 243 Ida. *Nature* 374(6525):783–785
- Cheng AF, Izenberg N, Chapman CR, Zuber MT (2002) Pondered deposits on asteroid 433 Eros. *Meteoritics and Planetary Science* 37:1095–1105, DOI 10.1111/j.1945-5100.2002.tb00880.x
- Cheng AF, Michel P, Jutzi M, Rivkin AS, Stickle A, Barnouin O, Ernst C, Atchison J, Pravec P, Richardson DC, et al (2016) Asteroid Impact & Deflection Assessment mission: Kinetic impactor. *Planetary and Space Science* 121:27–35
- Chujo T, Mori O, Kawaguchi J, Yano H (2017) Categorization of Brazil nut effect and its reverse under less-convective conditions for microgravity geology. *Monthly Notices of the Royal Astronomical Society* 474(4):4447–4459
- Cizeau P, Makse HA, Stanley HE (1999) Mechanisms of granular spontaneous stratification and segregation in two-dimensional silos. *Physical Review E* 59:4408
- Cleary PW, Sawley ML (2002) DEM modelling of industrial granular flows: 3D case studies and the effect of particle shape on hopper discharge. *Applied Mathematical Modelling* 26:89–111
- Clement E, Rajchenbach J, Duran J (1995) Mixing of a granular material in a bidimensional rotating drum. *Europhysics Letters* 30:1
- Comito C (2012) Numerical n-body approach to binary asteroid formation and evolution. PhD thesis, Università degli studi di Torino; Université Nice Sophia Antipolis
- Comito C, Thirouin A, Campo Bagatin A, Tanga P, Ortiz JL, Richardson DC (2011) Deformation and splitting of asteroids by YORP spin-up. In: EPSC-DPS Joint Meeting 2011, p 420
- Cotto-Figueroa D, Statler TS, Richardson DC, Tanga P (2015) Coupled spin and shape evolution of small rubble-pile asteroids: Self-limitation of the yorp effect. *The Astrophysical Journal* 803(1):25
- Cundall PA, Strack OD (1979) A discrete numerical model for granular assemblies. *Geotechnique* 29(1):47–65
- Daniels KE (2013) Rubble-pile near earth objects: insights from granular physics. In: Asteroids, Springer, pp 271–286
- Delbo M, Mueller M, Emery JP, Rozitis B, Capria MT (2015) Asteroid thermophysical modeling. In: Asteroids IV (P Michel, F DeMeo, W Bottke, eds) pp 107–128
- DeMeo F, Alexander C, Walsh K, Chapman C, Binzel R (2015) The compositional structure of the asteroid belt. In: Michel P, DeMeo F, Bottke W (eds) Asteroids IV, University of Arizona Press, Tucson, AZ, pp 13–41, DOI 10.2458/azu_uapress_9780816532131-ch002
- Dijksman JA, Rietz F, Lrincz KA, van Hecke M, Losert W (2012) Invited article: Refractive index matched scanning of dense granular materials. *Review of Scientific Instruments* 83(1):011,301, DOI 10.1063/1.3674173
- Dobrovolskis AR (1982) Internal stresses in Phobos and other triaxial bodies. *Icarus* 52(1):136–148
- Dohnanyi J (1969) Collisional model of asteroids and their debris. *Journal of Geophysical Research* 74(10):2531–2554
- Donev A, Cisse I, Sachs EA, Dand Variano, Stillinger FH, Connelly R, Torquato S, Chaikin PM (2004) Improving the density of jammed disordered packings using ellipsoids. *Science* 303:990–993
- Dove A, Anderson S, Gomer G, Fraser M, John K, Fries M (2018) Regolith stratification and migration in an asteroid-like environment. In: Lunar and Planetary Science Conference, Lunar and Planetary Science Conference, vol 49, p 2993
- Drube L, Harris AW, Engel K, Falke A, Johann U, Eggl S, Cano JL, vila JM, Schwartz SR, Michel P (2016) The NEOT ω IST mission (Near-Earth Object Transfer of angular momentum spin test). *Acta Astronautica* 127:103–111, DOI 10.1016/j.actaastro.2016.05.009
- Durda DD, Bagatin AC, Alemañ RA, Flynn GJ, Strait MM, Clayton AN, Patmore EB (2015) The shapes of fragments from catastrophic disruption events: Effects of target shape and impact speed. *Planetary and Space Science* 107:77–83, DOI 10.1016/j.pss.2014.10.006, {VIII} Workshop on Catastrophic Disruption in the Solar System
- Durech J, Carry B, Delbo M, Kaasalainen M, Viikinkoski M (2015) Asteroid models from multiple data sources. In: Michel P, DeMeo F, Bottke W (eds) Asteroids IV, University of Arizona Press, Tucson, AZ, pp 183–202, DOI 10.2458/azu_uapress_9780816532131-ch010

- Dziugys A, Navakas R (2009) The role of friction in mixing and segregation of granular material. *Granular Matter* 11:403–416
- Eggl S, Hestroffer D, Thuillot W, Bancelin D, Cano JL, Cichocki F (2015) Post mitigation impact risk analysis for asteroid deflection demonstration missions. *Advances in Space Research* 56:528–548, DOI 10.1016/j.asr.2015.02.030
- Eggl S, Hestroffer D, Cano JL, Ávila JM, Drube L, Harris AW, Falke A, Johann U, Engel K, Schwartz SR, Michel P (2016) Dealing with uncertainties in asteroid deflection demonstration missions: NEOT ω IST. In: Chesley SR, Morbidelli A, Jedicke R, Farnocchia D (eds) *IAU Symposium*, IAU Symposium, vol 318, pp 231–238, DOI 10.1017/S1743921315008698, 1601.02103
- Falcon E, Wunenburger R, Evesque P, Fauve S, Chabot C, Garrabos Y, Beysens D (1999) Cluster formation in a granular medium fluidized by vibrations in low gravity. *Physical Review Letters* 83:440
- Falcon E, Bacri JC, Laroche C (2013) Equation of state of a granular gas homogeneously driven by particle rotations. *Europhysical Letters* 103:64,004
- Fall A, Weber B, Pakpour M, Lenoir N, Shahidzadeh N, Fiscina J, Wagner C, Bonn D (2014) Sliding friction on wet and dry sand. *Phys Rev Lett* 112:175,502, DOI 10.1103/PhysRevLett.112.175502
- Fan Y, Boukerkour Y, Blanc T, Umbanhowar P, Ottino JM, Lueptow RM (2012) Stratification, segregation, and mixing of granular materials in quasi-two-dimensional bounded heaps. *Physical Review E* 86:051,305
- Faraday M (1831) On a peculiar class of acoustical figures; and on certain forms assumed by groups of particles upon vibrating elastic surfaces. *Philosophical Transactions of the Royal Society of London* 121:299–340, URL <http://www.jstor.org/stable/107936>
- Farinella P, Paolicchi P, Tedesco E, Zappala V (1981) Triaxial equilibrium ellipsoids among the asteroids? *Icarus* 46(1):114–123
- Félix G, Thomas N (2004) Relation between dry granular flow regimes and morphology of deposits: formation of levées in pyroclastic deposits. *Earth and Planetary Science Letters* 221(1-4):197–213
- Ferrellec JF, McDowell GR (2010) A method to model realistic particle shape and inertia in DEM. *Granular Matter* 12:459–467
- Festou M, Keller HU, Weaver HA (2004) *Comets II*. University of Arizona Press
- Finger T, von Rűling F, Lévay S, Szabó B, Börzsőnyi T, Stannarius R (2016) Segregation of granular mixtures in a spherical tumbler. *Physical Review E* 93:032,903
- Fischer D, Finger T, Angenstein F, Stannarius R (2009) Diffusive and subdiffusive axial transport of granular material in rotating mixers. *Physical Review E* 80:061,302
- Fries M, Abell P, Brisset J, Britt D, Colwell J, Durda D, Dove A, Graham L, Hartzell C, John K, Leonard M, Love S, Sánchez DP, Scheeres DJ (2016) Strata-1: An International Space Station Experiment into Fundamental Regolith Properties in Microgravity. In: *Lunar and Planetary Science Conference, Lunar and Planetary Science Conference*, vol 47, p 2799
- Fries M, Abell P, Brisset J, Britt D, Colwell J, Dove A, Durda D, Graham L, Hartzell C, Hrovat K, John K, Karrer D, Leonard M, Love S, Morgan J, Poppin J, Rodriguez V, Sánchez-Lana P, Scheeres D, Whizin A (2018) The Strata-1 experiment on small body regolith segregation. *Acta Astronautica* 142:87 – 94, DOI 10.1016/j.actaastro.2017.10.025
- Fujiwara A, Kawaguchi J, Yeomans D, Abe M, Mukai T, Okada T, Saito J, Yano H, Yoshikawa M, Scheeres D, et al (2006) The rubble-pile asteroid Itokawa as observed by Hayabusa. *Science* 312(5778):1330–1334
- Gaia-Collaboration, Spoto F, Tanga P, Mignard F, Berthier J, Carry B, Cellino A (2018) Gaia Data Release 2: Observations of solar system objects. *Astronomy and Astrophysics* in press, DOI 10.1051/0004-6361/201832900
- Galache J, Graps A, Asime 2016 Contributors (2017) ASIME 2016 white paper: Answers to questions from the asteroid miners. In: *European Planetary Science Congress*, vol 11, p 985
- Gao Z, Zhao J, Li XS, Dafalias YF (2014) A critical state sand plasticity model accounting for fabric evolution. *Int J Numer Anal Meth Geomech* 38:370–390
- Garcimartin A, Maza D, Iquimiche JL, Zuriguel I (2002) Convective motion in a vibrated granular layer. *Physical Review E* 65:031,303
- Gehrels T, Matthews MS (eds) (1979) *Asteroids*
- Gehrels T, Drummond J, Levenson N (1987) The absence of satellites of asteroids. *Icarus* 70(2):257–263

- Godoy S, Risso D, Soto R, Cordero P (2008) Rise of a Brazil nut: A transition line. *Physical Review E* 78:031,301
- Goldhirsch I, Zanetti G (1993) Clustering instability in dissipative gases. *Physical Review Letters* 70:1619
- Goldreich P, Tremaine S (1978) The velocity dispersion in Saturn's rings. *Icarus* 34(2):227–239
- Gray JNMT, Ancey C (2011) Multi-component particle size-segregation in shallow granular avalanches. *Journal of Fluid Mechanics* 678:535–558
- Gray JNMT, Thornton AR (2005) A theory for particle size segregation in shallow granular free-surface flows. *Proceedings of the Royal Society of London A* 461:1447
- Greengard L, Rokhlin V (1987) A fast algorithm for particle simulations. *Journal of Computational Physics* 73(2):325–348, DOI 10.1016/0021-9991(87)90140-9
- Guillard F, Forterre Y, Pouliquen O (2014) Lift forces in granular media. *Physics of Fluids* 26(4):043,301
- Gundlach B, Blum G (2013) A new method to determine the grain size of planetary regolith. *Icarus* 223:479–492
- Güttler C, von Borstel I, Schräpler R, Blum J (2013) Granular convection and the Brazil nut effect in reduced gravity. *Physical Review E* 87:044,201
- Haff P (1983) Grain flow as a fluid-mechanical phenomenon. *Journal of Fluid Mechanics* 134:401–430
- Hajra SK, Khakhar DV (2011) Radial segregation of ternary granular mixtures in rotating cylinders. *Granular Matter* 13(4):475–486
- Harrington M, Weijs JH, Losert W (2013) Suppression and emergence of granular segregation under cyclic shear. *Phys Rev Lett* 111:078,001, DOI 10.1103/PhysRevLett.111.078001
- Harrington M, Lin M, Nordstrom KN, Losert W (2014) Experimental measurements of orientation and rotation of dense 3D packings of spheres. *Granular Matter* 16(2):185–191, DOI 10.1007/s10035-013-0474-0
- Harris A, Barucci M, Cano J, Fitzsimmons A, Fulchignoni M, Green S, Hestroffer D, Lappas V, Lork W, Michel P, et al (2013) The European Union funded NEOShield project: a global approach to near-Earth object impact threat mitigation. *Acta Astronautica* 90(1):80–84
- Harris AW (1996) The Rotation Rates of Very Small Asteroids: Evidence for 'Rubble Pile' Structure. In: *Lunar and Planetary Science Conference, Lunar and Planetary Science Conference*, vol 27
- Harris AW, Lagerros JS (2002) Asteroids in the thermal infrared. In: Bottke W, Cellino A, Paolicchi P, Binzel R (eds) *Asteroids III*, University of Arizona Press, Tucson, AZ
- Harris AW, Fahnestock EG, Pravec P (2009) On the shapes and spins of 'rubble pile' asteroids. *Icarus* 199(2):310–318
- Harth K, Kornek U, Trittel T, Strachauer U, Hme S, Will K, Stannarius R (2013) Granular gases of rod-shaped grains in microgravity. *Physical Review Letters* 110:144,102
- Harthong B, Jérrier JF, Dorémus P, Imbault D, Donzé FV (2009) Modeling of high-density compaction of granular materials by the discrete element method. *International Journal of Solids and Structures* 46(18):3357–3364
- Hartzell C, Carter D (2017) Electrostatic forces on grains near asteroids and comets. In: *EPJ Web of Conferences*, EDP Sciences, vol 140, p 14009
- Heisselmann D, Blum J, Fraser HJ, Wolling K (2010) Microgravity experiments on the collisional behavior of Saturnian ring particles. *Icarus* 206:424–430
- Henych T, Holsapple KA (2018) Interpretations of family size distributions: The Datura example. *Icarus* 304:127–134, DOI 10.1016/j.icarus.2017.05.018, asteroids and Space Debris
- Hérique A, Agnus B, Asphaug E, Barucci A, Beck P, Bellerose J, Biele J, Bonal L, Bousquet P, Bruzzone L, et al (2018) Direct observations of asteroid interior and regolith structure: Science measurement requirements. *Advances in Space Research* in press
- Hernquist L (1987) Performance characteristics of tree codes. *Astrophys J Suppl Ser* 64:715–734, DOI 10.1086/191215
- Herrmann H, Luding S (1998) Modeling granular media on the computer. *Continuum Mechanics and Thermodynamics* 10:189–231, DOI 10.1007/s001610050089
- Hestroffer D (1998) Photocentre displacement of minor planets: analysis of HIPPARCOS astrometry. *Astron Astrophys* 336:776–781
- Hestroffer D, DellOro A, Cellino A, Tanga P (2010) The Gaia mission and the asteroids. In: *Dynamics of Small Solar System Bodies and Exoplanets*, Springer, pp 251–340
- Hestroffer D, Agnan M, Segret B, Quinsac G, Vannitsen J, Rosenblatt P, Miau JJ (2017) BIRDY - Interplanetary CubeSat for planetary geodesy of Small Solar System Bodies

- (SSSB). AGU Fall Meeting Abstracts
- Hestroffer D, Bagatín AC, Losert W, Opsomer E, Sánchez P, Scheeres DJ, Staron L, Taberlet N, Yano H, Eggl S, et al (2017) Small solar system bodies as granular systems. In: EPJ Web of Conferences, EDP Sciences, vol 140, p 14011
- Hill KM, Gioia G, Amaravadi D (2004) Radial segregation patterns in rotating granular mixtures: Waviness selection. *Physical Review Letters* 93:224,301
- Hilton JL (2002) Asteroid masses and densities. In: *Asteroids III*, pp 103–112
- Hirabayashi M (2014) Structural failure of two-density-layer cohesionless biaxial ellipsoids. *Icarus* 236(0):178–180, DOI 10.1016/j.icarus.2014.02.024
- Hirabayashi M, Scheeres DJ (2014) Analysis of Asteroid (216) Kleopatra Using Dynamical and Structural Constraints. *The Astrophysical Journal* 780(2):160
- Hirabayashi M, Scheeres DJ (2015) Stress and Failure Analysis of Rapidly Rotating Asteroid (29075) 1950DA. *The Astrophysical Journal Letters* 798(1):L8
- Hirabayashi M, Scheeres DJ, Sánchez DP, Gabriel T (2014) Constraints on the Physical Properties of Main Belt Comet P/2013 R3 from its Breakup Event. *The Astrophysical Journal Letters* 789(1):L12
- Hirabayashi M, Sánchez DP, Scheeres DJ (2015) Internal structure of asteroids having surface shedding due to rotational instability. *The Astrophysical Journal* 808(1):63
- Hirabayashi M, Scheeres DJ, Chesley SR, Marchi S, McMahon JW, Steckloff J, Mottola S, Naidu SP, Bowling T (2016) Fission and reconfiguration of bilobate comets as revealed by 67P/Churyumov–Gerasimenko. *Nature* 534(7607):352
- Hockney RW, Eastwood JW (1988) *Computer simulation using particles*. crc Press
- Holsapple K, Giblin I, Housen K, Nakamura A, Ryan E (2002) Asteroid Impacts: Laboratory Experiments and Scaling Laws. In: Bottke W, Cellino A, Paolicchi P, Binzel R (eds) *Asteroids III*, pp 443–462
- Holsapple KA (2001) Equilibrium configurations of solid cohesionless bodies. *Icarus* 154(2):432–448, DOI 10.1006/icar.2001.6683
- Holsapple KA (2004) Equilibrium figures of spinning bodies with self-gravity. *Icarus* 172:272–303, DOI 10.1016/j.icarus.2004.05.023
- Holsapple KA (2007) Spin limits of Solar System bodies: From the small fast-rotators to 2003 EL61. *Icarus* 187(2):500–509, DOI 10.1016/j.icarus.2006.08.012
- Hong DC, Quinn PV, Luding S (2001) Reverse brazil nut problem: Competition between percolation and condensation. *Physical Review Letters* 86:15
- Hou M, Liu R, Zhai G, Sun Z, Lu K, Garrabos Y, Evesque P (2008) Velocity distribution of vibration-driven granular gas in Knudsen regime in microgravity. *Microgravity Science and Technology* 20:73–80
- Hsieh HH, Jewitt D (2006) A population of comets in the main asteroid belt. *Science* 312(5773):561–563
- van der Hucht KA (2008) *Proceedings of the Twenty Sixth General Assembly Prague 2006: Transactions of the International Astronomical Union XXVIB*, vol 26. Cambridge University Press
- Huerta DA, Ruiz-Suarez JC (2004) Vibration-induced granular segregation: A phenomenon driven by three mechanisms. *Physical Review Letters* 92:11
- Hut P, Makino J, McMillan S (1995) Building a better leapfrog. *The Astrophysical Journal* 443:L93–L96
- Jacobson R, Spitale J, Porco C, Beurle K, Cooper N, Evans M, Murray C (2007) Revised orbits of Saturn’s small inner satellites. *The Astronomical Journal* 135(1):261
- Jacobson SA, Scheeres DJ (2011) Dynamics of rotationally fissioned asteroids: Source of observed small asteroid systems. *Icarus* 214:161–178, DOI 10.1016/j.icarus.2011.04.009
- Jaeger HM, Nagel SR, Behringer RP (1996a) Granular solids, liquids, and gases. *Rev Mod Phys* 68:1259–1273, DOI {10.1103/RevModPhys.68.1259}
- Jaeger HM, Nagel SR, Behringer RP (1996b) The physics of granular materials. *Physics Today* 49(4):32–38
- Jean M (1999) The non-smooth contact dynamics method. *Computer methods in applied mechanics and engineering* 177(3-4):235–257
- Jenkins J, Richman M (1985) Kinetic theory for plane flows of a dense gas of identical, rough, inelastic, circular disks. *The Physics of Fluids* 28(12):3485–3494
- Jenkins JT, Savage SB (1983) A theory for the rapid flow of identical, smooth, nearly elastic, spherical particles. *Journal of fluid mechanics* 130:187–202

- Jewitt D (2009) The Active Centaurs. *The Astronomical Journal* 137:4296–4312, DOI 10.1088/0004-6256/137/5/4296
- Jewitt D, Agarwal J, Weaver H, Mutchler M, Larson S (2013) The Extraordinary Multi-tailed Main-belt Comet P/2013 P5. *The Astrophysical Journal Letters* 778(1):L21
- Jewitt D, Agarwal J, Li J, Weaver H, Mutchler M, Larson S (2014) Disintegrating Asteroid P/2013 R3. *The Astrophysical Journal Letters* 784(1):L8
- Jewitt D, Hsieh H, Agarwal J (2015) The Active Asteroids. In: Michel P, DeMeo FE, Bottke WF (eds) *Asteroids IV*, pp 221–241, DOI 10.2458/azu_uapress_9780816532131-ch012
- John KK, Saucedo VL, Fisher KR, Fries MD, Dove AR, Leonard MJ, Graham LD, Abell PA (2018) Hermes Microgravity Research Facility on the ISS. In: *Lunar and Planetary Science Conference, Lunar and Planetary Science Conference*, vol 49, p 1790
- Jones R, Chesley S, Connolly A, Harris A, Ivezić Z, Knezevic Z, Kubica J, Milani A, Trilling D, Collaboration LSST, et al (2009) *Solar System science with LSST. Earth, Moon, and Planets* 105(2-4):101–105
- Jop P, Forterre Y, Pouliquen O (2006) A constitutive law for dense granular flows. *Nature* 441(7094):727
- Jutzi M, Michel P (2014) Hypervelocity impacts on asteroids and momentum transfer I. Numerical simulations using porous targets. *Icarus* 229:247–253
- Jutzi M, Michel P, Benz W, Richardson DC (2010) Fragment properties at the catastrophic disruption threshold: The effect of the parent body’s internal structure. *Icarus* 207(1):54–65, DOI 10.1016/j.icarus.2009.11.016
- Kaasalainen M, Torppa J (2001) Optimization Methods for Asteroid Lightcurve Inversion. I. Shape Determination. *Icarus* 153:24–36
- Kaasalainen M, Torppa J, Muinonen K (2001) Optimization Methods for Asteroid Lightcurve Inversion. II. The Complete Inverse Problem. *Icarus* 153:37–51
- Kaasalainen M, Mottola S, Fulchignoni M (2002) Asteroid models from disk-integrated data. In: Bottke W, Cellino A, Paolicchi P, Binzel R (eds) *Asteroids III*, University of Arizona Press, Tucson, AZ, pp 139–150
- Kay JP, Dombard AJ (2018) Formation of the bulge of Iapetus through long-wavelength folding of the lithosphere. *Icarus* 302:237–244
- Kelfoun K (2011) Suitability of simple rheological laws for the numerical simulation of dense pyroclastic flows and long-runout volcanic avalanches. *Journal of Geophysical Research: Solid Earth* 116(B8)
- Kerr RA (1985) Could an asteroid be a comet in disguise; two asteroids of the inner solar system are strong candidates for once-active comets that now masquerade as inert hunks of rock. *Science* 227:930–932
- Khakhar DV, McCarthy J, Shinbrot T, Ottino JM (1997) Radial segregation of granular mixtures in rotating cylinders. *Physics of Fluids* 9:3600
- Khakhar DV, Orpe VA, Hajra SK (2003) Segregation of granular materials in rotating cylinders. *Physica A* 318:126
- Kim T, Nam J, Yun J, Lee K, You S (2009) Relationship between cohesion and tensile strength in wet sand at low normal stresses. In: *Proc. of 17th Intern. Conf. on Soil Mechanics and Geotechnical Engineering*, Olexandria, 2009, JOS Press, Amsterdam, Berlin, Tokyo, Washington, vol 367, p 364
- Klypin A (2017) Methods for cosmological N-body simulations. URL <http://www.skiesanduniverses.org/resources/KlypinNbody.pdf>
- Knight JB, Jaeger HM, Nagel S (1993) Vibration-induced size separation in granular media: The convection connection. *Physical Review Letters* 70:24
- Koepppe JP, Enz M, Kakalios J (1998) Phase diagram for avalanche stratification of granular media. *Physical Review E* 58:R4104
- Kok JF, Parteli EJR, Michaels TI, Karam DB (2012) The physics of wind-blown sand and dust. *Reports on Progress in Physics* 75(10):106,901
- Konopliv AS, Miller JK, Owen WM, Yeomans DK, Giorgini JD, Garmier R, Barriot JP (2002) A global solution for the gravity field, rotation, landmarks, and ephemeris of Eros. *Icarus* 160(2):289–299
- Kou B, Cao Y, Li J, Xia C, Li Z, Dong H, Zhang A, Zhang J, Kob W, Wang Y (2017) Granular materials flow like complex fluids. *Nature* 551(7680):360
- Kudrolli A (2004a) Size separation in vibrated granular matter. *Reports on Progress in Physics* 67(3):209

- Kudrolli A (2004b) Size separation in vibrated granular matter. *Reports on Progress in Physics* 67:209
- Lagrée PY, Staron L, Popinet S (2011) The granular column collapse as a continuum: validity of a two-dimensional Navier–Stokes model with a μ (I)-rheology. *Journal of Fluid Mechanics* 686:378–408
- Landau LD, Lifshitz E (1986) Theory of elasticity, vol. 7. *Course of Theoretical Physics* 3:109
- Lecante M, Garrabos Y, Falcon E, Lecoutre-Chabot C, Palencia F, Evesque P, Beysens D (2006) Microgravity experiments on vibrated granular gases in a dilute regime: non-classical statistics. *Journal of Statistical Mechanics: Theory and Experiment* 7:07,012
- Lee V, Waitukaitis SR, Miskin MZ, Jaeger HM (2015) Direct observation of particle interactions and clustering in charged granular streams. *Nature Physics* 11:733–737
- Leinhardt ZM, Richardson DC, Quinn T (2000) Direct N-body simulations of rubble pile collisions. *Icarus* 146:133–151, DOI 10.1006/icar.2000.6370, astro-ph/9908221
- Levasseur-Regourd AC, Rotundi A, Bentley M, Della Corte V, Fulle M, Hadamcik E, Hilchenbach M, Hines D, Lasue J, Merouane S, et al (2015) Physical properties of dust particles in cometary comae: from clues to evidence with the Rosetta mission. In: *European Planetary Science Congress 2015*, vol 10, pp EPSC2015–932
- Li JY, Helfenstein P, Buratti B, Takir D, Clark BE (2015) Asteroid photometry. In: Michel P, DeMeo F, Bottke W (eds) *Asteroids IV*, University of Arizona Press, Tucson, AZ, pp 129–150, DOI 10.2458/azu_uapress_9780816532131-ch007
- Liu AJ, Nagel SR (1998) Nonlinear dynamics: Jamming is not just cool any more. *Nature* 396(6706):21–22
- Liu C, Nagel SR, Schecter DA, Coppersmith SN, Majumdar S, Narayan O, Witten TA (1995) Force Fluctuations in Bead Packs. *Science* 269:513
- Lohse D, Bergmann R, Mikkelsen R, Zeilstra C, van der Meer D, Versluis M, van der Weele K, van der Hoef M, Kuipers H (2004) Impact on soft sand: Void collapse and jet formation. *Phys Rev Lett* 93:198,003, DOI 10.1103/PhysRevLett.93.198003
- Losert W, Cooper DGW, Delour A, Jand Kudrolli, Gollub JP (1999) Velocity statistics in excited granular media. *Chaos* 9:682
- Louge MY, Jenkins JT, Xu H, Arnarson BO (2012) Granular segregation in collisional shearing flows. In: Aref H, Phillips JW (eds) *Mechanics for a New Millennium*
- Lu M, McDowell GR (2007) The importance of modelling ballast particle shape in the discrete element method. *Granular Matter* 9:69
- Lu XP, Cellino A, Hestroffer D, Ip WH (2016) Cellinoid shape model for hipparcos data. *Icarus* 267:24–33
- Lubachevsky BD (1991) How to simulate billiards and similar systems. *Journal of Computational Physics* 94(2):255 – 283, DOI 10.1016/0021-9991(91)90222-7
- Lucas A, Mangeney A, Ampuero JP (2014) Frictional velocity-weakening in landslides on Earth and on other planetary bodies. *Nature Communications* 5:3417
- Lucchitta BK (1979) Landslides in Valles Marineris, Mars. *Journal of Geophysical Research: Solid Earth* 84(B14):8097–8113
- Ludewig F, Vandewalle N (2012) Strong interlocking of nonconvex particles in random packings. *Physical Review E* 85:051,307
- Maaß CC, Isert N, Maret G, Aegerter CM (2008) Experimental investigation of the freely cooling granular gas. *Physical Review Letters* 100:248,001
- Makse HA (1999) Continuous avalanche segregation of granular mixtures in thin rotating drums. *Physical Review Letters* 83:3186
- Makse HA, Ball RC, Stanley HE, Warr S (1998) Dynamics of granular stratification. *Physical Review Letters* 58:3357
- Marchis F, Hestroffer D, Descamps P, Berthier J, Bouchez AH, Campbell RD, Chin JC, Van Dam MA, Hartman SK, Johansson EM, et al (2006) A low density of 0.8 g cm^{-3} for the Trojan binary asteroid 617 Patroclus. *Nature* 439(7076):565
- Margot JL, Pravec P, Taylor P, Carry B, Jacobson S (2015) Asteroid systems: binaries, triples, and pairs. In: Michel P, DeMeo F, Bottke W (eds) *Asteroids IV*, University of Arizona Press, Tucson, AZ, pp 355–374
- Masiero JR, Grav T, Mainzer AK, Nugent CR, Bauer JM, Stevenson R, Sonnett S (2014) Main-belt Asteroids with WISE/NEOWISE: Near-infrared Albedos. *The Astrophysical Journal* 791:121, DOI 10.1088/0004-637X/791/2/121, 1406.6645
- Masiero JR, Nugent C, Mainzer AK, Wright EL, Bauer JM, Cutri RM, Grav T, Kramer E, Sonnett S (2017) NEOWISE Reactivation Mission Year Three: Asteroid Diameters and

- Albedos. *The Astronomical Journal* 154(4):168
- Matsumura S, Richardson DC, Michel P, Schwartz SR, Ballouz RL (2014) The Brazil nut effect and its application to asteroids. *Monthly Notices of the Royal Astronomical Society* 443:3368–3380
- Mattson W, Rice BM (1999) Near-neighbor calculations using a modified cell-linked list method. *Computer Physics Communications* 119(2-3):135–148
- Maurel C, Ballouz RL, Richardson DC, Michel P, Schwartz SR (2017) Numerical simulations of oscillation-driven regolith motion: Brazil-nut effect. *Mon Not R Astron Soc* 464:2866–2881, DOI 10.1093/mnras/stw2641
- McCarthy DF, McCarthy DF (1977) *Essentials of soil mechanics and foundations*. Reston Publishing Company
- McMahon J, Scheeres D, Hesar S, Farnocchia D, Chesley S, Lauretta D (2018) The OSIRIS-REx radio science experiment at Bennu. *Space Science Reviews* 214(1):43
- McNamara S, Young WR (1994) Inelastic collapse in two dimensions. *Phys Rev E* 50:R28–R31, DOI 10.1103/PhysRevE.50.R28
- Meech KJ, Weryk R, Micheli M, Kleyna JT, Hainaut OR, Jedicke R, Wainscoat RJ, Chambers KC, Keane JV, Petric A, et al (2017) A brief visit from a red and extremely elongated interstellar asteroid. *Nature* 552(7685):378
- Mehta A (2012) *Granular Matter: an interdisciplinary approach*. Springer
- Merline WJ, Close L, Dumas C, Chapman C, Roddier F, Menard F, Slater D, Duvert G, Shelton C, Morgan T (1999) Discovery of a moon orbiting the asteroid 45 Eugenia. *Nature* 401(6753):565
- Merline WJ, Weidenschilling SJ, Durda DD, Margot JL, Pravec P, Storrs AD (2002) Asteroids do have satellites. In: *Asteroids III* (WF Bottke, A Cellino, P Paolicchi, RP Binzel, eds) 1:289–312
- Metzger MJ, Remy B, Glasser BJ (2011) All the Brazil nuts are not on top: Vibration induced granular size segregation of binary, ternary and multi-sized mixtures. *Powder technology* 205:42–51
- Michel P, Richardson DC (2013) Collision and gravitational reaccumulation: Possible formation mechanism of the asteroid Itokawa. *Astron Astrophys* 554:L1
- Michel P, Benz W, Tanga P, Richardson DC (2001) Collisions and gravitational reaccumulation: Forming asteroid families and satellites. *Science* 294(5547):1696–1700
- Michel P, Tanga P, Benz W, Richardson DC (2002) Formation of asteroid families by catastrophic disruption: Simulations with fragmentation and gravitational reaccumulation. *Icarus* 160:10–23, DOI 10.1006/icar.2002.6948
- Michel P, DeMeo FE, Bottke WF (eds) (2015) *Asteroids IV*. DOI 10.2458/azu_uapress_9780816532131
- Michel P, Richardson DC, Durda DD, Jutzi M, Asphaug E (2015) Collisional formation and modeling of asteroid families. In: Michel P, DeMeo F, Bottke W (eds) *Asteroids IV*, University of Arizona Press, Tucson, AZ, pp 341–354
- Michel P, Kueppers M, Sierks H, Carnelli I, Cheng AF, Mellab K, Granvik M, Kestilä A, Kohout T, Muinonen K, et al (2018) European component of the AIDA mission to a binary asteroid: Characterization and interpretation of the impact of the DART mission. *Advances in Space Research in press*
- MiDi-GDR (2004) On dense granular flows. *The European Physical Journal E* 14(4):341–365
- Miyamoto H, Yano H, Scheeres DJ, Abe S, Barnouin-Jha O, Cheng AF, Demura H, Gaskell RW, Hirata N, Ishiguro M, Michikami T, Nakamura AM, Nakamura R, Saito J, Sasaki S (2007) Regolith Migration and Sorting on Asteroid Itokawa. *Science* DOI 10.1126/science.1134390
- Möbius M, Lauderdale BE, Nagel SR, Jaeger HM (2001) Brazil-nut effect: Size separation of granular particles. *Nature* 414:270
- Morbidelli A, Chambers J, Lunine J, Petit JM, Robert F, Valsecchi G, Cyr K (2000) Source regions and timescales for the delivery of water to the Earth. *Meteoritics & Planetary Science* 35(6):1309–1320
- Moreau JJ (1994) Some numerical methods in multibody dynamics: application to granular materials. *European journal of mechanics A Solids* 13:93–114
- Mouret S, Hestroffer D, Mignard F (2007) Asteroid masses and improvement with Gaia. *Astron Astrophys* 472:1017–1027, DOI 10.1051/0004-6361:20077479
- Movshovitz N, Asphaug E, Korycansky D (2012) Numerical modeling of the disruption of comet D/1993 F2 Shoemaker-Levy 9 representing the progenitor by a gravitationally bound

- assemblage of randomly shaped polyhedra. *The Astrophysical Journal* 759(2):93
- Müller T, Marciniak A, Kiss C, Duffard R, Alí-Lagoa V, Bartczak P, Butkiewicz-Bak M, Dudziński G, Fernández-Valenzuela E, Marton G, et al (2017) Small Bodies Near and Far (SBNF): a benchmark study on physical and thermal properties of small bodies in the Solar System. *Advances in Space Research* in press
- Murdoch N, Michel P, Richardson DC, Nordstrom K, Berardi CR, Green SF, Losert W (2012) Numerical simulations of granular dynamics II: Particle dynamics in a shaken granular material. *Icarus* 219(1):321–335
- Murdoch N, Rozitis B, Green S, Lophem TL, Michel P, Losert W (2013a) Granular shear flow in varying gravitational environments. *Granular Matter* 15(2):129–137, DOI 10.1007/s10035-013-0395-y
- Murdoch N, Rozitis B, Green S, Michel P, de Lophem TL, Losert W (2013b) Simulating regoliths in microgravity. *Monthly Notices of the Royal Astronomical Society* 433(1):506–514
- Murdoch N, Rozitis B, Nordstrom K, Green S, Michel P, de Lophem T, Losert W (2013) Granular Convection in Microgravity. *Physical Review Letters* 110(1):018307, DOI 10.1103/PhysRevLett.110.018307
- Murdoch N, Sánchez P, Schwartz SR, Miyamoto H (2015) Asteroid Surface Geophysics. In: Michel P, DeMeo F, Bottke W (eds) *Asteroids IV*, University of Arizona Press, Tucson, AZ, pp 767–792, DOI 10.2458/azu_uapress_9780816532131-ch039
- Murdoch N, Cadu A, Mimoun D, Karatekin O, Garcia R, Carrasco J, Garcia de Quiros J, Vasseur H, Ritter B, Eubanks M, et al (2016) Investigating the surface and subsurface properties of the Didymos binary asteroid with a landed CubeSat. In: *EGU General Assembly Conference Abstracts*, vol 18, p 12140
- Murdoch N, Avila Martinez I, Sunday C, Zenou E, Cherrier O, Cadu A, Gourinat Y (2017a) An experimental study of low-velocity impacts into granular material in reduced gravity. *Monthly Notices of the Royal Astronomical Society* 468(2):1259–1272
- Murdoch N, Hempel S, Pou L, Cadu A, Garcia RF, Mimoun D, Margerin L, Karatekin O (2017b) Probing the internal structure of the asteroid Didymoon with a passive seismic investigation. *Planetary and Space Science* 144:89–105
- Neuffer D, Schultz R (2006) Mechanisms of slope failure in Valles Marineris, Mars. *Quarterly Journal of Engineering Geology and Hydrogeology* 39(3):227–240
- Noirhomme M, Ludewig N, Fand Vandewalle, Opsomer E (2017) Cluster growth in driven granular gases. *Physical Review E* 95:022,905
- Noll KS, Grundy WM, Chiang EI, Margot JL, Kern SD (2008) Binaries in the Kuiper Belt. In: Barucci, M A, Boehnhardt, H, Cruikshank, D P, & Morbidelli, A (ed) *The Solar System Beyond Neptune*, pp 345–363
- O’Brien DP, Walsh KJ, Morbidelli A, Raymond SN, Mandell AM (2014) Water delivery and giant impacts in the ‘Grand Tack’ scenario. *Icarus* 239:74–84
- Ogawa S (1978) Multitemperature theory of granular materials. In: *Proc. of the US-Japan Seminar on Continuum Mechanical and Statistical Approaches in the Mechanics of Granular Materials*, Gakajutsu Bunken Fukyu-Kai, pp 208–217
- Opsomer E, Ludewig F, Vandewalle N (2011) Phase transitions in vibrated granular systems in microgravity. *Physical Review E* 84:051,306
- Opsomer E, Vandewalle N, Noirhomme M, Ludewig F (2014) Clustering and segregation in driven granular fluids. *The European Physical Journal Special Topics* 37:115
- Opsomer E, Noirhomme M, Vandewalle N, Falcon E, Merminod S (2017) Segregation and pattern formation in dilute granular media under microgravity conditions. *npj Microgravity* 3:1
- Orpe A, Khakhar DV (2001) Scaling relations for granular flow in quasi-two-dimensional rotating cylinders. *Physical Review E* 64:031,302
- Ortiz JL, Santos-Sanz P, Sicardy B, Benedetti-Rossi G, Bérard D, Morales N, Duffard R, Braga-Ribas F, Hopp U, Ries C, et al (2017) The size, shape, density and ring of the dwarf planet Haumea from a stellar occultation. *Nature* 550(7675):219
- Ostro SJ, Hudson RS, Benner LAM, Giorgini JD, Magri C, Margot JL, Nolan MC (2002) Asteroid Radar Astronomy. In: *Asteroids III* (WF Bottke, A Cellino, P Paolicchi, RP Binzel, eds) pp 151–168
- Ottino JM, Khakhar DV (2000) Mixing and segregation of granular materials. *Annual Review of Fluid Mechanics* 32(1):55–91, DOI 10.1146/annurev.fluid.32.1.55

- Oyama Y (1939) Axial segregation of granular materials. *Bull Inst Phys Chem Res (Tokyo)*, Rept 5 18:600
- Paetzold M (2017) Mass Determination of Small Bodies in the Solar System. AGU Fall Meeting Abstracts
- Pächtz T, Herrmann HJ, Shinbrot T (2010) Why do particle clouds generate electric charges? *Nature Physics* 6:364, DOI 10.1038/nphys1631
- Patrick R, Nicodemi M, Delannay R, Ribiere P, Bideau D (2005) Slow relaxation and compaction of granular systems. *Nature materials* 4(2):121
- Peale S, Canup R (2015) The origin of the natural satellites. in *Planets and Moons, Treatise on Geophys*, vol 10, pp 559–604
- Pelton JN, Allahdadi F (eds) (2015) *Handbook of Cosmic Hazards and Planetary Defense*. Springer
- Pena AA, Garcia-Rojo R, Herrmann HJ (2007) Influence of particle shape on sheared dense granular media. *Granular Matter* 9:279–291
- Perera V, Jackson AP, Asphaug E, Ballouz RL (2016) The spherical Brazil nut effect and its significance to asteroids. *Icarus* 278:194–203
- Perna D, Dotto E, Ieva S, Barucci MA, Bernardi F, Fornasier S, De Luise F, Perozzi E, Rossi A, Mazzotta Epifani E, Micheli M, Deshapriya JDP (2016) Grasping the nature of potentially hazardous asteroids. *The Astronomical Journal* 151:11, DOI 10.3847/0004-6256/151/1/11
- Pletser V (2004) Short duration microgravity experiments in physical and life sciences during parabolic flights: the first 30 ESA campaigns. *Acta Astronautica* 55:829–854
- Polishook D, Moskovitz N, Binzel R, Burt B, DeMeo F, Hinkle M, Lockhart M, Mommert M, Person M, Thirouin A, et al (2016) A 2 km-size asteroid challenging the rubble-pile spin barrier—A case for cohesion. *Icarus* 267:243–254
- Pöschel T, Brilliantov NV (2003) *Granular gas dynamics*. Springer
- Pouliquen O, Cassar C, Jop P, Forterre Y, Nicolas M (2006) Flow of dense granular material: towards simple constitutive laws. *Journal of Statistical Mechanics: Theory and Experiment* 2006(07):P07,020
- Poux M, Fayote P, Bertrand J, Bridons D, Bousquet J (1991) Strong interlocking of nonconvex particles in random packings. *Powder Technology* 8:63
- Pravec P, Harris AW (2000) Fast and slow rotation of asteroids. *Icarus* 148(1):12–20, DOI 10.1006/icar.2000.6482
- Pravec P, Harris AW, Michalowski T (2002) Asteroid rotations. In: Bottke W, Cellino A, Paolicchi P, Binzel R (eds) *Asteroids III*, pp 113–122
- Pravec P, Vokrouhlický D, Polishook D, Scheeres DJ, Harris AW, Galad A, Vaduvescu O, Pozo F, Barr A, Longa P, et al (2010) Formation of asteroid pairs by rotational fission. *Nature* 466(7310):1085
- Procopio AT, Zavaliangos A (2005) Simulation of multi-axial compaction of granular media from loose to high relative densities. *Journal of the Mechanics and Physics of Solids* 53(7):1523–1551
- Radjai F (2015) Modeling force transmission in granular materials. *Comptes Rendus Physique* 16:3–9
- Radjai F, Dubois F (2011) *Discrete-element modeling of granular materials*. Wiley-ISTE
- Radjai F, Richefeu V (2009a) Bond anisotropy and cohesion of wet granular materials. *Phil Trans R Soc A* 367:5123–5138
- Radjai F, Richefeu V (2009b) Contact dynamics as a nonsmooth discrete element method. *Mechanics of Materials* 41(6):715–728
- Radjai F, Jean M, Moreau JJ, Roux S (1996) Force Distributions in Dense Two-Dimensional Granular Systems. *Phys Rev Lett* 77(2):274–
- Radjai F, Schäfer J, Dippel S, Wolf D (1997) Collective friction of an array of particles: A crucial test for numerical algorithms. *Journal de Physique I* 7(9):1053–1070
- Richardson D, Michel P, Walsh K, Flynn K (2009) Numerical simulations of asteroids modelled as gravitational aggregates with cohesion. *Planetary and Space Science* 57(2):183–192
- Richardson DC (1993) A new tree code method for simulation of planetesimal dynamics. *Mon Not R Astron Soc* 261:396–414, DOI 10.1093/mnras/261.2.396
- Richardson DC (1994) Tree code simulations of planetary rings. *Mon Not R Astron Soc* 269:493, DOI 10.1093/mnras/269.2.493
- Richardson DC, Quinn T, Stadel J, Lake G (2000) Direct Large-Scale N-Body Simulations of Planetesimal Dynamics. *Icarus* 143(1):45–59, DOI 10.1006/icar.1999.6243

- Richardson DC, Leinhardt ZM, Melosh HJ, Bottke WF Jr, Asphaug E (2002) Gravitational aggregates: Evidence and evolution. In: Bottke W, Cellino A, Paolicchi P, Binzel R (eds) *Asteroids III*, pp 501–515
- Richardson DC, Walsh KJ, Murdoch N, Michel P (2011) Numerical simulations of granular dynamics: I. Hard-sphere discrete element method and tests. *Icarus* 212(1):427–437
- Richardson JE, Melosh HJ, Greenberg R (2004) Impact-induced seismic activity on Asteroid 433 Eros: A surface modification process. *Science* 306(5701):1526–1529
- Rietz F, Stannarius R (2008) On the brink of jamming: Granular convection in densely filled containers. *Physical Review Letters* 100(7):078002, DOI 10.1103/PhysRevLett.100.078002
- Rivkin AS, Emery JP (2010) Detection of ice and organics on an asteroidal surface. *Nature* 464(7293):1322
- Rocchetti N, Frascarelli D, Nesmachnow S, Tancredi G (2017) Performance improvements of a parallel multithreading self-gravity algorithm. In: *Latin American High Performance Computing Conference*, Springer, pp 291–306
- Rognon P, Einav I (2010) Thermal transients and convective particle motion in dense granular materials. *Phys Rev Lett* 105:218,301, DOI 10.1103/PhysRevLett.105.218301
- Rosato A, Starndburg KJ, Prinz F, Swendsen RH (1987) Why the brazil nuts are on the top: Size segregation of particulate matter by shaking. *Physical Review Letters* 58:1038
- Rosenblatt P, Lainey V, Le Maistre S, Marty J, Dehant V, Pätzold M, Van Hoolst T, Häusler B (2008) Accurate Mars Express orbits to improve the determination of the mass and ephemeris of the Martian moons. *Planetary and Space science* 56(7):1043–1053
- Rosenblatt P, Charnoz S, Dunseath KM, Terao-Dunseath M, Trinh A, Hyodo R, Genda H, Toupin S (2016) Accretion of Phobos and Deimos in an extended debris disc stirred by transient moons. *Nature Geoscience* 9(8):581–583
- Roux S, Radjai F (1998) Texture-dependent rigid-plastic behavior. In: *Physics of dry granular media*, Springer, pp 229–236
- Rouyer F, Menon N (2000) Velocity Fluctuations in a Homogeneous 2D Granular Gas in Steady State. *Physical Review Letters* 85:3676
- Rozitis B, MacLennan E, Emery JP (2014) Cohesive forces prevent the rotational breakup of rubble-pile asteroid (29075) 1950 DA. *Nature* 512(7513):174–176, DOI 10.1038/nature13632
- Rubincam DP (2000) Radiative spin-up and spin-down of small asteroids. *Icarus* 148(1):2–11, DOI 10.1006/icar.2000.6485
- Russell HN (1906) On the light-variations of asteroids and satellites. *Astrophys J* 24:1–18, DOI 10.1086/141361
- Saito J, Miyamoto H, Nakamura R, Ishiguro M, Michikami T, Nakamura A, Demura H, Sasaki S, Hirata N, Honda C, et al (2006) Detailed images of asteroid 25143 Itokawa from Hayabusa. *Science* 312(5778):1341–1344
- Salo H (2001) Numerical simulations of the collisional dynamics of planetary rings. In: *Granular Gases*, Springer, pp 328–349
- Sánchez DP, Scheeres DJ (2012) DEM simulation of rotation-induced reshaping and disruption of rubble-pile asteroids. *Icarus* 218(2):876–894, DOI 10.1016/j.icarus.2012.01.014
- Sánchez DP, Scheeres DJ (2015) Scaling Rule Between Cohesive Forces and the Size of a Self-Gravitating Aggregate. In: *Lunar and Planetary Science Conference, Lunar and Planetary Science Conference*, vol 46, p 2556
- Sánchez P (2015) Asteroid evolution: Role of geotechnical properties”, doi = ”10.1017/s1743921315008583. *Proceedings of the International Astronomical Union* 10(S318):111–121
- Sánchez P, Scheeres DJ (2009) Granular Mechanics in Asteroid Regolith: Simulating and Scaling the Brazil Nut Effects. In: *Lunar and Planetary Science Conference, Lunar and Planetary Science Conference*, vol 40, p 2228
- Sánchez P, Scheeres DJ (2011) Simulating asteroid rubble piles with a self-gravitating soft-sphere distinct element method model. *The Astrophysical Journal* 727(2):120
- Sánchez P, Scheeres DJ (2014) The strength of regolith and rubble pile asteroids. *Meteoritics & Planetary Science* 49(5):788–811, DOI 10.1111/maps.12293
- Sánchez P, Scheeres DJ (2016) Disruption patterns of rotating self-gravitating aggregates: A survey on angle of friction and tensile strength. *Icarus* 271:453–471, DOI 10.1016/j.icarus.2016.01.016
- Sánchez P, Scheeres DJ (2018) Rotational evolution of self-gravitating aggregates with cores of variable strength. *Planetary and Space Science* DOI 10.1016/j.pss.2018.04.001

- Sanchez P, Colombo C, Vasile M, Radice G (2009) Multicriteria comparison among several mitigation strategies for dangerous near-Earth objects. *Journal of Guidance, Control, and Dynamics* 32(1):121–142
- Savage S, Lun C (1988) Particle size segregation in inclined chute flow of dry cohesionless granular solids. *Journal of Fluid Mechanics* 189:311–335
- Savage SB (1989) Flow of granular materials. In: *Theoretical and applied mechanics*, Elsevier, pp 241–266
- Savage SB (1993) *Disorder and Granular Media*. D. Bideau, North-Holland, Amsterdam
- Scheeres D (2015) Landslides and mass shedding on spinning spheroidal asteroids. *Icarus* 247(0):1–17, DOI 10.1016/j.icarus.2014.09.017
- Scheeres D, Hartzell C, Sánchez P, Swift M (2010) Scaling forces to asteroid surfaces: The role of cohesion. *Icarus* 210(2):968–984, DOI 10.1016/j.icarus.2010.07.009
- Scheeres DJ, Britt D, Carry B, Holsapple KA (2015) Asteroid interiors and morphology. In: Michel P, DeMeo F, Bottke W (eds) *Asteroids IV*, University of Arizona Press, Tucson, AZ, pp 745–766, DOI 10.2458/azu_uapress_9780816532131-ch038
- Scholz C, Pöschel T (2017) Velocity distribution of a homogeneously driven two-dimensional granular gas. *Physical Review Letters* 118:198,003
- Schorghofer N (2008) The lifetime of ice on main belt asteroids. *The Astrophysical Journal* 682(1):697
- Schröter M, Ulrich S, Kerft J, Swift JB, Swinney HL (2006) Mechanisms in the size segregation of a binary granular mixture. *Physical Review E* 74:011,307
- Schwamb ME, Jones RL, Chesley SR, Fitzsimmons A, Fraser WC, Holman MJ, Hsieh H, Ragozzine D, Thomas CA, Trilling DE, Brown ME, Bannister MT, Bodewits D, de Val-Borro M, Gerdes D, Granvik M, Kelley MSP, Knight MM, Seaman RL, Ye QZ, Young LA (2018) Large synoptic survey telescope solar system science roadmap. ArXiv e-prints 1802.01783
- Schwartz SR, Richardson DC, Michel P (2012) An implementation of the soft-sphere discrete element method in a high-performance parallel gravity tree-code. *Granular Matter* 14(3):363–380
- Schwartz SR, Michel P, Richardson DC (2013) Numerically simulating impact disruptions of cohesive glass bead agglomerates using the soft-sphere discrete element method. *Icarus* 226(1):67–76
- Schwartz SR, Michel P, Jutzi M, Marchi S, Zhang Y, Richardson DC (2018) Catastrophic disruptions as the origin of bilobate comets. *Nature Astronomy* pp 379–382
- Seiden G, Thomas PJ (2011) Complexity, segregation, and pattern formation in rotating-drum flows. *Review of Modern Physics* 83:1323
- Serero D, Noskowitz SH, Tan ML, Goldhirsch I (2009) Binary granular gas mixtures: Theory, layering effects and some open questions. *The European Physical Journal Special Topics* 179:221–247
- Shäfer J, Dippel S, Wolf D (1996) Force schemes in simulations of granular materials. *Journal de Physique I* 6(1):5–20
- Sharma I (2013) Structural stability of rubble-pile asteroids. *Icarus* 223(1):367–382
- Sharma I, Jenkins JT, Burns JA (2009) Dynamical passage to approximate equilibrium shapes for spinning, gravitating rubble asteroids. *Icarus* 200(1):304–322, DOI 10.1016/j.icarus.2008.11.003
- Shimokawa M, Suetsugu Y, Hiroshige R, Hirano T, Sakaguchi H (2015) Pattern formation in a sandpile of ternary granular mixtures. *Physical Review E* 91:062,205
- Shinbrot T, Duong NH, Kwan L, Alvarez MM (2004) Dry granular flows can generate surface features resembling those seen in Martian gullies. *Proceedings of the National Academy of Sciences* 101(23):8542–8546, DOI 10.1073/pnas.0308251101
- Sierks H, Lamy P, Barbieri C, Koschny D, Rickman H, Rodrigo R, A’Hearn MF, Angrilli F, Barucci MA, Bertaux JL, Bertini I, Besse S, Carry B, Cremonese G, Da Deppo V, Davidsson B, Debei S, De Cecco M, De Leon J, Ferri F, Fornasier S, Fulle M, Hviid SF, Gaskell RW, Groussin O, Gutierrez P, Ip W, Jorda L, Kaasalainen M, Keller HU, Knollenberg J, Kramm R, Kührt E, Küppers M, Lara L, Lazzarin M, Leyrat C, Moreno JLL, Magrin S, Marchi S, Marzari F, Massironi M, Michalik H, Moissl R, Naletto G, Preusker F, Sabau L, Sabolo W, Scholten F, Snodgrass C, Thomas N, Tubiana C, Vernazza P, Vincent JB, Wenzel KP, Andert T, Pätzold M, Weiss BP (2011) Images of Asteroid 21 Lutetia: A Remnant Planetesimal from the Early Solar System. *Science* 334:487, DOI 10.1126/science.1207325

- Spahn F, Schmidt J (2006) Hydrodynamic description of planetary rings. *GAMM-Mitteilungen* 29(1):118–143
- Stadel JG (2001) Cosmological N-body simulations and their analysis. PhD thesis, University of Washington, Washington, DC
- Stansberry J, Grundy W, Brown M, Cruikshank D, Spencer J, Trilling D, Margot J (2008) Physical Properties of Kuiper Belt and Centaur Objects: Constraints from the Spitzer Space Telescope. In: Barucci MA, Boehnhardt H, Cruikshank DP, Morbidelli A (eds) *The Solar System Beyond Neptune*, pp 161–179
- Staron L (2016) Segregation mechanisms in granular systems: role of gravity and velocity fluctuations. In: *EGU General Assembly Conference Abstracts*, vol 18, p 8047
- Staron L, Lagrée PY, Popinet S (2012) The granular silo as a continuum plastic flow: The hour-glass vs. the clepsydra. *Physics of Fluids* 24(10):103,301
- Sugimoto Y, Radice G, Ceriotti M, Sanchez JP (2014) Hazardous near Earth asteroid mitigation campaign planning based on uncertain information on fundamental asteroid characteristics. *Acta Astronautica* 103:333–357, DOI 10.1016/j.actaastro.2014.02.022
- Sunday C, Murdoch N, Cherrier O, Serrano SM, Nardi CV, Janin T, Martinez IA, Gourinat Y, Mimoun D (2016) A novel facility for reduced-gravity testing: A setup for studying low-velocity collisions into granular surfaces. *Review of Scientific Instruments* 87(8):084,504, DOI 10.1063/1.4961575
- Syal MB, Dearborn DS, Schultz PH (2013) Limits on the use of nuclear explosives for asteroid deflection. *Acta Astronautica* 90(1):103–111, DOI 10.1016/j.actaastro.2012.10.025, nEO Planetary Defense: From Threat to Action - Selected Papers from the 2011 IAA Planetary Defense Conference
- Tancredi G, Maciel A, Heredia L, Richeri P, Nesmachnow S (2012) Granular physics in low-gravity environments using discrete element method. *Monthly Notices of the Royal Astronomical Society* 420(4):3368–3380
- Tanga P, Comito C, Paolicchi P, Hestroffer D, Cellino A, Dell’Oro A, Richardson DC, Walsh K, Delbo M (2009a) Rubble-pile reshaping reproduces overall asteroid shapes. *The Astrophysical Journal Letters* 706(1):L197
- Tanga P, Hestroffer D, Delbo M, Richardson DC (2009b) Asteroid rotation and shapes from numerical simulations of gravitational re-accumulation. *Planetary and Space Science* 57(2):193–200
- Tanga P, Campo Bagatin A, Thirouin A, Cellino A, Comito C, Ortiz J, Hestroffer D, Richardson D (2013) Possible routes to spin up fission for the formation of asteroid binaries and pairs. In: *European Planetary Science Congress*, vol 8
- Tardivel S, Sánchez P, Scheeres DJ (2018) Equatorial cavities on asteroids, an evidence of fission events. *Icarus* 304:192–208, DOI 10.1016/j.icarus.2017.06.037
- Tatsumi S, Murayama Y, Hayakawa H, Sano M (2009) Experimental study on the kinetics of granular gases under microgravity. *Journal of Fluid Mechanics* 641:521–539
- Taylor PA, Howell ES, Nolan MC, Thane AA (2012) The Shape and Spin Distributions of Near-Earth Asteroids Observed with the Arecibo Radar System. In: *American Astronomical Society Meeting Abstracts #220*, American Astronomical Society Meeting Abstracts, vol 220, p 128.02
- Terzaghi K, Peck RB, Mesri G (1996) *Soil mechanics in engineering practice*. John Wiley & Sons
- Thomas P (2010) Sizes, shapes, and derived properties of the Saturnian satellites after the Cassini nominal mission. *Icarus* 208(1):395–401
- Thomas PA, Bray JD (1999) Capturing nonspherical shape of granular media with disk clusters. *Journal of Geotechnical and Geoenvironmental Engineering* 125:169–178
- Thuillet F, Maurel C, Michel P, Biele J, Ballouz RL, Richardson DC (2017) Numerical Simulations of Surface Package Landing on a Low-Gravity Granular Surface: Application to the Landing of MASCOT onboard Hayabusa2. In: *Lunar and Planetary Science Conference, Lunar and Planetary Science Conference*, vol 48, p 1810
- Tiscareno MS, Murray CD (2018) *Planetary Ring Systems: Properties, Structure, and Evolution*, vol 19. Cambridge University Press
- Toiya M, Stambaugh J, Losert W (2004) Transient and oscillatory granular shear flow. *Physical Review Letters* 93(8):088,001
- Verlet L (1967) Computer ‘experiments’ on classical fluids. I. Thermodynamical properties of Lennard-Jones molecules. *Physical Review* 159(1):98

- Von Kampen P, Kaczmarczik U, Rath HJ (2006) The new Drop Tower catapult system. *Acta Astronautica* 59:278–283
- Vu-Quoc L, Zhang X, Walton OR (2000) A 3-D discrete-element method for dry granular flows of ellipsoidal particles. *Computer methods in applied mechanics and engineering* 187:483–528
- Wadsley JW, Stadel J, Quinn T (2004) Gasoline: a flexible, parallel implementation of TreeSPH. *New Astronomy* 9:137–158, DOI 10.1016/j.newast.2003.08.004, [astro-ph/0303521](#)
- Walker R, Binns D, Carnelli I, Kueppers M, Galvez A (2016) CubeSat Opportunity Payload Intersatellite Network Sensors (COPINS) on the ESA Asteroid Impact Mission (AIM). In: 5th Interplanetary CubeSat Workshop (iCubeSat)
- Walsh K, Richardson D (2008) A steady-state model of NEA binaries formed by tidal disruption of gravitational aggregates. *Icarus* 193(2):553–566
- Walsh KJ, Jacobson SA (2015) Formation and evolution of binary asteroids. In: Michel P, DeMeo F, Bottke W (eds) *Asteroids IV*, University of Arizona Press, Tucson, AZ, pp 375–393
- Walsh KJ, Richardson DC (2006) Binary near-Earth asteroid formation: Rubble pile model of tidal disruptions. *Icarus* 180(1):201–216
- Walsh KJ, Richardson DC, Michel P (2008) Rotational breakup as the origin of small binary asteroids. *Nature* 454(7201):188
- Walsh KJ, Richardson DC, Michel P (2012) Spin-up of rubble-pile asteroids: Disruption, satellite formation, and equilibrium shapes. *Icarus* 220(2):514–529, DOI <http://dx.doi.org/10.1016/j.icarus.2012.04.029>
- Warner BD, Harris AW, Pravec P (2009) The asteroid lightcurve database. *Icarus* 202:134–146, DOI 10.1016/j.icarus.2009.02.003, updated 2018 June 24. <http://www.minorplanet.info/lightcurvedatabase.html>
- Weidenschilling SJ, Paolicchi P, Zappala V (1989) Do asteroids have satellites? In: *Asteroids II* (RP Binzel, T Gehrels, MS Matthews, eds) pp 643–658
- Weissman PR, A’Hearn MF, McFadden L, Rickman H (2002) Evolution of comets into asteroids. In: Bottke W, Cellino A, Paolicchi P, Binzel R (eds) *Asteroids III*, University of Arizona Press, Tucson, AZ, pp 669–686
- Wilkening LL, Matthews MS (1982) *Comets*. University of Arizona Press
- Will CM (2014) The confrontation between general relativity and experiment. *Living Reviews in Relativity* 17:4, DOI 10.12942/lrr-2014-4, 1403.7377
- Wisdom J, Tremaine S (1988) Local simulations of planetary rings. *The Astronomical Journal* 95:925–940
- Yeomans D, Antreasian P, Barriot JP, Chesley S, Dunham D, Farquhar R, Giorgini J, Helfrich C, Konopliv A, McAdams J, et al (2000) Radio science results during the NEAR-Shoemaker spacecraft rendezvous with Eros. *Science* 289(5487):2085–2088
- Yeomans DK, Barriot JP, Dunham D, Farquhar R, Giorgini J, Helfrich C, Konopliv A, McAdams J, Miller J, Owen W, et al (1997) Estimating the mass of asteroid 253 Mathilde from tracking data during the NEAR flyby. *Science* 278(5346):2106–2109
- Yu Y, Richardson DC, Michel P (2017) Structural analysis of rubble-pile asteroids applied to collisional evolution. *Astrodynamics* 1(1):57–69
- Zhang Y, Richardson DC, Barnouin OS, Michel P, Schwartz SR, Ballouz RL (2018) Rotational failure of rubble-pile bodies: Influences of shear and cohesive strengths. *The Astrophysical Journal* 857(1):15
- Zik O, Levine D, Lipson SG, Shtrikman S, Stavans J (1994) Rotationally induced segregation of granular materials. *Physical Review Letters* 73:644
- Zuriguel I, Gray JMNT, Peixinho J, Mullin T (2006) Pattern selection by a granular wave in a rotating drum. *Physical Review E* 73:061,302