



Polytopal realizations of finite type g-vector fans

Christophe Hohlweg, Vincent Pilaud, Salvatore Stella

► To cite this version:

Christophe Hohlweg, Vincent Pilaud, Salvatore Stella. Polytopal realizations of finite type g-vector fans. *Advances in Mathematics*, 2018, 328, pp.713-749. 10.1016/j.aim.2018.01.019 . hal-02343570

HAL Id: hal-02343570

<https://hal.science/hal-02343570>

Submitted on 2 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POLYTOPAL REALIZATIONS OF FINITE TYPE \mathbf{g} -VECTOR FANS

CHRISTOPHE HOHLWEG, VINCENT PILAUD, AND SALVATORE STELLA

ABSTRACT. This paper shows the polytopality of any finite type \mathbf{g} -vector fan, acyclic or not. In fact, for any finite Dynkin type Γ , we construct a universal associahedron $\text{Asso}_{\text{un}}(\Gamma)$ with the property that any \mathbf{g} -vector fan of type Γ is the normal fan of a suitable projection of $\text{Asso}_{\text{un}}(\Gamma)$.

1. INTRODUCTION

A *generalized associahedron* is a polytope which realizes the cluster complex of a finite type cluster algebra of S. Fomin and A. Zelevinsky [FZ03b, FZ02, FZ03a]. Generalized associahedra were first constructed by F. Chapoton, S. Fomin and A. Zelevinsky [CFZ02] using the \mathbf{d} -vector fans of [FZ03b]. Further realizations were obtained by C. Hohlweg, C. Lange and H. Thomas [HLT11] based on the Cambrian lattices of N. Reading [Rea06] and Cambrian fans of N. Reading and D. Speyer [RS09]. These constructions were later revisited by S. Stella [Ste13] using an approach similar to the original one of [CFZ02], and by V. Pilaud and C. Stump [PS15a] via brick polytopes.

The realizations of [HLT11] start from an acyclic initial exchange matrix B_o , and construct a generalized associahedron $\text{Asso}(B_o)$ whose normal fan is the \mathbf{g} -vector fan of the cluster algebra $\mathcal{A}_{\text{pr}}(B_o)$ with principal coefficients at B_o . They rely on a combinatorial understanding of the \mathbf{g} -vector fans as Cambrian fans [RS09]. A major obstruction in dropping the acyclicity assumption in this approach is that this combinatorial description is only partially available beyond acyclic cases [RS15]. Therefore, it remained a challenging open problem, since the appearance of generalized associahedra, to construct polytopal realizations of all finite type \mathbf{g} -vector fans including the cyclic cases. This paper answers this problem.

Theorem 1. *For any finite type initial exchange matrix B_o , the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(B_o)$ with respect to B_o is the normal fan of a generalized associahedron $\text{Asso}(B_o)$.*

When we start from an acyclic initial exchange matrix, our construction precisely recovers the associahedra of [HLT11, Ste13, PS15a]. These can all be obtained by deleting inequalities from the facet description of the permutahedron of the corresponding finite reflection group. The main difficulty to extend the previous approach to arbitrary initial exchange matrices lies in the fact that this property, intriguing as it might be, is essentially a coincidence. First, the hyperplane arrangement \mathcal{H} supporting the \mathbf{g} -vector fan is no longer the Coxeter arrangement of a finite reflection group. Even worse, we prove that the generalized associahedron $\text{Asso}(B_o)$ usually cannot be obtained by deleting inequalities in the facet description of any zonotope whose normal fan is \mathcal{H} . This behavior already appears in type D_5 .

To overcome this situation, we develop an alternative approach based on a uniform understanding of the linear dependences among adjacent cones in the \mathbf{g} -vector fan. In fact, not only we cover uniformly all finite type initial exchange matrices, but we actually can treat them simultaneously with a universal object.

Theorem 2. *For any given finite Dynkin type Γ , there exists a *universal associahedron* $\text{Asso}_{\text{un}}(\Gamma)$ such that, for any initial exchange matrix B_o of type Γ , the generalized associahedron $\text{Asso}(B_o)$ is a suitable projection of the universal associahedron $\text{Asso}_{\text{un}}(\Gamma)$. In particular, all \mathbf{g} -vector fans of type Γ are sections of the normal fan of the universal associahedron $\text{Asso}_{\text{un}}(\Gamma)$.*

CH was supported by NSERC Discovery grant *Coxeter groups and related structures*. SS is a Marie Curie - Cofund Fellow at INdAM. VP was partially supported by the French ANR grant SC3A (15 CE40 0004 01).

This universal associahedron provides a tool to study simultaneously geometric properties of all generalized associahedra of a given finite Dynkin type. For example, it is known that the vertex barycenters of all generalized associahedra of [HLT11] lie at the origin. In type A , this property was observed by F. Chapoton for J.-L. Loday's realization of the classical associahedron [Lod04], conjectured for all associahedra of C. Hohlweg and C. Lange in [HL07], proved by C. Hohlweg, J. Lortie and A. Raymond [HLR10] and revisited by C. Lange and V. Pilaud in [LP13]. For arbitrary acyclic finite types, it was conjectured by C. Hohlweg, C. Lange and H. Thomas in [HLT11] and proved by V. Pilaud and C. Stump using the brick polytope approach [PS15b]. In the present paper, we use the universal associahedron to extend this surprising property to all generalized associahedra $\text{Asso}(\mathbf{B}_o)$.

Theorem 3. *The origin is the vertex barycenter of the universal associahedron $\text{Asso}_{\text{un}}(\Gamma)$, and thus of all generalized associahedra $\text{Asso}(\mathbf{B}_o)$.*

The paper is organized as follows. In Section 2, we collect all the definitions and properties of finite type cluster algebras needed in this paper. In Section 3, we recall convenient criteria to check that a collection of cones forms a polyhedral fan and that a simplicial fan is the normal fan of a polytope. Based on a precise understanding of the linear dependences in \mathbf{g} -vectors of adjacent cones described in Section 4, we prove the polytopality of all finite type \mathbf{g} -vector fans in Section 5. Section 6 is devoted to two special cases: that of acyclic initial exchange matrices for which our construction yields the same generalized associahedra as [HLT11, Ste13, PS15a], and that of type A which presents several remarkable features. In particular we prove that the facet description of the associahedron $\text{Asso}(\mathbf{B}_o)$ is contained in the facet description of the corresponding zonotope $\text{Zono}(\mathbf{B}_o)$ for any initial exchange matrix \mathbf{B}_o of type A . Further properties of our generalized associahedra are explored in Section 7, including their connection to green mutations (Section 7.1), the construction of the universal associahedron (Section 7.2), its vertex barycenter (Section 7.3), and a discussion on the relation between $\text{Asso}(\mathbf{B}_o)$ and $\text{Zono}(\mathbf{B}_o)$ (Section 7.4).

2. FINITE TYPE CLUSTER ALGEBRAS

We begin by recalling some standard notions on cluster algebras simplifying, whenever possible, our notations to deal with the case at hand. This section can be used as a compendium of the results concerning finite type that are scattered through the literature. We refer to [FZ07] for a general treatment of cluster algebras.

2.1. Cluster algebras. We will be working in the ambient field $\mathbb{Q}(x_1, \dots, x_n, p_1, \dots, p_m)$ of rational expressions in $n + m$ variables with coefficients in \mathbb{Q} and we denote by \mathbb{P}_m its abelian multiplicative subgroup generated by the elements $\{p_i\}_{i \in [m]}$. Given $p = \prod_{i \in [m]} p_i^{a_i} \in \mathbb{P}_m$ we will write

$$\{p\}_+ := \prod_{i \in [m]} p_i^{\max(a_i, 0)} \quad \text{and} \quad \{p\}_- := \prod_{i \in [m]} p_i^{-\min(a_i, 0)}$$

so that $p = \{p\}_+ \{p\}_-^{-1}$.

A *seed* Σ is a triple $(\mathbf{B}, \mathbf{P}, \mathbf{X})$ consisting of an exchange matrix, a coefficient tuple, and a cluster:

- the *exchange matrix* \mathbf{B} is an integer $n \times n$ skew-symmetrizable matrix, *i.e.*, such that there exist a diagonal matrix \mathbf{D} with $-\mathbf{B}\mathbf{D} = (\mathbf{B}\mathbf{D})^T$,
- the *coefficient tuple* \mathbf{P} is any subset of n elements of \mathbb{P}_m ,
- the *cluster* \mathbf{X} is a set of *cluster variables*, n rational functions in the ambient field that are algebraically independent over $\mathbb{Q}(p_1, \dots, p_m)$.

To shorten our notation we think of rows and columns of \mathbf{B} , as well as elements of \mathbf{P} , as being labelled by the elements of \mathbf{X} : we write $\mathbf{B} = (b_{xy})_{x, y \in \mathbf{X}}$ and $\mathbf{P} = \{p_x\}_{x \in \mathbf{X}}$. Moreover we say that a cluster variable x (resp. a coefficient p) belongs to $\Sigma = (\mathbf{B}, \mathbf{P}, \mathbf{X})$ to mean $x \in \mathbf{X}$ (resp. $p \in \mathbf{P}$).

Given a seed $\Sigma = (\mathbf{B}, \mathbf{P}, \mathbf{X})$ and a cluster variable $x \in \Sigma$, we can construct a new seed $\mu_x(\Sigma) = \Sigma' = (\mathbf{B}', \mathbf{P}', \mathbf{X}')$ by *mutation* in direction x , where:

- the new cluster X' is obtained from X by replacing x with the cluster variable x' defined by the following *exchange relation*:

$$xx' = \{p_x\}_+ \prod_{\substack{y \in X \\ b_{xy} > 0}} y^{b_{xy}} + \{p_x\}_- \prod_{\substack{y \in X \\ b_{xy} < 0}} y^{-b_{xy}}$$

and leaving the remaining cluster variables unchanged so that $X \setminus \{x\} = X' \setminus \{x'\}$.

- the row (resp. column) of B' indexed by x' is the negative of the row (resp. column) of B indexed by x , while all the other entries satisfy:

$$b'_{yz} = b_{yz} + \frac{1}{2}(|b_{yx}|b_{xz} + b_{yx}|b_{xz}|),$$

- the elements of the new coefficient tuple P' are

$$p'_y = \begin{cases} p_x^{-1} & \text{if } y = x', \\ p_y \{p_x\}_-^{b_{xy}} & \text{if } y \neq x' \text{ and } b_{xy} \leq 0, \\ p_y \{p_x\}_+^{b_{xy}} & \text{if } y \neq x' \text{ and } b_{xy} > 0, \end{cases}$$

A straightforward computation shows that mutations are involutions, *i.e.*, $\mu_{x'}(\mu_x(\Sigma)) = \Sigma$ so they define an equivalence relation on the collection of all seeds.

Fix a seed $\Sigma_o = (B_o, P_o, X_o)$ and call it *initial*. Up to an automorphism of the ambient field we will assume that $X_o = \{x_1, \dots, x_n\}$ and drop X_o from our notation.

Definition 4 ([FZ07, Def. 2.11]). The (geometric type) *cluster algebra* $\mathcal{A}(B_o, P_o)$ is the $\mathbb{Z}\mathbb{P}_m$ -subring of the ambient field generated by all the cluster variables in all the seeds mutationally equivalent to the initial seed Σ_o .

Example 5. The simplest possible choice of coefficient tuple in the initial seed, namely $m = 0$ and $P_o = \{1\}_{i \in [n]}$, gives rise to the *cluster algebra without coefficients* which we will denote by $\mathcal{A}_{\text{fr}}(B_o)$. Note that this algebra, up to an automorphism of the ambient field $\mathbb{Q}(x_1, \dots, x_n)$, depends only on the mutation class of B_o and not on the exchange matrix itself. The appearance of B_o in the notation $\mathcal{A}_{\text{fr}}(B_o)$ is just to fix the embedding inside the ambient field.

2.2. Finite type. We will only be dealing with cluster algebras of *finite type* *i.e.*, cluster algebras having only a finite number of cluster variables. As it turns out, being of finite type is a property that depends only on the exchange matrix in the initial seed and not on the coefficient tuple.

The *Cartan companion* of an exchange matrix B is the symmetrizable matrix $A(B)$ given by:

$$a_{xy} = \begin{cases} 2 & \text{if } x = y, \\ -|b_{xy}| & \text{otherwise.} \end{cases}$$

Theorem 6 ([FZ03a, Thm. 1.4]). *The cluster algebra $\mathcal{A}(B_o, P_o)$ is of finite type if and only if there exists an exchange matrix B obtained by a sequence of mutations from B_o such that its Cartan companion is a Cartan matrix of finite type. Moreover the type of $A(B)$ is uniquely determined by B_o : if B' is any other exchange matrix obtained by mutation from B_o and such that $A(B')$ is a finite type Cartan matrix then $A(B')$ and $A(B)$ are related by a simultaneous permutation of rows and columns.*

In accordance with the above statement, when talking about the (*cluster*) *type* of $\mathcal{A}(B_o, P_o)$ or B_o we will refer to the Cartan type of $A(B)$. We reiterate that the Cartan type of $A(B_o)$ need not be finite: being of finite type is a property of the mutation class.

For a finite type cluster algebra $\mathcal{A}(B_o, P_o)$, we will consider the root system of $A(B_o)$. To avoid any confusion later on let us state clearly the conventions we use in this paper: for us simple roots $\{\alpha_x\}_{x \in X_o}$ and fundamental weights $\{\omega_x\}_{x \in X_o}$ are two basis of the same vector space V ; the matrix relating them is the Cartan matrix $A(B_o)$. Fundamental weights are the dual basis to simple coroots $\{\alpha_x^\vee\}_{x \in X_o}$, while simple roots are the dual basis to fundamental coweights $\{\omega_x^\vee\}_{x \in X_o}$; coroots and coweights are two basis of the dual space V^\vee and they are related by the transpose of the Cartan matrix. This set of conventions is the standard one in Lie theory but it is not the one generally used in the setting of Coxeter groups [BB05, Chap. 4].

A finite type exchange matrix B_o is said to be *acyclic* if $A(B_o)$ is itself a Cartan matrix of finite type and *cyclic* otherwise. An acyclic finite type exchange matrix is said to be *bipartite* if each of its rows (or equivalently columns) consists either of non-positive or non-negative entries.

2.3. Principal coefficients and g- and c-vectors. Among all the cluster algebras having a fixed initial exchange matrix, a central role is played by those with principal coefficients. Indeed, thanks to the results in [FZ07], they encode enough informations to understand all the other possible choices of coefficients.

Definition 7 ([FZ07, Def. 3.1]). A cluster algebra is said to have *principal coefficients* (at the initial seed) if its ambient field is $\mathbb{Q}(x_1, \dots, x_n, p_1, \dots, p_n)$ and the initial coefficient tuple consists of the generators of \mathbb{P}_n i.e., $P_o = \{p_i\}_{i \in [n]}$. In this case we will write $\mathcal{A}_{\text{pr}}(B_o)$ for $\mathcal{A}(B_o, \{p_i\}_{i \in [n]})$, and we reindex the generators $\{p_i\}_{i \in [n]}$ of \mathbb{P}_n by $\{p_x\}_{x \in X_o}$.

In the above definition, it is important to specify that principal coefficients are with respect to a specific exchange matrix, even though it is usually omitted. In other words $\mathcal{A}_{\text{pr}}(B_o)$ and $\mathcal{A}_{\text{pr}}(B'_o)$ are in general different cluster algebras even when B_o and B'_o are related by mutations.

A notable property of cluster algebras with principal coefficients is that they are \mathbb{Z}^n -graded (in the basis $\{\omega_x\}_{x \in X_o}$ of V). The degree function $\deg(B_o, \cdot)$ on $\mathcal{A}_{\text{pr}}(B_o)$ is obtained by setting

$$\deg(B_o, x) := \omega_x \quad \text{and} \quad \deg(B_o, p_x) := \sum_{y \in X_o} -b_{yx} \omega_y$$

for any $x \in X_o$. This assignment makes all exchange relations and all cluster variables in $\mathcal{A}_{\text{pr}}(B_o)$ homogeneous [FZ07] and it justifies the definition of the following family of integer vectors associated to cluster variables.

Definition 8 ([FZ07]). The *g-vector* of a cluster variable $x \in \mathcal{A}_{\text{pr}}(B_o)$ is its degree

$$\mathbf{g}(B_o, x) := \deg(B_o, x) \in V.$$

We denote by $\mathbf{g}(B_o, \Sigma) := \{\mathbf{g}(B_o, x) \mid x \in \Sigma\}$ the set of *g-vectors* of the cluster variable in the seed Σ of $\mathcal{A}_{\text{pr}}(B_o)$.

The next definition gives another family of integer vectors, introduced implicitly in [FZ07], that are relevant in the structure of $\mathcal{A}_{\text{pr}}(B_o)$.

Definition 9. Given a seed Σ in $\mathcal{A}_{\text{pr}}(B_o)$, the *c-vector* of a cluster variable $x \in \Sigma$ is the vector

$$\mathbf{c}(B_o, x \in \Sigma) := \sum_{y \in X_o} c_{yx} \alpha_y \in V$$

of exponents of $p_x = \prod_{y \in X_o} (p_y)^{c_{yx}}$. Let $\mathbf{c}(B_o, \Sigma) := \{\mathbf{c}(B_o, x \in \Sigma) \mid x \in \Sigma\}$ denote the set of *c-vectors* of a seed Σ . Finally, let $\mathbf{C}(B_o) := \bigcup_{\Sigma} \mathbf{c}(B_o, \Sigma)$ denote the set of all *c-vectors* in $\mathcal{A}_{\text{pr}}(B_o)$.

It is worth spending few words here to emphasize the fact that, contrary to what happens for *g-vectors*, *c-vectors* are not attached to cluster variables *per se* but depends on the seed in which the given cluster variable sits.

An important features of *c-vectors* is that their entries weekly agree in sign. This is one of the various reformulation of the *sign-coherence* conjecture of [FZ07] recently established in full generality by [GHKK14]. In the setting of finite type cluster algebras, this result can also be deduced in several ways from earlier works: one proof is to combine [DWZ10] with [Dem10]; another one is to use surfaces [FST08, FT12] and orbifolds [FST12] to study types A_n, B_n, C_n and D_n , and to deal with exceptional types by direct inspection. Here we prefer to observe it as a corollary of the following theorem that will be useful later on to justify our notation.

Theorem 10 ([NS14, Thm. 1.3]). *The c-vectors of the finite type cluster algebra $\mathcal{A}_{\text{pr}}(B_o)$ are roots in the root system whose Cartan matrix is $A(B_o)$.*

Again note that, since $A(B_o)$ may be not of finite type, the root system in this statement is, in general, not finite. For example, for the cyclic type A_3 exchange matrix, the Cartan companion $A(B_o)$ is of affine type $A_2^{(1)}$, see its Coxeter arrangement in Figure 5 (top right). More precisely,

it is finite if and only if B_\circ is acyclic. We will discuss in more details the relation of \mathbf{c} -vectors with root systems in Remark 21.

Another consequence of [NS14, Thm. 1.3] is that in a cluster algebra of finite type, there are nh distinct \mathbf{c} -vectors where h is the Coxeter number of the given finite type. We remind to our reader that the same algebra has $(h+2)n/2$ distinct \mathbf{g} -vectors, one for each cluster variable.

Our next task in this section is to discuss a duality relation in between \mathbf{c} -vectors and \mathbf{g} -vectors. A first step is to recall the notion of the *cluster complex* of $\mathcal{A}(B_\circ, P_\circ)$: it is the abstract simplicial complex whose vertices are the cluster variables of $\mathcal{A}(B_\circ, P_\circ)$ and whose facets are its clusters. As it turns out, at least in the finite type cases, this complex is independent of the choice of coefficients, see [FZ03a, Thm. 1.13] and [FZ07, Conj. 4.3]. In particular this means that, up to isomorphism, there is only one cluster complex for each finite type: the one associated to $\mathcal{A}_{\text{fr}}(B_\circ)$. We will use this remark later on to relate cluster variables of different cluster algebras of the same finite type. Note also that, again when $\mathcal{A}(B_\circ, P_\circ)$ is of finite type, the cluster complex is a *pseudomanifold* [FZ03a].

For a skew-symmetrizable exchange matrix B_\circ , the matrix $B_\circ^\vee := -B_\circ^T$ is still skew-symmetrizable. The cluster algebras $\mathcal{A}_{\text{pr}}(B_\circ)$ and $\mathcal{A}_{\text{pr}}(B_\circ^\vee)$ can be thought as *dual* to each other. Indeed their types are Langlands dual of each other. Moreover their cluster complexes are isomorphic: by performing the same sequence of mutations we can identify any cluster variable x of $\mathcal{A}_{\text{pr}}(B_\circ)$ with a cluster variable x^\vee of $\mathcal{A}_{\text{pr}}(B_\circ^\vee)$, and any seed Σ in $\mathcal{A}_{\text{pr}}(B_\circ)$ with a seed Σ^\vee in $\mathcal{A}_{\text{pr}}(B_\circ^\vee)$. More importantly the following crucial property holds.

Theorem 11 ([NZ12, Thm. 1.2]). *For any seed Σ of $\mathcal{A}_{\text{pr}}(B_\circ)$, let Σ^\vee be its dual in $\mathcal{A}_{\text{pr}}(B_\circ^\vee)$. Then the \mathbf{g} -vectors $\mathbf{g}(B_\circ, \Sigma)$ of the cluster variables in Σ and the \mathbf{c} -vectors $\mathbf{c}(B_\circ^\vee, \Sigma^\vee)$ of the cluster variables in Σ^\vee are dual bases, i.e.,*

$$\langle \mathbf{g}(B_\circ, x) \mid \mathbf{c}(B_\circ^\vee, y^\vee \in \Sigma^\vee) \rangle = \delta_{x=y}$$

for any two cluster variables $x, y \in \Sigma$.

In view of the above results, and since $A(B_\circ^\vee) = A(B_\circ)^T$, the \mathbf{c} -vectors of a finite type cluster algebra $\mathcal{A}_{\text{pr}}(B_\circ^\vee)$ can be understood as coroots for $A(B_\circ)$ so that the \mathbf{g} -vectors of $\mathcal{A}_{\text{pr}}(B_\circ)$ become weights. This justifies our choice to define \mathbf{g} -vectors in the weight basis.

2.4. Coefficient specialization and universal cluster algebra. We now want to relate, within a given finite type, cluster algebras with different choices of coefficients. Pick a finite type exchange matrix B_\circ and let $\mathcal{A}(B_\circ, P_\circ) \subset \mathbb{Q}(x_1, \dots, x_n, p_1, \dots, p_m)$ and $\mathcal{A}(B_\circ, \overline{P}_\circ) \subset \mathbb{Q}(\overline{x}_1, \dots, \overline{x}_n, \overline{p}_1, \dots, \overline{p}_\ell)$ be any two cluster algebras having B_\circ as exchange matrix in their initial seed. As we said, cluster variables and seeds in these two algebras are in bijection because their cluster complexes are isomorphic. Let us write

$$x \longleftrightarrow \overline{x} \quad \text{and} \quad \Sigma \longleftrightarrow \overline{\Sigma}$$

for this bijection. We will say that $\mathcal{A}(B_\circ, \overline{P}_\circ)$ is obtained from $\mathcal{A}(B_\circ, P_\circ)$ by a *coefficient specialization* if there exist a map of abelian groups $\eta : \mathbb{P}_m \rightarrow \mathbb{P}_\ell$ such that, for any p_x in some seed Σ of $\mathcal{A}(B_\circ, P_\circ)$

$$\eta(\{p_x\}_+) = \{\overline{p_x}\}_+ \quad \text{and} \quad \eta(\{p_x\}_-) = \{\overline{p_x}\}_-$$

and which extends in a unique way to a map of algebras that satisfy

$$\eta(x) = \overline{x}.$$

Note that this is not the most general definition (see [FZ07, Def. 12.1 and Prop. 12.2]) but it will suffice here. Armed with the notion of coefficient specialization we can now introduce the last kind of cluster algebra of finite type we will need.

Definition 12 ([FZ07, Def. 12.3 and Thm. 12.4]). Pick a finite type exchange matrix B_\circ . The *cluster algebra with universal coefficients* $\mathcal{A}_{\text{un}}(B_\circ)$ is the unique (up to canonical isomorphism) cluster algebra such that any other cluster algebra of the same type as B_\circ can be obtained from it by a unique coefficient specialization.

Let us insist on the fact that, in view of the universal property it satisfies, $\mathcal{A}_{\text{un}}(\mathbf{B}_o)$ depends only on the type of \mathbf{B}_o and not on the exchange matrix \mathbf{B}_o itself. We again keep \mathbf{B}_o in the notation only to fix an embedding into the ambient field.

Rather than proving the existence and explaining the details of how such a universal algebra is built, we will recall here one of its remarkable properties that follows directly from the \mathbf{g} -vector recursion [NZ12, Prop. 4.2 (v)] and that we will need in our proofs later on. Denote by $\mathcal{X}(\mathbf{B}_o)$ the set of all cluster variables in $\mathcal{A}_{\text{un}}(\mathbf{B}_o)$ and let $\{p[x]\}_{x \in \mathcal{X}(\mathbf{B}_o)}$ be the generators¹ of $\mathbb{P}_{|\mathcal{X}(\mathbf{B}_o)|}$.

Theorem 13 ([Rea14, Theo. 10.12]). *The cluster algebra $\mathcal{A}_{\text{un}}(\mathbf{B}_o)$ can be realized over $\mathbb{P}_{|\mathcal{X}(\mathbf{B}_o)|}$. The coefficient tuple $\mathbf{P} = \{p_x\}_{x \in X}$ at each seed $\Sigma = (\mathbf{B}, \mathbf{P}, X)$ of $\mathcal{A}_{\text{un}}(\mathbf{B}_o)$ is given by the formula*

$$p_x = \prod_{y \in \mathcal{X}(\mathbf{B}_o)} (p[y])^{[\mathbf{g}(\mathbf{B}^T, y^T); x^T]}$$

where we denote by $[\mathbf{v}; x]$ the x -th coefficient of a vector \mathbf{v} in the weight basis $(\omega_x)_{x \in X}$. The bijection of the elements of $\mathcal{X}(\mathbf{B}_o)$ with the cluster variables of $\mathcal{A}_{\text{pr}}(\mathbf{B}^T)$, appearing in the formula, is given by an isomorphism of the corresponding cluster complexes similar to the one discussed above.

Remark 14. In view of this result, it is straightforward to produce the coefficient specialization morphism to get any cluster algebra with principal coefficients of type \mathbf{B}_o from $\mathcal{A}_{\text{un}}(\mathbf{B}_o)$. Namely, for any seed $\Sigma_\star = (\mathbf{B}_\star, \mathbf{P}_\star, X_\star)$ of $\mathcal{A}_{\text{un}}(\mathbf{B}_o)$, we obtain $\mathcal{A}_{\text{pr}}(\mathbf{B}_\star)$ by evaluating to 1 all the coefficients $p[y]$ corresponding to cluster variables y not in Σ_\star .

We conclude this review giving an example: the cluster algebra of type B_2 with universal coefficients. We will do so in terms of “tall” rectangular matrices to help readers, not familiar with the language we adopt here, recognize an hopefully more familiar setting. Indeed, to pass from our seeds to the one consisting of extended exchange matrices and clusters with frozen variables, it suffices to observe that each $p \in \mathbb{P}_m$ can be encoded in a vector. In this way any n -tuple of elements of \mathbb{P}_m corresponds to a $m \times n$ integer matrix and one gets an extended exchange matrix by glueing it below the exchange matrix of the seed. The frozen variables are the generators of \mathbb{P}_m and the rules of mutations we discussed become then the usual mutations of extended exchange matrices. We prefer to use the notation we set up here following [FZ03a, FZ07] because it makes more evident the distinction in between coefficients and cluster variables, and because it is more suited to deal with coefficient specializations.

Example 15. Consider the exchange matrix

$$\mathbf{B}_o = \begin{bmatrix} 0 & 1 \\ -2 & 0 \end{bmatrix}.$$

Any cluster algebra built from this matrix will contain 6 cluster variables. We will call them $\mathcal{X}(\mathbf{B}_o) = \{x_1, x_2, x_3, x_4, x_5, x_6\}$. The corresponding cluster algebra with universal coefficients will then be a subring of

$$\mathbb{Q}(x_1, x_2, p[x_1], p[x_2], p[x_3], p[x_4], p[x_5], p[x_6]).$$

Namely it will be the cluster algebra

$$\mathcal{A}_{\text{un}}(\mathbf{B}_o) = \mathcal{A}\left(\mathbf{B}_o, \left\{ \frac{p[x_1]p[x_5]p[x_6]^2}{p[x_3]}, \frac{p[x_2]}{p[x_4]p[x_5]p[x_6]} \right\}\right)$$

Figure 1 shows the exchange graph of this algebra listing all the seeds in terms of extended exchange matrices.

One final computational remark: there is a simple algorithm to compute one of the rectangular exchange matrices appearing in a cluster algebra of finite type with universal coefficients. Let \mathbf{B} be an exchange matrix of the given finite type having only non-negative entries above its diagonal; it is acyclic and, by [YZ08, Eqn. (1.4)], it corresponds to the *coxeter element* $c = s_1 \cdots s_n$ in the associated Weyl group (note that the labelling of simple roots may not be the standard one

¹A note to the reader that might be scared of a circular reasoning here: the set of generators $\{p[x]\}_{x \in \mathcal{X}(\mathbf{B}_o)}$ is just a collection of symbols and its cardinality can be precomputed: it depends only on the type of the algebra and not on its coefficients.




FIGURE 1. The exchange graph of type B_2 with attached the rectangular matrices giving universal coefficients. Highlighted in blue are the entries of the coefficient part that give the principal coefficient cluster algebra at the seed attached to the red node.

here). Let $w_\circ(c)$ denote the c -sorting word for w_\circ , that is the lexicographically minimal reduced expression of w_\circ that appears as a subword of c^∞ . For each s_i obtained reading from left to right the word $cw_\circ(c)$, repeat the following two steps:

- add a row to the bottom of B whose only non-zero entry is a 1 in column i
- replace B by its mutation in direction i .

The matrix obtained at the end of the procedure is the desired one.

3. POLYHEDRAL GEOMETRY AND FANS

The second ingredient of this paper is discrete geometry of polytopes and fans. We refer to [Zie98] for a textbook on this topic.

3.1. Polyhedral fans. A *polyhedral cone* is a subset of the vector space V defined equivalently as the positive span of finitely many vectors or as the intersection of finitely many closed linear halfspaces. Throughout the paper, we write $\mathbb{R}_{\geq 0}\mathbf{\Lambda}$ for the positive span of a set $\mathbf{\Lambda}$ of vectors of V .

The *faces* of a cone C are the intersections of C with its supporting hyperplanes. In particular, the 1-dimensional (resp. codimension 1) faces of C are called *rays* (resp. *facets*) of C . A cone is *simplicial* if it is generated by a set of independent vectors.

A *polyhedral fan* is a collection \mathcal{F} of polyhedral cones of V such that

- if $C \in \mathcal{F}$ and F is a face of C , then $F \in \mathcal{F}$,
- the intersection of any two cones of \mathcal{F} is a face of both.

A fan is *simplicial* if all its cones are, and *complete* if the union of its cones covers the ambient space V . For a simplicial fan \mathcal{F} with rays \mathcal{X} , the collection $\{X \subseteq \mathcal{X} \mid \mathbb{R}_{\geq 0}X \in \mathcal{F}\}$ of generating sets of the cones of \mathcal{F} defines a *pseudomanifold* (in other words, a pure and thin simplicial complex, *i.e.*, with a notion of flip). The following statement characterizes which pseudomanifolds are complete simplicial fans. A formal proof can be found *e.g.*, in [DRS10, Coro. 4.5.20].

Proposition 16. *Consider a pseudomanifold Δ with vertex set \mathcal{X} and a set of vectors $\{\mathbf{r}(x)\}_{x \in \mathcal{X}}$ of V . For $X \in \Delta$, let $\mathbf{r}(X) := \{\mathbf{r}(x) \mid x \in X\}$. Then the collection of cones $\{\mathbb{R}_{\geq 0}\mathbf{r}(X) \mid X \in \Delta\}$ forms a complete simplicial fan if and only if*

- (1) *there exists a facet X of Δ such that $\mathbf{r}(X)$ is a basis of V and such that the open cones $\mathbb{R}_{>0}\mathbf{r}(X)$ and $\mathbb{R}_{>0}\mathbf{r}(X')$ are disjoint for any facet X' of Δ distinct from X ;*
- (2) *for any two adjacent facets X, X' of Δ with $X \setminus \{x\} = X' \setminus \{x'\}$, there is a linear dependence*

$$\gamma \mathbf{r}(x) + \gamma' \mathbf{r}(x') + \sum_{y \in X \cap X'} \delta_y \mathbf{r}(y) = 0$$

on $\mathbf{r}(X \cup X')$ where the coefficients γ and γ' have the same sign. (When these conditions hold, these coefficients do not vanish and the linear dependence is unique up to rescaling.)

3.2. Polytopes and normal fans. A *polytope* is a subset P of V^\vee defined equivalently as the convex hull of finitely many points or as a bounded intersection of finitely many closed affine halfspaces. The *faces* of P are the intersections of P with its supporting hyperplanes. In particular, the dimension 0 (resp. dimension 1, resp. codimension 1) faces of P are called *vertices* (resp. *edges*, resp. *facets*) of P . The (outer) *normal cone* of a face F of P is the cone in V generated by the outer normal vectors of the facets of P containing F . The (outer) *normal fan* of P is the collection of the (outer) normal cones of all its faces. We say that a complete polyhedral fan in V is *polytopal* when it is the normal fan of a polytope in V^\vee . The following statement provides a characterization of polytopality of complete simplicial fans. It is a reformulation of regularity of triangulations of vector configurations, introduced in the theory of secondary polytopes [GKZ08], see also [DRS10]. We present here a convenient formulation from [CFZ02, Lem. 2.1].

Proposition 17. *Consider a pseudomanifold Δ with vertex set \mathcal{X} and a set of vectors $\{\mathbf{r}(x)\}_{x \in \mathcal{X}}$ of V such that $\mathcal{F} := \{\mathbb{R}_{\geq 0}\mathbf{r}(X) \mid X \in \Delta\}$ forms a complete simplicial fan. Then the following are equivalent:*

- (1) *\mathcal{F} is the normal fan of a simple polytope in V^\vee ;*
- (2) *There exists a map $h : \mathcal{X} \rightarrow \mathbb{R}_{>0}$ such that for any two adjacent facets X, X' of Δ with $X \setminus \{x\} = X' \setminus \{x'\}$, we have*

$$\gamma h(x) + \gamma' h(x') + \sum_{y \in X \cap X'} \delta_y h(y) > 0,$$

where

$$\gamma \mathbf{r}(x) + \gamma' \mathbf{r}(x') + \sum_{y \in X \cap X'} \delta_y \mathbf{r}(y) = 0$$

is the unique (up to rescaling) linear dependence with $\gamma, \gamma' > 0$ between the rays of $\mathbf{r}(X \cup X')$.

Under these conditions, \mathcal{F} is the normal fan of the polytope defined by

$$\{\mathbf{v} \in V^\vee \mid \langle \mathbf{r}(x), \mathbf{v} \rangle \leq h(x) \text{ for all } x \in \mathcal{X}\}.$$

4. THE \mathbf{g} -VECTOR FAN

We first recast a well known fact concerning the cones spanned by the \mathbf{g} -vectors of a finite type cluster algebra with principal coefficients. We insist on the fact that the following statement is valid for any finite type exchange matrix B_\circ , acyclic or not.

Theorem 18. *For any finite type exchange matrix B_\circ , the collection of cones*

$$\mathcal{F}^{\mathbf{g}}(B_\circ) := \{ \mathbb{R}_{\geq 0} \mathbf{g}(B_\circ, \Sigma) \mid \Sigma \text{ seed of } \mathcal{A}_{\text{pr}}(B_\circ) \},$$

together with all their faces, forms a complete simplicial fan, called the \mathbf{g} -vector fan of B_\circ .

There are several ways to find or deduce Theorem 18 from the literature. First, it was established in the acyclic case in [RS09, YZ08, Ste13] (see Example 22). As already observed by N. Reading in [Rea14, Thm. 10.6], the general case then follows from the initial seed recursion on \mathbf{g} -vectors [NZ12, Prop. 4.2 (v)], valid thanks to sign-coherence. A second proof would be to use the unique cluster expansion property of any vector in the weight lattice (following from the fact that cluster monomials are a basis of $\mathcal{A}_{\text{pr}}(B_\circ)$ in finite type), and to use approximation by this lattice to show that any vector is covered exactly once by the interiors of the cones of the \mathbf{g} -vector fan. Note that contrarily to what sometimes appears in the literature, this approximation argument is subtle as it relies on the integrity of the \mathbf{g} -vectors². Finally, another possible proof is to use Proposition 16: the first point is a simplified version of the unique expansion property, and the second point is a consequence of the following description of the linear dependence between the \mathbf{g} -vectors of two adjacent clusters, which will be crucial in the next section.

Lemma 19. *For any finite type exchange matrix B_\circ and any adjacent seeds (B, P, X) and (B', P', X') in $\mathcal{A}_{\text{pr}}(B_\circ)$ with $X \setminus \{x\} = X' \setminus \{x'\}$, the \mathbf{g} -vectors of $X \cup X'$ satisfy precisely one of the following two linear dependences*

$$\mathbf{g}(B_\circ, x) + \mathbf{g}(B_\circ, x') = \sum_{\substack{y \in X \cap X' \\ b_{xy} < 0}} -b_{xy} \mathbf{g}(B_\circ, y) \quad \text{or} \quad \mathbf{g}(B_\circ, x) + \mathbf{g}(B_\circ, x') = \sum_{\substack{y \in X \cap X' \\ b_{xy} > 0}} b_{xy} \mathbf{g}(B_\circ, y).$$

Proof. This is a straightforward consequence of the definition of \mathbf{g} -vectors together with sign coherence. Indeed all exchange relations in $\mathcal{A}_{\text{pr}}(B_\circ)$ are homogeneous and

$$xx' = \{p_x\}_+ \prod_{\substack{y \in X \\ b_{xy} > 0}} y^{b_{xy}} + \{p_x\}_- \prod_{\substack{y \in X \\ b_{xy} < 0}} y^{-b_{xy}}$$

means that

$$\begin{aligned} \deg(B_\circ, x) + \deg(B_\circ, x') &= \deg(B_\circ, \{p_x\}_+) + \sum_{\substack{y \in X \cap X' \\ b_{xy} > 0}} b_{xy} \deg(B_\circ, y) \\ &= \deg(B_\circ, \{p_x\}_-) + \sum_{\substack{y \in X \cap X' \\ b_{xy} < 0}} -b_{xy} \deg(B_\circ, y). \end{aligned}$$

Now, by sign-coherence, exactly one of $\{p_x\}_+$ and $\{p_x\}_-$ is 1 so that its degree is 0. \square

Remark 20. Note that which of the two possible linear dependences is satisfied by the \mathbf{g} -vectors of $X \cup X'$ depends on the initial exchange matrix B_\circ . In particular, the geometry of the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(B_\circ)$ changes as B_\circ varies within a given mutation class.

²For an illustration of the subtlety, consider the 8 cones in \mathbb{R}^3 defined by the coordinate hyperplanes, rotate the 4 cones with $x \geq 0$ around the x -axis by $\pi/4$, and finally rotate all the cones around the origin by an irrational rotation so that each rational direction belongs to the interior of one of the 8 resulting cones. Then any vector in \mathbb{Z}^3 belongs to a single cone, but the resulting cones do not form a fan (since the cones with $x \geq 0$ intersect improperly those with $x \leq 0$).




FIGURE 2. The dual \mathbf{c} -vector fan $\mathcal{F}^{\mathbf{c}}(\mathbf{B}_\circ^\vee)$ (thin red) and \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(\mathbf{B}_\circ)$ (bold blue) for the type A_3 (left) and type C_3 (right) cyclic initial exchange matrices. Each 3-dimensional fan is intersected with the unit sphere and stereographically projected to the plane from the pole $(-1, -1, -1)$. We use this drawing convention for all 3-dimensional fans in this paper.

For any finite type exchange matrix \mathbf{B}_\circ , the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(\mathbf{B}_\circ)$ can be seen as a coarsening of two other fans naturally associated to $\mathbf{A}(\mathbf{B}_\circ)$. Denote by $\mathcal{F}^{\mathbf{Cox}}(\mathbf{B}_\circ^\vee)$ the *dual Coxeter fan* i.e., the fan of regions of the hyperplane arrangement given by the root system of $\mathbf{A}(\mathbf{B}_\circ^\vee)$. Similarly let $\mathcal{F}^{\mathbf{c}}(\mathbf{B}_\circ^\vee)$ be the *dual c-vector fan* i.e., the fan of regions of the arrangement of hyperplanes orthogonal to all the \mathbf{c} -vector of $\mathcal{A}_{\text{pr}}(\mathbf{B}_\circ^\vee)$. By Theorem 11, $\mathcal{F}^{\mathbf{g}}(\mathbf{B}_\circ)$ coarsens $\mathcal{F}^{\mathbf{c}}(\mathbf{B}_\circ^\vee)$ which, in turn, coarsens $\mathcal{F}^{\mathbf{Cox}}(\mathbf{B}_\circ^\vee)$ by Theorem 10. See Figure 5 for examples of these fans for different exchange matrices of type A_3 .

Remark 21. By further inspecting [NS14, Thm. 1.3] we can say more about $\mathcal{F}^{\mathbf{c}}(\mathbf{B}_\circ^\vee)$. Indeed its supporting hyperplane arrangement contains the hyperplanes associated to small roots in the root system of $\mathbf{A}(\mathbf{B}_\circ^\vee)$. Therefore, it turns out that the dual \mathbf{c} -vector fan $\mathcal{F}^{\mathbf{c}}(\mathbf{B}_\circ^\vee)$ intersected with the Tits cone contains the *Shi arrangement* for the root system of $\mathbf{A}(\mathbf{B}_\circ^\vee)$ (see [HNW16, Sect. 3.6 & Def. 3.18] for a review on the topic). In order to see that, it is enough to compare the description of the possible supports of the \mathbf{c} -vectors (in terms of simple roots) given in [NS14, Thm. 1.3] with the description of the possible supports of small roots given in [Bri98].

Example 22. When the exchange matrix \mathbf{B}_\circ is acyclic, the \mathbf{g} -vector fan is the Cambrian fan constructed by N. Reading and D. Speyer [RS09], while the dual \mathbf{c} -vector fan is the type $\mathbf{A}(\mathbf{B}_\circ^\vee)$ Coxeter fan. Section 6.1 provides a detailed discussion of the acyclic case.

Example 23. Figure 2 illustrates the \mathbf{g} -vector fans for the initial exchange matrices

$$\begin{bmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 0 & -1 & 2 \\ 1 & 0 & -2 \\ -1 & 1 & 0 \end{bmatrix}$$

(type A_3 cyclic) (type C_3 cyclic)

Note that these matrices are the only examples of 3-dimensional cyclic exchange matrices (up to duality and simultaneous permutations of rows and columns).

5. POLYTOPALITY

In this section, we show that the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(\mathbf{B}_\circ)$ is polytopal for any finite type exchange matrix \mathbf{B}_\circ . As discussed in Section 6.1, this result was previously known for acyclic finite type exchange matrices [HLT11, Ste13, PS15a]. The proof of this paper, although similar in spirit to that of [Ste13], actually simplifies the previous approaches.

We first consider some convenient functions which will be used later in Theorem 26 to lift the \mathbf{g} -vector fan. The existence of such functions will be discussed in Proposition 28.

Definition 24. A positive function h on the cluster variables of $\mathcal{A}(\mathbf{B}_\circ, \mathbf{P}_\circ)$ is *exchange submodular*³ if, for any pair of adjacent seeds $(\mathbf{B}, \mathbf{P}, X)$ and $(\mathbf{B}', \mathbf{P}', X')$ with $X \setminus \{x\} = X' \setminus \{x'\}$, it satisfies

$$h(x) + h(x') > \max \left(\sum_{\substack{y \in X \cap X' \\ b_{xy} < 0}} -b_{xy} h(y), \sum_{\substack{y \in X \cap X' \\ b_{xy} > 0}} b_{xy} h(y) \right).$$

Definition 25. Let h be a positive function on the cluster variables of $\mathcal{A}_{\text{pr}}(\mathbf{B}_\circ)$ we define:

(i) a point

$$\mathbf{p}^h(\mathbf{B}_\circ, \Sigma) := \sum_{x \in \Sigma} h(x) \mathbf{c}(\mathbf{B}_\circ^\vee, x^\vee \in \Sigma^\vee) \in V^\vee$$

for each seed Σ of $\mathcal{A}_{\text{pr}}(\mathbf{B}_\circ)$,

(ii) a halfspace $\mathbf{H}_{\leq}^h(\mathbf{B}_\circ, x)$ and a hyperplane $\mathbf{H}_{=}^h(\mathbf{B}_\circ, x)$ by

$$\begin{aligned} \mathbf{H}_{\leq}^h(\mathbf{B}_\circ, x) &:= \{ \mathbf{v} \in V^\vee \mid \langle \mathbf{g}(\mathbf{B}_\circ, x) \mid \mathbf{v} \rangle \leq h(x) \} \\ \text{and } \mathbf{H}_{=}^h(\mathbf{B}_\circ, x) &:= \{ \mathbf{v} \in V^\vee \mid \langle \mathbf{g}(\mathbf{B}_\circ, x) \mid \mathbf{v} \rangle = h(x) \} \end{aligned}$$

for each cluster variable x of $\mathcal{A}_{\text{pr}}(\mathbf{B}_\circ)$.

The following statement is the central result of this paper. We refer again to Proposition 28 for the existence of exchange submodular functions.

Theorem 26. For any finite type exchange matrix \mathbf{B}_\circ and any exchange submodular function h , the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(\mathbf{B}_\circ)$ is the normal fan of the \mathbf{B}_\circ -associahedron $\text{Asso}^h(\mathbf{B}_\circ) \subseteq V^\vee$ equivalently defined as

- (i) the convex hull of the points $\mathbf{p}^h(\mathbf{B}_\circ, \Sigma)$ for all seeds Σ of $\mathcal{A}_{\text{pr}}(\mathbf{B}_\circ)$, or
- (ii) the intersection of the halfspaces $\mathbf{H}_{\leq}^h(\mathbf{B}_\circ, x)$ for all cluster variables x of $\mathcal{A}_{\text{pr}}(\mathbf{B}_\circ)$.

Proof. Consider two adjacent seeds $\Sigma = (\mathbf{B}, \mathbf{P}, X)$ and $\Sigma' = (\mathbf{B}', \mathbf{P}', X')$ with $X \setminus \{x\} = X' \setminus \{x'\}$. By Lemma 19, the linear dependence between the \mathbf{g} -vectors of $X \cup X'$ is of the form

$$\mathbf{g}(\mathbf{B}_\circ, x) + \mathbf{g}(\mathbf{B}_\circ, x') = \sum_{\substack{y \in X \cap X' \\ \varepsilon(\mathbf{B}_\circ, \Sigma, \Sigma') b_{xy} > 0}} \varepsilon(\mathbf{B}_\circ, \Sigma, \Sigma') b_{xy} \mathbf{g}(\mathbf{B}_\circ, y)$$

for some $\varepsilon(\mathbf{B}_\circ, \Sigma, \Sigma') \in \{\pm 1\}$ (depending on the initial exchange matrix \mathbf{B}_\circ). However, by Definition 24, the function h satisfies

$$h(x) + h(x') > \max \left(\sum_{\substack{y \in X \cap X' \\ b_{xy} < 0}} -b_{xy} h(y), \sum_{\substack{y \in X \cap X' \\ b_{xy} > 0}} b_{xy} h(y) \right) \geq \sum_{\substack{y \in X \cap X' \\ \varepsilon(\mathbf{B}_\circ, \Sigma, \Sigma') b_{xy} > 0}} \varepsilon(\mathbf{B}_\circ, \Sigma, \Sigma') b_{xy} h(y).$$

Applying the characterization of Proposition 17, we thus immediately obtain that the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(\mathbf{B}_\circ)$ is the normal fan of the polytope defined as the intersection of the halfspaces $\mathbf{H}_{\leq}^h(\mathbf{B}_\circ, x)$ for all cluster variables x of $\mathcal{A}_{\text{pr}}(\mathbf{B}_\circ)$.

³The term “exchange submodular” is inspired from a particular situation in type A (namely, when \mathbf{B}_\circ has 1’s on the upper subdiagonal and -1 ’s on the lower subdiagonal), where such functions really correspond to the classical submodular functions, *i.e.*, functions $h : 2^{[n]} \rightarrow \mathbb{R}$ such that $h(X) + h(Y) > h(X \cup Y) + h(X \cap Y)$ for any distinct non-trivial subsets X, Y of $[n]$.

Finally, to show the vertex description, we just need to observe that, for any seed Σ of $\mathcal{A}_{\text{pr}}(\mathbf{B}_o)$, the point $\mathbf{p}^h(\mathbf{B}_o, \Sigma)$ is the intersection of the hyperplanes $\mathbf{H}_{\Sigma}^h(\mathbf{B}_o, x)$ for all $x \in \Sigma$. Indeed, since $\mathbf{g}(\mathbf{B}_o, \Sigma)$ and $\mathbf{c}(\mathbf{B}_o^\vee, \Sigma^\vee)$ form dual bases by Theorem 11, we have for any $x \in \Sigma$:

$$\langle \mathbf{g}(\mathbf{B}_o, x) | \mathbf{p}^h(\mathbf{B}_o, \Sigma) \rangle = \sum_{y \in \Sigma} h(y) \langle \mathbf{g}(\mathbf{B}_o, x) | \mathbf{c}(\mathbf{B}_o^\vee, y^\vee \in \Sigma^\vee) \rangle = \sum_{y \in \Sigma} h(y) \delta_{x=y} = h(x). \quad \square$$

Remark 27. By definition $\text{Asso}^h(\mathbf{B}_o)$ fulfills the following properties:

- the normal vectors are the \mathbf{g} -vectors of all cluster variables of $\mathcal{A}_{\text{pr}}(\mathbf{B}_o)$,
- for any two adjacent seeds $\Sigma = (\mathbf{B}, \mathbf{P}, \mathbf{X})$ and $\Sigma' = (\mathbf{B}', \mathbf{P}', \mathbf{X}')$ with $\mathbf{X} \setminus \{x\} = \mathbf{X}' \setminus \{x'\}$, the edge of $\text{Asso}^h(\mathbf{B}_o)$ joining the vertex $\mathbf{p}^h(\mathbf{B}_o, \Sigma)$ to the vertex $\mathbf{p}^h(\mathbf{B}_o, \Sigma')$ is a negative multiple of the dual \mathbf{c} -vector $\mathbf{c}(\mathbf{B}_o^\vee, x^\vee \in \Sigma^\vee) = -\mathbf{c}(\mathbf{B}_o^\vee, x'^\vee \in \Sigma'^\vee)$. More precisely,

$$\mathbf{p}^h(\mathbf{B}_o, \Sigma') - \mathbf{p}^h(\mathbf{B}_o, \Sigma) = \left(\sum_y b_{xy} h(y) - h(x') - h(x) \right) \mathbf{c}(\mathbf{B}_o^\vee, x^\vee \in \Sigma^\vee),$$

where the sum runs over the variables $y \in \Sigma \cap \Sigma'$ such that b_{xy} has the same sign as the \mathbf{c} -vector $\mathbf{c}(\mathbf{B}_o^\vee, x^\vee \in \Sigma^\vee)$. Note that $\sum_y b_{xy} h(y) - h(x') - h(x) < 0$ since h is exchange sumodular.

Our next step is to show the existence of exchange submodular functions for any finite type cluster algebra with principal coefficients. The important observation here is that the definition of exchange submodular function does not involve in any way the coefficients of $\mathcal{A}_{\text{pr}}(\mathbf{B}_o)$ so that it suffices to construct one in the coefficient free cases. Indeed, if h is exchange submodular for $\mathcal{A}_{\text{fr}}(\mathbf{B}_o)$, and η is the coefficient specialization morphism given by

$$\begin{array}{ccc} \eta : \mathcal{A}_{\text{pr}}(\mathbf{B}_o) & \longrightarrow & \mathcal{A}_{\text{fr}}(\mathbf{B}_o) \\ p_i & \longmapsto & 1 \end{array}$$

one gets the desired map by setting

$$h(x) := h(\eta(x))$$

for any cluster variable x of $\mathcal{A}_{\text{pr}}(\mathbf{B}_o)$.

Recall that, up to an obvious automorphism of the ambient field, there exists a unique cluster algebra without coefficients for each given finite type [FZ07]. We can therefore, without loss of generality, assume that \mathbf{B}_o is bipartite and directly deduce our result from [Ste13, Prop. 8.3] obtained as an easy consequence of [CFZ02, Lem. 2.4] which we recast here in our current setting.

When \mathbf{B}_o is acyclic, the Weyl group of $\mathbf{A}(\mathbf{B}_o)$ is finite and has a longest element w_o . A point $\lambda^\vee := \sum_{x \in \mathbf{X}_o} \lambda_x^\vee \omega_x^\vee$ in the interior of the fundamental Weyl chamber of $\mathbf{A}(\mathbf{B}_o^\vee)$ (that is to say $\lambda_x^\vee > 0$ for all $x \in \mathbf{X}_o$) is *fairly balanced* if $w_o(\lambda^\vee) = -\lambda^\vee$.

Proposition 28. *Let $\mathcal{A}_{\text{fr}}(\mathbf{B}_o)$ be any finite type cluster algebra without coefficients and assume that \mathbf{B}_o is bipartite. To each fairly balanced point λ^\vee corresponds an exchange submodular function h_{λ^\vee} on $\mathcal{A}_{\text{fr}}(\mathbf{B}_o)$.*

Proof. To define h_{λ^\vee} , recall from the construction in [FZ03b, FZ03a] that the set of cluster variables in $\mathcal{A}_{\text{fr}}(\mathbf{B}_o)$ is acted upon by a dihedral group generated by the symbols τ_+ and τ_- .

Each $\langle \tau_+, \tau_- \rangle$ -orbit of cluster variables meets the initial seed in either 1 or 2 elements. We will define h_{λ^\vee} to be constant on the orbits of this action: on any element in the orbit of the initial cluster variable x the function h_{λ^\vee} evaluates to the x -th coordinate of λ^\vee when written in the basis of simple coroots. The requirement that λ^\vee is fairly balanced, as explained in the proof of [Ste13, Thm. 6], is tantamount to say that $h_{\lambda^\vee}(x) = h_{\lambda^\vee}(y)$ if x and y are initial cluster variables in the same $\langle \tau_+, \tau_- \rangle$ -orbit.

The fact that the function h_{λ^\vee} defined in this way is exchange submodular is then the content of [CFZ02, Lem. 2.4] together with the computation following [Ste13, Prop. 8]. The only minor thing to observe is that, there, this function appears as a piecewise linear function on the ambient space of the root lattice and thus, instead of writing the cluster expansions of the two monomials in the right hand side of the exchange relations as we do here, their total denominator vectors appear. \square

Remark 29. Reading through [Ste13] one may have the impression, as the author did at the time, that for a given λ^\vee a different exchange submodular function has been constructed for each choice of acyclic initial seed. This is not the case. Indeed, if one unravels the definitions, it is easy to see that the functions defined there only differ because the set of \mathbf{g} -vectors they use as domain are different. Taken as function on the cluster variables in the respective cluster algebras with principal coefficients, under the specialization maps η , they all correspond to the same function on $\mathcal{A}_{\text{fr}}(\mathbf{B}_o)$.

A particular example of fairly balanced point is the point

$$\rho^\vee := \sum_{x \in X_o} \omega_x^\vee.$$

Note that ρ^\vee is both the sum of the fundamental coweights and the half sum of all positive coroots of the root system of finite type $A(\mathbf{B}_o)$. In particular h_{ρ^\vee} is the *half compatibility sum* of x , *i.e.*, the half sum of the compatibility degrees

$$h_{\rho^\vee}(x) := \frac{1}{2} \sum_{y \neq x} (y \parallel x)$$

over all cluster variables distinct from x . (See [FZ03b, CP15] for the definition and discussion of the relevant properties of compatibility degrees.) The point ρ^\vee is particularly relevant in representation theory and its role in this context has already been observed in [CFZ02, Rem. 1.6]. We call *balanced \mathbf{B}_o -associahedron* and denote by $\text{Asso}(\mathbf{B}_o)$ the \mathbf{B}_o -associahedron $\text{Asso}^{h_{\rho^\vee}}(\mathbf{B}_o)$ for the exchange submodular function h_{ρ^\vee} .

Example 30. When \mathbf{B}_o is acyclic, the \mathbf{B}_o -associahedron $\text{Asso}(\mathbf{B}_o)$ was already constructed in [HLT11, Ste13, PS15a]. It is then obtained by deleting inequalities from the facet description of the permutahedron of the Coxeter group of type $A(\mathbf{B}_o)$. Section 6.1 provides a detailed discussion of the acyclic case.

Example 31. Following Example 23, we have represented in Figure 3 the \mathbf{B}_o -associahedra $\text{Asso}(\mathbf{B}_o)$ for the same two initial exchange matrices

$$\begin{bmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 0 & -1 & 2 \\ 1 & 0 & -2 \\ -1 & 1 & 0 \end{bmatrix}$$

(type A_3 cyclic) (type C_3 cyclic)

Note that the leftmost associahedron of Figure 3 appeared as a mysterious realization of the associahedron in [CSZ15].

We sum up by stating the main result of this paper.

Corollary 32. *For any finite type exchange matrix \mathbf{B}_o , the \mathbf{g} -vector fan $\mathcal{F}^\mathbf{g}(\mathbf{B}_o)$ is polytopal.*

6. TWO FAMILIES OF EXAMPLES

Before investigating more combinatorial and geometric properties of the \mathbf{B}_o -associahedron $\text{Asso}(\mathbf{B}_o)$, we take a moment to study two specific families of examples, corresponding to initial exchange matrices that are either acyclic (Section 6.1), or of type A (Section 6.2).

6.1. Acyclic case. We first consider an acyclic initial seed, *i.e.*, with exchange matrix \mathbf{B}_o whose Cartan companion $A(\mathbf{B}_o)$ is itself a Cartan matrix of finite type. We denote by $W(\mathbf{B}_o)$ the Weyl group of type $A(\mathbf{B}_o)$ and by w_o its longest element. Note that the choice of an acyclic seed is equivalent to the choice of a Coxeter element c of $W(\mathbf{B}_o)$.

We gather in the following list some relevant facts from the literature (some of which were already observed earlier in the text). We refer to [Hoh12] for a detailed survey on these properties.

Roots and weights:

- the cluster variables of $\mathcal{A}_{\text{pr}}(\mathbf{B}_o)$ are in bijection with the almost positive roots of $A(\mathbf{B}_o)$,
- the dual \mathbf{c} -vectors of $\mathcal{A}_{\text{pr}}(\mathbf{B}_o)$ are all the coroots of $A(\mathbf{B}_o)$,




FIGURE 3. The associahedra $\text{Asso}(\text{B}_o)$ for the type A_3 (left) and type C_3 (right) cyclic initial exchange matrices.

- the \mathbf{g} -vectors of $\mathcal{A}_{\text{pr}}(\text{B}_o)$ are some weights of $A(\text{B}_o)$.

Cambrian fan and Coxeter fan:

- the dual \mathbf{c} -vector fan $\mathcal{F}^c(\text{B}_o^\vee)$ coincides with the dual Coxeter fan $\mathcal{F}^{\text{Cox}}(\text{B}_o^\vee)$,
- the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(\text{B}_o)$ coincides with the c -Cambrian fan of N. Reading and D. Speyer [RS09].

HLT associahedron and permutahedron: Consider any fairly balanced point $\lambda^\vee := \sum_{x \in X_o} \lambda_x^\vee \omega_x^\vee$ in the fundamental chamber of $A(\text{B}_o^\vee)$. Then the B_o -associahedron $\text{Asso}^{h_{\lambda^\vee}}(\text{B}_o)$ coincides with the c -associahedron $\text{Asso}^{\lambda^\vee}(c)$ constructed by C. Hohlweg, C. Lange and H. Thomas [HLT11] and later revisited by S. Stella [Ste13] and by V. Pilaud and C. Stump [PS15a] in the context of brick polytopes. In particular, $\text{Asso}^{\lambda^\vee}(c)$ is defined by the inequalities normal to the \mathbf{g} -vectors of $\mathcal{A}_{\text{pr}}(\text{B}_o)$ in the facet description of the B_o -permutahedron

$$\begin{aligned} \text{Perm}^{\lambda^\vee}(\text{B}_o) &:= \text{conv} \{w(\lambda^\vee) \mid w \in W(\text{B}_o)\} \\ &= \{\mathbf{v} \in V^\vee \mid \langle w(\omega_x) \mid \mathbf{v} \rangle \leq \lambda_x^\vee \text{ for all } x \in X_o, w \in W(\text{B}_o)\}. \end{aligned}$$

See [Hoh12] for more details on the relation between $\text{Perm}^{\lambda^\vee}(\text{B}_o)$ and $\text{Asso}^{\lambda^\vee}(c)$.

Cambrian lattice and weak order: When oriented in the direction $-\sum_{x \in X_o} \omega_x$,

- the graph of the B_o -permutahedron is the Hasse diagram of the weak order on $W(\text{B}_o)$,
- the graph of the B_o -associahedron is the Hasse diagram of the c -Cambrian lattice of N. Reading [Rea06], which is a lattice quotient and a sublattice of the weak order on $W(\text{B}_o)$.

Vertex barycenter: For any fairly balanced point λ^\vee , the origin is the vertex barycenter of both the B_o -permutahedron $\text{Perm}^{\lambda^\vee}(\text{B}_o)$ and the B_o -associahedron $\text{Asso}^{\lambda^\vee}(\text{B}_o)$:

$$\sum_{w \in W(\text{B}_o)} w(\lambda^\vee) = \sum_{\Sigma} \mathbf{p}^{\lambda^\vee}(\text{B}_o, \Sigma) = \mathbf{0}.$$

In type A , this property was observed by F. Chapoton for J.-L. Loday's realization of the classical associahedron [Lod04] and conjectured for arbitrary Coxeter element by C. Hohlweg and C. Lange in [HL07] in the balanced case. It was later proved by C. Hohlweg, J. Lortie and A. Raymond [HLR10] and revisited by C. Lange and V. Pilaud in [LP13]. Both proofs use an orbit refinement of this property. For arbitrary finite types, it was conjectured by C. Hohlweg, C. Lange and H. Thomas in [HLT11] and proved by V. Pilaud and C. Stump using the brick polytope approach [PS15b].

We will see in Section 7 how these properties of the B_o -associahedron $\text{Asso}^{h_{\lambda^\vee}}(\text{B}_o)$ for finite type acyclic initial exchange matrices B_o extend to arbitrary initial exchange matrices.

6.2. Type A. We now consider an initial exchange matrix of type A . It is well known that the vertices of the type A associahedron are counted by the Catalan numbers and are therefore in bijection with all Catalan families. We use in this paper the classical model by triangulations of a convex polygon [FZ03b], that we briefly recall now.

Triangulation model for type A cluster algebras. Consider a convex $(n+3)$ -gon Ω_n . A *dissection* is a set of pairwise non-crossing internal diagonals of Ω_n , and a *triangulation* is a maximal dissection (thus decomposing Ω_n into triangles). A triangulation T defines a matrix $B(T) = (b_{\gamma\delta})$ whose rows and columns are indexed by the diagonals of T and where

$$b_{\gamma\delta} = \begin{cases} 1 & \text{if } \gamma \text{ follows } \delta \text{ in counter-clockwise order around a triangle of } T, \\ -1 & \text{if } \gamma \text{ precedes } \delta \text{ in counter-clockwise order around a triangle of } T, \\ 0 & \text{otherwise.} \end{cases}$$

A *flip* in a triangulation T consists in exchanging an internal diagonal γ by the other diagonal γ' of the quadrilateral formed by the two triangles of T containing it. The reader can observe that the exchange matrix $B(T')$ of the resulting triangulation $T' = T \triangle \{\gamma, \gamma'\}$ is obtained from the exchange matrix $B(T)$ by a mutation in direction γ (as defined in Section 2).

Moreover, note that if the triangulation T has no internal triangle, then we can order linearly its diagonals $\gamma_1, \dots, \gamma_n$ such that $b_{\gamma_i, \gamma_{i+1}} = \pm 1$ and $b_{\gamma_i, \gamma_j} = 0$ if $|i - j| \neq 1$, so that the Cartan companion $A(B(T))$ is precisely the type A Cartan matrix. The reciprocal statement clearly holds as well: any exchange matrix whose Cartan companion is the type A Cartan matrix is the exchange matrix $B(T)$ of a triangulation T with no internal triangle. Therefore, the flip graph on triangulations of Ω_n completely encodes the combinatorics of mutations in type A .

The choice of a type A initial exchange matrix B_\circ is thus equivalent to the choice of an initial triangulation T_\circ of Ω_n . The cluster algebra $\mathcal{A}_{\text{fr}}(B(T_\circ))$ has a cluster variable x_δ for each (internal) diagonal of the polygon Ω_n . Recall that if δ, δ' are the two diagonals of a quadrilateral with edges $\kappa, \lambda, \mu, \nu$ in cyclic order, then the corresponding cluster variables are related by the exchange relation $x_\delta x_{\delta'} = x_\kappa x_\mu + x_\lambda x_\nu$. (Here and elsewhere it is understood that if κ is a boundary edge, then $x_\kappa = 1$ and similarly for λ, μ , and ν .)

The compatibility degree in type A is very simple: for $\gamma \neq \delta$, we have $(x_\gamma \parallel x_\delta) = 1$ if γ and δ cross, and 0 otherwise. In particular, the function h_ρ is given by $h_\rho(x_\gamma) = i(n-2-i)/2$ for a diagonal γ with i vertices of Ω_n on one side and $n-2-i$ on the other side.

Finally, note that since the exchange matrix $B(T_\circ)$ is skew-symmetric, the algebra $\mathcal{A}_{\text{pr}}(B(T_\circ))$ and its dual $\mathcal{A}_{\text{pr}}(B(T_\circ)^\vee)$ coincide. We therefore take the freedom to omit mentioning duals in all this type A discussion.

Shear coordinates for \mathbf{g} - and \mathbf{c} -vectors. We now provide a combinatorial interpretation for the \mathbf{g} - and \mathbf{c} -vectors in a type A cluster algebra in terms of the triangulation model. Our presentation is a light version (for the special case of triangulations of a disk) of the shear coordinates of [FST08, FT12] developed to provide combinatorial models for cluster algebras from surfaces.

We consider $2n+6$ points on the unit circle labeled clockwise by $1_\circ, 2_\bullet, \dots, (2n+5)_\circ, (2n+6)_\bullet$. We say that $1_\circ, 3_\circ, \dots, (2n+5)_\circ$ are the *hollow vertices* and that $2_\bullet, 4_\bullet, \dots, (2n+6)_\bullet$ are the *solid vertices*. We simultaneously consider *hollow triangulations* (based on hollow vertices) and *solid triangulations* (based on solid vertices), but never mix hollow and solid vertices in our triangulations. To help distinguishing them, hollow vertices and diagonals appear red while solid vertices and diagonals appear blue in all pictures. See *e.g.*, Figure 4.

Let T be a hollow (resp. solid) triangulation, let $\delta \in T$, and let γ be a solid (resp. hollow) diagonal. We denote by $Q(\delta \in T)$ the quadrilateral formed by the two triangles of T incident to δ . When γ crosses δ , we define $\varepsilon(\delta \in T, \gamma)$ to be 1, -1, or 0 depending on whether γ crosses $Q(\delta \in T)$ as a Σ , as a Δ , or in a corner. If γ and δ do not cross, then we set $\varepsilon(\delta \in T, \gamma) = 0$.

Fix once and for all a *reference triangulation* T_\circ of the hollow polygon and let $(\omega_{\delta_\circ})_{\delta_\circ \in T_\circ}$ and $(\alpha_{\delta_\bullet})_{\delta_\bullet \in T_\bullet}$ denote dual bases of \mathbb{R}^{T_\circ} . The reference triangulation T_\circ of the hollow polygon defines an initial triangulation $T_\bullet^- := \{(i-1)_\bullet, (j-1)_\bullet \mid (i_\circ, j_\circ) \in T_\circ\}$ of the solid polygon, with $B(T_\bullet^-) = B(T_\circ)$. The cluster algebra $\mathcal{A}_{\text{pr}}(B(T_\bullet^-))$ has one cluster variable x_{δ_\bullet} for each solid internal diagonal δ_\bullet . For a diagonal δ_\bullet and a triangulation T_\bullet with $\delta_\bullet \in T_\bullet$, we write $\mathbf{g}(T_\bullet, \delta_\bullet)$




FIGURE 4. Examples of twisted red and blue triangulations.

and $\mathbf{c}(T_o, \delta_\bullet \in T_\bullet)$ for the \mathbf{g} - and \mathbf{c} -vectors of the variable x_{δ_\bullet} computed from the initial seed triangulation T_o^- . We denote by $\mathbf{C}(T_o)$ the set of all \mathbf{c} -vectors of $\mathcal{A}_{\text{pr}}(\mathcal{B}(T_o))$.

The following statement provides a combinatorial interpretation of the \mathbf{g} - and \mathbf{c} -vectors.

Proposition 33. *For any diagonal δ_\bullet and a triangulation T_\bullet with $\delta_\bullet \in T_\bullet$, we have*

$$\mathbf{g}(T_o, \delta_\bullet) := \sum_{\delta_o \in T_o} \varepsilon(\delta_o \in T_o, \delta_\bullet) \omega_{\delta_o} \quad \text{and} \quad \mathbf{c}(T_o, \delta_\bullet \in T_\bullet) := \sum_{\delta_o \in T_o} -\varepsilon(\delta_\bullet \in T_\bullet, \delta_o) \alpha_{\delta_o}.$$

Intuitively, the \mathbf{g} -vector of δ_\bullet is given by alternating ± 1 along the *zigzag* of δ_\bullet in T_o (the diagonals of T_o that cross opposite edges of $Q(\delta_o \in T_o)$) and the \mathbf{c} -vector of δ_\bullet in T_\bullet is up to a sign the characteristic vector of the diagonals of T_o that cross opposite edges of $Q(\delta_\bullet \in T_\bullet)$.

For example, the solid diagonal $2_\bullet 10_\bullet$ in the triangulation T_o^ℓ of Figure 4 (left) has \mathbf{g} -vector $\mathbf{g}(T_o, 2_\bullet 10_\bullet) = \omega_{3_\circ 15_\circ} - \omega_{9_\circ 15_\circ}$ and \mathbf{c} -vector $\mathbf{c}(T_o, 2_\bullet 10_\bullet \in T_\bullet^\ell) = -\alpha_{5_\circ 15_\circ} - \alpha_{9_\circ 15_\circ}$, while the blue diagonal $4_\bullet 14_\bullet$ in the triangulation T_o^r of Figure 4 (right) has \mathbf{g} -vector $\mathbf{g}(T_o, 4_\bullet 14_\bullet) = \omega_{5_\circ 15_\circ}$ and \mathbf{c} -vector $\mathbf{c}(T_o, 4_\bullet 14_\bullet \in T_\bullet^r) = \alpha_{5_\circ 15_\circ} + \alpha_{9_\circ 15_\circ}$.

Note that there is one \mathbf{g} -vector $\mathbf{g}(T_o, \delta_\bullet)$ for each internal diagonal δ_\bullet . In contrast, many $\delta_\bullet \in T_\bullet$ give the same \mathbf{c} -vector $\mathbf{c}(T_o, \delta_\bullet \in T_\bullet)$. For a diagonal $\delta_\bullet = u_\bullet v_\bullet$, let A_o^- (resp. A_o^+) denote the edges of T_o crossed by δ_\bullet and not incident to the vertices $u_o + 1$ or $v_o + 1$ (resp. $u_o - 1$ or $v_o - 1$). Define $\mathbf{c}^-(\delta_\bullet) := -\sum_{\delta_o \in A_o^-} \omega_{\delta_o}$ and $\mathbf{c}^+(\delta_\bullet) := \sum_{\delta_o \in A_o^+} \omega_{\delta_o}$. The negative (resp. positive) \mathbf{c} -vectors of $\mathbf{C}(T_o)$ are then precisely given by the vectors $\mathbf{c}^-(\delta_\bullet)$ (resp. $\mathbf{c}^+(\delta_\bullet)$) for all diagonals δ_\bullet not in $T_o^- := \{(i-1)_\bullet(j-1)_\bullet \mid (i_\circ, j_\circ) \in T_o\}$ (resp. $T_o^+ := \{(i+1)_\bullet(j+1)_\bullet \mid (i_\circ, j_\circ) \in T_o\}$).

Specializing Theorem 18, the simplicial complex of dissections of Ω_n is realized by the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(T_o) := \{\mathbf{g}(T_o, D_\bullet) \mid D_\bullet \text{ dissection of } \Omega_n\}$, which coarsens the \mathbf{c} -vector fan $\mathcal{F}^{\mathbf{c}}(T_o)$ (defined by the arrangement of hyperplanes orthogonal to the \mathbf{c} -vectors of $\mathbf{C}(T_o)$), which in turn coarsens the Coxeter fan $\mathcal{F}^{\text{Cox}}(T_o)$ (defined by the Coxeter arrangement for the Cartan matrix $A(\mathcal{B}(T_o))$). These fans are illustrated in Figure 5 for various initial hollow triangulations T_o .

T_o -zonotope, T_o -associahedron and T_o -parallelepiped. Using these \mathbf{g} - and \mathbf{c} -vectors, we now consider three polytopes associated to T_o :

- (1) The *T_o -zonotope* $\text{Zono}(T_o)$ is the Minkowski sum of all \mathbf{c} -vectors:

$$\text{Zono}(T_o) := \sum_{\mathbf{c} \in \mathbf{C}(T_o)} \mathbf{c}.$$

Its normal fan is the fan given by the arrangement of the hyperplanes normal to the \mathbf{c} -vectors of $\mathbf{C}(T_o)$.

- (2) The *T_o -associahedron* $\text{Asso}(T_o)$ is the polytope defined equivalently as
 - (i) the convex hull of the points $\mathbf{p}(T_o, T_\bullet) := \sum_{\delta_\bullet \in T_\bullet} h_\rho(\delta_\bullet) \mathbf{c}(T_o, \delta_\bullet \in T_\bullet)$ for all solid triangulations T_\bullet ,
 - (ii) the intersection of the hyperplanes $\mathbf{H}_\leq(T_o, \delta_\bullet) := \{\mathbf{v} \in \mathbb{R}^{T_o} \mid \langle \mathbf{g}(T_o, \delta_\bullet) \mid \mathbf{v} \rangle \leq h_\rho(\delta_\bullet)\}$ for all solid diagonal δ_\bullet .

The normal fan of $\text{Asso}(T_o)$ is the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(T_o)$.




FIGURE 5. Stereographic projections of the Coxeter fan $\mathcal{F}^{\text{Cox}}(T_o)$, the \mathbf{c} -vector fan $\mathcal{F}^c(T_o)$, and the \mathbf{g} -vector fans $\mathcal{F}^g(T_o)$ for various reference triangulations T_o . The 3-dimensional fan is intersected with the unit sphere and stereographically projected to the plane from the pole in direction $(-1, -1, -1)$.

(3) The T_\circ -*parallelepiped* $\text{Para}(T_\circ)$ is the parallelepiped defined as

$$\text{Para}(T_\circ) := \{\mathbf{v} \in \mathbb{R}^{T_\circ} \mid |\langle \omega_{\delta_\circ} | \mathbf{v} \rangle| \leq h_\rho((i-1)_\bullet(j-1)_\bullet) \text{ for all } \delta_\circ = i_\circ j_\circ \in T_\circ\}.$$

Its normal fan is the coordinate fan, defined by the coordinate hyperplanes.

These polytopes are illustrated in Figure 6 for various initial hollow triangulations T_\circ . Note that the T_\circ -zonotope $\text{Zono}(T_\circ)$ is not simple in general. To simplify the presentation, we restricted the definition to the balanced exchange submodular function h_ρ , but similar definitions would of course hold with any exchange submodular function.

These three polytopes are strongly related: not only their normal fans, but even their inequality descriptions, refine each other, as stated in our next proposition. This property is quite specific to type A as shown in Section 7.4.

Proposition 34. *All facet defining inequalities of $\text{Para}(T_\circ)$ are facet defining inequalities of $\text{Asso}(T_\circ)$, and all facet defining inequalities of $\text{Asso}(T_\circ)$ are facet defining inequalities of $\text{Zono}(T_\circ)$.*

Proof. The first part of the sentence is immediate since for any initial diagonal $\delta_\circ = i_\circ j_\circ$, we have

$$\mathbf{g}(T_\circ, (i-1)_\bullet(j-1)_\bullet) = \omega_{\delta_\circ}, \quad \text{and} \quad \mathbf{g}(T_\circ, (i+1)_\bullet(j+1)_\bullet) = -\omega_{\delta_\circ}.$$

For the second part of the statement, let $k(\gamma_\bullet)$ denote the maximum of $\langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{v} \rangle$ over $\text{Zono}(T_\circ)$. As $\text{Zono}(T_\circ)$ is the Minkowski sum of all \mathbf{c} -vectors, we have

$$k(\gamma_\bullet) = \sum_{\substack{\mathbf{c} \in \mathbf{C}(T_\circ) \\ \langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c} \rangle > 0}} \langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c} \rangle.$$

To compute this sum, recall that the \mathbf{g} -vector $\mathbf{g}(T_\circ, \gamma_\bullet)$ has alternating ± 1 along the zigzag Z_\circ of γ_\bullet in T_\circ . Choose a \mathbf{c} -vector $\mathbf{c} \in \mathbf{C}(T_\circ)$ and let $\delta_\bullet \in T_\bullet$ be such that $\mathbf{c} = \mathbf{c}(T_\circ, \delta_\bullet \in T_\bullet)$. Since all diagonals of Z_\circ that traverse $Q(\delta_\bullet \in T_\bullet)$ cross it in the same way (either all as Σ or all as Z), we have $\langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c} \rangle \in \{-1, 0, 1\}$. We thus want to count the \mathbf{c} -vectors $\mathbf{c} \in \mathbf{C}(T_\circ)$ for which $\langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c} \rangle > 0$. It actually turns out that it is more convenient and equivalent (since $\mathbf{C}(T_\circ) = -\mathbf{C}(T_\circ)$) to count the \mathbf{c} -vectors $\mathbf{c} \in \mathbf{C}(T_\circ)$ for which $\langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c} \rangle < 0$.

Decompose the zigzag Z_\circ into $Z_\circ = Z_\circ^+ \sqcup Z_\circ^-$ such that $\mathbf{g}(T_\circ, \gamma_\bullet) = \sum_{\delta_\circ \in Z_\circ^+} \omega_{\delta_\circ} - \sum_{\delta_\circ \in Z_\circ^-} \omega_{\delta_\circ}$. For a diagonal $\delta_\bullet = u_\bullet v_\bullet$, let A_\circ^- (resp. A_\circ^+) denote the edges of T_\circ crossed by δ_\bullet and not incident to $u_\circ + 1$ or $v_\circ + 1$ (resp. $u_\circ - 1$ or $v_\circ - 1$). Let $\mathbf{c}^-(\delta_\bullet) := -\sum_{\delta_\circ \in A_\circ^-} \omega_{\delta_\circ}$ and $\mathbf{c}^+(\delta_\bullet) := \sum_{\delta_\circ \in A_\circ^+} \omega_{\delta_\circ}$. Recall that the negative (resp. positive) \mathbf{c} -vectors of $\mathbf{C}(T_\circ)$ are then precisely given by the vectors $\mathbf{c}^-(\delta_\bullet)$ (resp. $\mathbf{c}^+(\delta_\bullet)$) for all diagonals δ_\bullet not in $T_\bullet^- := \{(i-1)_\bullet(j-1)_\bullet \mid (i_\circ, j_\circ) \in T_\circ\}$ (resp. $T_\bullet^+ := \{(i+1)_\bullet(j+1)_\bullet \mid (i_\circ, j_\circ) \in T_\circ\}$). We leave it to the reader to check that:

- (i) If γ_\bullet and δ_\bullet do not cross and have no common endpoint, both $|Z_\circ \cap A_\circ^-|$ and $|Z_\circ \cap A_\circ^+|$ are even. Thus $\langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c}^-(\delta_\bullet) \rangle = \langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c}^+(\delta_\bullet) \rangle = 0$.
- (ii) If γ_\bullet and δ_\bullet have a common endpoint, and $\gamma_\bullet \delta_\bullet$ form a counterclockwise angle, then $|Z_\circ \cap A_\circ^-|$ is even while $Z_\circ \cap A_\circ^+$ is empty or starts and ends in Z_\circ^+ . Thus $\langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c}^-(\delta_\bullet) \rangle = 0$ while $\langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c}^+(\delta_\bullet) \rangle \geq 0$. The situation is similar if $\gamma_\bullet \delta_\bullet$ form a clockwise angle.
- (iii) If γ_\bullet and δ_\bullet cross, $Z_\circ \cap A_\circ^-$ and $Z_\circ \cap A_\circ^+$ are empty or start and end both in Z_\circ^- or both in Z_\circ^+ . Thus, either $\langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c}^-(\delta_\bullet) \rangle < 0$ and $\langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c}^+(\delta_\bullet) \rangle \geq 0$ or conversely.

This shows that there are as many \mathbf{c} -vectors $\mathbf{c} \in \mathbf{C}(T_\circ)$ for which $\langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{c} \rangle < 0$ as diagonals δ_\bullet crossing γ_\bullet . In other words, $k(\gamma_\bullet) = h_\rho(\gamma_\bullet)$.

Finally, we obtained that the inequality $\langle \mathbf{g}(T_\circ, \gamma_\bullet) | \mathbf{v} \rangle \leq k(\gamma_\bullet)$ defines a face $\mathbf{F}(\gamma_\bullet)$ of the zonotope $\text{Zono}(T_\circ)$. This face $\mathbf{F}(\gamma_\bullet)$ is the Minkowski sum of the \mathbf{c} -vectors of $\mathbf{C}(T_\circ)$ orthogonal to $\mathbf{g}(T_\circ, \gamma_\bullet)$. Theorem 11 ensures that any triangulation T_\bullet containing γ_\bullet already provides $n-1$ linearly independent such \mathbf{c} -vectors $\mathbf{c}(T_\circ, \delta_\bullet \in T_\bullet)$ for $\delta_\bullet \in T_\bullet \setminus \{\gamma_\bullet\}$. We obtain that $\mathbf{F}(\gamma_\bullet)$ has dimension $n-1$ and is therefore a facet of the zonotope $\text{Zono}(T_\circ)$. \square

Vertex barycenter. We now use this (type A) interpretation of the T_\circ -associahedron $\text{Asso}(T_\circ)$ to show that its vertex barycenter is also at the origin. Our approach is independent, and somewhat simpler than the previous proofs of this property for type A acyclic associahedra [HLR10, LP13].




FIGURE 6. The zonotope $\text{Zono}(T_o)$, associahedron $\text{Asso}(T_o)$ and parallelepiped $\text{Para}(T_o)$ for different reference triangulations T_o . The first column is J.-L. Loday's associahedron [Lod04], the second column is one of C. Hohlweg and C. Lange's associahedra [HL07], the third column appeared in a discussion in C. Ceballos, F. Santos and G. Ziegler's survey on associahedra [CSZ15, Fig. 3].

Proposition 35. *For any initial triangulation T_o , the origin is the vertex barycenter of the T_o -zonotope $\text{Zono}(T_o)$, the T_o -associahedron $\text{Asso}(T_o)$ and the T_o -parallelepiped $\text{Para}(T_o)$.*

Proof. Assume that we started from a regular $(2n + 6)$ -gon with alternative hollow and solid vertices. Consider a diagonal δ_o of T_o and let Ψ denote the reflexion of the plane which stabilizes δ_o . Note that Ψ sends solid diagonals (resp. triangulations) onto solid diagonals (resp. triangulations). Moreover, a diagonal δ_\bullet crosses δ_o if and only if its image $\Psi(\delta_\bullet)$ crosses δ_o . Since Ψ reverses the orientation, we therefore obtain that

$$\varepsilon(\delta_\bullet \in T_\bullet, \delta_o) = -\varepsilon(\Psi(\delta_\bullet) \in \Psi(T_\bullet), \delta_o),$$

for any $\delta_\bullet \in T_\bullet$. Finally, Ψ preserves the length of a diagonal δ_\bullet , so that $h_\rho(\Psi(\delta_\bullet)) = h_\rho(\delta_\bullet)$. Summing over all diagonals in all solid triangulations, we obtain that the δ_o -coordinate of the vertex barycenter of $\text{Asso}(T)$ is given by

$$\left(\sum_{T_\bullet} \mathbf{p}(T_o, T_\bullet) \right)_{\delta_o} = \sum_{T_\bullet} \sum_{\delta_\bullet \in T_\bullet} (h_\rho(\delta_\bullet) \mathbf{c}(T_o, \delta_\bullet \in T_\bullet))_{\delta_o} = - \sum_{T_\bullet} \sum_{\delta_\bullet \in T_\bullet} h_\rho(\delta_\bullet) \varepsilon(\delta_\bullet \in T_\bullet, \delta_o) = 0$$

since the contribution of $\delta_\bullet \in T_\bullet$ is balanced by that of $\Psi(\delta_\bullet) \in \Psi(T_\bullet)$. Since this holds for any $\delta_o \in T_o$, we conclude that the vertex barycenter of $\text{Asso}(T_o)$ is the origin. It is immediate for the other two polytopes $\text{Zono}(T_o)$ and $\text{Para}(T_o)$ since they are centrally symmetric. \square

Proposition 35 will be generalized in Section 7.3 for arbitrary seeds in arbitrary finite types and for arbitrary fairly balanced point λ .

7. FURTHER PROPERTIES OF $\text{Asso}(B_o)$

In this section, we discuss further geometric properties of the B_o -associahedron $\text{Asso}^h(B_o)$, motivated by the specific families of examples presented in Section 6. We also introduce the universal associahedron mentioned in Theorem 2, a high dimensional polytope which simultaneously contains the associahedra $\text{Asso}^h(B_o)$ for all exchange matrices B_o of a given finite type.

7.1. Green mutations. Motivated by Section 6.1, we consider a natural orientation of mutations introduced by B. Keller in the context of quantum dilogarithm identities [Kel11]. For two adjacent seeds $\Sigma = (B, P, X)$ and $\Sigma' = (B', P', X')$ of $\mathcal{A}_{\text{pr}}(B_o)$ with $X \setminus \{x\} = X' \setminus \{x'\}$, the mutation $\Sigma \rightarrow \Sigma'$ is a *green mutation* when the dual \mathbf{c} -vector $\mathbf{c}(B_o^\vee, x^\vee \in \Sigma^\vee) = -\mathbf{c}(B_o^\vee, x'^\vee \in \Sigma'^\vee)$ is positive. The directed graph $\mathcal{G}(B_o)$ of green mutations is known to be acyclic, and even the Hasse diagram of a lattice when the type of B_o is simply laced, see [GM15, Coro. 4.7] and the references therein. Further lattice theoretic properties of $\mathcal{G}(B_o)$ are discussed in [GM15].

Example 36. For an acyclic initial exchange matrix B_o , the lattice of green mutations is the c -Cambrian lattice of N . Reading [Rea06] (where c is the Coxeter element corresponding to B_o).

It turns out that this green mutation digraph $\mathcal{G}(B_o)$ is apparent in the B_o -associahedron. Indeed, the following statement is a direct consequence of Remark 27.

Proposition 37. *For any finite type exchange matrix B_o , the graph of the associahedron $\text{Asso}^h(B_o)$, oriented in the linear direction $-\sum_{x \in X_o} \omega_x$, is the graph $\mathcal{G}(B_o)$ of green mutations in $\mathcal{A}_{\text{pr}}(B_o)$.*

Proof. Consider two adjacent seeds $\Sigma = (B, P, X)$ and $\Sigma' = (B', P', X')$ of $\mathcal{A}_{\text{pr}}(B_o)$ with $X \setminus \{x\} = X' \setminus \{x'\}$. By Remark 27, we have

$$\mathbf{p}^h(B_o, \Sigma') - \mathbf{p}^h(B_o, \Sigma) = -\gamma \mathbf{c}(B_o^\vee, x^\vee \in \Sigma^\vee)$$

for some positive $\gamma \in \mathbb{R}_{>0}$. Therefore,

$$\left\langle - \sum_{x \in X_o} \omega_x \mid \mathbf{p}^h(B_o, \Sigma') - \mathbf{p}^h(B_o, \Sigma) \right\rangle = \gamma \left\langle \sum_{x \in X_o} \omega_x \mid \mathbf{c}(B_o^\vee, x^\vee \in \Sigma^\vee) \right\rangle$$

is positive if and only if $\mathbf{c}(B_o^\vee, x^\vee \in \Sigma^\vee)$ is a positive \mathbf{c} -vector. \square

7.2. Universal associahedron. For each initial exchange matrix B_o of a given type, we constructed in Section 5 a generalized associahedron $\text{Asso}^h(B_o)$ by lifting the \mathbf{g} -vector fan using an exchange submodular function h on the cluster variables of $\mathcal{A}_{\text{pr}}(B_o)$. As already observed though, the function h is independent of the coefficients of $\mathcal{A}_{\text{pr}}(B_o)$, so that all \mathbf{g} -vector fans can be lifted with the same function h . This motivates the definition of a universal associahedron.

For this, consider the finite type cluster algebra $\mathcal{A}_{\text{un}}(B_o)$ with universal coefficients, and let $\mathcal{X}(B_o)$ denote its set of cluster variables. Consider a $|\mathcal{X}(B_o)|$ -dimensional vector space U with basis $\{\beta_x\}_{x \in \mathcal{X}(B_o)}$ and its dual space U^\vee with basis $\{\beta_{x^\vee}^\vee\}_{x^\vee \in \mathcal{X}(B_o^\vee)}$. As before, the cluster variables of $\mathcal{A}_{\text{un}}(B_o)$ and $\mathcal{A}_{\text{un}}(B_o^\vee)$ are related by $x \leftrightarrow x^\vee$. For $X \subseteq \mathcal{X}(B_o)$, we denote by \mathbf{H}_X the coordinate subspace of U spanned by $\{\beta_x\}_{x \in X}$.

Given a seed Σ in $\mathcal{A}_{\text{un}}(B_o)$, the **u-vector** of a cluster variable $x \in \Sigma$ is the vector

$$\mathbf{u}(B_o, x \in \Sigma) := \sum_{y \in \mathcal{X}(B_o)} u_{yx} \beta_y \in U$$

of exponents of $p_x = \prod_{y \in \mathcal{X}(B_o)} (p[y])^{u_{yx}}$. Remark 14 then reformulates geometrically in terms of **u**- and **c**-vectors as follows. Choose a seed $\Sigma_\star = (B_\star, P_\star, X_\star)$ in $\mathcal{A}_{\text{un}}(B_o)$ that you want to make initial. Then, for any cluster variable x in a seed Σ , the **c**-vector $\mathbf{c}(B_\star, x \in \Sigma)$ is the orthogonal projection of the **u**-vector $\mathbf{u}(B_o, x \in \Sigma)$ on the coordinate subspace \mathbf{H}_{X_\star} . (Here and elsewhere we identify \mathbf{H}_{X_\star} with V and $\mathbf{H}_{X_\star^\vee}$ with V^\vee in the obvious way.)

We are now ready to define the universal associahedron.

Definition 38. For any finite type exchange matrix B_o and any exchange submodular function h , the **universal B_o -associahedron** is the polytope $\text{Asso}_{\text{un}}^h(B_o)$ in U^\vee defined as the convex hull of the points

$$\mathbf{p}_{\text{un}}^h(B_o, \Sigma) := \sum_{x \in \Sigma} h(x) \mathbf{u}(B_o^\vee, x^\vee \in \Sigma^\vee) \in U^\vee$$

for each seed Σ of $\mathcal{A}_{\text{un}}(B_o)$.

Note that $\text{Asso}_{\text{un}}^h(B_o)$ does not depend on B_o but only on its cluster type. We keep B_o in the notation since it fixes the indexing of the spaces U and U^\vee .

Example 39. We illustrate Definition 38 on the type C_2 exchange matrix:

$$B_o = \begin{bmatrix} 0 & 2 \\ -1 & 0 \end{bmatrix}.$$

The cluster algebra $\mathcal{A}_{\text{un}}(B_o)$ has 6 cluster variables that we denote by $\mathcal{X}(B_o) = \{x_1, x_2, x_3, x_4, x_5, x_6\}$. It is straightforward to verify that to the point $2\rho^\vee$ corresponds the function $h_{2\rho^\vee}$ with value 3 on x_1, x_3, x_5 and 4 on x_2, x_4, x_6 . The **u**-vectors we need to compute our polytope are those associated to the algebra $\mathcal{A}_{\text{un}}(B_o^\vee)$ that appears in Example 15. (Notationally one should think of cluster variables in Example 15 as $\{x_i^\vee\}$.) Using Figure 1 we then get that $\text{Asso}_{\text{un}}^{h_{2\rho^\vee}}(B_o)$ is the convex hull of the 6 points

$$\begin{array}{lll} (3, 4, -3, -4, -1, 2), & (3, -4, -3, -2, 1, 4), & (1, 4, 3, -4, -3, -2), \\ (-3, -4, -1, 2, 3, 4), & (-1, 2, 3, 4, -3, -4), & (-3, -2, 1, 4, 3, -4). \end{array}$$

These are in general position so $\text{Asso}_{\text{un}}^{h_{2\rho^\vee}}(B_o)$ is a 5-dimensional simplex embedded in \mathbb{R}^6 .

Our interest in $\text{Asso}_{\text{un}}^h(B_o)$ comes from the following property.

Theorem 40. Fix a finite type exchange matrix B_o and an exchange submodular function h . For any seed $(B_\star, P_\star, X_\star)$ of $\mathcal{A}_{\text{un}}(B_o)$, the orthogonal projection of the universal associahedron $\text{Asso}_{\text{un}}^h(B_o)$ on the coordinate subspace $\mathbf{H}_{X_\star^\vee}$ of U^\vee spanned by $\{\beta_{x^\vee}^\vee\}_{x^\vee \in X_\star^\vee}$ is the B_\star -associahedron $\text{Asso}^h(B_\star)$.

Proof. Denote by $\pi_{X_\star^\vee}$ the orthogonal projection on $\mathbf{H}_{X_\star^\vee}$. We already observed that

$$\mathbf{c}(B_\star^\vee, x^\vee \in \Sigma^\vee) = \pi_{X_\star^\vee}(\mathbf{u}(B_o^\vee, x^\vee \in \Sigma^\vee))$$

n	dimension of ambient space	dimension	# vertices	# facets	# vertices / facet	# facets / vertex
1	2	1	2	2	1	1
2	5	4	5	5	4	4
3	9	8	14	60	$9 \leq \cdot \leq 10$	$30 \leq \cdot \leq 42$
4	14	13	42	8960	$14 \leq \cdot \leq 28$	$3463 \leq \cdot \leq 4244$

TABLE 1. Some statistics for the universal associahedron of type A_n for $n \in [4]$.

for any cluster variable x in any seed Σ . It follows that

$$\mathbf{p}^h(B_\star, \Sigma) = \pi_{X_\star^\vee}(\mathbf{p}_{\text{un}}^h(B_\circ, \Sigma))$$

for any seed Σ . We conclude that

$$\begin{aligned} \text{Asso}^h(B_\circ) &:= \text{conv} \{ \mathbf{p}^h(B_\star, \Sigma) \mid \Sigma \text{ seed in } \mathcal{A}_{\text{pr}}(B_\circ) \} \\ &= \text{conv} \{ \pi_{X_\star^\vee}(\mathbf{p}_{\text{un}}^h(B_\star, \Sigma)) \mid \Sigma \text{ seed in } \mathcal{A}_{\text{un}}(B_\circ) \} \\ &= \pi_{X_\star^\vee}(\text{conv} \{ \mathbf{p}_{\text{un}}^h(B_\star, \Sigma) \mid \Sigma \text{ seed in } \mathcal{A}_{\text{un}}(B_\circ) \}) =: \pi_{X_\star^\vee}(\text{Asso}_{\text{un}}^h(B_\circ)). \quad \square \end{aligned}$$

Remark 41. Consider the normal fan \mathcal{F} of the universal B_\circ -associahedron $\text{Asso}_{\text{un}}^h(B_\circ)$. Then for any seed $\Sigma_\star = (B_\star, P_\star, X_\star)$ in $\mathcal{A}_{\text{un}}(B_\circ)$, the section of \mathcal{F} by the coordinate subspace \mathbf{H}_{X_\star} of U spanned by $\{\beta_x\}_{x \in X_\star}$ is the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(B_\star)$. We therefore call *universal \mathbf{g} -vector fan* the normal fan $\mathcal{F}_{\text{un}}^{\mathbf{g}}(B_\circ)$ of the universal B_\circ -associahedron $\text{Asso}_{\text{un}}(B_\circ)$.

We now gather some observations on the universal B_\circ -associahedron $\text{Asso}_{\text{un}}^h(B_\circ)$:

- The universal associahedron $\text{Asso}_{\text{un}}^h(B_\circ)$ is *a priori* defined in U^\vee . However, computer experiments indicate that it has codimension 1. The linear space containing $\text{Asso}_{\text{un}}^h(B_\circ)$ seems to be expressed naturally in terms of the Cartan matrix of type B_\circ and h .
- As an immediate consequence of Theorem 40, we obtain that the vertices of the universal associahedron $\text{Asso}_{\text{un}}^h(B_\circ)$ are precisely the points $\mathbf{p}_{\text{un}}^h(B_\circ, \Sigma)$ for all seeds Σ of $\mathcal{A}_{\text{un}}(B_\circ)$, and that the mutation graph of the cluster algebra $\mathcal{A}_{\text{un}}(B_\circ)$ is a subgraph of the graph of $\text{Asso}_{\text{un}}^h(B_\circ)$. However, this inclusion is strict in general.
- In general $\text{Asso}_{\text{un}}^h(B_\circ)$ is neither simple nor simplicial. Table 1 presents some statistics for the number of vertices per facet and facets per vertex in the universal associahedron of type A_n for $n \in [4]$.
- Computer experiments indicate that the face lattice (and thus the f -vector) of the universal B_\circ -associahedron $\text{Asso}_{\text{un}}^h(B_\circ)$ is independent of h .

To conclude, let us insist on the fact that Theorem 40 describes the projection of the universal associahedron $\text{Asso}_{\text{un}}^h(B_\circ)$ on coordinate subspaces corresponding to clusters of $\mathcal{A}_{\text{un}}(B_\circ)$. It turns out that the projections on coordinate subspaces corresponding to all faces (not necessarily facets) of the cluster complex create relevant simplicial complexes, fans and polytopes [Cha16, GM16, MP17]. This naturally raises the question to understand all coordinate projections of the universal associahedron $\text{Asso}_{\text{un}}^h(B_\circ)$.

7.3. Vertex barycenter. As mentioned in Section 6.1, the vertex barycenters of all associahedra constructed by C. Hohlweg, C. Lange and H. Thomas in [HLT11] coincide with the origin. This intriguing property observed in [HLT11, Conj. 5.1] was proved by V. Pilaud and C. Stump [PS15b]. We show in this section that it also extends to all associahedra $\text{Asso}^h(B_\circ)$ for any initial exchange matrix B_\circ and any exchange submodular function h . In fact, it is a consequence of the following stronger statement.

Theorem 42. *For any finite type exchange matrix B_\circ and any exchange submodular function h , the origin is the vertex barycenter of the universal B_\circ -associahedron $\text{Asso}_{\text{un}}^h(B_\circ)$.*

Our proof of this theorem relies on its validity in the acyclic case [PS15b], and on the following observation.

Lemma 43. *Fix a finite type exchange matrix B_o and an exchange submodular function h . For any seed $(B_\star, P_\star, X_\star)$ of $\mathcal{A}_{\text{un}}(B_o)$, the vertex barycenter of the B_\star -associahedron $\text{Asso}^h(B_\star)$ is the image of the vertex barycenter of the universal associahedron $\text{Asso}_{\text{un}}^h(B_o)$ by the orthogonal projection on the coordinate subspace $\mathbf{H}_{X_\star^\vee}$ of U^\vee spanned by $\{\beta_{x^\vee}^\vee\}_{x^\vee \in X_\star^\vee}$.*

Proof. Denote by $\pi_{X_\star^\vee}$ the orthogonal projection on $\mathbf{H}_{X_\star^\vee}$. Since $\pi_{X_\star^\vee}$ is linear, we have

$$\sum_{\Sigma} \mathbf{p}^h(B_\star, \Sigma) = \sum_{\Sigma} \pi_{X_\star^\vee}(\mathbf{p}_{\text{un}}^h(B_o, \Sigma)) = \pi_{X_\star^\vee}\left(\sum_{\Sigma} \mathbf{p}_{\text{un}}^h(B_o, \Sigma)\right),$$

where Σ runs over all seeds of $\mathcal{A}_{\text{un}}(B_o)$. The result follows since these seeds index the vertices of both $\text{Asso}^h(B_\star)$ and $\text{Asso}_{\text{un}}^h(B_o)$. \square

Proof of Theorem 42. Consider a cluster variable x of $\mathcal{A}_{\text{un}}(B_o)$. Let $(B_\star, P_\star, X_\star)$ be an acyclic seed of $\mathcal{A}_{\text{un}}(B_o)$ containing x (such a seed exists, we could even require that it is bipartite). Since by [PS15b] the vertex barycenter of $\text{Asso}^h(B_\star)$ is at the origin, we obtain by Lemma 43 that the x -coordinate of the vertex barycenter of the universal associahedron $\text{Asso}_{\text{un}}^h(B_o)$ vanishes. Applying the same argument independently for all cluster variables x of $\mathcal{A}_{\text{un}}(B_o)$ concludes the proof. \square

Corollary 44. *For any finite type exchange matrix B_o and any exchange submodular function h , the origin is the vertex barycenter of the B_o -associahedron $\text{Asso}^h(B_o)$.*

Proof. This is an immediate consequence of Theorem 42 and Lemma 43. \square

7.4. Zonotope. Motivated by the specific families presented in Section 6, it is natural to investigate whether there exists a zonotope $\text{Zono}^h(B_o)$ whose facet description contains all inequalities of the associahedron $\text{Asso}^h(B_o)$. In this section, we show that such a zonotope does not always exist in general.

A first naive option. Mimicking Section 6, the natural choice is to consider the zonotope $\text{Zono}(B_o) := \sum_{\mathbf{c} \in \mathbf{C}(B_o^\vee)} \mathbf{c}$. Indeed, we have seen in Section 6 that all inequalities of $\text{Asso}(B_o)$ are inequalities of $\text{Zono}(B_o)$ when B_o is either acyclic or of type A.

However, this property already fails for the type C_3 cyclic initial exchange matrix

$$B_o = \begin{bmatrix} 0 & -1 & 2 \\ 1 & 0 & -2 \\ -1 & 1 & 0 \end{bmatrix}.$$




FIGURE 7. The zonotope counter-example in type C_3 . The B_o -associahedron $\text{Asso}(B_o)$ (left), the B_o -zonotope $\text{Zono}(B_o)$ (middle), and their superposition (right) for the type C_3 cyclic initial exchange matrix B_o . The hyperplanes supporting the shaded facets of $\text{Asso}(B_o)$ are parallel to but do not coincide with the hyperplanes supporting the shaded facets of $\text{Zono}(B_o)$.




FIGURE 8. The polytope defined by the inequalities of $\text{Zono}(\mathbf{B}_o)$ defining facets whose normal vectors are \mathbf{g} -vectors of $\mathcal{A}_{\text{pr}}(\mathbf{B}_o)$ (left), and a zonotope whose facet description contains all inequalities of $\text{Asso}(\mathbf{B}_o)$ (right).

One indeed checks that in the direction of the \mathbf{g} -vectors $(-1, 1, 0)$ and $(1, -1, 0)$, the right hand sides of the inequalities are 4 in $\text{Asso}(\mathbf{B}_o)$ and 3 in $\text{Zono}(\mathbf{B}_o)$. This is visible in Figure 7 where the facets of $\text{Asso}(\mathbf{B}_o)$ and $\text{Zono}(\mathbf{B}_o)$ orthogonal to the \mathbf{g} -vectors $(-1, 1, 0)$ and $(1, -1, 0)$ are shaded.

Figure 8 (left) shows that the polytope defined by the inequalities of $\text{Zono}(\mathbf{B}_o)$ defining facets whose normal vectors are \mathbf{g} -vectors of $\mathcal{A}_{\text{pr}}(\mathbf{B}_o)$ is not an associahedron of type C_3 (it is not even a simple polytope).

Note by the way that the two \mathbf{g} -vectors $(-1, 1, 0)$ and $(1, -1, 0)$ are opposite (thus correspond to parallel facets of $\text{Asso}(\mathbf{B}_o)$). This should sound unusual as the only pairs of opposite \mathbf{g} -vectors in both situations of Section 6 are the pairs of opposite coordinate vectors $\{\omega_x, -\omega_x\}$ for $x \in X_o$.

General approach. For any tuple $\gamma := (\gamma_{\mathbf{c}})_{\mathbf{c} \in \mathbf{C}(\mathbf{B}_o^\vee)}$ of positive coefficients, we consider the zonotope

$$\text{Zono}^\gamma(\mathbf{B}_o) := \sum_{\mathbf{c} \in \mathbf{C}(\mathbf{B}_o^\vee)} \gamma_{\mathbf{c}} \mathbf{c}$$

By definition, its normal fan is $\mathcal{F}^\mathbf{c}(\mathbf{B}_o^\vee)$. For any ray \mathbf{g} of this fan, the inequality defining the facet normal to \mathbf{g} is given by $\langle \mathbf{g} | \mathbf{v} \rangle \leq k(\mathbf{g})$ where

$$k(\mathbf{g}) := \sum_{\substack{\mathbf{c} \in \mathbf{C}(\mathbf{B}_o^\vee) \\ \langle \mathbf{g} | \mathbf{c} \rangle > 0}} \gamma_{\mathbf{c}} \langle \mathbf{g} | \mathbf{c} \rangle.$$

The facet description of the zonotope $\text{Zono}^\gamma(\mathbf{B}_o)$ thus contains all inequalities of the associahedron $\text{Asso}^h(\mathbf{B}_o)$ if and only if $\gamma = \gamma(h)$ is a positive solution to the system of linear equations

$$\sum_{\substack{\mathbf{c} \in \mathbf{C}(\mathbf{B}_o^\vee) \\ \langle \mathbf{g}(\mathbf{B}_o, x) | \mathbf{c} \rangle > 0}} \gamma_{\mathbf{c}} \langle \mathbf{g}(\mathbf{B}_o, x) | \mathbf{c} \rangle = h(x) \quad \text{for all cluster variable } x \text{ of } \mathcal{A}_{\text{pr}}(\mathbf{B}_o).$$

For example, such a solution exists for the type C_3 cyclic initial exchange matrix, as illustrated in Figure 8 (right). Note that we had to pick different coefficients for different elements of $\mathbf{C}(\mathbf{B}_o^\vee)$ (leading to unnaturally narrow faces) since we already observed in Figure 7 that constant coefficients are not suitable.

In contrast, a quick computer experiment shows that this system has no solution for the type D_5 cyclic initial exchange matrix

$$\mathbf{B}_o = \begin{bmatrix} 0 & 1 & 0 & 0 & -1 \\ -1 & 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 & 1 \\ 1 & 0 & 0 & -1 & 0 \end{bmatrix}$$




FIGURE 9. An obstruction to the existence of a centrally symmetric zonotope whose facet description would contain that of the associahedron $\text{Asso}(\mathbf{B}_o)$ for a cyclic initial seed \mathbf{B}_o in type D_5 . The two arcs γ_ℓ (left) and γ_r (right) have opposite \mathbf{g} -vectors but different compatibility sums.

and the balanced exchange submodular function h_{ρ^\vee} . Rather than showing the detailed linear system, we prefer to convince the reader that there is no solution under the natural assumption that $\gamma_{\mathbf{c}} = \gamma_{-\mathbf{c}}$ for any $\mathbf{c} \in \mathbf{C}(\mathbf{B}_o^\vee)$, *i.e.*, that the zonotope $\text{Zono}^\gamma(\mathbf{B}_o)$ is centrally symmetric. To see that no such solution exists, consider the classical punctured pentagon model for the type D_5 cluster algebra [FST08]. Recall that

- (i) cluster variables correspond to internal arcs up to isotopy (with the subtlety that an arc incident to the puncture can be tagged or not),
 - (ii) \mathbf{g} -vectors can be read by shear coordinates [FT12] in a similar way as in Section 6.2, and
 - (iii) the compatibility degree between two arcs is the (minimal) number of crossings between them.
- We therefore obtain that the arcs γ_ℓ connecting 1_\bullet to 3_\bullet in Figure 9 (left) and the arc γ_r connecting 4_\bullet to 5_\bullet in Figure 9 (right) satisfy:

$$\begin{aligned} \mathbf{g}(\mathbf{B}_o, x_{\gamma_\ell}) &= (-1, 0, 1, 0, 0) \quad \text{and} \quad h_{\rho^\vee}(x_{\gamma_\ell}) = 7, \\ \mathbf{g}(\mathbf{B}_o, x_{\gamma_r}) &= (1, 0, -1, 0, 0) \quad \text{and} \quad h_{\rho^\vee}(x_{\gamma_r}) = 9. \end{aligned}$$

In other words, we exhibited two cluster variables with opposite \mathbf{g} -vectors while belonging to distinct $\langle \tau_+, \tau_- \rangle$ -orbits. Since a centrally symmetric zonotope would have the same right-hand-sides on opposite normal vectors, this shows that no such polytope exists.

ACKNOWLEDGEMENTS

We thank N. Reading for helpful discussions on this topic and comments on a preliminary version of this paper. We are also grateful to A. Garver and T. McConville for bibliographic inputs. This work answers one of the questions A. Zelevinsky posed to the third author as possible Ph.D. problems. He would therefore like to express once again gratitude to his advisor for the guidance received.

REFERENCES

- [BB05] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter groups*, volume 231 of *Graduate Texts in Mathematics*. Springer, New York, 2005.
- [Bri98] Brigitte Brink. The set of dominance-minimal roots. *J. Algebra*, 206(2):371–412, 1998.
- [CFZ02] Frédéric Chapoton, Sergey Fomin, and Andrei Zelevinsky. Polytopal realizations of generalized associahedra. *Canad. Math. Bull.*, 45(4):537–566, 2002.
- [Cha16] Frédéric Chapoton. Stokes posets and serpent nests. *Discrete Math. Theor. Comput. Sci.*, 18(3), 2016.
- [CP15] Cesar Ceballos and Vincent Pilaud. Denominator vectors and compatibility degrees in cluster algebras of finite type. *Trans. Amer. Math. Soc.*, 367(2):1421–1439, 2015.
- [CSZ15] Cesar Ceballos, Francisco Santos, and Günter M. Ziegler. Many non-equivalent realizations of the associahedron. *Combinatorica*, 35(5):513–551, 2015.

- [Dem10] Laurent Demonet. Mutations of group species with potentials and their representations. Applications to cluster algebras. Preprint, [arXiv:1003.5078](#), to appear in *Combinatorica*, 2010.
- [DRS10] Jesus A. De Loera, Jörg Rambau, and Francisco Santos. *Triangulations: Structures for Algorithms and Applications*, volume 25 of *Algorithms and Computation in Mathematics*. Springer Verlag, 2010.
- [DWZ10] Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky. Quivers with potentials and their representations II: applications to cluster algebras. *J. Amer. Math. Soc.*, 23(3):749–790, 2010.
- [FST08] Sergey Fomin, Michael Shapiro, and Dylan Thurston. Cluster algebras and triangulated surfaces I. Cluster complexes. *Acta Math.*, 201(1):83–146, 2008.
- [FST12] Anna Felikson, Michael Shapiro, and Pavel Tumarkin. Cluster algebras and triangulated orbifolds. *Adv. Math.*, 231(5):2953–3002, 2012.
- [FT12] Sergey Fomin and Dylan Thurston. Cluster algebras and triangulated surfaces. part II: Lambda lengths. Preprint, [arXiv:1210.5569](#), 2012.
- [FZ02] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. I. Foundations. *J. Amer. Math. Soc.*, 15(2):497–529, 2002.
- [FZ03a] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. II. Finite type classification. *Invent. Math.*, 154(1):63–121, 2003.
- [FZ03b] Sergey Fomin and Andrei Zelevinsky. Y-systems and generalized associahedra. *Ann. of Math. (2)*, 158(3):977–1018, 2003.
- [FZ07] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. IV. Coefficients. *Compos. Math.*, 143(1):112–164, 2007.
- [GHKK14] Mark Gross, Paul Hacking, Sean Keel, and Maxim Kontsevich. Canonical bases for cluster algebras. Preprint, [arXiv:1411.1394](#), 2014.
- [GKZ08] Israel Gelfand, Mikhail Kapranov, and Andrei Zelevinsky. *Discriminants, resultants and multidimensional determinants*. Modern Birkhäuser Classics. Birkhäuser Boston Inc., Boston, MA, 2008. Reprint of the 1994 edition.
- [GM15] Alexander Garver and Thomas McConville. Lattice properties of oriented exchange graphs and torsion classes. Preprint, [arXiv:1507.04268](#), 2015.
- [GM16] Alexander Garver and Thomas McConville. Oriented flip graphs and noncrossing tree partitions. Preprint, [arXiv:1604.06009](#), 2016.
- [HL07] Christophe Hohlweg and Carsten Lange. Realizations of the associahedron and cyclohedron. *Discrete Comput. Geom.*, 37(4):517–543, 2007.
- [HLR10] Christophe Hohlweg, Jonathan Lortie, and Annie Raymond. The centers of gravity of the associahedron and of the permutahedron are the same. *Electron. J. Combin.*, 17(1):Research Paper 72, 14, 2010.
- [HLT11] Christophe Hohlweg, Carsten Lange, and Hugh Thomas. Permutahedra and generalized associahedra. *Adv. Math.*, 226(1):608–640, 2011.
- [HNW16] Christophe Hohlweg, Philippe Nadeau, and Nathan Williams. Automata, reduced words and garside shadows in coxeter groups. *Journal of Algebra*, 457:431 – 456, 2016.
- [Hoh12] Christophe Hohlweg. Permutahedra and associahedra: generalized associahedra from the geometry of finite reflection groups. In *Associahedra, Tamari lattices and related structures*, volume 299 of *Prog. Math. Phys.*, pages 129–159. Birkhäuser/Springer, Basel, 2012.
- [Kel11] Bernhard Keller. On cluster theory and quantum dilogarithm identities. In *Representations of algebras and related topics*, EMS Ser. Congr. Rep., pages 85–116. Eur. Math. Soc., Zürich, 2011.
- [Lod04] Jean-Louis Loday. Realization of the Stasheff polytope. *Arch. Math. (Basel)*, 83(3):267–278, 2004.
- [LP13] Carsten Lange and Vincent Pilaud. Associahedra via spines. Preprint, [arXiv:1307.4391](#), to appear in *Combinatorica*, 2013.
- [MP17] Thibault Manneville and Vincent Pilaud. Geometric realizations of the accordion complex of a dissection. In preparation, 2017.
- [NS14] Tomoki Nakanishi and Salvatore Stella. Diagrammatic description of c -vectors and d -vectors of cluster algebras of finite type. *Electron. J. Combin.*, 21(1):Paper 1.3, 107, 2014.
- [NZ12] Tomoki Nakanishi and Andrei Zelevinsky. On tropical dualities in cluster algebras. In *Algebraic groups and quantum groups*, volume 565 of *Contemp. Math.*, pages 217–226. Amer. Math. Soc., Providence, RI, 2012.
- [PS15a] Vincent Pilaud and Christian Stump. Brick polytopes of spherical subword complexes and generalized associahedra. *Adv. Math.*, 276:1–61, 2015.
- [PS15b] Vincent Pilaud and Christian Stump. Vertex barycenter of generalized associahedra. *Proc. Amer. Math. Soc.*, 143(6):2623–2636, 2015.
- [Rea06] Nathan Reading. Cambrian lattices. *Adv. Math.*, 205(2):313–353, 2006.
- [Rea14] Nathan Reading. Universal geometric cluster algebras. *Math. Z.*, 277(1-2):499–547, 2014.
- [RS09] Nathan Reading and David E. Speyer. Cambrian fans. *J. Eur. Math. Soc.*, 11(2):407–447, 2009.
- [RS15] Nathan Reading and David E. Speyer. A Cambrian framework for the oriented cycle. *Electron. J. Combin.*, 22(4):Paper 4.46, 21, 2015.
- [Ste13] Salvatore Stella. Polyhedral models for generalized associahedra via Coxeter elements. *J. Algebraic Combin.*, 38(1):121–158, 2013.

- [YZ08] Shih-Wei Yang and Andrei Zelevinsky. Cluster algebras of finite type via Coxeter elements and principal minors. *Transform. Groups*, 13(3-4):855–895, 2008.
- [Zie98] Günter M. Ziegler. *Lectures on Polytopes*, volume 152 of *Graduate texts in Mathematics*. Springer-Verlag, New York, 1998.

(Christophe Hohlweg) LACIM, UNIVERSITÉ DU QUÉBEC À MONTRÉAL (UQAM)

E-mail address: `hohlweg.christophe@uqam.ca`

URL: <http://hohlweg.math.uqam.ca/>

(Vincent Pilaud) CNRS & LIX, ÉCOLE POLYTECHNIQUE, PALAISEAU

E-mail address: `vincent.pilaud@lix.polytechnique.fr`

URL: <http://www.lix.polytechnique.fr/~pilaud/>

(Salvatore Stella) UNIVERSITÀ DEGLI STUDI DI ROMA “LA SAPIENZA”

E-mail address: `stella@mat.uniroma1.it`

URL: <http://www1.mat.uniroma1.it/people/stella/>